



Adhesin activity of *Leptospira interrogans* lipoprotein identified by *in vivo* and *in vitro* shotgun phage display

Swiany Silveira Lima^{a,b,1}, Ana Tung Ching Ching^{a,c,1}, Regiane Degan Fávaro^a, Josefa B. Da Silva^a, Maria L.S. Oliveira^a, Eneas Carvalho^a, Patrícia A.E. Abreu^d, Sílvia A. Vasconcellos^e, Paulo Lee Ho^{a,b,c,*}

^a Centro de Biotecnologia, Instituto Butantan, CEP 05503-900 São Paulo, Brazil

^b Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, CEP 05508-000 São Paulo, Brazil

^c Programa de Pós-Graduação Interunidades em Biotecnologia USP/IPT/Instituto Butantan, Instituto de Ciências Biomédicas da USP, CEP 05508-900 São Paulo, Brazil

^d Laboratório de Bacteriologia, Instituto Butantan, CEP 05503-900 São Paulo, Brazil

^e Departamento de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia da USP, CEP 05508 270 São Paulo, Brazil

ARTICLE INFO

Article history:

Received 13 December 2012

Available online 3 January 2013

Keywords:

Leptospira interrogans

Adhesin

Lipoprotein

Phage display

Vaccine

ABSTRACT

Leptospira interrogans causes leptospirosis, one of the most common zoonotic diseases in the world. This pathogenic spirochete is able to bind to extracellular matrix, to express virulent factors and to cause host death. Until now, there is no effective human vaccine for the disease. Shotgun phage display genomic libraries of *L. interrogans* were constructed and used for *in vivo* biopanning in hamsters and screened for ligands able to bind to LLC-PK1 epithelial cells. In both panning procedures, clones coding for the putative lipoprotein LIC12976 were identified and, in order to confirm its adhesin activity, a recombinant protein was produced in *Escherichia coli* and showed to interact with A31 fibroblasts, LLC-PK1 and Vero epithelial cells *in vitro*. Moreover, rLIC12976 was shown to bind to laminin, indicating an adhesin function. This protein was also detected in extracts of *L. interrogans* from different serovars and it was found to be conserved among pathogenic leptospires. Further, the protein was tested as vaccine candidate and immunization of hamsters with LIC12976 did not confer protection against a lethal challenge with the homologous *L. interrogans* serovar Copenhageni. Nevertheless, LIC12976 seems to act as an adhesin, and may be important for the host–pathogen interaction, so that its study can contribute to the understanding of the virulence mechanisms in pathogenic leptospires.

© 2013 Elsevier Inc. Open access under the [Elsevier OA license](http://creativecommons.org/licenses/by/3.0/).

1. Introduction

Leptospira is the etiologic agent of leptospirosis, a zoonosis recognized as an emerging infectious disease that is extremely spread throughout the world [1,2]. The disease transmission, directly or indirectly from animals to humans, is through cuts and abrasions in the skin or via the conjunctiva, and may also occur in intact skin after prolonged exposure to water contaminated with urine of infected animals [1].

The knowledge on the mechanism of pathogenicity of *Leptospira* is very limited [1,3]. Nowadays, the genomes of nine strains of the genus *Leptospira* were sequenced: *L. interrogans* serovar Lai [4], *L. interrogans* strain Fiocruz L1–130 serovar Copenhageni [5], JB197 and L550 strains from *Leptospira borgpetersenii* serovar Hardjo [6], *Leptospira santarosai* serovar Shermani [7], VAR010 and MMD0835 strains from *Leptospira licerasiae* serovar Varillal [8]

and *Leptospira biflexa* serovar Patoc strains Paris and Ames [9]. Comparative genomic studies can provide knowledge about mechanisms of pathogenicity and thus help to establish strategies to control the disease and to identify potential candidates for drugs or vaccine development [2,5].

Bacteria interacting with eukaryotic hosts express adhesive molecules on their surfaces that mediate adhesion to a receptor on the target cell surface or with soluble macromolecules and what can head the pathogen to a specific location in the host [10]. Pathogenic bacteria utilize numerous mechanisms to cause disease in humans. The first step in host–pathogen interaction and tissue colonization is the adherence to the surface of the cell or to extracellular matrix proteins. Once attached, pathogens are able to initiate specific biochemical processes that include proliferation, secretion of toxins, immune evasion responses and cell invasion, culminating in the appearance of disease symptoms [11].

Jacobsson and Frykberg [12] were pioneers to identify adhesins using the shotgun phage display. In this technique, the phage library is constructed from random fragments from genomic DNA. Shotgun phage display has identified a number of genes coding for adhesins involved in host–pathogen interactions [13–15].

* Corresponding author at: Centro de Biotecnologia, Instituto Butantan, CEP 05503-900 São Paulo, Brazil. Fax: +55 (11) 3726 1505.

E-mail address: hoplee@butantan.gov.br (P.L. Ho).

¹ Both authors contributed equally to this work.

In the present study, a new adhesin, LIC12976, was identified through *in vivo* and *in vitro* screening a shotgun phage display library constructed using genomic DNA from *L. interrogans* serovar Copenhageni. In agreement, we showed that the recombinant protein rLIC12976 is able to bind to mammalian cells, including kidney epithelial cells and laminin.

2. Materials and methods

2.1. Microorganisms and culture condition

L. interrogans serovar Copenhageni Fiocruz L1–130 was obtained from ATCC catalog BAA-1198. It was cultivated as described [16]. Virulence was maintained by iterative passages in Golden Syrian hamsters.

2.2. Library construction, biopanning and sequence analysis

The libraries construction and the *in vivo* biopanning were done as described [17–19]. Briefly, hamsters were anesthetized intraperitoneally with 2.5% Avertin [20] and the thoracic cavity was opened in order to expose the heart. 1×10^9 phages/mL (200 μ L/animal) from the library were injected directly into the animal heart. After 5 min of phage circulation, the animal was perfused and the phages were recovered by removing the kidneys. The panning procedure was repeated for four rounds. This study has the approval of the Ethics Committee for Animal Use from Instituto Butantan, protocol number 564/08. The *in vitro* biopanning was performed according the BRASIL methodology [21] with modifications. Briefly, 1×10^{12} phages/mL (100 μ L/tube) were incubated with 1×10^6 pig kidney epithelial LLC-PK1 cells. After centrifugation, the phages in the precipitate were eluted by low pH, neutralized with Tris–HCl pH 8.0, centrifuged after the addition of 1% BSA-DMEM to remove cellular debris and the supernatant was again incubated with LLC-PK1 cells. After panning procedures, clones of each round were analyzed as described [18].

2.3. Cloning, expression, purification of rLIC12976 and antisera production

Leptospira's genomic DNAs were extracted using DNAzol Reagent (Invitrogen). The gene *lic12976* was amplified by PCR using specific primers forward 5' TAGGCTCGAGGAACGTCATTCTCTTC 3' and reverse 5' ACCAAGCTTCTAAAAACGTTCCGAATT 3' containing *Xho* I and *Hind* III restriction sites (bold). The amplified product from *L. interrogans* serovar Copenhageni genomic DNA was cloned into the pGEM-T easy vector (Promega) and subcloned into the pAE vector [22], as described [18]. *E. coli* BL21 Star (DE3) pLysS was used for protein expression overnight at 20 °C and the purification of the recombinant protein was performed as described [23]. The presence of the recombinant protein was evaluated by 12% SDS-PAGE, the fractions of interest were dialysed against PBS and the protein was quantified by Bradford (Bio-Rad). The antisera against rLIC12976 and immunoblot assay were performed as described [23,24], using BSA 2% in PBST during incubation with antiserum.

2.4. Bioinformatic analysis

The PSORT program (<http://psort.nibb.ac.jp/>) was used to predict the sub cellular localization of the protein. Public and custom sequence-specific search algorithms were used for identification of sequence motifs including signal peptides, lipoprotein cleavages sites and transmembrane domains (<http://www.cbs.dtu.dk/services/TMHMM>) and (<http://www.cbs.dtu.dk/services/SignalP>), as described [25].

2.5. Binding of rLIC12976 to mammalian cells and extracellular matrix components

A31 (mouse BALB/c3T3 clone A31 fibroblast), Vero (african green monkey kidney epithelial cell – *Cercopithecus aethiops*) and LLC-PK1 (pig kidney epithelial cell – *Sus scrofa*) cells were cultivated and fixed as described [18]. The binding procedure was performed according [16] with modifications. Briefly, Nunc MaxiSorp microtiter plates were coated with 1.0 μ g of plasma fibronectin, fibrinogen, BSA, laminin, cellular fibronectin and monosialoganglioside GM1 (all purchased from Sigma–Aldrich) and were incubated at 4 °C overnight and washed three times with PBST (0.1% Tween). The cells and the ECM proteins were blocked with 200 μ L of 1% BSA in PBST and 10% BSA in PBST, respectively. The incubation were proceed at 37 °C for 1 h and then at 4 °C for 16 h. After three washes with PBST, 1.0 μ g of rLIC12976 protein was diluted in PBS, incubated for 1 h and 30 min at 37 °C and washed six times with PBST. The detection was performed by incubation with polyclonal antisera anti-LIC12976 1:5000 dilution for 1 h at 37 °C followed by three washes with PBST and incubation with anti-mouse IgG peroxidase conjugate 1:10,000 dilution. After three washes with PBST, the plate was developed as described [23]. For competition experiments, rLIC12976 was incubated with laminin (0, 0.5, 1.0 and 5.0 μ g) for 1 h and 30 min before the incubation with the cells.

2.6. Immunization of hamsters and leptospiral challenge

All procedures were approved by the Committee for the Use of Experimental Animals of the Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, São Paulo, SP, Brazil. Immunization and challenge were done as described [26].

3. Results

3.1. LIC12976 was selected by *in vivo* and *in vitro* biopanning and it is conserved in pathogenic leptospiral genomes

After four rounds of *in vivo* biopanning through hamster perfusion combined with the recovering of the phages from kidneys in parallel with *in vitro* biopanning against kidney epithelial LLC-PK1 cells, two and one, respectively, phage clones in frame coding for a partial sequence of LIC12976 were selected. LIC12976 encodes a putative lipoprotein with 47.7 kDa of molecular mass corresponding to 418 amino acids residues and pI 8.12. This protein possesses a signal peptide, but functionally motifs besides the lipobox, were not identified (Supplementary Fig. 1). A PCR using primers designed on *L. interrogans* serovar Copenhageni strain Fiocruz L1–130 DNA sequence was used to investigate the presence of this gene in other *Leptospira* serovars. Indeed, this gene was amplified in all pathogenic serovars assayed of the species *L. interrogans*, comprising a fragment of approximately 1.2 kb, but not in the saprophytic *L. biflexa* serovar Patoc (Fig. 1), confirming that *lic12976* is conserved in pathogenic leptospires species. The recombinant LIC12976 (rLIC12976) with 45.8 kDa corresponding to amino acid 29–418 was expressed and purified.

3.2. rLIC12976 is expressed in pathogenic *Leptospira* serovars

The polyclonal antiserum against rLIC12976 was able to specifically recognize the recombinant protein (Fig. 2A) but not other *L. interrogans* recombinant proteins. Immunoblot analysis were also performed on a panel of *Leptospira* serovar extracts and the antiserum against rLIC12976 reacted with a band corresponding to a molecular mass of LIC12976 in all pathogenic serovar samples




Fig. 1. PCR amplification of the gene encoding the LIC12976 in *L. interrogans* serovars Copenhageni, Autumnalis, Canicola, Hardjo, Icterohaemorrhagiae, Pomona and Pyrogenes and *L. biflexa* serovar Patoc. M is the molecular marker 1 kb plus (Invitrogen).




Fig. 2. (A) Polyclonal serum raised against LIC12976 is specific against the homologous recombinant protein. Other recombinant proteins were tested, such as LigA [37], LipL32 [24], TlyC [27], LIC11399 (Hashimoto V.L., unpublished results), LIC13435 [28], LIC13418 (Lima S.S., unpublished results). (B) Presence of LIC12976 protein in *L. interrogans* serovars. Sera from mice immunized with LIC12976 reacted with the native protein from whole cell extracts of pathogenic *L. interrogans* serovars Australis, Bratislava, Canicola, Pyrogenes, Hardjo and Copenhageni but failed to recognize the *L. biflexa* serovar Patoc. Sera also react with the rLIC12976.

(Australis, Bratislava, Canicola, Pyrogenes, Hardjo and Copenhageni). In accordance to the result that *lic12976* presence is conserved in pathogenic leptospires species, no reaction was observed with nonpathogenic saprophytic *L. biflexa* serovar Patoc (Fig. 2B) indicating that LIC12976 is only expressed by pathogenic leptospires.

3.3. rLIC12976 binds to mammalian cells and laminin in vitro

Since LIC12976 was identified after panning against kidney (*in vivo*) and LLC-PK1 epithelial kidney cells (*in vitro*), the recombinant protein rLIC12976 was subjected to binding assays to mammalian cells, what showed that rLIC12976 was able to adhere to all tested cells including LLC-PK1 (Fig. 3A). In order to identify possible cellular targets of LIC12976, plasma and ECM purified proteins were also assayed. We showed that rLIC12976 was able to bind to laminin (Fig. 3B). TlyC was used as positive [27] and LIC13435 as negative control [28], respectively (data not shown).

Competition assays were performed to investigate if laminin was able to inhibit the binding of rLIC12976 to mammalian cells (A31, LLC-PK1 or Vero). Nevertheless, using different concentrations of laminin (0, 0.5, 1.0 and 5.0 μ g), higher binding of the recombinant protein to the cells was observed. When we probed the experiment with anti-laminin antibodies, we observed that the laminin added to the cultures was able to bind to the cells (Fig. 3C and D). In the binding assay, rLIC12976 different concentrations were tested and showed that the interaction with mammalian cells and laminin is rLIC12976 concentration dependent; when higher concentration is used, more ligation is observed (data not shown).

3.4. rLIC12976 was not protective against a lethal leptospiral challenge in hamsters

In order to evaluate the potential of LIC12976 as a vaccine antigen against leptospirosis, hamsters were immunized with rLIC12976 and challenged with virulent *L. interrogans* serovar Copenhageni. It was observed that LIC12976 was not a protective antigen, in contrast to LigAC [29], a *L. interrogans* immunoglobulin-like protein used as a positive control (Fig. 4). To date, the most promising vaccine candidate is the C-terminal portion of LigA, which although not being able to completely eliminate leptospires from kidneys, confers protection in a hamster model of leptospirosis [26].

4. Discussion

L. interrogans shotgun phage display libraries were constructed in order to characterize new adhesins. Many adhesins have been identified in pathogenic bacteria using this technique [14,18,19,30–32].

In this work, LIC12976 was identified after *in vivo* and *in vitro* biopanning. It codes for a lipoprotein and our results showed that it was expressed in all tested *L. interrogans* pathogenic serovars. We also observed that rLIC12976 was able to bind to mammalian cells and specifically to laminin. Interestingly, addition of laminin, instead of competing with rLIC12976 for cell binding, enhanced the interaction of rLIC12976 to the cells, indicating that laminin mediates the binding of rLIC12976 to the cells, what probably led to higher binding levels of rLIC12976 to the cells.

Lipoproteins are a set of bacterial membrane proteins with many different functions. These proteins are required for virulence of many pathogenic bacteria, and potentially play a key role in host–pathogen interactions ranging from adhesion to the surface of the host cells with the induction of inflammatory processes to the translocation of virulence factors into the host cell cytoplasm [33]. In *Leptospira*, lipoproteins have important structural and functional roles; they may be enzymes, transporters, adhesins, toxins and other proteins essential for bacterial virulence [34].

The immunogenic proteins, specially the outer membrane surface proteins, may be effective as candidates for a new vaccine. The identification of proteins, which are conserved among pathogenic leptospires and can crossprotect against various serovars, has become a major focus of leptospiral research [25]. In the tested conditions, rLIC12976 did not present protective properties as a vaccine antigen, despite the fact that this recombinant protein is immunogenic. The lack of protection may be related to the presence of many different adhesins in leptospires (see references below). This suggest that there is a probable redundant system for bacterial adhesion to host, based on the diversity of these proteins, that is not significantly affected when just one adhesin activity is impaired. Furthermore, future studies combining rLIC12976 with other protein antigens and also with novel adjuvants could be per-




Fig. 3. Adhesion of the rLIC12976 to cells, extracellular matrix components (ECM) and plasma proteins. (A) 96-well plates were plated with A31, LLC-PK1 and Vero cells. (B) ELISA plates wells were coated with 1 µg of BSA, GM1, plasma fibronectin, cellular fibronectin, fibrinogen and laminin. One microgram of rLIC12976 protein was added per well and further revealed with anti-rLIC12976. In control, A31, LLC-PK1 and Vero cells were incubated with the primary and secondary antibodies only. (C) and (D) Competition assay was performed with different concentrations of laminin and rLIC12976 incubated with A31, LLC-PK1 and Vero cells and revealed with anti-LIC12976 or anti-laminin, respectively.




Fig. 4. Survival of hamsters immunized with rLIC12976 and challenged with virulent *L. interrogans* serovar Copenhageni str. Fiocruz L1-130.

formed to test and improve the protection as a new vaccinal candidate.

The ability of *L. interrogans* to cause disease is complex and multifactorial. Studies have shown that this bacterium is able to enter into phagocytic and non-phagocytic cells, and interact with cultured fibroblasts, epithelial cells, endothelial cells and microglial cells, indicating that pathogen surface molecules have a role in adhesion and invasion of the host tissue [2].

So far, several *Leptospira* adhesins have been identified due to the ability to bind to host such as the LipL32 specifically binds to fibronectin and type IV collagen [24,35]; Lsa24/LenA and Lsa 21, the first specifically binds to laminin and the second is capable of binding to laminin, type IV collagen and fibronectin [16,36]; LigA e LigB, both with adhesion proprieties to immobilized ECM and plasma proteins [37] and heparin [18]. LenB, LenC, LenD, LenE e LenF (all paralogs of LenA) that exhibit affinity for fibronectin [38]. The adhesin TlyC binds to laminin, collagen IV and plasma fibronectin [27]. Lsa20 binds to laminin and to plasminogen [39]. Lsa66 and Lsa30 bind to laminin and plasma fibronectin [40,41]. The large amount of adhesins identified up to now in pathogenic leptospires is intriguing. It may represent redundant processes to bind to the host to assure the rate of the infection success.

This is the first report about a leptospiral lipoprotein identified by panning against epithelial kidney cells and also from hamster's kidneys. The identification of a same putative protein in both independent techniques strongly suggests that this putative lipoprotein, LIC12976, could be a new adhesin. Together with the observation that *lic12976* gene is expressed in pathogenic leptospires, our data suggest that this protein may be important for the host interaction process. Binding to the host cell by adhesin is one of the most important skills for the onset of the disease. Thus, research involving the characterization of adhesins may contribute to the knowledge of the first molecular steps involved in host-pathogen interaction. In addition, the search for adhesins

may contribute to the development of vaccines that would block the early stages of infection, such as cell adhesion, preventing bacterial infection [42–44].

Acknowledgments

The authors wish to thank the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação Butantan for financial support.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bbrc.2012.12.095>.

References

- [1] P.N. Levett, Leptospirosis, Clin. Microbiol. Rev. 14 (2001) 296–326.
- [2] R.U. Palaniappan, S. Ramanujam, Y.F. Chang, Leptospirosis: pathogenesis, immunity, and diagnosis, Curr. Opin. Infect. Dis. 20 (2007) 284–292.
- [3] B. Adler, M. Lo, T. Seemann, G.L. Murray, Pathogenesis of leptospirosis: the influence of genomics, Vet. Microbiol. 153 (2011) 73–81.
- [4] S.X. Ren, G. Fu, X.G. Jiang, R. Zeng, Y.G. Miao, H. Xu, Y.X. Zhang, H. Xiong, G. Lu, L.F. Lu, H.Q. Jiang, J. Jia, Y.F. Tu, J.X. Jiang, W.Y. Gu, Y.Q. Zhang, Z. Cai, H.H. Sheng, H.F. Yin, Y. Zhang, G.F. Zhu, M. Wan, H.L. Huang, Z. Qian, S.Y. Wang, W. Ma, Z.J. Yao, Y. Shen, B.Q. Qiang, Q.C. Xia, X.K. Guo, A. Danchin, I. Saint Girons, R.L. Somerville, Y.M. Wen, M.H. Shi, Z. Chen, J.G. Xu, G.P. Zhao, Unique physiological and pathogenic features of *Leptospira interrogans* revealed by whole-genome sequencing, Nature 422 (2003) 888–893.
- [5] A.L. Nascimento, A.I. Ko, E.A. Martins, C.B. Monteiro-Vitorello, P.L. Ho, D.A. Haake, S. Verjovski-Almeida, R.A. Hartskeerl, M.V. Marques, M.C. Oliveira, C.F. Menck, L.C. Leite, H. Carrer, L.L. Coutinho, W.M. Degraive, O.A. Dellagostin, H. El-Dorry, E.S. Ferro, M.I. Ferro, L.R. Furlan, M. Gamberini, E.A. Giglioti, A. Goes-Neto, G.H. Goldman, M.H. Goldman, R. Harakava, S.M. Jeronimo, I.L. Junqueira-de-Azevedo, E.T. Kimura, E.E. Kuramae, E.G. Lemos, M.V. Lemos, C.L. Marino, L.R. Nunes, R.C. de Oliveira, G.G. Pereira, M.S. Reis, A. Schrieffer, W.J. Siqueira, P. Sommer, S.M. Tsai, A.J. Simpson, J.A. Ferro, L.E. Camargo, J.P. Kitajima, J.C. Setubal, M.A. Van Sluys, Comparative genomics of two *Leptospira interrogans* serovars reveals novel insights into physiology and pathogenesis, J. Bacteriol. 186 (2004) 2164–2172.
- [6] D.M. Bulach, R.L. Zuerner, P. Wilson, T. Seemann, A. McGrath, P.A. Cullen, J. Davis, M. Johnson, E. Kuczek, D.P. Alt, B. Peterson-Burch, R.L. Coppel, J.I. Rood, J.K. Davies, B. Adler, Genome reduction in *Leptospira borgpetersenii* reflects limited transmission potential, Proc. Natl. Acad. Sci. USA 103 (2006) 14560–14565.
- [7] L.F. Chou, Y.T. Chen, C.W. Lu, Y.C. Ko, C.Y. Tang, M.J. Pan, Y.C. Tian, C.H. Chiu, C.C. Hung, C.W. Yang, Sequence of *Leptospira santarosai* serovar Shermani genome and prediction of virulence-associated genes, Gene 511 (2012) 364–370.
- [8] J.N. Ricaldi, D.E. Fouts, J.D. Selengut, D.M. Harkins, K.P. Patra, A. Moreno, J.S. Lehmann, J. Purushe, R. Sanka, M. Torres, N.J. Webster, J.M. Vinetz, M.A. Mathias, Whole genome analysis of *Leptospira licerasiae* provides insight into leptospiral evolution and pathogenicity, PLoS Negl. Trop. Dis. 6 (2012) e1853.
- [9] M. Picardeau, D.M. Bulach, C. Bouchier, R.L. Zuerner, N. Zidane, P.J. Wilson, S. Creno, E.S. Kuczek, S. Bommezzadri, J.C. Davis, A. McGrath, M.J. Johnson, C. Boursaux-Eude, T. Seemann, Z. Rouy, R.L. Coppel, J.I. Rood, A. Lajus, J.K. Davies, C. Medigue, B. Adler, Genome sequence of the saprophyte *Leptospira biflexa* provides insights into the evolution of *Leptospira* and the pathogenesis of leptospirosis, PLoS One 3 (2008) e1607.
- [10] H.H. Niemann, W.D. Schubert, D.W. Heinz, Adhesins and invasins of pathogenic bacteria: a structural view, Microbes Infect. 6 (2004) 101–112.
- [11] K.A. Kline, S. Falker, S. Dahlberg, S. Normark, B. Henriques-Normark, Bacterial adhesins in host-microbe interactions, Cell Host Microbe 5 (2009) 580–592.
- [12] K. Jacobsson, L. Frykberg, Cloning of ligand-binding domains of bacterial receptors by phage display, Biotechniques 18 (1995) 878–885.
- [13] K. Jacobsson, A. Rosander, J. Bjerketorp, L. Frykberg, Shotgun phage display – selection for bacterial receptors or other exported proteins, Biol. Proced. Online 5 (2003) 123–135.
- [14] C. Beckmann, J.D. Waggoner, T.O. Harris, G.S. Tamura, C.E. Rubens, Identification of novel adhesins from Group B streptococci by use of phage display reveals that C5a peptidase mediates fibronectin binding, Infect. Immun. 70 (2002) 2869–2876.
- [15] M. Nilsson, J. Bjerketorp, B. Guss, L. Frykberg, A fibrinogen-binding protein of *Staphylococcus lugdunensis*, FEMS Microbiol. Lett. 241 (2004) 87–93.
- [16] A.S. Barbosa, P.A. Abreu, F.O. Neves, M.V. Atzingen, M.M. Watanabe, M.L. Vieira, Z.M. Morais, S.A. Vasconcellos, A.L. Nascimento, A newly identified leptospiral adhesin mediates attachment to laminin, Infect. Immun. 74 (2006) 6356–6364.
- [17] C.F. Barbas III, D.R. Burton, J.K. Scott, G.J. Silverman, Phage Display: A Laboratory Manual, first ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, 2001.
- [18] A.T. Ching, R.D. Fávoro, S.S. Lima, A. de Alencar Muniz Chaves, M.A. de Lima, H.B. Nader, P.A. Abreu, P.L. Ho, *Leptospira interrogans* shotgun phage display identified LigB as a heparin-binding protein, Biochem. Biophys. Res. Commun. 427 (2012) 774–779.
- [19] S. Antonara, R.M. Chafel, M. LaFrance, J. Coburn, *Borrelia burgdorferi* adhesins identified using in vivo phage display, Mol. Microbiol. 66 (2007) 262–276.
- [20] V.E. Papaioannou, J.G. Fox, Efficacy of tribromoethanol anesthesia in mice, Lab. Anim. Sci. 43 (1993) 189–192.
- [21] R.J. Giordano, M. Cardo-Vila, J. Lahdenranta, R. Pasqualini, W. Arap, Biopanning and rapid analysis of selective interactive ligands, Nat. Med. 7 (2001) 1249–1253.
- [22] C.R. Ramos, P.A. Abreu, A.L. Nascimento, P.L. Ho, A high-copy T7 *Escherichia coli* expression vector for the production of recombinant proteins with a minimal N-terminal His-tagged fusion peptide, Braz. J. Med. Biol. Res. 37 (2004) 1103–1109.
- [23] P. Hauk, S. Negrotto, E.C. Romero, S.A. Vasconcellos, M.E. Genovez, R.J. Ward, M. Schattner, R.M. Gómez, P.L. Ho, Expression and characterization of HlyX hemolysin from *Leptospira interrogans* serovar Copenhageni: potentiation of hemolytic activity by LipL32, Biochem. Biophys. Res. Commun. 333 (2005) 1341–1347.
- [24] P. Hauk, F. Macedo, E.C. Romero, S.A. Vasconcellos, Z.M. de Moraes, A.S. Barbosa, P.L. Ho, In LipL32, the major leptospiral lipoprotein, the C terminus is the primary immunogenic domain and mediates interaction with collagen IV and plasma fibronectin, Infect. Immun. 76 (2008) 2642–2650.
- [25] M. Gamberini, R.M. Gomez, M.V. Atzingen, E.A. Martins, S.A. Vasconcellos, E.C. Romero, L.C. Leite, P.L. Ho, A.L. Nascimento, Whole-genome analysis of *Leptospira interrogans* to identify potential vaccine candidates against leptospirosis, FEMS Microbiol. Lett. 244 (2005) 305–313.
- [26] E.F. Silva, M.A. Medeiros, A.J. McBride, J. Matsunaga, G.S. Esteves, J.G. Ramos, C.S. Santos, J. Croda, A. Homma, O.A. Dellagostin, D.A. Haake, M.G. Reis, A.I. Ko, The terminal portion of leptospiral immunoglobulin-like protein LigA confers protective immunity against lethal infection in the hamster model of leptospirosis, Vaccine 25 (2007) 6277–6286.
- [27] E. Carvalho, A.S. Barbosa, R.M. Gomez, A.M. Cianciarullo, P. Hauk, P.A. Abreu, L.C. Fiorini, M.L. Oliveira, E.C. Romero, A.P. Goncalves, Z.M. Morais, S.A. Vasconcellos, P.L. Ho, Leptospiral TlyC is an extracellular matrix-binding protein and does not present hemolysin activity, FEBS Lett. 583 (2009) 1381–1385.
- [28] V.L. Hashimoto, Z.M. Moraes, A.P. Gonçalves, S.A. Vasconcellos, P.L. Ho, Failure of LIC13435 protein of *Leptospira interrogans* serovar Copenhageni to confer protection in immunized hamsters, J. Vaccines Vaccin. 02 (2011) 120.
- [29] M.M. Castiblanco-Valencia, T.R. Fraga, L.B. Silva, D. Monaris, P.A. Abreu, S. Strobel, M. Jozsi, L. Isaac, A.S. Barbosa, Leptospiral immunoglobulin-like proteins interact with human complement regulators factor H, FHL-1, FHR-1, and C4BP, J. Infect. Dis. 205 (2012) 995–1004.
- [30] M. Nilsson, L. Frykberg, J.I. Flock, L. Pei, M. Lindberg, B. Guss, A fibrinogen-binding protein of *Staphylococcus epidermidis*, Infect. Immun. 66 (1998) 2666–2673.
- [31] J. Bjerketorp, M. Nilsson, A. Ljungh, J.I. Flock, K. Jacobsson, L. Frykberg, A novel von Willebrand factor binding protein expressed by *Staphylococcus aureus*, Microbiology 148 (2002) 2037–2044.
- [32] L.M. Mullen, S.P. Nair, J.M. Ward, A.N. Rycroft, R.J. Williams, B. Henderson, Comparative functional genomic analysis of Pasteurellaceae adhesins using phage display, Vet. Microbiol. 122 (2007) 123–134.
- [33] A. Kovacs-Simon, R.W. Titball, S.L. Michell, Lipoproteins of bacterial pathogens, Infect. Immun. 79 (2011) 548–561.
- [34] M. Madan Babu, K. Sankaran, DOLOP–database of bacterial lipoproteins, Bioinformatics 18 (2002) 641–643.
- [35] D.E. Hoke, S. Egan, P.A. Cullen, B. Adler, LipL32 is an extracellular matrix-interacting protein of *Leptospira* spp. and *Pseudomonas aeruginosa*, Infect. Immun. 76 (2008) 2063–2069.
- [36] M.V. Atzingen, A.S. Barbosa, T. De Brito, S.A. Vasconcellos, Z.M. de Moraes, D.M. Lima, P.A. Abreu, A.L. Nascimento, Lsa21, a novel leptospiral protein binding adhesive matrix molecules and present during human infection, BMC Microbiol. 8 (2008) 70.
- [37] H.A. Choy, M.M. Kelley, T.L. Chen, A.K. Moller, J. Matsunaga, D.A. Haake, Physiological osmotic induction of *Leptospira interrogans* adhesion: LigA and LigB bind extracellular matrix proteins and fibrinogen, Infect. Immun. 75 (2007) 2441–2450.
- [38] B. Stevenson, H.A. Choy, M. Pinne, M.L. Rotondi, M.C. Miller, E. Demoll, P. Kraicz, A.E. Cooley, T.P. Creamer, M.A. Suchard, C.A. Brissette, A. Verma, D.A. Haake, *Leptospira interrogans* endostatin-like outer membrane proteins bind host fibronectin, laminin and regulators of complement, PLoS One 2 (2007) e1188.
- [39] R.S. Mendes, M. Von Atzingen, Z.M. de Moraes, A.P. Goncalves, S.M. Serrano, A.F. Asega, E.C. Romero, S.A. Vasconcellos, A.L. Nascimento, The novel leptospiral surface adhesin Lsa20 binds laminin and human plasminogen and is probably expressed during infection, Infect. Immun. 79 (2011) 4657–4667.
- [40] R. Oliveira, Z.M. de Moraes, A.P. Goncalves, E.C. Romero, S.A. Vasconcellos, A.L. Nascimento, Characterization of novel OmpA-like protein of *Leptospira interrogans* that binds extracellular matrix molecules and plasminogen, PLoS One 6 (2011) e21962.

- [41] N.M. Souza, M.L. Vieira, I.J. Alves, Z.M. de Morais, S.A. Vasconcellos, A.L. Nascimento, Lsa30, a novel adhesin of *Leptospira interrogans* binds human plasminogen and the complement regulator C4bp, *Microb. Pathog.* 53 (2012) 125–134.
- [42] T.M. Witzemann, J.E. Adamou, S. Langermann, Adhesins as targets for vaccine development, *Emerg. Infect. Dis.* 5 (1999) 395–403.
- [43] M. Scarselli, M.M. Giuliani, J. Adu-Bobie, M. Pizza, R. Rappuoli, The impact of genomics on vaccine design, *Trends Biotechnol.* 23 (2005) 84–91.
- [44] G. Rajam, J.M. Anderton, G.M. Carlone, J.S. Sampson, E.W. Ades, Pneumococcal surface adhesin A (PsaA): a review, *Crit. Rev. Microbiol.* 34 (2008) 163–173.