



Contents lists available at [SciVerse ScienceDirect](#)

Biochemical and Biophysical Research Communications

journal homepage: www.elsevier.com/locate/ybbrc



Corrigendum

Corrigendum to “Modeling the Zn^{2+} and Cd^{2+} metalation mechanism in mammalian metallothionein 1a” [Biochem. Biophys. Res. Commun. 426 (2012) 601–607]

Duncan E.K. Sutherland^a, Kelly L. Summers^b, Martin J. Stillman^{a,b,*}

^a Department of Chemistry, The University of Western Ontario, London, Ontario, Canada N6A 5B7

^b Department of Biology, The University of Western Ontario, London, Ontario, Canada N6A 5B7

The authors regret an error in [Fig. 3](#). A corrected version of figure has been provided.

DOI of original article: <http://dx.doi.org/10.1016/j.bbrc.2012.08.136>

* Corresponding author. Address: Department of Chemistry, The University of Western Ontario, Richmond St., London, Ontario, Canada N6A 5B7. Fax: +1 519 661 3022.
E-mail address: martin.stillman@uwo.ca (M.J. Stillman).




Fig. 3. Experimental and theoretical data showing Cd²⁺ (A, B, C) and Zn²⁺ (D, E, F) speciation during metalation of the full MT protein, as well as the α- and β-domains. Modeled data: (G, H, I) A model using noncooperativity rules (declining K_{as}) of ESI-MS data. (J, K, L) A model using weakly cooperativity rules (equal K_{as}). (M, N, O) A model using strongly cooperativity rules (increasing K_{as}). Each line corresponds to a different metalation state: Zn⁰ (), Zn¹ (), Zn² (), Zn³ (), Zn⁴ (), Zn⁵ (), Zn⁶ () and Zn⁷ ().