

Current Topics

Delivery of Retinoid-Based Therapies To Target Tissues[†]

Alexander R. Moise,^{*,‡} Noa Noy,[‡] Krzysztof Palczewski,[‡] and William S. Blaner^{§,||}

Department of Pharmacology, School of Medicine, Case Western Reserve University, Cleveland, Ohio 44106-4965,
and Institute of Human Nutrition and Department of Medicine, College of Physicians and Surgeons,
Columbia University, New York, New York 10032

Received February 13, 2007; Revised Manuscript Received March 11, 2007

ABSTRACT: Through its various metabolites, vitamin A controls essential physiological functions. Both naturally occurring metabolites and novel retinoid analogues have shown effectiveness in many clinical settings that include skin diseases and cancer, and in animal models of human conditions affecting vision. In this review, we analyze several potential retinoid-based therapies from the point of view of drug metabolism and transport to target tissues. We focus on the endogenous factors that affect the absorption, transport, and metabolism of retinoids by taking into account data obtained from the analysis of animal models that lack the enzymes or proteins involved in the storage and absorption of retinoids. We also discuss findings of toxicity associated with retinoids in an effort to improve the outcome of retinoid-based therapies. In this context, we review evidence that esterification of retinol and retinol-based drugs within target tissues provides one of the most efficient means to improve the absorption and to reduce the toxicity associated with pharmacological doses of retinoids. Future retinoid-based therapeutic strategies could involve targeted delivery mechanisms leading to lower toxicity and improved effectiveness of retinoids.

Vitamin A, retinol, plays essential roles in many biological processes including vision, immunity, growth, development, and cellular differentiation. The various functions of vitamin A are carried out by several known physiologically active metabolites including 11-*cis*-retinal, the visual chromophore (1), and all-*trans*-retinoic acid (RA¹), which controls gene

expression through the RA receptor (RAR) (2, 3). Other retinol metabolites that have shown biological activities *in vitro* are the 9-*cis*-isomer of RA, which activates both retinoid X receptors (RXRs) and RARs (4), and the *retro*-retinoids, anhydroretinol (AR), and 14-hydroxy-4,14-*retro*-retinol (14-HRR), which are involved in the regulation of lymphocyte proliferation (5–7). All-*trans*-13,14-dihydroxy-retinol was shown to have activity comparable to that of 14-HRR in supporting the proliferation of promyelocytic HL-60 cells and in the activation of T cells (8). The conversion of all-*trans*-retinol to all-*trans*-13,14-dihydroretinol by the recently described enzyme all-*trans*-retinol–all-*trans*-13,14-dihydroretinol saturase (retinol saturase or RetSat) enzyme leads to the formation of all-*trans*-13,14-dihydroretinol (9), which can be further oxidized to all-*trans*-13,14-dihydroretinoic acid (10). The metabolism of vitamin A in relation to

[†] This research was supported in part by U.S. Public Health Service Grants EY01730, EY015399, and EY08061 from the National Eye Institute, National Institutes of Health, Bethesda, MD (to K.P.) and Grant DK068437 from the National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD (to W.S.B.).

^{*} To whom correspondence should be addressed. Phone: 216-368-1284. Fax: 216-368-1300. E-mail: ram50@case.edu.

[‡] Case Western Reserve University.

[§] Institute of Human Nutrition, Columbia University.

^{||} Department of Medicine, Columbia University.




FIGURE 1: Retinoid metabolism and bioactive retinoids in higher vertebrates. Known endogenous bioactive retinoids are highlighted by yellow boxes and include all-*trans*-retinoic acid (RA), which regulates gene expression via RAR, 11-*cis*-retinal, which is the chromophore of the photoreceptor molecule rhodopsin, all-*trans*-13,14-dihydroxyretinol, and the *retro*-retinoids, AR and 14-hydroxy-*retro*-retinol, which control cellular proliferation. All-*trans*-retinol can be esterified by the LRAT or by the ARAT enzyme to produce retinyl esters that can be hydrolyzed back to all-*trans*-retinol by retinyl ester hydrolase (REH) (reaction 1). The LRAT enzyme uses phospholipid (PL) as acyl donors and prefers retinol complexed to CRBP as the substrate. The oxidation of all-*trans*-retinol by short chain dehydrogenase/reductase (SDR) or ADH enzymes leads to the formation of all-*trans*-retinal (reaction 2), which in turn is oxidized by retinaldehyde dehydrogenases (RALDH) to all-*trans*-RA (reaction 3). All-*trans*-RA can be converted to more polar metabolites through oxidation by cytochrome P450 CYP26 enzymes (reaction 4) or through glucuronidation by UDP-glucuronosyltransferase (UGT) enzymes (reaction 5). All-*trans*-retinol can be saturated by RetSat and leads to the formation of all-*trans*-13,14-dihydroretinol (reaction 6), which can be oxidized by the same enzymes as those of all-*trans*-retinol to produce all-*trans*-13,14-dihydroretinoic acid. The production of 11-*cis*-retinal occurs in RPE cells and through esterification, isomerization, and oxidation reactions, with isomerization involving the RPE65 enzyme (reaction 7). An alternate pathway for the production of the visual chromophore for cones operates independently of RPE65 (reaction 8). The enzymes involved in the production of all-*trans*-13,14-dihydroxyretinol, AR, and 14-hydroxy-*retro*-retinol in vertebrates are not currently known, though a retinol dehydratase from insects was shown to catalyze the formation of AR.

the formation of known or potential endogenous bioactive retinoids is shown in Figure 1.

Therapeutic applications for retinoids have made use of both naturally occurring retinoids and synthetic agonists and antagonists of RAR and RXR. Supplementation with all-*trans*-retinol is used in developing countries to correct vitamin A deficiency (VAD). This treatment is one of the

safest and most efficacious therapeutic uses of retinoids and has enormous worldwide impact, saving millions of lives every year at a cost of pennies per treated individual (11). All-*trans*-RA (tretinoin, Retin-A) and 13-*cis*-RA (isotretinoin, Accutane) are clinically used in the treatment of severe acne and other dermatological conditions (12). Retinoids like all-*trans*-RA, 13-*cis*-RA, *N*-(4-hydroxyphenyl)retinamide (4-HPR or fenretinide), and non-retinoid synthetic subtype specific RAR and RXR-agonists and modulators are used in the treatment and chemoprevention of various forms of cancer (13–16). With respect to vision, there are several proposed retinoid-based therapies with potential in the treatment of autosomal dominant retinitis pigmentosa (ADRP), Leber congenital amaurosis (LCA), macular degeneration, or Stargardt disease (17). Supplementation of the substitute chromophore 9-*cis*-retinal can reconstitute the visual pigment in animal models of LCA (18–21). Inhibitors of the visual cycle, such as retinylamine (22) and 13-*cis*-RA (23), and reduction of serum retinol using 4-HPR (24) have shown effectiveness in animal models of Stargardt disease and

¹ Abbreviations: A2E, pyridinium bis-retinoid; ADH, medium-chain alcohol dehydrogenases; ADRP, autosomal dominant retinitis pigmentosa; apoE, apolipoprotein E; AR, anhydroretinol; ARAT, acyl-CoA-retinol acyltransferase; CRBP, cellular retinol-binding protein; DGAT1, diacylglycerol acyltransferase 1; 4-HPR, *N*-(4-hydroxyphenyl)retinamide (or fenretinide); 14-HRR, 14-hydroxy-4,14-*retro*-retinol; LCA, Leber congenital amaurosis; LPL, lipoprotein lipase; LRAT, lecithin-retinol acyltransferase; RA, retinoic acid; 13-*cis*-RA, 13-*cis*-retinoic acid (or isotretinoin, Accutane); RALDH, retinaldehyde dehydrogenase; RAR, retinoic acid receptor; RBP, retinol-binding protein; RetSat, all-*trans*-retinol—all-*trans*-13,14-dihydroretinol saturase (or retinol saturase); RPE65, retinal pigmented epithelial-specific 65 kDa protein; RXR, retinoid-X receptor; SDR, short-chain dehydrogenase/reductase; STRA6, stimulated by retinoic acid 6; TTR, transthyretin; UGT1, UDP-glucuronosyltransferase 1; vitamin A, all-*trans*-retinol; VAD, vitamin A deficiency; VLDL, very low-density lipoprotein.




FIGURE 2: Targets of retinoid treatments. Both all-*trans*-retinol and all-*trans*-retinyl esters (palmitate) are effective in the treatment and prevention of vitamin A deficiency (VAD). The reconstitution of rhodopsin with 9-*cis*-retinal forms isorhodopsin and is an effective means of restoring visual function in several models of Leber congenital amaurosis (LCA). Inhibitors of the visual cycle, retinylamine and 13-*cis*-RA, can slow the rate of 11-*cis*-retinal formation in cases where excessive activation of rhodopsin can lead to phototoxicity and accumulation of all-*trans*-retinal and toxic metabolites as seen in Stargardt disease and age-related macular degeneration. A similar result can also be obtained with *N*-(4-hydroxyphenyl)retinamide (4-HPR or fenretinide), which binds RBP and leads to its excretion through glomerular filtration. Retinylamine can be amidated by the LRAT enzyme, allowing for the storage of the drug in an inactive form with lower toxicity. Independently of RAR, 4-HPR can lead to the apoptosis of tumor cells. RAR can be modulated by all-*trans*-RA, 9-*cis*-RA, and many synthetic agonists and antagonists. The retinoid X receptor can be modulated by 9-*cis*-RA and several synthetic agonists and antagonists. Both 13-*cis*- and 9-*cis*-RA can isomerize to all-*trans*-RA *in vivo*. The drug targets are shown in bold font, and the retinoids used in various therapies are shown in red font.

macular degeneration by slowing the rate of the visual cycle and preventing the accumulation of all-*trans*-retinal and the production of phototoxic byproducts such as pyridinium bis-retinoid (A2E) (17). Some of the currently investigated retinoid-based drugs and therapeutic targets are depicted in Figure 2.

Retinoid therapies have demonstrated clinical efficacy as treatments for many debilitating diseases; however, one must overcome significant obstacles associated with the well-established teratogenic and toxic effects of retinoids. Many of these obstacles could be overcome if the delivery mechanism is more specific. In this review, we focus on factors affecting the delivery of retinol and other retinoids to specific target tissues.

ABSORPTION OF RETINOIDS

Retinoids delivered orally are efficiently absorbed by the intestine. Retinyl esters are hydrolyzed to retinol within the intestinal lumen by the actions of pancreatic triglyceride lipase (25) or at the intestinal brush border by the membrane-bound enzyme phospholipase B (26, 27). The newly generated retinol as well as the retinol obtained from the diet or supplements is taken into the enterocyte as retinol. Because of the lipophilic character of retinol, absorption is enhanced by a fatty meal (28). Within the enterocyte, newly absorbed retinol is re-esterified through the actions of the enzyme lecithin retinol-acyl transferase (LRAT), which catalyzes the transacylation of a fatty acyl group from lecithin to retinol (29–33). Cellular retinol-binding protein II (CRBP-II), a member of the family of intracellular lipid-binding proteins, which binds retinol with high affinity, helps facilitate retinol uptake into the body providing the retinol as a substrate for

LRAT (34, 35). Although LRAT is the predominant enzyme for catalyzing retinyl ester formation within the intestine, a study of *Lrat*^{−/−} mice indicates that an acyl-CoA-dependent enzyme, referred to as acyl-CoA-retinol acyltransferase (ARAT), may also be involved in retinyl ester formation in the liver, mammary gland, or adipose tissue (32, 36). Ultimately, the retinyl ester newly synthesized by LRAT and/or ARAT is incorporated along with other dietary lipids into nascent chylomicrons that are then secreted from the enterocyte into the lymphatic system (37).

TRANSPORT OF RETINOIDS TO TARGET TISSUES

Role of Chylomicrons and Their Remnants. Nascent chylomicrons in the lymphatic system enter the general circulation through the thoracic duct where they undergo metabolism to form chylomicron remnants. The conversion of chylomicrons to chylomicron remnants involves the actions of lipoprotein lipase (LPL), which hydrolyzes chylomicron triglycerides to free fatty acids, resulting in a smaller particle and facilitating the recruitment to the lipolyzed particle of several apolipoproteins, especially apolipoprotein E (apoE), that are needed for receptor-mediated uptake of the remnants by the liver (38, 39). About 66–75% of retinyl esters contained in chylomicrons are taken up by the liver and resecreted into the circulation as retinol bound to serum retinol-binding protein (RBP) or stored as esters in hepatic stellate cells (also called Ito cells, lipocytes, or fat-storing cells) (40, 41). The remaining 25–33% of the retinyl esters from chylomicrons are taken up by target tissues that include adipose tissue, heart, muscle, lungs, reproductive organs, and bone marrow. The clearance of chylomicron

retinoids appears to involve the actions of LPL (42). This reaction most likely involves LPL-catalyzed retinyl ester hydrolysis but possibly also the well-characterized actions of LPL in facilitating whole particle uptake by cells (42, 43). The importance of chylomicron-derived retinoid toward meeting target tissue needs is evidenced by RBP-deficient mice, which, aside from a mild vision phenotype, are phenotypically normal (44–46). As long as RBP-deficient mice are provided sufficient quantities of vitamin A in the diet, they are able to maintain normal retinoid-dependent functions and even to acquire retinyl ester stores in target tissues. When these mutant mice are deprived of dietary retinoid, they are at increased risk of developing symptoms of VAD because they are not able to mobilize hepatic retinyl ester stores through the actions of RBP (44–46). Thus, the delivery of retinoids by chylomicrons and their remnants can be an important route for retinoid delivery to target tissues.

Role of RBP in the Delivery of Retinol. As mentioned earlier, unesterified retinol is transported in the blood as retinol bound to RBP (47). RBP is primarily synthesized and secreted from the liver, where the majority of the body's retinoid is stored (48, 49). However, RBP also is synthesized and secreted in lesser amounts by extrahepatic tissues, including the eye, adipose tissue, kidney, lungs, and heart (50). RBP forms a one-to-one protein–protein complex with another blood protein transthyretin (TTR), which prevents renal clearance of the retinol–RBP by the glomerulus (51–53). Hepatic secretion of holo-RBP is needed for the efficient export of retinol from liver stores (44–46). There is some controversy regarding how retinol is taken up by cells from the retinol–RBP–TTR complex (54). Retinol rapidly dissociates from RBP and rapidly flip–flops across the plasma membrane (55). Moreover, the rate of dissociation of retinol from RBP is not rate limiting for retinol uptake by cells, which argues that it is not necessary to invoke the actions of a cell surface receptor for RBP to account for retinol uptake into cells (54, 56, 57). Yet, other studies suggest that retinol uptake into cells involves a cell surface receptor for RBP (54, 58–62). One possible receptor is coded by a retinoic acid-responsive gene, *STRA6* (stimulated by retinoic acid-6) (63). The *STRA6* protein is a multispinning membrane protein that confers RBP binding to transfected cells and mediates the uptake of retinol (62). *STRA6* is expressed in several blood–organ barriers, such as Sertoli cells, the yolk sac, chorioallantoic placenta, choroid plexi (64), and RPE (62, 64). Mutations in *STRA6* cause severe malformations, such as anophthalmia, congenital heart defects, diaphragmatic hernia, alveolar capillary dysplasia, lung hypoplasia, and mental retardation (65). However, patients affected by deleterious mutations in the RBP gene suffer from night-blindness early in life but are otherwise unaffected (66). A similar phenotype is seen in *Rbp*^{−/−} mice, manifested as visual impairment early in life (44). These observations argue for additional roles for the *STRA6* protein during development that are independent of its role as a receptor for RBP (67).

The retinoid needs for all tissues aside from the eye can be fully met through chylomicron delivery of retinoid or retinoid bound to serum albumin, as is evidenced by the visual phenotype of the *Rbp*^{−/−} mice (44). Thus, the eye may rely on the *STRA6* receptor to ensure sufficient retinoid uptake from RBP to support the visual cycle. However,

because the complete absence of RBP does not impair retinoid-dependent functions in other tissues, a cell surface receptor for RBP may not be needed to ensure the normal uptake of retinol by such tissues in times of sufficient dietary vitamin A intake.

RA Delivery by Serum Albumin. RA is far less hydrophobic than retinol or retinyl ester, and it is not absorbed in chylomicrons but rather transported via the portal system bound to albumin because binding to RBP hinders its interaction with TTR (68). Pharmacokinetic studies of many of the synthetic analogues of RA indicate that they are absorbed primarily via the portal system like the natural acidic retinoid forms (69). However, depending on the lipophilic nature of a synthetic retinoid, some of its absorption may involve uptake via nascent chylomicrons, or it may be taken up solely via the portal route. This absorption will depend on the chemical properties of the particular retinoid.

In well-nourished humans and animal models, the circulating levels of all-*trans*-RA and 13-*cis*-RA bound to albumin are approximately 0.1 to 0.4% of those of retinol–RBP during fasting (70). Yet, this circulating RA can contribute significantly to tissue pools of RA in well-nourished animals (71). Tracer kinetic turnover studies in rats have shown that, for most tissues, 5–30% of the RA present within the tissue is derived from the circulating RA pool, whereas the brain and liver derive greater than 80% of their RA from the circulating RA pool (71). Thus, despite the low levels of all-*trans*- and 13-*cis*-RA in circulation, these acids are rapidly taken up by cells and tissues. Albumin-bound RA can be an important delivery pathway through which cells/tissues normally acquire retinoid. Like retinol, RA undergoes dissociation from albumin prior to traversing the membrane and entering into the cell, and RA is known to flip–flop readily across membrane bilayers. The delivery of synthetic RA analogues via albumin is pharmacologically important, and presumably, this process mimics RA uptake. However, there is little information available in the literature regarding how synthetic retinoids enter cells/tissues from the circulation.

Other Routes of Retinoid Delivery. In humans as well as in animal models, retinyl esters are normally secreted at low levels (1–2% of those of retinol–RBP) from the liver bound to nascent very low-density lipoprotein (VLDL) (72). The physiologic significance of VLDL retinyl ester has not been systematically investigated, but, invoking similarities between VLDL and chylomicron metabolism, it seems likely that the retinyl ester contained within VLDL is destined for uptake by extrahepatic tissues. The circulation also contains low levels of retinol- β -glucuronides and retinoyl- β -glucuronides (levels similar to those of all-*trans*- and 13-*cis*-RA) that are fully water soluble (54, 73, 74), hence more easily excreted. Alternatively, these retinoid- β -glucuronides may be used by the body to satisfy tissue retinoid needs because they can be readily hydrolyzed to retinol and RA (54, 73, 74).

ROLE OF LRAT IN THE ABSORPTION AND STORAGE OF RETINOL

As is true for the liver and RPE, LRAT is quantitatively the most important enzyme for catalyzing retinyl ester formation in most tissues (20, 32, 75). Consequently, LRAT is essential for the accumulation of retinoids within the liver

and eye. In the liver, retinyl esters are found predominantly in hepatic stellate cells, which represent the main cellular retinoid storage site in the body. RPE cells also represent an important storage of retinyl esters reserved mostly for use in the production of the visual chromophore. The importance of LRAT in allowing tissues to accumulate and store retinol is evidenced by the complete absence of retinyl esters in nearly all of the tissues that have been examined in *Lrat*^{-/-} mice. The one exception to this general statement is adipose tissue, where retinyl ester levels are 2- to 3-fold elevated in *Lrat*^{-/-} mice compared to those in age- and gender-matched wild-type mice (20, 32). Although adipocytes have long been known to be important cellular sites of retinyl ester accumulation in the body accounting for the second largest retinoid store after the liver, little is known regarding how retinol is processed and becomes esterified within this cell type (76–80). LRAT is not needed to catalyze retinyl ester formation in adipose tissue but it is not presently clear which enzyme is responsible for retinyl ester formation within this tissue. The diacylglycerol acyltransferase 1 (DGAT1) enzyme, which catalyzes the formation of triglyceride from diacylglycerol and an acyl-CoA, catalyzes acyl-CoA-dependent retinol esterification *in vitro*, hence exhibiting ARAT activity (32, 81, 82). However, DGAT1, although highly expressed in adipose tissue, does not seem to be the only enzyme involved in catalyzing retinyl ester formation, because adipose tissue retinyl ester levels are the same for *Lrat*^{-/-} mice and *Lrat*^{-/-}/*Dgat1*^{-/-} double knockout mice (Blaner, W. et al., unpublished observations). Thus, it remains unclear which other enzymes can catalyze retinyl ester formation and facilitate the accumulation of retinoid stores within adipose tissue.

In addition to allowing for retinoid accumulation within tissues, the sequestration of retinol as retinyl ester also serves to limit the generation of RA during development (83) and acts as a protective mechanism following the consumption of high doses of dietary vitamin A (84). Although the levels of serum retinol and RA are within the normal range, *Lrat*^{-/-} mice are not capable of sequestering retinol as retinyl esters, thus exhibiting signs of RA toxicity when maintained on a vitamin A-sufficient chow diet. This observation is also supported by the upregulation in *Lrat*^{-/-} mice of Cyp26 genes that are known to have central roles in the detoxification (catabolism) of RA (75). In addition, when we employed microarray analysis to identify other differentially expressed genes in the livers of *Lrat*^{-/-} mice and their wild-type counterparts, we found that among the genes upregulated in the liver of *Lrat*^{-/-} mice maintained on a standard chow diet were several isoforms of the UDP-glucuronosyltransferase 1 (UGT1) family of enzymes (85) (Table 1). The UGT1 locus is known to be subject to regulation by RA (86). As mentioned above, β -glucuronidation products of all-*trans*-RA, 13-*cis*-RA, 4-oxo-RA, and 5,6-epoxy-RA are normally present at low levels in the blood and bile and at greater concentrations following supplementation with large doses of retinol or RA (87–89). Elevated expression levels of UGT1 in the livers of *Lrat*^{-/-} mice could be taken to suggest that the liver is sensing excessive levels of RA, leading to the induction of detoxification mechanisms.

Esterification by LRAT leads to active recruitment of the retinoid from circulation as well as the establishment of a long-lived storage pool. This property of LRAT can be

manipulated for therapeutic advantage for the delivery of retinol and other retinoids with an alcohol functional group. Another potential retinoid-based therapeutic retinylamine has a reactive amine group instead of alcohol, yet it can also be recruited and stored via amidation by LRAT (90) (Figure 2). The conversion of retinylamine to pharmacologically inactive retinylamides occurs *in vitro* and *in vivo* in the liver and RPE and leads to the safe storage of this potent inhibitor, which can be released after hydrolysis back to free retinylamine. This transformation leads to a prolonged therapeutic effect and lower toxicity of this compound in comparison with those of other proposed visual cycle inhibitors such as 13-*cis*-RA (90). In summary, the presence of LRAT in the target tissue can ensure the efficient and active uptake of retinol and other retinoids with a functional hydroxyl group as well as retinoids that can be amidated by LRAT.

ROLE OF CRBPS IN THE UPTAKE OF RETINOID BY SPECIFIC TISSUES

Three cellular-retinol-binding proteins, CRBP-I, II, and III, are known to facilitate the esterification of retinoids by LRAT. The physiologic roles of LRAT and the CRBPs in the absorption and storage of retinol have been best studied in knockout animal models. CRBP-I is expressed in the liver, kidney, and eye, and *Crbp-I*^{-/-} mice display lower levels of hepatic retinyl esters and reduced numbers of lipid droplets in hepatic stellate cells than wild-type mice. *Crbp-I*^{-/-} mice succumb more easily to symptoms of VAD than wild-type mice if they are maintained on a vitamin A-restricted diet (91). CRBP-II is expressed primarily in the intestine, and its lack in *Crbp-II*^{-/-} mice results in decreased levels of stored retinoids and increased neonatal mortality, especially during dietary vitamin A restriction (92). CRBP-III is expressed in the heart, muscle, adipose, and mammary tissues (93). *Crbp-III*^{-/-} mice have difficulty incorporating retinol into milk. In some cases, the loss of one form of CRBP can be partly compensated for by the upregulation of other CRBPs, as is the case of the upregulation of CRBP-III in the absence of CRBP-I and vice versa (94). Although the studies of *Crbp*^{-/-} mice demonstrate that the absence of these proteins results in a much less severe metabolic phenotype than that observed with *Lrat*^{-/-} mice, these studies do show that like LRAT, the CRBPs have an important role in facilitating retinol uptake, esterification, and storage by cells (20, 32, 91–93).

The critical role of LRAT in catalyzing the esterification and thus the uptake and storage of retinol suggests that this enzyme may be a useful target for assuring the specific uptake of a retinoid therapeutic agent by a target tissue. Coupled with the hydrolysis of the esters, this potentially provides a way to localize release of the drug within a specific cell type or tissue. The release of stored retinoid drugs from their ester pools depends on the activity of retinyl ester hydrolases in the respective target tissues. Manipulating this reaction may provide a means for a controlled delivery of therapeutic retinoids, which, in principle, would allow for the usage of lower administered doses and hence lower toxicity. Moreover, the expression of LRAT is subject to regulation by RA (95, 96). STRA6 was described as an RA-responsive gene (63), and modulation of the levels of STRA6 in WiDr colon adenocarcinoma cells by RA leads to the increased uptake of retinol in these cells (62). As a result,

Table 1: Differentially Expressed Genes in the Liver of *Lrat*^{-/-} Mice vs WT^a

gene symbol	gene description	accession no.	fold change <i>Lrat</i> ^{-/-} /WT
upregulated in <i>Lrat</i> ^{-/-} mice vs WT			
<i>Ugt1a9</i>	UDP-glucuronosyltransferase 1 family A9	S57479	7.9
<i>Apoa4</i>	apolipoprotein A-IV	M64249	6.0
<i>Mt1</i>	metallothionein 1	BC027262	4.4
<i>Alas1</i>	aminolevulinate synthase 1	M63245	4.4
<i>Gdpd3</i>	Glycerophosphodiester phosphodiesterase domain containing 3	NM_024228	3.6
<i>Cxcl14</i>	chemokine (C-X-C motif) ligand 14	AF352785	3.3
<i>Hspa8</i>	heat shock protein 8	X05837	3.1
<i>Serpina4-ps1</i>	serpin A4, pseudogene 1	BC024071	3.0
<i>Hsp110</i>	heat shock protein 110	AK075731	3.0
<i>Dhtkd1</i>	dehydrogenase E1 and transketolase domain containing 1	AK050057	2.7
<i>Hspa1b</i>	heat shock protein 1B	M12573	2.5
<i>Lcn13</i>	lipocalin 13	NM_153558	2.5
<i>Cap1</i>	adenylate cyclase- associated protein 1	NM_007598	2.4
<i>Tppp</i>	tubulin polymerization- promoting protein	NM_182839	2.3
<i>Tsc22d3</i>	TSC22-related inducible leucine zipper 3	AF024519	2.3
downregulated in <i>Lrat</i> ^{-/-} mice vs WT			
<i>Clec2h</i>	C-type lectin domain family 2, member h	NM_053165	0.49
<i>MGC60813</i>	unknown	BC058613	0.45
<i>Tubb2a</i>	tubulin, beta 2 a	NM_009450	0.44
<i>Cml5</i>	camello-like 5	AK007530	0.43
<i>Cg10671</i>	Cg10671 like	BC048823	0.43
<i>Mm.2121</i>	unknown	NM_026790	0.38
<i>Tiam2</i>	T-cell lymphoma invasion and metastasis 2	NM_011878	0.38
<i>Cebpe</i>	CCAAT/enhancer binding protein epsilon	NM_207131	0.37
<i>Hsd3b5</i>	hydroxy-delta-5-steroid dehydrogenase	NM_008295	0.32
<i>Camk2b</i>	calcium/calmodulin- dependent protein kinase II, beta	AK083344	0.19
<i>Gadd45g</i>	growth arrest and DNA-damage- inducible 45 gamma	AK002237	0.15

^a Total RNA from the three livers of 3 month old *Lrat*^{-/-} (or WT) mice maintained on a standard diet were pooled to generate one sample. We used two different pools of hepatic RNA from *Lrat*^{-/-} mice to compare them with two different pools of WT mice. Each sample of pooled RNA was detection-labeled and hybridized in duplicate on a mouse genomic microarray using a service provided by NimbleGen System, Inc. (Madison, WI) as described previously (139). The microarray contained 37,364 genes and covered the entire mouse transcriptome as represented by the University of California, Santa Cruz database (build HG 17), using a minimum of 11 probes per gene. The expression of genes was normalized according to the probe signal, and the average signal for each gene was normalized for each sample replicate. Array data for samples across the whole study were normalized by NimbleGen Systems, Inc. (Madison, WI), using the robust multichip analysis feature of the data analysis package found at <http://www.bioconductor.org>. Project-wide spreadsheets of robust multichip analysis results were exported to Microsoft Excel. Expression level ratios for all of the possible pairwise comparisons, comprising a four-way comparison of two *Lrat*^{-/-} and two WT samples, were calculated. These pairwise ratios were imported to Microsoft Access and mined for credible fold changes. Changes in gene expression greater than or equal to 2-fold in at least three of the four comparisons were considered significant.

induction of the expression of LRAT or STRA6 using RA could increase the delivery of retinol to target tissues. Indeed, co-administration of RA effectively increases the delivery of retinol to lungs (97, 98). In addition, the expression of CRBP-I is also upregulated via RAR action (99), whereas CRBP-II expression is controlled via RXR (100). In fact, agonists of RXR have been shown to affect the esterification of retinol in certain cell types (101). Possibly, ectopic expression of STRA6, LRAT, and/or CRBP via gene therapy

or transcriptional regulation could help ensure the delivery of retinoids to specific target tissues or cell types. Both LRAT and CRBP are downregulated in many tumor cell types (102–117); therefore, less retinol is available for oxidation to produce RA in cancer cells. Upregulation of the retinoid pool in tumor cells might lead to higher endogenous levels of RA at the tumor site enabling the anticarcinogenic activities of this agent. Future studies could examine the feasibility and therapeutic effectiveness of increasing the

intracellular retinoid pool through the upregulation of the enzymes and accessory proteins involved in the generation of retinyl esters.

RETINOL TOXICITY AND THE SIDE EFFECTS ASSOCIATED WITH RETINOID TREATMENT

Excess levels of vitamin A lead to a wide range of deleterious effects on health. Retinoids present a serious teratogenic risk and can lead to craniofacial, cardiac, thymic, and central nervous system malformations (118), which are the most severe manifestations of toxicity associated with retinoid therapy. Teratogenic effects are seen for both natural and synthetic retinoids used in clinical trials (119, 120), but these are less pronounced for preformed retinol supplements (121). The administration of retinyl palmitate to rats, mice, and hamsters led to exencephaly, with frequent occurrences of cleft palate, spina bifida, eye defects, hydrocephaly, and shortening of the mandible and maxilla (122–125). The teratogenic effects are believed to stem mainly from the transcriptional activities of retinoids, mediated via the activation of RAR or RXR. The teratogenic consequences of retinoid supplementation during pregnancy depend on the stage of development, the nature of the retinoid metabolite, and the genetic and environmental factors (120). Caution must be exercised in prescribing retinoids to ensure that pregnancy is avoided during treatment (126). Additionally, the administration of retinoids to young animals and children results in delayed growth as a result of premature epiphyseal plate closure (127, 128).

Less severe consequences of retinoid supplementation result in acute and chronic forms of retinoid toxicity. Acute intoxication with vitamin A occurs following the ingestion of large amounts of vitamin A over a short period of time, but cases of acute toxicity are relatively rare. The clinical effective dose for vitamin A administration in the case of degenerative eye diseases such as retinitis pigmentosa (15,000 IU/day) is below the maximal recommended dose (25,000 IU/day) and was shown to be well tolerated after 12 years of treatment (129, 130). In the few documented cases, recovery from acute retinoid toxicity is rapid once intake is reduced (131, 132). Chronic effects are manifested as drying, ulceration and desquamation of the skin, anorexia, anemia, and bone loss. Liver damage leading to fibrosis and hepatic stellate cell activation are both seen in patients affected by hypervitaminosis A (133). Upon activation, hepatic stellate cells lose their capacity to store retinyl esters and secrete collagen, leading to cirrhosis (132, 134). Oxidation of retinol generates RA, which stimulates osteoclast formation and activity leading to bone resorption and hypercalcemia (135, 136). Bone loss could represent a health issue in the elderly, leading to osteopenia, osteoporosis, and increased risk of hip fracture (137, 138).

CONCLUSIONS

Vitamin A metabolites, including RAs and their synthetic derivatives, collectively known as retinoids, modulate the rates of transcription of the genes involved in the regulation of cell proliferation and differentiation and thus exhibit chemotherapeutic and chemopreventive activities in a number of human cancers. Another vitamin A metabolite, 11-*cis*-retinal, serves as the visual chromophore, and several

proposed retinoid-based therapies show promise in the treatment of eye diseases such as ADRP, LCA, and macular degeneration. Retinoid-based therapy is, however, often confounded by the toxicity of these compounds at pharmacological doses. Recent observations suggest that factors that regulate the uptake and storage of retinoids in cells may serve as targets for novel retinoid-based therapeutic strategies. Such factors include LRAT, the enzyme that catalyzes retinoid esterification and storage, and CRBPs, proteins that cooperate with LRAT in regulating retinoid uptake and metabolism. Manipulating the levels or activities of these proteins may allow for a better control of the levels of active retinoids in specific tissues, thereby reducing the amounts needed for efficacious therapy and targeting the agents to specific tissues.

REFERENCES

- Wald, G. (1968) Molecular basis of visual excitation, *Science* 162, 230–239.
- Chambon, P. (1996) A decade of molecular biology of retinoic acid receptors, *FASEB J.* 10, 940–954.
- Mark, M., Ghyselinck, N. B., and Chambon, P. (2006) Function of retinoid nuclear receptors: lessons from genetic and pharmacological dissections of the retinoic acid signaling pathway during mouse embryogenesis, *Annu. Rev. Pharmacol. Toxicol.* 46, 451–480.
- Heyman, R. A., Mangelsdorf, D. J., Dyck, J. A., Stein, R. B., Eichele, G., Evans, R. M., and Thaller, C. (1992) 9-*cis* retinoic acid is a high affinity ligand for the retinoid X receptor, *Cell* 68, 397–406.
- Buck, J., Derguini, F., Levi, E., Nakanishi, K., and Hammerling, U. (1991) Intracellular signaling by 14-hydroxy-4,14-retro-retinol, *Science* 254, 1654–1656.
- Buck, J., Grun, F., Derguini, F., Chen, Y., Kimura, S., Noy, N., and Hammerling, U. (1993) Anhydroretinol: a naturally occurring inhibitor of lymphocyte physiology, *J. Exp. Med.* 178, 675–680.
- Derguini, F., Nakanishi, K., Hammerling, U., and Buck, J. (1994) Intracellular signaling activity of synthetic (14R)-, (14S)-, and (14RS)-14-hydroxy-4,14-retro-retinol, *Biochemistry* 33, 623–628.
- Derguini, F., Nakanishi, K., Hammerling, U., Chua, R., Eppinger, T., Levi, E., and Buck, J. (1995) 13,14-Dihydroxy-retinol, a new bioactive retinol metabolite, *J. Biol. Chem.* 270, 18875–18880.
- Moise, A. R., Kuksa, V., Imanishi, Y., and Palczewski, K. (2004) Identification of all-trans-retinol:all-trans-13,14-dihydroretinol saturase, *J. Biol. Chem.* 279, 50230–50242.
- Moise, A. R., Kuksa, V., Blaner, W. S., Baehr, W., and Palczewski, K. (2005) Metabolism and transactivation activity of 13,14-dihydroretinoic acid, *J. Biol. Chem.* 280, 27815–27825.
- Humphrey, J. H., West, K. P., Jr., and Sommer, A. (1992) Vitamin A deficiency and attributable mortality among under-5-year-olds, *Bull. W. H. O.* 70, 225–232.
- Chivot, M. (2005) Retinoid therapy for acne. A comparative review, *Am. J. Clin. Dermatol.* 6, 13–19.
- Chomienne, C., Ballerini, P., Balitrand, N., Amar, M., Bernard, J. F., Boivin, P., Daniel, M. T., Berger, R., Castaigne, S., and Degos, L. (1989) Retinoic acid therapy for promyelocytic leukaemia, *Lancet* 2, 746–747.
- Hong, W. K., and Sporn, M. B. (1997) Recent advances in chemoprevention of cancer, *Science* 278, 1073–1077.
- Simeone, A. M., and Tari, A. M. (2004) How retinoids regulate breast cancer cell proliferation and apoptosis, *Cell. Mol. Life Sci.* 61, 1475–1484.
- Hail, N., Jr., Kim, H. J., and Lotan, R. (2006) Mechanisms of fenretinide-induced apoptosis, *Apoptosis* 11, 1677–1694.
- Travis, G. H., Golczak, M., Moise, A. R., and Palczewski, K. (2007) Diseases caused by defects in the visual cycle: retinoids as potential therapeutic agents, *Annu. Rev. Pharmacol. Toxicol.* 47, 469–512.
- Van Hooser, J. P., Aleman, T. S., He, Y. G., Cideciyan, A. V., Kuksa, V., Pittler, S. J., Stone, E. M., Jacobson, S. G., and Palczewski, K. (2000) Rapid restoration of visual pigment and function with oral retinoid in a mouse model of childhood blindness, *Proc. Natl. Acad. Sci. U.S.A.* 97, 8623–8628.

19. Van Hooser, J. P., Liang, Y., Maeda, T., Kuksa, V., Jang, G. F., He, Y. G., Rieke, F., Fong, H. K., Detwiler, P. B., and Palczewski, K. (2002) Recovery of visual functions in a mouse model of Leber congenital amaurosis, *J. Biol. Chem.* 277, 19173–19182.
20. Batten, M. L., Imanishi, Y., Tu, D. C., Doan, T., Zhu, L., Pang, J., Glushakova, L., Moise, A. R., Baehr, W., Van Gelder, R. N., Hauswirth, W. W., Rieke, F., and Palczewski, K. (2005) Pharmacological and rAAV gene therapy rescue of visual functions in a blind mouse model of Leber congenital amaurosis, *PLoS Med.* 2, e333.
21. Maeda, A., Maeda, T., and Palczewski, K. (2006) Improvement in rod and cone function in mouse model of *Fundus albipunctatus* after pharmacologic treatment with 9-cis-retinal, *Invest. Ophthalmol. Visual Sci.* 47, 4540–4546.
22. Golczak, M., Kuksa, V., Maeda, T., Moise, A. R., and Palczewski, K. (2005) Positively charged retinoids are potent and selective inhibitors of the trans-cis isomerization in the retinoid (visual) cycle, *Proc. Natl. Acad. Sci. U.S.A.* 102, 8162–8167.
23. Radu, R. A., Mata, N. L., Nusinowitz, S., Liu, X., and Travis, G. H. (2004) Isotretinoin treatment inhibits lipofuscin accumulation in a mouse model of recessive Stargardt's macular degeneration, *Novartis Found. Symp.* 255, 51–63; discussion 63–67, 177–178.
24. Radu, R. A., Han, Y., Bui, T. V., Nusinowitz, S., Bok, D., Lichter, J., Widder, K., Travis, G. H., and Mata, N. L. (2005) Reductions in serum vitamin A arrest accumulation of toxic retinal fluorophores: a potential therapy for treatment of lipofuscin-based retinal diseases, *Invest. Ophthalmol. Visual Sci.* 46, 4393–4401.
25. van Bennekum, A. M., Fisher, E. A., Blaner, W. S., and Harrison, E. H. (2000) Hydrolysis of retinyl esters by pancreatic triglyceride lipase, *Biochemistry* 39, 4900–4906.
26. Rigrup, K. M., Kakkad, B., and Ong, D. E. (1994) Purification and partial characterization of a retinyl ester hydrolase from the brush border of rat small intestine mucosa: probable identity with brush border phospholipase B, *Biochemistry* 33, 2661–2666.
27. Harrison, E. H. (2005) Mechanisms of digestion and absorption of dietary vitamin A, *Annu. Rev. Nutr.* 25, 87–103.
28. Tso, P., Lee, T., and DeMichele, S. J. (2001) Randomized structured triglycerides increase lymphatic absorption of tocopherol and retinol compared with the equivalent physical mixture in a rat model of fat malabsorption, *J. Nutr.* 131, 2157–2163.
29. MacDonald, P. N., and Ong, D. E. (1988) A lecithin:retinol acyltransferase activity in human and rat liver, *Biochem. Biophys. Res. Commun.* 156, 157–163.
30. MacDonald, P. N., and Ong, D. E. (1988) Evidence for a lecithin:retinol acyltransferase activity in the rat small intestine, *J. Biol. Chem.* 263, 12478–12482.
31. Ruiz, A., Winston, A., Lim, Y. H., Gilbert, B. A., Rando, R. R., and Bok, D. (1999) Molecular and biochemical characterization of lecithin retinol acyltransferase, *J. Biol. Chem.* 274, 3834–3841.
32. O'Byrne, S. M., Wongsiriroj, N., Libien, J., Vogel, S., Goldberg, I. J., Baehr, W., Palczewski, K., and Blaner, W. S. (2005) Retinoid absorption and storage is impaired in mice lacking lecithin:retinol acyltransferase (LRAT), *J. Biol. Chem.* 280, 35647–35657.
33. Batten, M. L., Imanishi, Y., Maeda, T., Tu, D. C., Moise, A. R., Bronson, D., Possin, D., Van Gelder, R. N., Baehr, W., and Palczewski, K. (2004) Lecithin-retinol acyltransferase is essential for accumulation of all-trans-retinyl esters in the eye and in the liver, *J. Biol. Chem.* 279, 10422–10432.
34. Herr, F. M., and Ong, D. E. (1992) Differential interaction of lecithin-retinol acyltransferase with cellular retinol binding proteins, *Biochemistry* 31, 6748–6755.
35. Ong, D. E. (1994) Cellular transport and metabolism of vitamin A: roles of the cellular retinoid-binding proteins, *Nutr. Rev.* 52, S24–S31.
36. Randolph, R. K., Winkler, K. E., and Ross, A. C. (1991) Fatty acyl CoA-dependent and -independent retinol esterification by rat liver and lactating mammary gland microsomes, *Arch. Biochem. Biophys.* 288, 500–508.
37. Blomhoff, R., Green, M. H., Berg, T., and Norum, K. R. (1990) Transport and storage of vitamin A, *Science* 250, 399–404.
38. Cooper, A. D. (1997) Hepatic uptake of chylomicron remnants, *J. Lipid Res.* 38, 2173–2192.
39. Redgrave, T. G. (2004) Chylomicron metabolism, *Biochem. Soc. Trans.* 32, 79–82.
40. Goodman, D. W., Huang, H. S., and Shiratori, T. (1965) Tissue distribution and metabolism of newly absorbed vitamin A in the rat, *J. Lipid Res.* 6, 390–396.
41. Packer, L. (2005) *Carotenoids and Retinoids: Molecular Aspects and Health Issues*, AOC Press, Champaign, IL.
42. van Bennekum, A. M., Kako, Y., Weinstock, P. H., Harrison, E. H., Deckelbaum, R. J., Goldberg, I. J., and Blaner, W. S. (1999) Lipoprotein lipase expression level influences tissue clearance of chylomicron retinyl ester, *J. Lipid Res.* 40, 565–574.
43. Otarod, J. K., and Goldberg, I. J. (2004) Lipoprotein lipase and its role in regulation of plasma lipoproteins and cardiac risk, *Curr. Atheroscler. Rep.* 6, 335–342.
44. Quadro, L., Blaner, W. S., Salchow, D. J., Vogel, S., Piantedosi, R., Gouras, P., Freeman, S., Cosma, M. P., Colantuoni, V., and Gottesman, M. E. (1999) Impaired retinal function and vitamin A availability in mice lacking retinol-binding protein, *EMBO J.* 18, 4633–4644.
45. Vogel, S., Piantedosi, R., O'Byrne, S. M., Kako, Y., Quadro, L., Gottesman, M. E., Goldberg, I. J., and Blaner, W. S. (2002) Retinol-binding protein-deficient mice: biochemical basis for impaired vision, *Biochemistry* 41, 15360–15368.
46. Quadro, L., Hamberger, L., Gottesman, M. E., Wang, F., Colantuoni, V., Blaner, W. S., and Mendelsohn, C. L. (2005) Pathways of vitamin A delivery to the embryo: insights from a new tunable model of embryonic vitamin A deficiency, *Endocrinology* 146, 4479–4490.
47. Kanai, M., Raz, A., and Goodman, D. S. (1968) Retinol-binding protein: the transport protein for vitamin A in human plasma, *J. Clin. Invest.* 47, 2025–2044.
48. Blomhoff, R., Rasmussen, M., Nilsson, A., Norum, K. R., Berg, T., Blaner, W. S., Kato, M., Mertz, J. R., Goodman, D. S., Eriksson, U., et al. (1985) Hepatic retinol metabolism. Distribution of retinoids, enzymes, and binding proteins in isolated rat liver cells, *J. Biol. Chem.* 260, 13560–13565.
49. Blaner, W. S., Hendriks, H. F., Brouwer, A., de Leeuw, A. M., Knook, D. L., and Goodman, D. S. (1985) Retinoids, retinoid-binding proteins, and retinyl palmitate hydrolase distributions in different types of rat liver cells, *J. Lipid Res.* 26, 1241–1251.
50. Soprano, D. R., and Blaner, W. S. (1994) Plasma Retinol Binding Protein, in *The Retinoids: Biology, Chemistry, and Medicine*, 2nd ed., Raven Press, New York.
51. Peterson, P. A. (1971) Studies on the interaction between prealbumin, retinol-binding protein, and vitamin A, *J. Biol. Chem.* 246, 44–49.
52. Melhus, H., Nilsson, T., Peterson, P. A., and Rask, L. (1991) Retinol-binding protein and transthyretin expressed in HeLa cells form a complex in the endoplasmic reticulum in both the absence and the presence of retinol, *Exp. Cell Res.* 197, 119–124.
53. van Bennekum, A. M., Wei, S., Gamble, M. V., Vogel, S., Piantedosi, R., Gottesman, M., Episkopou, V., and Blaner, W. S. (2001) Biochemical basis for depressed serum retinol levels in transthyretin-deficient mice, *J. Biol. Chem.* 276, 1107–1113.
54. Vogel, S., Gamble, M. V., and Blaner, W. S. (1999) Retinoid Uptake, Metabolism and Transport, in *The Handbook of Experimental Pharmacology: Retinoids: The Biochemical and Molecular Basis of Vitamin A and Retinoid Action*, Vol. 139, Springer, New York.
55. Noy, N., and Xu, Z. J. (1990) Kinetic parameters of the interactions of retinol with lipid bilayers, *Biochemistry* 29, 3883–3888.
56. Noy, N. (2000) Retinoid-binding proteins: mediators of retinoid action, *Biochem. J.* 348, 481–495.
57. Noy, N., and Blaner, W. S. (1991) Interactions of retinol with binding proteins: studies with rat cellular retinol-binding protein and with rat retinol-binding protein, *Biochemistry* 30, 6380–6386.
58. Heller, J. (1975) Interactions of plasma retinol-binding protein with its receptor. Specific binding of bovine and human retinol-binding protein to pigment epithelium cells from bovine eyes, *J. Biol. Chem.* 250, 3613–3619.
59. Rask, L., and Peterson, P. A. (1976) In vitro uptake of vitamin A from the retinol-binding plasma protein to mucosal epithelial cells from the monkey's small intestine, *J. Biol. Chem.* 251, 6360–6366.
60. International Agency for Research on Cancer. (2000) *IARC Handbooks of Cancer Prevention, Vol. 4. Retinoids*. World Health Organization Press, Geneva.
61. Sundaram, M., Sivaprasadarao, A., DeSousa, M. M., and Findlay, J. B. (1998) The transfer of retinol from serum retinol-binding protein to cellular retinol-binding protein is mediated by a membrane receptor, *J. Biol. Chem.* 273, 3336–3342.
62. Kawaguchi, R., Yu, J., Honda, J., Hu, J., Whitelegge, J., Ping, P., Wiita, P., Bok, D., and Sun, H. (2007) A membrane receptor for retinol binding protein mediates cellular uptake of vitamin A, *Science* 315, 820–825.

63. Bouillet, P., Oulad-Abdelghani, M., Vicaire, S., Garnier, J. M., Schuhbaur, B., Dolle, P., and Chambon, P. (1995) Efficient cloning of cDNAs of retinoic acid-responsive genes in P19 embryonal carcinoma cells and characterization of a novel mouse gene, *Stra1* (mouse LERK-2/Eplg2), *Dev. Biol.* 170, 420–433.
64. Bouillet, P., Sapin, V., Chazaud, C., Messaddeq, N., Decimo, D., Dolle, P., and Chambon, P. (1997) Developmental expression pattern of *Stra6*, a retinoic acid-responsive gene encoding a new type of membrane protein, *Mech. Dev.* 63, 173–186.
65. Pasutto, F., Sticht, H., Hammersen, G., Gillesen-Kaesbach, G., Fitzpatrick, D. R., Nurnberg, G., Brasch, F., Schirmer-Zimmermann, H., Tolmie, J. L., Chitayat, D., Houge, G., Fernandez-Martinez, L., Keating, S., Mortier, G., Hennekam, R. C., von der Wense, A., Slavotinek, A., Meinecke, P., Bitoun, P., Becker, C., Nurnberg, P., Reis, A., and Rauch, A. (2007) Mutations in *STRA6* cause a broad spectrum of malformations including anophthalmia, congenital heart defects, diaphragmatic hernia, alveolar capillary dysplasia, lung hypoplasia, and mental retardation, *Am. J. Hum. Genet.* 80, 550–560.
66. Seeliger, M. W., Biesalski, H. K., Wissinger, B., Gollnick, H., Gielen, S., Frank, J., Beck, S., and Zrenner, E. (1999) Phenotype in retinol deficiency due to a hereditary defect in retinol binding protein synthesis, *Invest. Ophthalmol. Visual Sci.* 40, 3–11.
67. Blaner, W. S. (2007) *STRA6*, a cell-surface receptor for retinol-binding protein: the plot thickens, *Cell Metab.* 5, 164–166.
68. Noy, N., Slosberg, E., and Scarlata, S. (1992) Interactions of retinol with binding proteins: studies with retinol-binding protein and with transthyretin, *Biochemistry* 31, 11118–11124.
69. Retinoic Acid. (2000) *IARC Handbooks of Cancer Prevention*, Vol. 4.
70. Blaner, W. S., and Olson, J. A. (1994) Retinol and Retinoic acid Metabolism, in *The Retinoids: Biology, Chemistry, and Medicine*, 2nd ed., Raven Press, New York.
71. Kurlandsky, S. B., Gamble, M. V., Ramakrishnan, R., and Blaner, W. S. (1995) Plasma delivery of retinoic acid to tissues in the rat, *J. Biol. Chem.* 270, 17850–17857.
72. Krasinski, S. D., Cohn, J. S., Russell, R. M., and Schaefer, E. J. (1990) Postprandial plasma vitamin A metabolism in humans: a reassessment of the use of plasma retinyl esters as markers for intestinally derived chylomicrons and their remnants, *Metabolism* 39, 357–365.
73. Goswami, B. C., Reida, A. K., Ivanoff, K. D., Barua, A. B., and Olson, J. A. (2003) Intestinal absorption and metabolism of retinoyl beta-glucuronide in humans, and of 15-[14C]-retinoyl beta-glucuronide in rats of different vitamin A status, *J. Nutr. Biochem.* 14, 703–709.
74. Ueltschy, A., Gunning, D. B., Barua, A. B., and Olson, J. A. (2002) Effects of subcutaneously injected graded doses of all-trans retinoic acid and all-trans retinoyl beta-glucuronide on the outcome of pregnancy in Sprague-Dawley rats, *Int. J. Vitam. Nutr. Res.* 72, 229–235.
75. Liu, L., and Gudas, L. J. (2005) Disruption of the lecithin:retinol acyltransferase gene makes mice more susceptible to vitamin A deficiency, *J. Biol. Chem.* 280, 40226–40234.
76. Tsutsumi, C., Okuno, M., Tannous, L., Piantadosi, R., Allan, M., Goodman, D. S., and Blaner, W. S. (1992) Retinoids and retinoid-binding protein expression in rat adipocytes, *J. Biol. Chem.* 267, 1805–1810.
77. Zovich, D. C., Orologa, A., Okuno, M., Kong, L. W., Talmage, D. A., Piantadosi, R., Goodman, D. S., and Blaner, W. S. (1992) Differentiation-dependent expression of retinoid-binding proteins in BFC-1 beta adipocytes, *J. Biol. Chem.* 267, 13884–13889.
78. Blaner, W. S., Obunike, J. C., Kurlandsky, S. B., al-Haideri, M., Piantadosi, R., Deckelbaum, R. J., and Goldberg, I. J. (1994) Lipoprotein lipase hydrolysis of retinyl ester. Possible implications for retinoid uptake by cells, *J. Biol. Chem.* 269, 16559–16565.
79. Okuno, M., Caraveo, V. E., Goodman, D. S., and Blaner, W. S. (1995) Regulation of adipocyte gene expression by retinoic acid and hormones: effects on the gene encoding cellular retinol-binding protein, *J. Lipid Res.* 36, 137–147.
80. Wei, S., Lai, K., Patel, S., Piantadosi, R., Shen, H., Colantuoni, V., Kraemer, F. B., and Blaner, W. S. (1997) Retinyl ester hydrolysis and retinol efflux from BFC-1beta adipocytes, *J. Biol. Chem.* 272, 14159–14165.
81. Yen, C. L., Monetti, M., Burri, B. J., and Farese, R. V., Jr. (2005) The triacylglycerol synthesis enzyme DGAT1 also catalyzes the synthesis of diacylglycerols, waxes, and retinyl esters, *J. Lipid Res.* 46, 1502–1511.
82. Orland, M. D., Anwar, K., Cromley, D., Chu, C. H., Chen, L., Billheimer, J. T., Hussain, M. M., and Cheng, D. (2005) Acyl coenzyme A dependent retinol esterification by acyl coenzyme A: diacylglycerol acyltransferase 1, *Biochim. Biophys. Acta* 1737, 76–82.
83. Isken, A., Holzschuh, J., Lampert, J. M., Fischer, L., Oberhauser, V., Palczewski, K., and von Lintig, J. (2006) Sequestration of retinyl esters is essential for retinoid signaling in the zebrafish embryo, *J. Biol. Chem.* 282, 1144–1151.
84. Molotkov, A., Ghyselinck, N. B., Chambon, P., and Duester, G. (2004) Opposing actions of cellular retinol-binding protein and alcohol dehydrogenase control the balance between retinol storage and retinoic acid synthesis, *Biochem. J.* 383, 295–302.
85. Guillemette, C. (2003) Pharmacogenomics of human UDP-glucuronosyltransferase enzymes, *Pharmacogenomics J.* 3, 136–158.
86. Brierley, C. H., Senafi, S. B., Clarke, D., Hsu, M. H., Johnson, E. F., and Burchell, B. (1996) Regulation of the human bilirubin UDP-glucuronosyltransferase gene, *Adv. Enzyme Regul.* 36, 85–97.
87. Zile, M. H., Schnoes, H. K., and DeLuca, H. F. (1980) Characterization of retinoyl beta-glucuronide as a minor metabolite of retinoic acid in bile, *Proc. Natl. Acad. Sci. U.S.A.* 77, 3230–3233.
88. Frolik, C. A., Swanson, B. N., Dart, L. L., and Sporn, M. B. (1981) Metabolism of 13-cis-retinoic acid: identification of 13-cis-retinoyl- and 13-cis-4-oxoretinoyl-beta-glucuronides in the bile of vitamin A-normal rats, *Arch. Biochem. Biophys.* 208, 344–352.
89. Napoli, J. L., Khalil, H., and McCormick, A. M. (1982) Metabolism of 5,6-epoxyretinoic acid in vivo: isolation of a major intestinal metabolite, *Biochemistry* 21, 1942–1949.
90. Golczak, M., Imanishi, Y., Kuksa, V., Maeda, T., Kubota, R., and Palczewski, K. (2005) Lecithin:retinol acyltransferase is responsible for amidation of retinylamine, a potent inhibitor of the retinoid cycle, *J. Biol. Chem.* 280, 42263–42273.
91. Ghyselinck, N. B., Bavik, C., Sapin, V., Mark, M., Bonnier, D., Hindelang, C., Dierich, A., Nilsson, C. B., Hakansson, H., Sauvant, P., Azais-Braesco, V., Frasson, M., Picaud, S., and Chambon, P. (1999) Cellular retinol-binding protein I is essential for vitamin A homeostasis, *EMBO J.* 18, 4903–4914.
92. E, X., Zhang, L., Lu, J., Tso, P., Blaner, W. S., Levin, M. S., and Li, E. (2002) Increased neonatal mortality in mice lacking cellular retinol-binding protein II, *J. Biol. Chem.* 277, 36617–36623.
93. Vogel, S., Mendelsohn, C. L., Mertz, J. R., Piantadosi, R., Waldburger, C., Gottesman, M. E., and Blaner, W. S. (2001) Characterization of a new member of the fatty acid-binding protein family that binds all-trans-retinol, *J. Biol. Chem.* 276, 1353–1360.
94. Piantadosi, R., Ghyselinck, N., Blaner, W. S., and Vogel, S. (2005) Cellular retinol-binding protein type III is needed for retinoid incorporation into milk, *J. Biol. Chem.* 280, 24286–24292.
95. Randolph, R. K., and Ross, A. C. (1991) Vitamin A status regulates hepatic lecithin:retinol acyltransferase activity in rats, *J. Biol. Chem.* 266, 16453–16457.
96. Zolfaghari, R., and Ross, A. C. (2002) Lecithin:retinol acyltransferase expression is regulated by dietary vitamin A and exogenous retinoic acid in the lung of adult rats, *J. Nutr.* 132, 1160–1164.
97. Ross, A. C., Ambalavanan, N., Zolfaghari, R., and Li, N. Q. (2006) Vitamin A combined with retinoic acid increases retinol uptake and lung retinyl ester formation in a synergistic manner in neonatal rats, *J. Lipid Res.* 47, 1844–1851.
98. Ross, A. C., Li, N. Q., and Wu, L. (2006) The components of VARA, a nutrient-metabolite combination of vitamin A and retinoic acid, act efficiently together and separately to increase retinyl esters in the lungs of neonatal rats, *J. Nutr.* 136, 2803–2807.
99. Vogel, S., Piantadosi, R., Frank, J., Lalazar, A., Rockey, D. C., Friedman, S. L., and Blaner, W. S. (2000) An immortalized rat liver stellate cell line (HSC-T6): a new cell model for the study of retinoid metabolism in vitro, *J. Lipid Res.* 41, 882–893.
100. Mangelsdorf, D. J., Umeson, K., Kliewer, S. A., Borgmeyer, U., Ong, E. S., and Evans, R. M. (1991) A direct repeat in the cellular retinol-binding protein type II gene confers differential regulation by RXR and RAR, *Cell* 66, 555–561.
101. Tang, X. H., Suh, M. J., Li, R., and Gudas, L. J. (2007) Cell proliferation inhibition and alterations in retinol esterification induced by phytanic acid and docosahexaenoic acid, *J. Lipid Res.* 48, 165–176.

102. Guo, X., Ruiz, A., Rando, R. R., Bok, D., and Gudas, L. J. (2000) Esterification of all-trans-retinol in normal human epithelial cell strains and carcinoma lines from oral cavity, skin and breast: reduced expression of lecithin:retinol acyltransferase in carcinoma lines, *Carcinogenesis* 21, 1925–1933.
103. Guo, X., Nanus, D. M., Ruiz, A., Rando, R. R., Bok, D., and Gudas, L. J. (2001) Reduced levels of retinyl esters and vitamin A in human renal cancers, *Cancer Res.* 61, 2774–2781.
104. Guo, X., Knudsen, B. S., Peehl, D. M., Ruiz, A., Bok, D., Rando, R. R., Rhim, J. S., Nanus, D. M., and Gudas, L. J. (2002) Retinol metabolism and lecithin:retinol acyltransferase levels are reduced in cultured human prostate cancer cells and tissue specimens, *Cancer Res.* 62, 1654–1661.
105. Simmons, D. P., Andreola, F., and De Luca, L. M. (2002) Human melanomas of fibroblast and epithelial morphology differ widely in their ability to synthesize retinyl esters, *Carcinogenesis* 23, 1821–1830.
106. Zhan, H. C., Gudas, L. J., Bok, D., Rando, R., Nanus, D. M., and Tickoo, S. K. (2003) Differential expression of the enzyme that esterifies retinol, lecithin:retinol acyltransferase, in subtypes of human renal cancer and normal kidney, *Clin. Cancer Res.* 9, 4897–4905.
107. Boorjian, S., Tickoo, S. K., Mongan, N. P., Yu, H., Bok, D., Rando, R. R., Nanus, D. M., Scherr, D. S., and Gudas, L. J. (2004) Reduced lecithin: retinol acyltransferase expression correlates with increased pathologic tumor stage in bladder cancer, *Clin. Cancer Res.* 10, 3429–3437.
108. Sheren-Manoff, M., Shin, S. J., Su, D., Bok, D., Rando, R. R., and Gudas, L. J. (2006) Reduced lecithin:retinol acyltransferase expression in human breast cancer, *Int. J. Oncol.* 29, 1193–1199.
109. Lotan, R., Ong, D. E., and Chytil, F. (1980) Comparison of the level of cellular retinoid-binding proteins and susceptibility to retinoid-induced growth inhibition of various neoplastic cell lines, *J. Natl. Cancer Inst.* 64, 1259–1262.
110. Fex, G., Ekelund, G., Leandoer, L., and Sternby, N. H. (1986) Cellular retinol-binding protein (CRBP) in human colorectal adenocarcinoma, *Br. J. Cancer* 53, 687–690.
111. Fex, G., Wahlberg, P., Biorklund, A., Wennerberg, J., and Willen, R. (1986) Studies of cellular retinol-binding protein (CRBP) in squamous-cell carcinomas of the head and neck region, *Int. J. Cancer* 37, 217–221.
112. Fex, G., Linell, F., and Ljungberg, O. (1985) Cellular retinol-binding protein in normal and neoplastic human mammary gland, *Breast Cancer Res. Treat.* 6, 131–136.
113. Kuppumbatti, Y. S., Bleiweiss, I. J., Mandeli, J. P., Waxman, S., and Mira, Y. L. R. (2000) Cellular retinol-binding protein expression and breast cancer, *J. Natl. Cancer Inst.* 92, 475–480.
114. Kuppumbatti, Y. S., Rexer, B., Nakajo, S., Nakaya, K., and Mira-y-Lopez, R. (2001) CRBP suppresses breast cancer cell survival and anchorage-independent growth, *Oncogene* 20, 7413–7419.
115. Arapshian, A., Bertran, S., Kuppumbatti, Y. S., Nakajo, S., and Mira-y-Lopez, R. (2004) Epigenetic CRBP downregulation appears to be an evolutionarily conserved (human and mouse) and oncogene-specific phenomenon in breast cancer, *Mol. Cancer* 3, 13.
116. Lotan, R. (2005) A crucial role for cellular retinol-binding protein I in retinoid signaling, *J. Natl. Cancer Inst.* 97, 3–4.
117. Farias, E. F., Ong, D. E., Ghyselinck, N. B., Nakajo, S., Kuppumbatti, Y. S., and Mira y Lopez, R. (2005) Cellular retinol-binding protein I, a regulator of breast epithelial retinoic acid receptor activity, cell differentiation, and tumorigenicity, *J. Natl. Cancer Inst.* 97, 21–29.
118. Lammer, E. J., Chen, D. T., Hoar, R. M., Agnish, N. D., Benke, P. J., Braun, J. T., Curry, C. J., Fernhoff, P. M., Grix, A. W., Jr., Lott, I. T., et al. (1985) Retinoic acid embryopathy, *N. Engl. J. Med.* 313, 837–841.
119. Soprano, D. R., and Soprano, K. J. (1995) Retinoids as teratogens, *Annu. Rev. Nutr.* 15, 111–132.
120. Collins, M. D., and Mao, G. E. (1999) Teratology of retinoids, *Annu. Rev. Pharmacol. Toxicol.* 39, 399–430.
121. Rothman, K. J., Moore, L. L., Singer, M. R., Nguyen, U. S., Mannino, S., and Milunsky, A. (1995) Teratogenicity of high vitamin A intake, *N. Engl. J. Med.* 333, 1369–1373.
122. Cohlan, S. Q. (1953) Excessive intake of vitamin A as a cause of congenital anomalies in the rat, *Science* 117, 535–536.
123. Cohlan, S. Q. (1954) Congenital anomalies in the rat produced by excessive intake of vitamin A during pregnancy, *Pediatrics* 13, 556–567.
124. Geelen, J. A., Langman, J., and Lowdon, J. D. (1980) The influence of excess vitamin A on neural tube closure in the mouse embryo, *Anat. Embryol.* 159, 223–234.
125. Collins, M. D., Tzimas, G., Hummler, H., Burgin, H., and Nau, H. (1994) Comparative teratology and transplacental pharmacokinetics of all-trans-retinoic acid, 13-cis-retinoic acid, and retinyl palmitate following daily administrations in rats, *Toxicol. Appl. Pharmacol.* 127, 132–144.
126. Abroms, L., Maibach, E., Lyon-Daniel, K., and Feldman, S. R. (2006) What is the best approach to reducing birth defects associated with isotretinoin? *PLoS Med.* 3, e483.
127. Standeven, A. M., Davies, P. J., Chandraratna, R. A., Mader, D. R., Johnson, A. T., and Thomazy, V. A. (1996) Retinoid-induced epiphyseal plate closure in guinea pigs, *Fundam. Appl. Toxicol.* 34, 91–98.
128. Prendiville, J., Bingham, E. A., and Burrows, D. (1986) Premature epiphyseal closure—a complication of etretinate therapy in children, *J. Am. Acad. Dermatol.* 15, 1259–1262.
129. Berson, E. L., Rosner, B., Sandberg, M. A., Hayes, K. C., Nicholson, B. W., Weigel-DiFranco, C., and Willett, W. (1993) A randomized trial of vitamin A and vitamin E supplementation for retinitis pigmentosa, *Arch. Ophthalmol.* 111, 761–772.
130. Sibulesky, L., Hayes, K. C., Pronczuk, A., Weigel-DiFranco, C., Rosner, B., and Berson, E. L. (1999) Safety of <7500 RE (<25000 IU) vitamin A daily in adults with retinitis pigmentosa, *Am. J. Clin. Nutr.* 69, 656–663.
131. Nagai, K., Hosaka, H., Kubo, S., Nakabayashi, T., Amagasaki, Y., and Nakamura, N. (1999) Vitamin A toxicity secondary to excessive intake of yellow-green vegetables, liver and laver, *J. Hepatol.* 31, 142–148.
132. Guarascio, P., Portmann, B., Visco, G., and Williams, R. (1983) Liver damage with reversible portal hypertension from vitamin A intoxication: demonstration of Ito cells, *J. Clin. Pathol.* 36, 769–771.
133. Nollevaux, M. C., Guiot, Y., Horsmans, Y., Leclercq, I., Rahier, J., Geubel, A. P., and Sempoux, C. (2006) Hypervitaminosis A-induced liver fibrosis: stellate cell activation and daily dose consumption, *Liver Int.* 26, 182–186.
134. Hautekeete, M. L., and Geerts, A. (1997) The hepatic stellate (Ito) cell: its role in human liver disease, *Virchows Arch.* 430, 195–207.
135. Scheven, B. A., and Hamilton, N. J. (1990) Retinoic acid and 1,25-dihydroxyvitamin D3 stimulate osteoclast formation by different mechanisms, *Bone* 11, 53–59.
136. Baxi, S. C., and Dailey, G. E., 3rd. (1982) Hypervitaminosis A. A cause of hypercalcemia, *West. J. Med.* 137, 429–431.
137. Genaro Pde, S., and Martini, L. A. (2004) Vitamin A supplementation and risk of skeletal fracture, *Nutr. Rev.* 62, 65–7.
138. Melhus, H., Michaelsson, K., Kindmark, A., Bergstrom, R., Holmberg, L., Mallmin, H., Wolk, A., and Ljunghall, S. (1998) Excessive dietary intake of vitamin A is associated with reduced bone mineral density and increased risk for hip fracture, *Ann. Intern. Med.* 129, 770–778.
139. Maeda, A., Maeda, T., Golczak, M., Imanishi, Y., Leahy, P., Kubota, R., and Palczewski, K. (2006) Effects of potent inhibitors of the retinoid cycle on visual function and photoreceptor protection from light damage in mice, *Mol. Pharmacol.* 70, 1220–1229.

BI7003069