

A Unique Insert of Leucyl-tRNA Synthetase Is Required for Aminoacylation and Not Amino Acid Editing[†]

Michael T. Vu and Susan A. Martinis*

Department of Biochemistry, Roger Adams Laboratory, University of Illinois at Urbana–Champaign, Box B4, 600 South Mathews Avenue, Urbana, Illinois 61801

Received October 5, 2006; Revised Manuscript Received February 2, 2007

ABSTRACT: Leucyl-tRNA synthetase (LeuRS) is a class I enzyme, which houses its aminoacylation active site in a canonical core that is defined by a Rossmann nucleotide binding fold. In addition, many LeuRSs bear a unique polypeptide insert comprised of about 50 amino acids located just upstream of the conserved KMSKS sequence. The role of this leucine-specific domain (LS-domain) remains undefined. We hypothesized that this domain may be important for substrate recognition in aminoacylation and/or amino acid editing. We carried out a series of deletion mutations and chimeric swaps within the leucine-specific domain of *Escherichia coli*. Our results support that the leucine-specific domain is critical for aminoacylation but not required for editing activity. Kinetic analysis determined that deletion of the LS-domain primarily impacts k_{cat} . Because of its proximity to the aminoacylation active site, we propose that this domain interacts with the tRNA during amino acid activation and/or tRNA aminoacylation. Although the leucine-specific domain does not appear to be important to the editing complex, it remains possible that it aids the dynamic translocation process that moves tRNA from the aminoacylation to the editing complex.

Aminoacylation is catalyzed by an ancient family of enzymes called the aminoacyl-tRNA synthetases (aaRS)¹ (1, 2). It is important to translation that the aaRS correctly link amino acids to the corresponding tRNA acceptors. Each of the aaRS is responsible for covalently attaching 1 of 20 standard amino acids to a set of cognate tRNA isoacceptors. The esterification of amino acid to tRNA occurs via a two-step reaction:



The first reversible step involves the activation of an amino acid concomitant with ATP hydrolysis to form an aminoacyl-adenylate intermediate. The amino acid is then transferred from the adenylate intermediate to the 3'-terminal adenosine moiety of the tRNA. The charged aminoacyl-tRNA binds to EF-Tu and is transported to the ribosome to extend the polypeptide chain.

Leucyl-tRNA synthetase (LeuRS) is a class I aaRS and is responsible for accurately aminoacylating leucine to its cognate tRNA^{Leu} isoacceptors (3). As is characteristic of all class I synthetases, the catalytic core is defined by a Rossmann nucleotide binding fold (4) that comprises the main body of the enzyme (5). This canonical class I core of

LeuRS contains inserts and appendages (5–7). For example, the enzyme has an amino acid editing function that resides within a large insert called the connective polypeptide 1 (CP1) (8). In addition, a novel C-terminal domain has been proposed to be important for tRNA interactions (9–12).

One unique small insert called the leucine-specific domain (LS-domain) is found in many bacterial LeuRSs (5). Its function remains undefined. This compact domain that is comprised of five β -strands and two short α -helices flanks the entrance of the aminoacylation active site (5). The LS-domain is inserted after the last β -strand of the Rossmann nucleotide binding fold and is linked to the conserved KMSKS sequence (Figure 1). Primary sequence alignments show that the LS-domain is not particularly conserved in sequence or length (Figure 1A). Indeed, it is completely missing in LeuRSs of many organisms including *Bacillus subtilis*, *Helicobacter pylori*, and *Pyrococcus horikoshii* (data not shown).

A large ensemble of LeuRS cocrystal structures that include tRNA complexes in the aminoacylation and editing conformation have failed to suggest a specific role for the LS-domain (5, 9, 13–16). The cocrystal structure of *P. horikoshii* LeuRS with the tRNA in the aminoacylation conformation has been solved (13). But *P. horikoshii* LeuRS does not possess a LS-domain. However, comparison between the apo *Thermus thermophilus* LeuRS and its complex with the tRNA in the editing conformation shows that the LS-domain is dynamic and moves approximately 19° (9). In the editing conformation of *T. thermophilus* LeuRS, the LS-domain does not make any contact with the tRNA. We hypothesized that this movement of the LS-domain may influence tRNA interactions or even interact

[†] This work was supported by the National Institutes of Health (Grant GM63789).

* To whom correspondence should be addressed. Phone: 217-244-2405. Fax: 217-244-5858. E-mail: martinis@life.uiuc.edu.

¹ Abbreviations: aaRS, aminoacyl-tRNA synthetase; LeuRS, leucyl-tRNA synthetase; LS-domain, leucine-specific domain; CP1, connective polypeptide 1.




FIGURE 1: Primary and tertiary structure of LeuRS enzymes. (A) Primary sequence alignment of LeuRSs from various organisms. Conserved and homologous residues are highlighted in black and gray, respectively. The bar above the sequence indicates the LS-domain sequence of *E. coli* LeuRS. Abbreviations are as follows: *Ec*, *E. coli*; *Pm*, *Pasteurella multocci*; *Hi*, *Haemophilus influenzae*; *Nm*, *Neisseria meningitidis*; *Pa*, *Pseudomonas aeruginosa*; *Xf*, *Xylella fastidiosa*; *Vc*, *Vibrio cholerae*; *Tt*, *Thermus thermophilus*. (B) Homology model of *E. coli* LeuRS (32). The protein is shown as a ribbon diagram. The LS-domain is highlighted in black, while the canonical class I core and CP1 editing domain are colored in gray. The bold arrow points to the Rossmann nucleotide binding fold that comprises the aminoacylation active site. (C) Sequence alignment of LeuRS with regions of ValRS and IleRS that correlate to the insertion site of the LS-domain. The shaded box highlights the two ValRS peptides that were used to make chimeric replacements of the LS-domain in *E. coli* LeuRS.

with the tRNA during its entry and/or exit. It is also possible that the LS-domain plays a role in tRNA translocation from the aminoacylation active site over 30 Å to the editing active site that is located in the CP1 domain.

To identify a function for the LS-domain, we carried out deletion and chimeric peptide replacement analysis. We tested each mutant LeuRS for alterations in aminoacylation and post-transfer editing activity. Our results support that the LS-domain is important to the overall aminoacylation reaction but does not significantly impact post-transfer editing of mischarged tRNA^{Leu}.

EXPERIMENTAL PROCEDURES

Materials and Resources. Oligonucleotide primers were synthesized by MWG Biotech (High Point, NC) or Integrated DNA Technologies (Coralville, IA). Tritium-labeled amino acids and [³²P]pyrophosphate were purchased from Perkin-Elmer (Boston, MA). Cloned *Pfu* DNA polymerase, dNTPs, and competent cells were obtained from Stratagene (La Jolla, CA). Structure predictions of mutant and chimeric proteins were generated using the DeepView/Swiss Pdb Viewer version 3.7 (17).

Mutagenesis and Purification of *Escherichia coli* LeuRS. Each 50 µL polymerase chain reaction (PCR) contained 63 ng of p15ec3-1 template, which encodes the *E. coli* *leuS* gene (18), 0.4 µM each of the forward and reverse primer that contained the mutation, 0.4 mM each of the dNTP, and 2.5 units of *Pfu* DNA polymerase in commercially prepared buffer. The mutant plasmid was amplified, screened, and confirmed by DNA sequencing as described previously (19). Plasmids expressing mutant and wild-type LeuRS were used to transform *E. coli* BL21(DE3). Protein expression was

induced with 1 mM isopropyl β-D-thiogalactopyranoside (IPTG) for 4 h, and LeuRS was purified by affinity chromatography via a fused N-terminal six-histidine tag (19). The final concentration was determined using a Bradford protein assay as described in the commercial protocol (Bio-Rad, Hercules, CA).

Isolation of in Vitro Transcribed tRNA^{Leu} and [³H]-Ile-tRNA^{Leu}. T7 RNA polymerase was purified (20), and tRNA^{Leu}_{UAA} (tRNA^{Leu}) was transcribed via in vitro runoff transcription (21). The plasmid containing the gene for *E. coli* tRNA^{Leu} (ptDNA^{Leu}) (22, 23) was digested with 25 units of *Bst*NI (Promega) in a 1 mL reaction at 60 °C overnight and then used as template for in vitro transcription as described previously (24). The ethanol-precipitated RNA was separated on a 10% acrylamide (19:1), 8 M urea gel. The tRNA^{Leu} band was detected by UV shadowing, excised, and recovered (19). The concentration was determined on the basis of the absorbance at 260 nm at 80 °C using the extinction coefficient 840700 L M⁻¹ cm⁻¹. Purified tRNA^{Leu} was refolded by denaturing at 80 °C for 1 min, followed by addition of 1 mM MgCl₂, and quickly cooled on ice. The tRNA was stored at -20 °C.

Mischarged [³H]-Ile-tRNA^{Leu} was generated in a reaction containing 60 mM Tris (pH 7.5), 10 mM MgCl₂, 1 mM DTT, 4 mM ATP, 22 µM [³H]isoleucine (150 µCi/mL), 4 µM tRNA^{Leu}, and 26 µM each of an editing-defective LeuRS (19). The reaction was incubated at room temperature for 3 h and then quenched with 4 µL of 10% acetic acid (25), followed by extraction using phenol/chloroform/isoamyl alcohol (25:24:1, pH 4.3). One-half volume of 4.6 M ammonium acetate, pH 5.0, and 3 µL of 25 mg/mL glycogen was added, followed by ethanol precipitation at -80 °C. The

tRNA was resuspended in 61 μL of 50 mM KH_2PO_4 (pH 5.0) and stored at -20°C .

Enzyme Activity Assays. Each aminoacylation reaction contained 60 mM Tris, pH 7.5, 10 mM MgCl_2 , 1 mM DTT, 22 μM [^3H]leucine (150 $\mu\text{Ci}/\text{mL}$), 4 μM tRNA^{Leu} , and 100 nM enzyme and was initiated with 4 mM ATP. Kinetic rate constants for aminoacylation were measured using six different concentrations of in vitro transcribed tRNA^{Leu} ranging from 0.2 to 20 μM . The apparent rate constants were determined on the basis of the average of three values. The post-transfer editing reaction consisted of 60 mM Tris, pH 7.5, 10 mM MgCl_2 , and approximately 6.5 μM [^3H]Ile- tRNA^{Leu} (150 $\mu\text{Ci}/\text{mL}$) and was initiated with 100 nM enzyme. Aliquots for either aminoacylation or post-transfer editing were quenched at specific time points by transferring 10 μL onto a Whatman filter pad that was presoaked with 5% trichloroacetic acid (TCA). The pads were washed, and radioactivity was quantitated as described previously (19).

Inorganic pyrophosphate (PP_i) exchange assays were carried out in 50 mM *N*-(2-hydroxyethyl)piperazine-*N'*-2-ethanesulfonic acid (HEPES), pH 8.0, 10 mM MgCl_2 , 1 mM DTT, 1 mM L-leucine, 1 mM [^{32}P] PP_i (78 mCi/mL), and 500 nM enzyme (26). The reaction was initiated with 1 mM ATP, and aliquots were quenched by spotting 2 μL onto a poly(ethylene imine) thin-layer chromatography (TLC) plate (Scientific Adsorbents Inc., Atlanta, GA) that had been prerun in water. ATP and PP_i were separated using 750 mM KH_2PO_4 (pH 3.5) and 4 M urea. Radiolabeled products were analyzed via phosphorimaging using a FUJIFILM BAS-MS 2040 imaging plate (FUJIFILM Medical System USA, Stanford, CT).

RNA Binding Assays. Nitrocellulose filter binding assays to measure tRNA binding to wild-type and mutant LeuRSs were carried out using 50 μL reactions that contained 60 mM Tris (pH 7.5), 10 mM MgCl_2 , 1 mM DTT, 20 μM L-leucine, 0.1 μM 5'- ^{32}P -labeled tRNA, and varying concentrations of enzyme ranging from 0.1 to 2 μM . The reaction was incubated at 37°C for 5 min and then quenched by spotting all 50 μL on a nitrocellulose membrane assembled on a MINIFOLD-1 Spot-Blot system (Schleicher & Schuell, Keene, NH). The membrane was washed with 200 μL of binding buffer [60 mM Tris (pH 7.5), 10 mM MgCl_2 , and 1 mM DTT] to remove unbound tRNA. Radiolabeled products were analyzed via phosphorimaging using a FUJIFILM BAS-MS 2040 imaging plate (FUJIFILM Medical System USA, Stanford, CT).

RESULTS

Deletion of the LS-Domain Abolishes Aminoacylation Activity but Not Post-Transfer Editing Activity. The LS-domain is found in many bacterial LeuRSs but not in the homologous IleRS and ValRS. We carried out a primary sequence alignment between LeuRS, IleRS, and ValRS to determine the end points of the LS-domain insert (Figure 1C). On the basis of structural and primary sequence alignments, we deleted the entire 49 amino acid insert, which spans Val 569 to Gly 618 in *E. coli* LeuRS. The LS-domain was replaced with either three or five alanines, which link the last β -strand of the Rossmann fold to the KMSKS conserved sequence. On the basis of the X-ray crystal structures of LeuRS, IleRS, and ValRS, we hypothesized that




FIGURE 2: Enzymatic activity of wild type and LeuRS LS-domain deletion mutants. (A) Aminoacylation reactions were carried out using 4 μM in vitro transcribed tRNA^{Leu} and 100 nM enzyme. (B) Post-transfer editing reactions included approximately 6.5 μM [^3H]Ile- tRNA^{Leu} and 100 nM enzyme. Symbols representing wild-type and mutant enzymes are as follows: no enzyme (\square), wild type (\circ), $\Delta\text{LSD-5A}$ (\triangle), and $\Delta\text{LSD-3A}$ (∇). Error bars represent the assay reproduced at least in triplicate and are presented for each point.

these alanine linker peptides would maintain the approximate distance between the two flanking β -strands that connect the LS-domain to the main body.

The LeuRS wild type and deletion mutants were purified by affinity chromatography. Aminoacylation activity of both mutant LeuRSs that lacked the LS-domain was abolished (Figure 2A). Interestingly, however, post-transfer editing activity of mischarged Ile- tRNA^{Leu} was largely unaffected and similar to the LeuRS wild-type enzyme (Figure 2B). Thus, the LS-domain plays a significant role in aminoacylation but is not required for amino acid editing.

Chimeric LeuRS and ValRS Mutants Restore Limited Aminoacylation Activity. We hypothesized that the short three or five alanine replacements of the LS-domain may have constrained movement of the conserved mobile KMSKS motif that is critical for aminoacylation activity (27–29). Sequence alignments with the homologous class I IleRS and ValRS enzymes identified a short peptide sequence that correlated with the location of the LS-domain insert (Figure 1C). The orientation of the short peptide in the X-ray crystal structure of *T. thermophilus* ValRS (30, 31) overlapped well with the *T. thermophilus* (5) and *E. coli* (32) LeuRS catalytic core. Thus, we introduced a seven-residue sequence (VLDEKGG) from *T. thermophilus* ValRS to replace the LeuRS LS-domain ($\Delta\text{LSD-valRStt}$). Likewise, we also swapped IRDDEGQ from *E. coli* ValRS into LeuRS ($\Delta\text{LSD-valRSec}$).

The chimeric LeuRS that contains the *T. thermophilus* ValRS peptide stimulated some aminoacylation, albeit significantly decreased compared to the LeuRS wild type (Figure 3). Kinetic analysis showed about a 30-fold decrease in the apparent $k_{\text{cat}}/K_{\text{m}}$ compared to the LeuRS wild-type enzyme (Table 1). In contrast, the *E. coli* ValRS peptide




FIGURE 3: Enzymatic activity of chimeric LeuRS mutants. (A) Aminoacylation reactions were carried out using 4 μ M in vitro transcribed tRNA^{Leu} and 100 nM enzyme. (B) Post-transfer editing reactions included about 6.5 μ M [³H]-Ile-tRNA^{Leu} and 100 nM enzyme. Symbols representing wild-type and mutant enzymes are as follows: no enzyme (\square), wild type (\circ), Δ LSD-valRStt (\blacktriangledown), and Δ LSD-valRSec (\blacktriangle). Error bars represent the assay reproduced at least in triplicate and are presented for each point.

Table 1: Apparent Kinetic Parameters for Aminoacylation^a

enzyme	K_m (μ M)	k_{cat} (s^{-1})	k_{cat}/K_m ($\mu M^{-1} s^{-1}$)	relative
wild-type LeuRS	0.73 ± 0.1	9.6 ± 2.7	13.0	1
Δ LSD-valRStt	2.0 ± 0.4	0.9 ± 0.1	0.45	0.03
R vec L	1.6 ± 0.2	0.43 ± 0.08	0.28	0.02
E vec K	1.3 ± 0.1	0.33 ± 0.05	0.25	0.02
RE vec LK	5.4 ± 0.6	2.0 ± 0.1	0.37	0.03
RED vec LKE	2.4 ± 0.4	1.4 ± 0.3	0.58	0.04
REI vec LKV	2.2 ± 0.7	1.4 ± 0.2	0.64	0.05

^a Apparent rate constants were determined using in vitro transcribed tRNA^{Leu} ranging in concentration from 0.2 to 20 μ M.

replacement failed to restore aminoacylation activity. Both chimeric mutants retained substantial post-transfer editing activity relative to that of the wild-type enzyme (Figure 3B). The K_D was measured for the Δ LSD-valRStt deletion mutant ($0.4 \pm 0.1 \mu$ M) using nitrocellulose filter binding assays and was similar to the wild-type protein (0.3μ M). Interestingly, the K_D was not measurable for the inactive Δ LSD-valRSec deletion mutant, suggesting that enzyme-tRNA interactions are very sensitive to the sequence of this peptide linker. This supports that the difference in leucylation activities may lie within the molecular details of the LS-domain rather than its overall global domain structure.

The two ValRS peptides from *T. thermophilus* and *E. coli* are highly similar. To identify specific amino acid determinants that might be important for aminoacylation activity, we systematically swapped different amino acids within this seven amino acid insert of the chimeric LeuRS mutants that contained the nonfunctional *E. coli* LeuRS peptide (Δ LSD-valRSec) (Figure 4A). In particular, we substituted an




FIGURE 4: Enzymatic activity of derivatives of Δ LSD-valRSec chimeric mutants. (A) Sequence of chimeric LeuRS mutants that contain a ValRS peptide insert and site-specific mutations within the inserted peptide of the Δ LSD-valRSec chimeras. Sites that were targeted by specific substitution are indicated with arrows. Residues within the chimeric sequence Δ LSD-valRSec that were substituted by the correlating *T. thermophilus* LeuRS residue are shaded. (B) Aminoacylation activity of chimeric mutants. The reaction was carried out using 4 μ M in vitro transcribed tRNA^{Leu} and 100 nM enzyme. An insert illustrates the wild-type activity relative to the mutants. The bold arrow within the insert identifies the region of the graph that has been blown up. (C) Post-transfer editing activity of single and double mutations of Δ LSD-valRSec. (D) Post-transfer editing activity of triple mutations of Δ LSD-valRSec. Post-transfer editing assays included about 6.5 μ M [³H]-Ile-tRNA^{Leu} and 100 nM enzyme. Symbols representing wild-type and mutant enzymes are as follows: no enzyme (\square), wild type (\circ), Δ LSD-valRStt (\blacktriangledown), Δ LSD-valRSec (\blacktriangle), R vec L (\blacklozenge), E vec K (\times), RE vec LK ($*$), RED vec LKE (\diamond), and REI vec LKV ($+$). Error bars represent the assay reproduced at least in triplicate and are presented for each point.

arginine for a leucine (R vec L) as well as a glutamic acid for a lysine (E vec K) in the *E. coli* ValRS peptide. Both single mutations stimulated aminoacylation activity of the Δ LSD-valRSec mutant (Figure 4B). As found for the LS-domain deletion mutants above, swapping these peptide

residue(s) had minimal, if any, effects on post-transfer editing (Figure 4C).

Interestingly, the R vec L and E vec K single mutations had K_m values of 1.6 and 1.3 μM , respectively, which were comparable to the wild-type enzyme (12). However, a large decrease in the apparent k_{cat} of about 25-fold results in substantially diminished activity. Combination of these two mutations (RE vec LK) showed further enhancement of aminoacylation activity compared to the single chimeric mutants (Figure 4B). While the double LeuRS mutant and chimeric $\Delta\text{LSD-valRStt}$ had a 5-fold increase in the apparent k_{cat} , compared to the single mutant, this enhanced activity was diminished somewhat by a corresponding increase in apparent K_m of approximately 5-fold.

We also tested whether leucylation activity of the inactive $\Delta\text{LSD-valRSec}$ could be enhanced further by swapping a third residue. Using the RE vec LK LeuRS chimeric mutant as a starting point, we swapped two homologous residues. One aspartic acid is conserved, while the second aspartic acid is replaced by a homologous glutamic acid (Figure 4A). This latter aspartic acid in the *E. coli* ValRS peptide was swapped with glutamic acid (RED vec LKE). In addition, the isoleucine was replaced with valine to yield a triple mutant, REI vec LKV. Both of the triple mutations had little, if any, affect on post-transfer editing (Figure 4D). Leucylation activity was slightly increased for the REI vec LKV mutant relative to RED vec LKE (Figure 4B). The RE vec LK double mutant exhibited lower leucylation activity compared to both the two triple mutants (RED vec LKE and REI vec LKV) and $\Delta\text{LSD-valRStt}$, which appeared to be primarily due to an increase in the apparent K_m to 5.4 μM for tRNA^{Leu} . All three mutants had comparable apparent k_{cat} parameters that were about 6-fold decreased compared to wild type (Table 1). Since the LeuRS mutants RED vec LKE, REI vec LKV, and $\Delta\text{LSD-valRStt}$ have similar apparent K_m and k_{cat} parameters, it appears that the key residues within the context of the ValRS peptide for stimulating leucylation activity are the conserved leucine and lysine residues.

Chimeric Mutations Selectively Impact Amino Acid Activation or Transfer to tRNA. Amino acid-dependent pyrophosphate exchange assays were carried out to test activity of the first step of the aminoacylation reaction. All of the chimeric LeuRS mutants, even $\Delta\text{LSD-valRSec}$, which was inactive in aminoacylation, activated amino acid (Figure 5). Interestingly, however, the triple mutations (REI vec LKV and RED vec LKE) as well as $\Delta\text{LSD-valRStt}$ were significantly more active in pyrophosphate exchange than the double and single mutations. Although the leucine-dependent pyrophosphate exchange activity of the triple mutations were decreased compared to the wild-type LeuRS, they were similar to the $\Delta\text{LSD-valRStt}$. This suggests that the peptide from ValRS is optimized for amino acid activation, but homologous valine and glutamic acid substitutions may actually impede transfer of the amino acid to the tRNA. Likewise, we hypothesize that although the available X-ray crystal structures fail to detect direct interactions between the LS-domain and the tRNA^{Leu} , this domain that is idiosyncratic to LeuRS influences tRNA aminoacylation during catalysis.




FIGURE 5: Pyrophosphate exchange activity of chimeric mutants. (A) The amino acid-dependent pyrophosphate exchange reaction was carried out in the presence of 1 mM leucine, 1 mM ATP, and 1 mM $[\text{P}^{32}]\text{PP}_i$. (B) Weakly active mutants in (A) are blown up. Symbols representing wild-type and mutant enzymes are as follows: wild type (\circ), $\Delta\text{LSD-valRStt}$ (\blacktriangledown), $\Delta\text{LSD-valRSec}$ (\blacktriangle), R vec L (\blacklozenge), E vec K (\times), RE vec LK ($*$), RED vec LKE (\diamond), and REI vec LKV ($+$). Error bars represent the assay reproduced at least in triplicate and are presented for each point.

DISCUSSION

The LS-domain is found in most bacteria and some eukaryotic LeuRSs. Although the X-ray crystal structure of *T. thermophilus* LeuRS (5) shows that the LS-domain folds into a discrete domain, it is not conserved in sequence or size (Figure 1). Thus, it may have evolved in a species-specific manner. Herein, we determined that the LS-domain is clearly important for aminoacylation but has little, if any, impact on amino acid editing. Although the X-ray crystal structure did not indicate any interactions between the LS-domain and the tRNA or propose obvious roles for this idiosyncratic domain during catalysis, our investigation suggests that it significantly influences tRNA interactions with the enzyme. The LS-domain may interact directly with tRNA^{Leu} during catalysis at a transient step that is not captured by the ensemble of X-ray cocrystal structures. This might include substrate binding, translocation, and/or product release.

The LS-domain may also influence critical neighboring sequences in aminoacylation. The more conserved KMSKS sequence is located in a loop just downstream of the LS-domain. The role of the mobile KMSKS motif in the class I tyrosyl-tRNA synthetase (TyrRS) has been extensively analyzed via structural and biochemical investigations at various stages of the reaction (33–36). Upon ATP binding, the KMSKS loop shifts to a “closed” conformation where the two lysines interact with the α - and β -phosphates of the ATP molecule, respectively, moving it closer to the active site (36). The α -phosphate group of the ATP molecule is juxtaposed to the carbonyl oxygen of the bound tyrosine ligand to facilitate adenylate bond formation (36, 37). Interestingly, however, the structure of the TyrRS–adenylate complex suggests that, in the closed conformation, the KMSKS would clash with the 3' adenosine of the tRNA

acceptor end (37). Thus, Kobayashi et al. have suggested that the flexible KMSKS loop moves to a "semiopen" conformation to facilitate entry of the tRNA acceptor stem (36). Once the 3' CCA end of the tRNA is in the active site, the KMSKS loop could again adopt the closed conformation and interact with the phosphate backbone of the tRNA 3' CCA end (36).

Deletion of the entire LS-domain may hinder movement of the KMSKS motif that is required for catalysis. Replacement of the LS-domain with a short chimeric sequence that is found in another class I enzyme could better facilitate the movement of the KMSKS loop to initiate activation but hinders KMSKS flexibility to efficiently revert to the semiopen conformation that allows entry of the tRNA acceptor end to the active site. Thus, the Δ LSD-valRSst and the two triple chimeric LeuRS mutants productively activated amino acids but failed to efficiently transfer the amino acid to tRNA during aminoacylation. In LeuRS, it is possible that both the LS-domain and the KMSKS loop work in tandem to sequentially and efficiently orchestrate substrate binding for catalysis and/or product release. Kinetic analysis showed that this peptide is exquisitely sensitive to changes in k_{cat} and K_m but in an idiosyncratic way. It is also possible that this sensitivity is amplified by the adjacent KMSKS conserved sequence that is required for aminoacylation (27, 29, 38, 39).

Structure predictions (17) of the ValRS peptides that were inserted into the two chimeric LeuRS mutants suggest that the inserted sequence of the chimeric LeuRS folds into a short α -helix. Interestingly, this correlates with the *P. horikoshii* TyrRS structure where the sequence just upstream of the KMSKS motif also forms a helix (34). This helix has been proposed to contribute to tRNA binding (34). Structures of TyrRS from *Archaeoglobus fulgidus*, *P. horikoshii*, and *Aeropyrum pernix* show that the KMSKS sequence has adopted different orientations among the three organisms (34). In the tyrosine-bound form, the KMSKS loop is positioned in a closed conformation for *A. pernix*. In contrast, the conformation of the KMSKS loop of *P. horikoshii* TyrRS correlates to the semiclosed form. The homologous KMSSS loop of the *A. fulgidus* TyrRS is positioned further away from the active site (34). Thus, the mobility of the KMSKS sequence required for catalysis may differ among LeuRSs with and without the LS-domain insert. Among the LeuRSs that contain the LS-domain, the mobility of the KMSKS sequence might be altered to permit cooperative movement that would enhance catalysis.

It still remains unclear why LeuRS of some organisms evolved to incorporate this compact LS-domain and others do not. Sequence alignments of corresponding LeuRSs lacking the LS-domain show that the peptide sequence before the KMSKS motif is relatively homologous among the ValRSs and IleRSs. Perhaps the LS-domain was inserted in some, but not all, LeuRSs to confer an evolutionary advantage to tRNA selection and/or catalysis by certain LeuRSs.

ACKNOWLEDGMENT

We thank Ms. Amy Williams as well as Drs. Richard Mursinna and Tommie Lincecum for technical advice and valuable discussions. We are grateful to Dr. Stephen Cusack

for providing coordinates of the *T. thermophilus* LeuRS cocrystal structure.

REFERENCES

1. Ibba, M., and Söll, D. (2000) Aminoacyl-tRNA synthesis, *Annu. Rev. Biochem.* 69, 617–650.
2. Martinis, S. A., Plateau, P., Cavarelli, J., and Florentz, C. (1999) Aminoacyl-tRNA synthetases: a family of expanding functions, *EMBO J.* 18, 4591–4596.
3. Lincecum, T. L., Jr., and Martinis, S. A. (2005) Leucyl-tRNA synthetase, in *The Aminoacyl-tRNA Synthetases* (Ibba, M., Francklyn, C., and Cusack, S., Eds.) pp 36–47, Landes Bioscience, Georgetown, TX.
4. Rossmann, M. G., Liljas, A., Branden, C. I., and Banaszak, L. J. (1975) Evolutionary and structural relationships among dehydrogenases, in *The Enzymes* (Boyer, P. D., Ed.) 3rd ed., pp 61–102, Academic Press, New York.
5. Cusack, S., Yaremchuk, A., and Tukalo, M. (2000) The 2 Å crystal structure of leucyl-tRNA synthetase and its complex with a leucyl-adenylate analogue, *EMBO J.* 19, 2351–2361.
6. Hou, Y. M., Shiba, K., Mottes, C., and Schimmel, P. (1991) Sequence determination and modeling of structural motifs for the smallest monomeric aminoacyl-tRNA synthetase, *Proc. Natl. Acad. Sci. U.S.A.* 88, 976–980.
7. Burbum, J. J., and Schimmel, P. (1991) Structural relationships and the classification of aminoacyl-tRNA synthetases, *J. Biol. Chem.* 266, 16965–16968.
8. Starzyk, R. M., Webster, T. A., and Schimmel, P. (1987) Evidence for dispensable sequences inserted into a nucleotide fold, *Science* 237, 1614–1618.
9. Tukalo, M., Yaremchuk, A., Fukunaga, R., Yokoyama, S., and Cusack, S. (2005) The crystal structure of leucyl-tRNA synthetase complexed with tRNA^{Leu} in the post-transfer-editing conformation, *Nat. Struct. Mol. Biol.* 12, 923–930.
10. Fukunaga, R., and Yokoyama, S. (2005) Crystal structure of leucyl-tRNA synthetase from the archaeon *Pyrococcus horikoshii* reveals a novel editing domain orientation, *J. Mol. Biol.* 346, 57–71.
11. Zheng, Y. G., Wei, H., Ling, C., Martin, F., Eriani, G., and Wang, E. D. (2004) Two distinct domains of the β subunit of *Aquifex Aeolicus* leucyl-tRNA synthetase are involved in tRNA binding as revealed by a three-hybrid selection, *Nucleic Acids Res.* 32, 3294–3303.
12. Hsu, J. L., Rho, S. B., Vanella, K. M., and Martinis, S. A. (2006) Functional divergence of a unique C-terminal domain of leucyl-tRNA synthetase to accommodate its splicing and aminoacylation roles, *J. Biol. Chem.* 281, 23075–23082.
13. Fukunaga, R., Ishitani, R., Nureki, O., and Yokoyama, S. (2005) Crystallization of leucyl-tRNA synthetase complexed with tRNA^{Leu} from the archaeon *Pyrococcus horikoshii*, *Acta Crystallogr., Sect. F: Struct. Biol. Cryst. Commun.* 61, 30–32.
14. Fukunaga, R., and Yokoyama, S. (2005) Aminoacylation complex structures of leucyl-tRNA synthetase and tRNA^{Leu} reveal two modes of discriminator-base recognition, *Nat. Struct. Mol. Biol.* 12, 915–922.
15. Lincecum, T. L., Jr., Tukalo, M., Yaremchuk, A., Mursinna, R. S., Williams, A. M., Sproat, B. S., Van Den Eynde, W., Link, A., Van Calenbergh, S., Gröthli, M., Martinis, S. A., and Cusack, S. (2003) Structural and mechanistic basis of pre- and post-transfer editing by leucyl-tRNA synthetase, *Mol. Cell* 11, 951–963.
16. Liu, Y., Liao, J., Zhu, B., Wang, E. D., and Ding, J. (2006) Crystal structures of the editing domain of *Escherichia coli* leucyl-tRNA synthetase and its complexes with met and ile reveal a lock-and-key mechanism for amino acid discrimination, *Biochem. J.* 394, 399–407.
17. Guex, N., and Peitsch, M. C. (1997) Swiss-Model and the Swiss-Pdbviewer: an environment for comparative protein modeling, *Electrophoresis* 18, 2714–2723.
18. Martinis, S. A., and Fox, G. E. (1997) Non-standard amino acid recognition by *Escherichia coli* leucyl-tRNA synthetase, *Nucleic Acids Symp. Ser.* 36, 125–128.
19. Zhai, Y., and Martinis, S. A. (2005) Two conserved threonines collaborate in the *Escherichia coli* leucyl-tRNA synthetase amino acid editing mechanism, *Biochemistry* 44, 15437–15443.
20. Grodberg, J., and Dunn, J. J. (1988) OmpT encodes the *Escherichia coli* outer membrane protease that cleaves T7 RNA polymerase during purification, *J. Bacteriol.* 170, 1245–1253.

21. Sampson, J. R., and Uhlenbeck, O. C. (1988) Biochemical and physical characterization of an unmodified yeast phenylalanine transfer RNA transcribed *in vitro*, *Proc. Natl. Acad. Sci. U.S.A.* 85, 1033–1037.
22. Normanly, J., Ogden, R. C., Horvath, S. J., and Abelson, J. (1986) Changing the identity of a transfer RNA, *Nature* 321, 213–219.
23. Tocchini-Valentini, G., Saks, M. E., and Abelson, J. (2000) tRNA leucine identity and recognition sets, *J. Mol. Biol.* 298, 779–793.
24. Mursinna, R. S., and Martinis, S. A. (2002) Rational design to block amino acid editing of a tRNA synthetase, *J. Am. Chem. Soc.* 124, 7286–7287.
25. Schreier, A. A., and Schimmel, P. R. (1972) Transfer ribonucleic acid synthetase catalyzed deacylation of aminoacyl transfer ribonucleic acid in the absence of adenosine monophosphate and pyrophosphate, *Biochemistry* 11, 1582–1589.
26. Lincecum, T. L., Jr., and Martinis, S. A. (2000) The tRNA synthetase proofreading and editing active site: a novel antibiotic drug, in *SAAS Bulletin Biochemistry and Biotechnology* (Ballal, S. K., Ed.) Vol. 13, pp 25–33, SAAS, Cookeville, TN.
27. Mechulam, Y., Dardel, F., Le, Corre, D., Blanquet, S., and Fayat, G. (1991) Lysine 335, part of the KMSKS signature sequence, plays a crucial role in the amino acid activation catalysed by the methionyl-tRNA synthetase from *Escherichia coli*, *J. Mol. Biol.* 217, 465–475.
28. Schmitt, E., Meinel, T., Blanquet, S., and Mechulam, Y. (1994) Methionyl-tRNA synthetase needs an intact and mobile 332KM-SKS336 motif in catalysis of methionyl adenylate formation, *J. Mol. Biol.* 242, 566–576.
29. First, E. A., and Fersht, A. R. (1995) Analysis of the role of the KMSKS loop in the catalytic mechanism of the tyrosyl-tRNA synthetase using multimeric cycles, *Biochemistry* 34, 5030–5043.
30. Fukai, S., Nureki, O., Sekine, S., Shimada, A., Vassilyev, D. G., and Yokoyama, S. (2003) Mechanism of molecular interactions for tRNA(val) recognition by valyl-tRNA synthetase, *RNA* 9, 100–111.
31. Fukai, S., Nureki, O., Sekine, S., Shimada, A., Tao, J., Vassilyev, D. G., and Yokoyama, S. (2000) Structural basis for double-sieve discrimination of L-valine from L-isoleucine and L-threonine by the complex of tRNA(Val) and valyl-tRNA synthetase, *Cell* 103, 793–803.
32. Lee, K. W., and Briggs, J. M. (2004) Molecular modeling study of the editing active site of *Escherichia coli* leucyl-tRNA synthetase: two amino acid binding sites in the editing domain, *Proteins* 54, 693–704.
33. Hountondji, C., Lederer, F., Dessen, P., and Blanquet, S. (1986) *Escherichia coli* tyrosyl- and methionyl-tRNA synthetases display sequence similarity at the binding site for the 3'-end of tRNA, *Biochemistry* 25, 16–21.
34. Kuratani, M., Sakai, H., Takahashi, M., Yanagisawa, T., Kobayashi, T., Murayama, K., Chen, L., Liu, Z. J., Wang, B. C., Kuroishi, C., Kuramitsu, S., Terada, T., Bessho, Y., Shirouzu, M., Sekine, S., and Yokoyama, S. (2006) Crystal structures of tyrosyl-tRNA synthetases from archaea, *J. Mol. Biol.* 355, 395–408.
35. Xin, Y., Li, W., and First, E. A. (2000) The “KMSKS” motif in tyrosyl-tRNA synthetase participates in the initial binding of tRNA(Tyr), *Biochemistry* 39, 340–347.
36. Kobayashi, T., Takimura, T., Sekine, R., Kelly, V. P., Kamata, K., Sakamoto, K., Nishimura, S., and Yokoyama, S. (2005) Structural snapshots of the KMSKS loop rearrangement for amino acid activation by bacterial tyrosyl-tRNA synthetase, *J. Mol. Biol.* 346, 105–117.
37. Yaremchuk, A., Krikilivyi, I., Tukalo, M., and Cusack, S. (2002) Class I tyrosyl-tRNA synthetase has a class II mode of cognate tRNA recognition, *EMBO J.* 21, 3829–3840.
38. Hountondji, C., Dessen, P., and Blanquet, S. (1993) The SKS of the KMSKS signature of class I aminoacyl-tRNA synthetases corresponds to the GKT/S sequence characteristic of the ATP-binding site of many proteins, *Biochimie* 75, 1137–1142.
39. Hountondji, C., Lazennec, C., Beauvallet, C., Dessen, P., Pernollet, J. C., Plateau, P., and Blanquet, S. (2002) Crucial role of conserved lysine 277 in the fidelity of tRNA aminoacylation by *Escherichia coli* valyl-tRNA synthetase, *Biochemistry* 41, 14856–14865.

BI062078J