

Subunit-Selective Interrogation of CO Recombination in Carbonmonoxy Hemoglobin by Isotope-Edited Time-Resolved Resonance Raman Spectroscopy[†]

Gurusamy Balakrishnan,^{#,||} Xiaojie Zhao,^{#,⊥} Edyta Podstawska,^{‡,§} Leonard M. Proniewicz,^{‡,§} James R. Kincaid,^{*,‡} and Thomas G. Spiro^{*,#,||}

Department of Chemistry, Princeton University, Princeton, New Jersey 08544, Department of Chemistry, Marquette University, Milwaukee, Wisconsin 53233, Department of Chemistry, Jagiellonian University, Krakow, Poland, Department of Chemistry, University of Washington, Seattle, Washington 98195, and Advanced Optowave Corporation, P.O. Box 530, Bohemia, New York 11716

Received November 27, 2008; Revised Manuscript Received February 25, 2009

ABSTRACT: Hemoglobin (Hb) is an allosteric tetrameric protein made up of $\alpha\beta$ heterodimers. The α and β chains are similar, but are chemically and structurally distinct. To investigate dynamical differences between the chains, we have prepared tetramers in which the chains are isotopically distinguishable, via reconstitution with ¹⁵N-heme. Ligand recombination and heme structural evolution, following HbCO dissociation, was monitored with chain selectivity by resonance Raman (RR) spectroscopy. For α but not for β chains, the frequency of the ν_4 porphyrin breathing mode increased on the microsecond time scale. This increase is a manifestation of proximal tension in the Hb T-state, and its time course is parallel to the formation of T contacts, as determined previously by UVRR spectroscopy. Despite the localization of proximal constraint in the α chains, geminate recombination was found to be equally probable in the two chains, with yields of $39 \pm 2\%$. We discuss the possibility that this equivalence is coincidental, in the sense that it arises from the evolutionary pressure for cooperativity, or that it reflects mechanical coupling across the $\alpha\beta$ interface, evidence for which has emerged from UVRR studies of site mutants.

Hemoglobin (Hb) continues to offer a compelling laboratory for the study of allostery in proteins. Ligand binding induces a shift in the Hb tetramer from a low-(T) to a high-(R) affinity structure (*I*–3), but analysis of this process is complicated by the fact that the tetramer is composed of heterodimers, $\alpha\beta$. The α and β chains are similar, but distinct. Much evidence points to differing constraints on ligand binding in the T state. In the α chains, the Fe atom is restrained by its proximal histidine ligand from readily approaching the heme plane in order to bind a distal ligand. This constraint is at the core of Perutz' stereochemical trigger mechanism for allostery (*4*). In the deoxy form, the Fe atom is five-coordinate and high-spin, and nonbonded forces drive it out of the heme plane toward the proximal ligand. Binding of the diatomic ligands O₂, NO, or CO requires the Fe to approach the heme plane, and it becomes low-spin. Constraint on this motion by the protein lowers the ligand affinity, while overcoming the constraint induces changes in the protein structure and alters contacts at the subunit interfaces, thereby destabilizing the T state.

However, the indicators of proximal constraint are attenuated in the β chain, which appear to be more flexible than the α chain (*5*). On the other hand, the distal binding pocket is more crowded in the T state for the β than for the α chain, leading Perutz to suggest an ordered sequence of ligand binding, first to the α and then to the β chains, the latter following the T–R conversion (*4*).

Since then there have been a number of studies attempting to distinguish α from β chain reactivity (*6*). All appear to show a surprising degree of functional similarity between the two kinds of subunits, even though structural and spectroscopic differences persist. However, most of these studies have employed hybrid constructs, in which α or β chains are substituted with metal ions that do not bind the ligand under study, in order to isolate the behavior of the functional Fe-containing chain (*6*–*19*). Although structural integrity is maintained in these hybrids, there are nevertheless questions about indirect effects of chemical substitution. Seeking to address the issue of $\alpha\beta$ equivalence in the *native* protein, we have employed chain-specific isotope labeling, reconstituting Hb with heme having ¹⁵N in the pyrrole rings. Time-resolved resonance Raman (TR3) spectroscopy was then used to monitor dynamics at the heme following HbCO photodissociation. The heme resonance Raman (RR) spec-

[#] Princeton University.

^{||} University of Washington.

[⊥] Advanced Optowave Corporation.

[‡] Marquette University.

[§] Jagiellonian University.

[†] This work was supported by NIH Grant GM 12526 (to T.G.S.) and DK35153 (to J.R.K.).

* Corresponding author. (T.G.S.) Phone: (206) 685-4964. Fax: (206) 685-8665. E-mail: spiro@chem.washington.edu. (J.R.K.) Phone: (414) 288-3539. Fax: (414) 288-7066. E-mail: james.kincaid@marquette.edu.

¹ Abbreviations: RR, resonance Raman; UVRR, ultraviolet resonance Raman; TR3, time-resolved resonance Raman; TLC, thin-layer chromatography; HbCO, carbonmonoxy hemoglobin; Hbdoxy, deligated hemoglobin; ¹⁵N–Fe^{III}(PPIX)Cl, ¹⁵N-ferritroporphyrin IX chloride; PMB, sodium *p*-(hydroxymercuri)benzoate; α_{N15}^{CO} , α_{N14}^{CO} , β_{N15}^{CO} , or β_{N14}^{CO} , α - or β -subunits containing CO bound heme with ¹⁴N or ¹⁵N isotopes at the pyrrole ring.

trum can distinguish α from β heme because the vibrational bands are isotope-sensitive.

We find that CO recombination rates and geminate recombination yields are indistinguishable for the α and β chains. However, the spectra also show a differing structural response of the two chains. We consider the possibility that the functional equivalence is a reflection of intradimer coupling between the two heme sites, or rather that chain-selective evolutionary changes lead to the same function for individual chains.

MATERIALS AND METHODS

Sample Preparation. ^{15}N -Ferriprotoporphyrin IX chloride (^{15}N -Fe^{III}(PPIX)Cl) was graciously provided by Professor Kevin Smith and used after checking for purity via thin-layer chromatography (TLC), proton NMR and electronic absorption spectroscopy (20–22). Sodium *p*-(hydroxymercuri)benzoate (PMB) and sodium perchlorate were purchased from Aldrich Chemical Co. and used without further purification. Biogel P-6DG (Bio-Rad Laboratories, Hercules, CA), CM cellulose (CM-52), and diethylaminoethyl cellulose (DE-53) (Whatman, Fairfield, NJ) were used for protein column chromatography.

Protein Preparation. Hb was isolated from red blood cells according to established procedures (23) and stored as the CO adduct. ApoHb was prepared by the acid–acetone method (24), and reconstitution with ^{15}N -isotopically labeled protoheme, to form ^{15}N -Hb, was accomplished as described previously (23, 25, 26); Hb and ^{15}N -Hb were dissociated into their subunits by reaction with PMB (27) and isolated as reported in the literature (25). Isotopically labeled hybrids were obtained as the carbon monoxide adducts by incubating the proper subunits, that is, $\alpha_{\text{N}15}^{\text{CO}}$ and $\beta_{\text{N}14}^{\text{CO}}$. The same was done for the $\alpha_{\text{N}14}^{\text{CO}}$ and $\beta_{\text{N}15}^{\text{CO}}$ subunits. Purification was accomplished by column chromatography (25–27).

Spectroscopy. Time-Resolved Resonance Raman (TR³). The TR³ spectroscopic setup has been described elsewhere (28, 29). Briefly, photolysis pulses at 419 nm (20 ns, 60 $\mu\text{J}/\text{pulse}$, 1 kHz) were generated by frequency doubling the output of a Ti:Sapphire laser (Photonics International Industries), which was pumped (527 nm, ~ 10 mJ/pulse, 70 ns, 1 kHz) by an intracavity frequency-doubled Nd:YLF laser (GM30, Photonics International Inc.). Probe pulses (20 ns, 1 kHz) at 426 nm (0.5 $\mu\text{J}/\text{pulse}$) were obtained by frequency doubling a second Ti:Sapphire laser. Scattered light was collected at 135° with a pair of fused quartz lenses, f-matched to a 1.26 m spectrograph (Spex 1269), which was equipped with a holographic grating (3600 groove mm^{-1}) and intensified photodiode-array detector (IPDA-1024 S/B, Roper Scientific). A time-delay signal generator (DG535, Stanford Research) was used to control the delay time between pump and probe pulses and synchronize the Raman scattered signals from the probe pulse with the gate of the detector. Interfering signals from the pump laser were eliminated by electronically gating the detector using a high voltage pulse generator (Roper Scientific, PG201). The time-resolution of the current set up is limited to 200 ns by the minimum gate-width of the IPDA intensifier at 1 kHz. The probe pulse wavelength, 426 nm, was selected to provide comparable resonance enhancement for the ν_4 RR bands of deoxyHb and HbCO.

HbCO samples having a concentration of ~ 0.25 mM in heme were prepared using CO-saturated phosphate buffer (pH 7.5, 50 mM). About 350 μL of sample solution was contained in a 5 mm glass NMR tube and spun around a stationary stirring wire to ensure vertical mixing of the sample. A CO atmosphere was maintained near the top surface of the sample during the entire experiment by continuous delivery of water saturated CO, under a vacuum exhaust. The diameters of the pump and probe laser spots at the samples were estimated to be ~ 80 and 60 μm , respectively, and the diameter of the NMR tube was 5 mm. To avoid spatial overlap mismatch between pump and probe laser spots, and to ensure that successive pulses excited a new sample volume, the spinning speed of the NMR tube (monitored online by a tachometer (08210, Cole Parmer)) was adjusted to 800 rpm for 200 ns to 3 μs time delays, 500 rpm for 6 to 50 μs delays, and 300 rpm for longer delay times. Sample integrity was checked by the reproducibility of Raman and absorption spectra.

RR Cross Section Measurement. The Raman setup used for static RR measurements was the same as above. Solutions of deoxyHb (60 μM in heme) were generated in 50 mM phosphate buffer (containing 0.2 M in NaClO_4 as a frequency and intensity standard) at pH 7.4 by irradiating a solution of HbCO with visible light at 4 °C under flowing ultrapure N_2 . The fraction of dimeric Hb is negligible at this concentration for deoxy Hb (30). About 0.4 mL of solution was transferred to an NMR tube under N_2 and the N_2 atmosphere was maintained during the RR spectral measurements.

Owing to the limited narrow spectral band-pass (220 cm^{-1}), the ν_4 heme band and the ClO_4^- reference band were measured in separate experiments without altering conditions. Identical results were obtained from three repeated measurements. These measurements were followed immediately by measuring the HbCO spectrum (CO atmosphere) under identical conditions, because the higher HbCO concentration (0.25 mM in heme) required to minimize the dimer dissociation, and lower laser power (0.3 mW) to minimize CO photodissociation, precluded the use of 0.2 M in NaClO_4 as the intensity standard owing to its very weak signal. The RR cross section of HbCO was calculated relative to deoxyHb from their relative spectral intensities. The RR cross section was obtained using the following equation (31)

$$\sigma_{\text{sample}} = \sigma_{\text{std}} \times \left(\frac{I_{\text{sample}}}{I_{\text{std}}} \right) \times \left(\frac{C_{\text{std}}}{C_{\text{sample}}} \right) \times \left(\frac{\epsilon_{\text{sample}} + \epsilon_0}{\epsilon_{\text{std}} + \epsilon_0} \right) \quad (1)$$

where σ is the RR cross section, I is the intensity, C is the molar concentration and ϵ_0 , ϵ_{sample} , and ϵ_{std} are the molar extinction coefficients (in $\text{M}^{-1} \text{cm}^{-1}$) of the sample at the laser excitation wavelength (ϵ_0) and at the Raman scattered wavelengths of the sample (ϵ_{sample}) and internal standard (ϵ_{std}), respectively.

The spectral data were processed with *Grams/AI* (7.0) software (Thermo Galactic). The HbCO spectra were deconvoluted using two Lorentzian bands, one for the CO bound form having a fixed bandwidth (8.6 cm^{-1}) and one for the photoproduct whose bandwidth was optimized. (It was not possible to use fixed bandwidth for the photoproduct, since as discussed below, the ν_4 band of the α subunits shifts to higher frequency during the conformational change from the R to T state, resulting in band broadening in the




FIGURE 1: Static RR spectra excited at 426 nm (0.3 mW) for the deoxy and CO-adducts of HbA and of the $(^{14}\alpha^{15}\beta)_2$ isotope hybrid.

composite traces.) In the case of the hybrids, the isotope splitting gives four resolved bands, permitting the use of four Lorentzian, two for the CO bound forms, both having a fixed bandwidth (8.6 cm^{-1}), and two for the deoxy components of the photoproducts, both having a fixed bandwidth (11.2 cm^{-1}). Band areas were used to calculate the RR cross sections.

RESULTS

The ν_4 RR band, arising from a breathing mode of the porphyrin ring, is the most prominent peak in RR spectra excited in resonance with the heme Soret absorption band, seen near 400 nm (32, 33). It is responsive to electron density in the porphyrin ring, and shifts up strongly when CO binds to deoxyHb (Figure 1), as electron density shifts from the ring to the CO ligand through back-donation of Fe electrons. This frequency separation permits documentation of the HbCO ligand saturation from measured peak intensities (34–36). In addition, ν_4 shifts down upon ^{15}N substitution at the pyrrole N atoms, for example, by 7 cm^{-1} in Ni-octaethyl porphyrin (37). A similar shift in Hb allows for chain-selective resolution of ν_4 , as seen in Figure 1. The peak heights reveal unequal cross-sections of the chains, higher and lower for α than β chains in the deoxy- and CO-forms, respectively (Table 1) (38, 39). These differences must arise from unequal resonance enhancements, perhaps stemming from slight differences in the Soret electronic bands of the two chains (13).

Figure 2 shows time-resolved RR spectra in the ν_4 region following a saturating photolysis pulse. Because of geminate recombination, a substantial HbCO peak is seen, even at the earliest time delay, and this peak grows at the expense of the deoxyHb peak at later times, reflecting bimolecular recombination. The bound CO fraction was determined at each delay, by deconvoluting the peaks and dividing their heights by those obtained from a probe-only spectrum (in order to compensate for the measurable photolysis induced by the probe pulse) (34–36). The resulting progress curves

Table 1: RR Cross Sections^a for the ν_4 Bands of HbA and of the $^{14}\alpha^{15}\beta$ Hybrid

	$^{426}\sigma_{\nu_4} \times 10^{-24}\text{ cm}^2\text{ mol}^{-1}\text{ str}^{-1}$	
	R_CO	T_deoxy
HbA ($\alpha + \beta$)	1.08	1.18
HbA (α -subunits)	0.50	0.78
HbA (β -subunits)	0.58	0.40

^a Calculated for CO adducts relative to the deoxy peaks, and for the deoxy hemes relative to ClO_4^- internal standard ($\sigma_{\text{ClO}_4^-} = 10.4 \times 10^{-30}\text{ cm}^2/\text{mol str}$) (38). The self-absorption correction was calculated from absorptivities: ($\epsilon_0 = 481\,840\text{ M}^{-1}\text{ cm}^{-1}$ at 426 nm, $\epsilon_{\text{sample}} = 58\,670\text{ M}^{-1}\text{ cm}^{-1}$ at deoxy ν_4 scattered wavelength of 452 nm, and $\epsilon_{\text{std}} = 298827\text{ M}^{-1}\text{ cm}^{-1}$ at ClO_4^- band scattered wavelength of 444 nm) (39).




FIGURE 2: Time-resolved RR spectra at the indicated time (μs) following photolysis of the CO adduct of HbA and of the $(^{15}\alpha^{14}\beta)_2$ and $(^{14}\alpha^{15}\beta)_2$ hybrids. Band positions and assignments are marked at the top. Samples were prepared at a heme concentration of $\sim 0.25\text{ mM}$ in CO saturated 50 mM sodium phosphate buffer [$\lambda_{\text{pump}} = 419\text{ nm}$, 65 mW, $\sim 20\text{ ns}$, 1 kHz; $\lambda_{\text{probe}} = 426\text{ nm}$, 0.5 mW, $\sim 20\text{ ns}$, 1 kHz].

(Figure 3) follow an initial rise to a plateau, representing the geminate yield, and a subsequent rise toward full saturation, due to recombination of CO from solution. The solid curves were constructed with the aid of previously determined (40) time constants for geminate recombination ($\tau_{\text{gem}} = 0.07\text{ }\mu\text{s}$) and for bimolecular recombination to the R and T states of Hb ($\tau_4, \tau_5 = 250$ and $1500\text{ }\mu\text{s}$ at 1 atm CO).

For the $^{15/14}\text{N}$ hybrids, the ν_4 envelope was decomposed into individual components via curve resolution. The quality of the curve fitting is illustrated in Figure 4. Essentially the same progress curves were obtained for both hybrids (Figure 3), and for native HbA, confirming the validity of the methodology. The important result is that the progress curves are indistinguishable for α and β chains, which have the same geminate yield, 0.39 ± 0.02 and similar recombination rates (the curves are too short to be precise about the rates).

However, the two chains do differ in their spectral response. Accurate fitting of the curves required a distinct ν_4 upshift with time for the α but not the β chains. As illustrated in Figure 5, among the four ν_4 components, only the α_{deoxy} peak shifts position with time, whether the chain contains ^{14}N or ^{15}N labeled heme. This shift was confirmed by taking difference spectra between early and late times,




FIGURE 3: The CO rebinding to photolyzed CO adducts of HbA and the indicated Hb hybrids, determined from the intensity of the ν_4 CO-heme band. The continuous lines correspond to modeling of the CO recombination reaction with the time constants (τ_0), τ_{gem} (geminate phase), τ_2 (R-state bimolecular rebinding), and τ_5 (T-state bimolecular rebinding).

which revealed a sigmoidal band only for the α_{deoxy} signal (Figure S1, Supporting Information). The time course for the α_{deoxy} ν_4 shift is shown in Figure 6, which also shows that the corresponding shift for unlabeled deoxyHb is half that of the α_{deoxy} peak, consistent with the HbA signal being a composite of the shifted α peak and the unshifted β peak. The solid curves are drawn to confirm to the time constants determined via time-resolved UVRR spectroscopy (34) for signals associated with changes in protein structure. Most of the α_{deoxy} ν_4 shift is associated with τ_2 (2 μs) and τ_3 (20 μs), the protein motions, which involve formation of the T state contacts (34).

DISCUSSION

The present results confirm a distinct structural difference between the two chains in the deoxy forms. We find that the ν_4 frequency shifts up with time in the α but not in the β chains. The largest changes (Figure 6) occur at ~ 2 and 20 μs after CO photodissociation, phases known from UVRR studies (34) to be associated with formation of the T state contacts at the “hinge” and “switch” regions of the $\alpha_1\beta_2$ interface.

The position of ν_4 is known to correlate negatively with the stretching frequency, $\nu_{\text{Fe-His}}$, of the bond between the heme Fe and the proximal histidine ligand, for a variety of deoxyHb constructs (41). A depressed $\nu_{\text{Fe-His}}$ is observed in the Hb T-state (216 cm^{-1} vs 222 cm^{-1} in deoxy-myoglobin or in relaxed forms of deoxyHb) (42, 43) and is the most direct evidence for tension at the proximal Fe-His connection, underlying the constraint to ligand binding that triggers the T-R transition in Perutz’ stereochemical mechanism (4). Moreover, the evolution of $\nu_{\text{Fe-His}}$ after HbCO photolysis was previously shown to track T state formation (44). Earlier RR

studies of hybrid Hb’s have shown that this tension is expressed in the α chains and the present results confirm that the tension develops in concert with the R-T conversion.

However, this tension is not manifest in the β chains. The β_{deoxy} ν_4 does not evolve in time, and the metal hybrid data indicate little change in $\nu_{\text{Fe-His}}$ between R and T states for the deoxy- β -chains (45).

Immediately after photolysis, neither chain is under tension, and the present results show them to be functionally equivalent. The CO rebinding rates and geminate yields are indistinguishable (Figure 3). UVRR studies have shown CO rebinding to be concurrent with the first concerted protein motion after photolysis (34, 36) which is detected as a loss of tertiary H-bond contacts, and interpreted as a rotation of the E and F helices that hold the heme like a clamshell (Figure 7). Displacement of the E helix toward the heme was suggested to impel the photodissociated CO to either rebind or to exit the heme pocket (34, 36).

This mechanism is presumed to be the same for both chains. Nevertheless, it is surprising that the branching ratio between CO exit and rebinding is exactly the same.

One possibility is that mechanical coupling across the $\alpha_1\beta_1$ dimer interface ensures equivalent functionality. In other words, rebinding to one-half of an $\alpha\beta$ dimer induces a structural change that is propagated to the other half, requiring equivalent rebinding at the opposite heme; the dimer then acts as a single dynamical unit. This hypothesis is supported by UVRR results showing equivalent changes in the protein dynamics when one of the anchoring interhelical H-bonds is abolished in one chain or another, or in both (46). Moreover, transmission of distal interactions across the $\alpha_1\beta_1$ interface was earlier demonstrated by Levy et al. (47), who found distinctive differences in the ligation of the distal histidine to Fe(III)-heme in valency hybrid Hbs, depending on whether ligated subunits were in the same dimer (symmetric hybrids) or in the opposite dimer (asymmetric hybrids). Molecular dynamics simulation of an $\alpha\beta$ dimer revealed spontaneous formation of a Fe(III)-distal histidine bond in the α subunit; this did not occur if the interface was frozen in the simulation, nor did it occur in isolated subunits (48).

Intradimer cooperativity in the T-state has long been propounded by Ackers and co-workers (49), based on extensive measurements of dimer assembly free energies. Analysis of several hybrid Hb’s revealed greater stability of doubly ligated tetramers when the two ligands are on the same dimer than when they are on opposite dimers. There has been controversy about the extent of the intradimer cooperativity, based on error analysis of different methodologies (50, 51), but the basic phenomenon seems well established (50).

The other possibility is that the α/β equivalence is intrinsic to the chains, and arises from evolutionary pressure to endow structurally inequivalent chains with the same functional properties. QM/MM computations indicate equal contribution to the R-T energy difference from structural change in the two chains (5). There is, of course, selective pressure for such a development, since deviations from binding equivalence reduce cooperativity, and therefore physiological efficiency, in the Hb tetramer (6). A variety of measurements have produced evidence for near-equivalence of the chains both in the R-state as well as in the T-state. Thus Bettati et




FIGURE 4: Example of RR band deconvolution at 3, 50, and 300 μs following CO photolysis, showing the experimental data (black) fitted spectra (green), deconvoluted deoxy bands (red), CO bands (blue), and the residuals (gray).




FIGURE 5: Deconvolution data showing that ν_4 shift with time only for α -subunits in the deoxy state.




FIGURE 6: Time evolution (log scale) for the deoxy ν_4 frequencies of HbA and for the α -subunits in $(^{15}\alpha^{14}\beta)_2$ and $(^{14}\alpha^{15}\beta)_2$ hybrids, following CO-photolysis. The continuous lines correspond to modeling with a series of time constants ($\tau_1 = 65$ ns; $\tau_2 = 0.74$ μs ; $\tau_3 = 20.5$ μs ; $\tau_{4,5} = 497$ μs) previously documented for protein structural changes from UVRR spectroscopy (34).

al. estimated the T-state O_2 affinities of α and β chains to differ by less than a factor of 3 from polarized spectral measurements of deoxyHb single crystals (52). The most comprehensive study of chain differences was carried out




FIGURE 7: Diagram of the EF clamshell rotation proposed (34, 36) to follow HbCO photodissociation. Displacement of the high-spin Fe from the heme plane impels the F helix away from the heme, while E helix movement toward the heme expels CO from the heme pocket or else induces geminate recombination.

by Unzai et al. (6), who analyzed CO and O_2 binding for hybrid Hb's in which one of the chains contained Mg(II) or Ni(II), which stabilize the T state, or Cr(III) and Mn(III), which stabilize the R state. They found similar affinities and binding rates for both chains in both states. There were up to 2-fold chain differences in CO and O_2 association and dissociation rates for R-state Hb and for CO association rates in the T-state (CO dissociation proved hard to measure); for O_2 the T-state association and dissociation rates were essentially the same for the two chains. While intradimer coupling could explain α/β functional equivalence in native Hb, it cannot do so for metal (symmetric) hybrids, since in that case either the α or β chains are nonfunctional in ligand binding.

The most relevant comparison of our results is with the rebinding measurements on Co/Fe hybrids by Hofrichter et al. (13), who found CO geminate rebinding times of 76 and 53 ns and geminate yields of 0.35 and 0.22 for β and α chains, respectively. The geminate yield is critical to the present argument since it is sensitive to steric factors in the heme pocket, which control the probability of rebinding vs escape (4, 53). These factors are different in the two chains.

The β pocket is more crowded than the α pocket, and this crowding might be expected to increase the rebinding probability. The 50% higher geminate yield of β than of α chains measured by Hofrichter et al. (13) is consistent with this difference. However, Samuni et al. have reported that the dissociated CO rapidly escapes into adjacent cavity in β -subunits of the $\alpha(\text{Zn})\beta(\text{FeCO})$ hybrid in the T-state and thus offsets the crowding effect, possibly favoring a higher geminate yield (54). In an event, the lower α yield was attributed by Hofrichter et al. (13) to a fraction of the molecules being in the T-state is, in which the geminate yield is known to be lowered. A variety of evidence supports the inference that the T-state is favored in α -Fe-containing, but not the β -Fe-containing Co/Fe hybrid (13). Thus it remains possible, though surprising, that the indistinguishability of the α and β geminate yields in native Hb is entirely ascribable to convergent evolution of the chains.

CONCLUSIONS

A variety of observations point to similar ligand affinities and binding rates in the two chains of Hb, despite well-characterized structural differences (6). This similarity can be viewed as the evolution of similar function by different intrachain mechanisms, driven by the requirement of high binding cooperativity for physiological efficiency of O₂ delivery by tetrameric Hb. However, the present results, in which rebinding is monitored for individual chains within native Hb, reveal identical α and β geminate yields, although a higher β geminate yield is expected for the more crowded β chains (4). A possible mechanism for abolition of chain differences is mechanical coupling that ensures equivalent binding in both halves of an $\alpha\beta$ dimer. A higher β than α geminate yield has been reported for Co/Fe hybrids (13), in which intradimer coupling would not be operative. However, this difference could be attributed to a population of T-state molecules in the α -Fe hybrid. Thus, while the present results firmly document equivalency in geminate rebinding, the precise mechanism for suppressing chain inequivalence remains an open issue.

SUPPORTING INFORMATION AVAILABLE

A detailed illustration of the deoxy ν_4 shift in the α subunits. This material is available free of charge via the Internet at <http://pubs.acs.org>.

REFERENCES

- Perutz, M. F. (1990) *Mechanisms of Cooperativity and Allosteric Regulation in Proteins*; Cambridge University Press: Cambridge.
- Baldwin, J., and Chothia, C. (1979) Hemoglobin - structural-changes related to ligand-binding and its allosteric mechanism. *J. Mol. Biol.* 129, 175.
- Monod, J., Wyman, J., and Changeux, J. P. (1965) On nature of allosteric transitions - A plausible model. *J. Mol. Biol.* 12, 88.
- Perutz, M. F. (1970) Stereochemistry of cooperative effects in haemoglobin. *Nature* 228, 726.
- Alcantara, R. E., Xu, C., Spiro, T. G., and Guallar, V. (2007) A quantum-chemical picture of hemoglobin affinity. *Proc. Nat. Acad. Sci. U. S. A.* 104, 18451-18455.
- Unzai, S., Eich, R., Shibayama, N., Olson, J. S., and Morimoto, H. (1998) Rate constants for O₂ and CO binding to the alpha and beta subunits within the R and T states of human hemoglobin. *J. Biol. Chem.* 273, 23150-23159.
- Brunori, M., Amiconi, G., Antonini, E., Wyman, J., and Winterha, Kh. (1970) Artificial intermediates in reaction of haemoglobin - functional and conformational properties of cyanmet intermediates. *J. Mol. Biol.* 49, 461.
- Cassoly, R., and Gibson, Q. H. (1972) Kinetics of ligand binding to hemoglobin valency hybrids and effect of anions. *J. Biol. Chem.* 247, 7332.
- Hoffman, B. M., Gibson, Q. H., Bull, C., Crepeau, R. H., Edelstein, S. J., Fisher, R. G., and McDonald, M. J. (1975) Manganese-substituted hemoglobin and myoglobin. *Ann. N.Y. Acad. Sci.* 244, 174-186.
- Ikedaaito, M., and Yonetani, T. (1980) Studies on cobalt myoglobins and hemoglobins. 11. The interaction of carbon-monoxide and oxygen with alpha-subunits and beta-subunits in iron-cobalt hybrid hemoglobins. *J. Mol. Biol.* 138, 845-858.
- Blough, N. V., and Hoffman, B. M. (1982) Equilibrium and kinetic CO binding parameters of T-state and R-state hemoglobin chains using iron-manganese hybrids. *J. Am. Chem. Soc.* 104, 4247-4250.
- Blough, N. V., and Hoffman, B. M. (1984) Carbon-monoxide binding to the ferrous chains of [Mn,Fe(II)] hybrid hemoglobins - pH-dependence of the chain affinity constants associated with specific hemoglobin ligation pathways. *Biochemistry* 23, 2875-2882.
- Hofrichter, J., Henry, E. R., Sommer, J. H., Deutsch, R., Ikedaaito, M., Yonetani, T., and Eaton, W. A. (1985) Nanosecond optical-spectra of iron cobalt hybrid hemoglobins - geminate recombination, conformational-changes, and intersubunit communication. *Biochemistry* 24, 2667-2679.
- Shibayama, N., Yonetani, T., Regan, R. M., and Gibson, Q. H. (1995) Mechanism of ligand-binding to Ni(II)-Fe(II) hybrid hemoglobins. *Biochemistry* 34, 14658-14667.
- Philo, J. S., Dreyer, U., and Lary, J. W. (1996) Quaternary structure dynamics and carbon monoxide binding kinetics of hemoglobin valency hybrids. *Biophys. J.* 70 1949-1965.
- Unzai, S., Hori, H., Miyazaki, G., Shibayama, N., and Morimoto, H. (1996) Oxygen equilibrium properties of chromium(III)-iron(II) hybrid hemoglobins. *J. Biol. Chem.* 271, 12451-12456.
- Ackers, G. K., Doyle, M. L., Myers, D., and Daugherty, M. A. (1992) Molecular code for cooperativity in hemoglobin. *Science* 255, 54-63.
- Mathews, A. J., and Olson, J. S. (1994) Assignment of Rate Constants for O₂ and CO Binding to Alpha-Subunit and Beta-Subunit within R-State and T-State Human Hemoglobin, in *Hemoglobins, Part C*, pp 363-386, Academic Press, New York.
- Mathews, A. J., and Olson, J. S. (1994) Assignment of rate constants for O₂ and CO binding to alpha-subunit and beta-subunit within R-state and T-state human hemoglobin. *Methods Enzymol.* 232, 363.
- Fuhrhop, J. H., and Smith, K. M., Eds. (1975) *Laboratory Methods in Porphyrin and Metalloporphyrin Research*, Elsevier Publishing Co., North Holland, Amsterdam.
- DiNello, R. K., and Chang, C. K. (1978) Isolation and modification of natural porphyrins, in *The Porphyrins: Structure and synthesis* (Dolphin, D., Ed.), pp 289-339, Academic Press, New York.
- Janson, T. R., and Katz, J. J. (1979) Nuclear magnetic resonance spectroscopy of diamagnetic porphyrins, in *The Porphyrins: Physical chemistry* (Dolphin, D., Ed.), pp 1-59, Academic Press, New York.
- Riggs, A. (1981) Preparation of blood hemoglobins of vertebrates. *Methods Enzymol.* 76, 5-29.
- Yip, Y. K., Waks, M., and Beychok, S. (1972) Influence of prosthetic groups on protein folding and subunit assembly. 1. Conformational differences between separated human alpha-globins and beta-globins. *J. Biol. Chem.* 247, 7237.
- Jeyarajah, S., and Kincaid, J. R. (1990) Resonance Raman studies of hemoglobins reconstituted with mesoheme - unperturbed iron histidine stretching frequencies in a functionally altered hemoglobin. *Biochemistry* 29, 5087-5094.
- Smith, K. M., Parish, D. W., and Inouye, W. S. (1986) Methyl deuteration reactions in vinylporphyrins - protoporphyrins IX, III, and XIII. *J. Org. Chem.* 51, 666-671.
- Yip, Y. K., Waks, M., and Beychok, S. (1977) Reconstitution of native human hemoglobin from separated globin chains and alloplex intermediates. *Proc. Nat. Acad. Sci. U. S. A.* 74, 64-68.
- Zhao, X. J., Chen, R. P., Tengroth, C., and Spiro, T. G. (1999) Solid-state tunable kHz ultraviolet laser for Raman applications. *Appl. Spectrosc.* 53, 1200-1205.
- Zhao, X. J., Tengroth, C., Chen, R. P., Simpson, W. R., and Spiro, T. G. (1999) Time-resolved Raman spectroscopy with a tunable ultraviolet kilohertz nanosecond laser. *J. Raman Spectrosc.* 30, 773-776.

30. Thomas, J. O., and Edelstein, S. J. (1973) Observation of dissociation of unliganded hemoglobin. 2. Effect of pH, salt, and dioxane. *J. Biol. Chem.* 248, 2901–2905.
31. Chang, S., Yang, W., and Spiro, T. G. (1990) Saturation effects on ultraviolet resonance Raman intensities - excimer/YAG laser comparison and aromatic amino-acid cross-sections. *J. Raman Spectrosc.* 21, 435–440.
32. Spiro, T. G., and Li, X. Y. (1987) Resonance Raman Spectra of Metalloporphyrins, in *Biological Application of Raman Spectroscopy* (Spiro, T. G., Ed.) pp 1–37, John Wiley & Sons, New York.
33. Spiro, T. G., and Streckas, T. C. (1974) Resonance Raman-spectra of heme proteins - effects of oxidation and spin state. *J. Am. Chem. Soc.* 96, 338–345.
34. Balakrishnan, G., Case, M. A., Pevsner, A., Zhao, X. J., Tengroth, C., McLendon, G. L., and Spiro, T. G. (2004) Time-resolved absorption and UV resonance Raman spectra reveal stepwise formation of T quaternary contacts in the allosteric pathway of hemoglobin. *J. Mol. Biol.* 340, 843–856.
35. Friedman, J. M., and Lyons, K. B. (1980) Transient Raman-study of CO-hemoprotein photolysis - origin of the quantum yield. *Nature* 284, 570–572.
36. Jayaraman, V., Rodgers, K. R., Mukerji, I., and Spiro, T. G. (1995) Hemoglobin allostery - resonance Raman-spectroscopy of kinetic intermediates. *Science* 269, 1843–1848.
37. Li, X. Y., Czernuszewicz, R. S., Kincaid, J. R., Stein, P., and Spiro, T. G. (1990) Consistent porphyrin force-field. 2. Nickel octaethylporphyrin skeletal and substituent mode assignments from N-15, meso-D4, and methylene-D16 Raman and infrared isotope shifts. *J. Phys. Chem.* 94, 47–61.
38. Dudik, J. M., Johnson, C. R., and Asher, S. A. (1985) Wavelength dependence of the preresonance Raman cross-sections of CH₃CN, SO₄²⁻, ClO₄⁻, and NO₃. *J. Chem. Phys.* 82, 1732–1740.
39. <http://omlc.ogi.edu/spectra/hemoglobin/summary.html>.
40. Hu, X. H., Frei, H., and Spiro, T. G. (1996) Nanosecond stepscan FTIR spectroscopy of hemoglobin: Ligand recombination and protein conformational changes. *Biochemistry* 35, 13001–13005.
41. Ondrias, M. R., Rousseau, D. L., Shelnutz, J. A., and Simon, S. R. (1982) Quaternary-transformation-induced changes at the heme in deoxyhemoglobins. *Biochemistry* 21, 3428–3437.
42. Friedman, J. M., Rousseau, D. L., and Ondrias, M. R. (1982) Time-resolved resonance Raman studies of hemoglobin. *Annu. Rev. Phys. Chem.* 33, 471–491.
43. Friedman, J. M., Rousseau, D. L., Ondrias, M. R., and Stepnoski, R. A. (1982) Transient Raman-study of hemoglobin - structural dependence of the iron-histidine linkage. *Science* 218, 1244–1246.
44. Jayaraman, V., and Spiro, T. G. (1996) Structural evolution of the heme group during the allosteric transition in hemoglobin: Insights from resonance Raman spectra of isotopically labeled heme. *Biospectroscopy* 2, 311–316.
45. Ondrias, M. R., Rousseau, D. L., Kitagawa, T., Ikedasaito, M., Inubushi, T., and Yonetani, T. (1982) Quaternary structure changes in iron-cobalt hybrid hemoglobins detected by resonance Raman-scattering. *J. Biol. Chem.* 257, 8766–8770.
46. Balakrishnan, G., Tsai, C. H., Wu, Q., Case, M. A., Pevsner, A., McLendon, G. L., Ho, C., and Spiro, T. G. (2004) Hemoglobin site-mutants reveal dynamical role of interhelical H-bonds in the allosteric pathway: Time-resolved UV resonance Raman evidence for intra-dimer coupling. *J. Mol. Biol.* 340, 857–868.
47. Levy, A., Sharma, V. S., Zhang, L., and Rifkind, J. M. (1992) A new mode for heme-heme interactions in hemoglobin associated with distal perturbations. *Biophys. J.* 61, 750–755.
48. Ramadas, N., and Rifkind, J. M. (1999) Molecular dynamics of human methemoglobin: The transmission of conformational information between subunits in an alpha beta dimer. *Biophys. J.* 76, 1796–1811.
49. Ackers, G. K., Holt, J. M., Huang, Y. W., Grinkova, Y., Klinger, A. L., and Denisov, I. (2000) Confirmation of a unique intra-dimer cooperativity in the human hemoglobin alpha(1)beta(1) half-oxygenated intermediate supports the symmetry rule model of allosteric regulation. *Prot. Str. Func. Gen.* 23–43.
50. Holt, J. M., Klinger, A. L., Yarian, C. S., Keelara, V., and Ackers, G. K. (2005) Asymmetric distribution of cooperativity in the binding cascade of normal human hemoglobin. 1. Cooperative and noncooperative oxygen binding in Zn-substituted hemoglobin. *Biochemistry* 44, 11925–11938.
51. Yun, K. M., Morimoto, H., and Shibayama, N. (2002) The contribution of the asymmetric alpha 1 beta 1 half-oxygenated intermediate to human hemoglobin cooperativity. *J. Biol. Chem.* 277, 1878–1883.
52. Bettati, S., and Mozzarelli, A. (1997) T state hemoglobin binds oxygen noncooperatively with allosteric effects of protons, inositol hexaphosphate, and chloride. *J. Biol. Chem.* 272, 32050–32055.
53. Dantsker, D., Roche, C., Samuni, U., Blouin, G., Olson, J. S., and Friedman, J. M. (2005) The position 68(E11) side chain in myoglobin regulates ligand capture, bond formation with heme iron, and internal movement into the xenon cavities. *J. Biol. Chem.* 280, 38740–38755.
54. Samuni, U., Juszczak, L., Dantsker, D., Khan, I., Friedman, A. J., Perez-Gonzalez-De-Apodaca, J., Bruno, S., Hui, H. L., Colby, J. E., Karasik, E., Kwiatkowski, L. D., Mozzarelli, A., Noble, R., and Friedman, J. M. (2003) Functional and spectroscopic characterization of half-liganded iron-zinc hybrid hemoglobin: Evidence for conformational plasticity within the T state. *Biochemistry* 42, 8272–8288.

BI802190F