

Structures of A β -Related Peptide–Monoclonal Antibody Complexes^{†,‡}

Anna Gardberg,^{§,○} Lezlee Dice,^{||,◇} Kathleen Pridgen,[§] Jan Ko,[⊥] Paul Patterson,[⊥] Susan Ou,[⊥] Ronald Wetzel,[#] and Chris Dealwis^{*,△}

[§]Department of Biochemistry and Cellular and Molecular Biology, University of Tennessee, Knoxville, Tennessee 37996,

^{||}Graduate School of Medicine, University of Tennessee, Knoxville, Tennessee 37920, [⊥]Division of Biology, California Institute of Technology, Pasadena, California 91125, [#]Structural Biology Department and Pittsburgh Institute for Neurodegenerative Diseases, Pittsburgh, Pennsylvania 15260, and [△]Department of Pharmacology and Center for Proteomics, School of Medicine, Case Western Reserve University, 10900 Euclid Avenue, Cleveland, Ohio 44106 [○]Current address: Center for Structural Molecular Biology, Oak Ridge National Laboratory, Oak Ridge, TN 37831-6142 [◇]Current address: Department of Molecular and Structural Biochemistry, North Carolina State University, Raleigh, NC 27607

Received January 26, 2009; Revised Manuscript Received April 17, 2009

ABSTRACT: Passive immunotherapy (PI) is being explored as a potential therapeutic against Alzheimer's disease. The most promising antibodies (Abs) used in PI target the EFRH motif of the A β N-terminus. The monoclonal anti-A β Ab PFA1 recognizes the EFRH epitope of A β . PFA1 has a high affinity for A β fibrils and protofibrils (0.1 nM), as well as good affinity for A β monomers (20 nM). However, PFA1 binds the toxic N-terminally modified pyroglutamate peptide pyro-Glu3-A β with a 77-fold loss in affinity compared to the WT A β (1–8). Furthermore, our earlier work illustrated PFA1's potential for cross-reactivity. The receptor tyrosine kinase Ror2, which plays a role in skeletal and bone formation, possesses the EFRH sequence. PFA1 Fab binds the Ror2(518–525) peptide sequence REEFRHEA with a 3-fold enhancement over WT A β (1–8). In this work, the crystal structures of the hybridoma-derived PFA1 Fab in complex with pyro-Glu3-A β peptide and with a cross-reacting peptide from Ror2 have been determined at resolutions of 1.95 and 2.7 Å, respectively. As with wild-type A β , these peptides bind to the Fab via a combination of charge- and shape-complementarity, hydrogen-bonding, and hydrophobic interactions. Comparison of the structures of the four peptides A β (1–8), Grip1, pyro-Glu3-A β (3–8), and Ror2 in complex with PFA1 shows that the greatest conformational flexibility occurs at residues 2 to 3 and 8 of the peptide. These structures provide a molecular basis of the specificity tolerance of PFA1 and its ability to recognize A β N-terminal heterogeneity. The structures provide clues to improving mAb specificity and affinity for pyroglutamate A β .

The deposition of amyloid aggregates of the amyloid β (A β)¹ peptide is a hallmark of the underlying pathology of Alzheimer's disease (AD). The A β peptides, A β (1–40) and A β (1–42), are degradation products of the amyloid precursor protein (APP) resulting from the sequential proteolysis by the secretases, β and γ , respectively (1). With two extra C-terminal residues, A β (1–42) is considered the more toxic species found in amyloid deposits of both humans and animal models of AD. The direct immunization of patients with A β fibrils in a phase II clinical trial was halted due

to the development of meningoencephalitis in 6% of the patients. Passive immunotherapy (PI) where the humanized IgGs are intravenously administered is potentially a safer treatment. At least three mechanisms for the effectiveness of A β immunotherapy in AD models have been proposed. Two mechanisms, microglial activation (2, 3) and catalytic dissolution (4), require that the antibody enter the brain. Another mechanism, sometimes called the peripheral sink hypothesis, does not have this requirement. In the peripheral sink hypothesis (5), Abs bind to A β in the blood and shift the distribution of A β between the brain and the peripheral circulatory system, leading to a net efflux of A β from the CNS to the peripheral circulatory system where it is degraded (5). All three mechanisms require Ab binding to A β in either monomer or aggregated forms. Since it is unclear which of these mechanisms contributes to AD therapy, it can be argued that the most appropriate Abs for therapy are those capable of recognizing all assembly forms of A β peptides.

In a recent study, we demonstrated the molecular basis of passive immunotherapy by determining the first X-ray crystal structures of the N-terminal epitope of A β (1–8) (sequence

[†]This work was supported by NIH Grants NS46356-04A1 and R01AG018416. The work of J. K., S. O. and P. H. P. was supported by grants from the McGrath Foundation and NIA/NIH.

[‡]The atomic coordinates have been deposited with RCSB ID 3EYU (PFA1-Ror2(518–525)) and 3EYS (PFA1-pyro-Glu3-A β (3–8)).

*Corresponding author. Phone: (216) 368-1652. Fax: (216) 368-1300. E-mail: chris.dealwis@case.edu.

Abbreviations: A β , amyloid β ; AD, Alzheimer's disease; PFA1, protofibril antibody 1; PFA2, protofibril antibody 2; SPR, surface plasmon resonance; GRIP1, glutamate receptor interacting protein 1; Ror2, receptor-related neurotrophic tyrosine kinase; AMPA, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionate; PEG3350, polyethylene glycol, average molecular weight 3350.

DAEFRHDS) bound to the monoclonal antibodies (mAbs) PFA1 and PFA2 (6). Briefly, the essential binding motif on A β is the EFRH sequence at positions 3–6; this motif is recognized by the Fabs' complementarity determining regions, notably a WWDD region within heavy chain CDR2 in both PFA1 and PFA2.

PFA1 and PFA2 mAbs bind A β monomers, protofibrils, and fibrils with nanomolar affinities (6) and hence show promise as potential immunotherapy agents. Besides universal binding to all known physical forms of A β (1–40) and A β (1–42), however, a desired antibody or antibodies for treating AD should theoretically exhibit certain properties in their binding profile. First, there are a variety of chemical degradation products of A β *in vivo*, some of which are of equal or greater concern, *vis à vis* aggregate formation and toxicity, than the intact A β sequence. Second, since the mAb's binding epitope is a relatively short sequence, it is possible that similar or identical sequences might exist elsewhere in the proteome and exhibit cross-reactivity to the mAb with potential toxic side effects. Thus, the ideal Ab for PI would exhibit good specificity for all toxic and aggregating forms of A β while at the same time avoiding high-affinity interactions with other proteins.

The analysis of neuritic and vascular plaques shows that there is substantial N-terminal heterogeneity of A β . For example, the A β peptide can have its D residues at positions 1 and 7 racemized or isomerized, and there are pyroglutaminyl (pyro-Glu) forms at residues 3 and 11 (7–10). In fact, there is evidence to suggest the major component of AD senile plaques is the N-terminal modified A β beginning at E3, which is posttranslationally modified to an N-terminal pyro-Glu(11–15). In some cases pyro-Glu forms of A β at residues 3 (pyro-Glu3-A β (3–8)) and 11 (pyro-Glu11-A β) constitute more than 50% of the A β in neuritic plaques (11). Interestingly, relative to age-matched controls, AD sufferers have an abundance of pyro-Glu-A β in their neuritic amyloid deposits, suggesting that pyro-Glu-A β may play an important role in the onset of AD.

Supporting evidence for a role for pyro-Glu-A β peptides in AD includes the following: (a) the pyro-Glu residue stabilizes the amyloid peptides against degradation by aminopeptidases (16, 17); (b) pyro-Glu forms of A β appear in early stages of the disease (18); (c) pyro-Glu-modified peptides form aggregates 250-fold faster than WT A β (19); (d) pyro-Glu-A β peptides display an enhanced cytotoxicity compared to WT A β (9, 14); (e) based on *in vitro* studies, pyro-Glu-modified A β peptides are potential seeding species for A β aggregate formation *in vivo* (20, 21); and (f) recently, inhibitors of the glutaminyl cyclase were shown to reduce plaque burden dramatically and improve cognition in AD mice (22). Unfortunately, given pyro-Glu3-A β 's potential importance in disease etiology, we found that PFA1 Fab binds pyro-Glu3-A β with a much reduced affinity (77-fold difference in K_d) when compared to A β (1–8) (6). We would like to explore the structural basis of this loss of affinity.

The shortness of the A β sequence epitope also raises possible specificity issues. As a preliminary investigation of the specificity of PFA1 and PFA2 with respect to the human genome, the WT A β (2–7) peptide AEFRHD and mutants derived from proteins listed in ref 6, AKFRHD, AEIRHD, AEFRSD, and REEF-RHEA, were synthesized, and their affinity for the two Fab fragments was determined by surface plasmon resonance (SPR) (Table 1). AEIRHD and AEFRSD showed no measurable binding to PFA1 and PFA2. However, we found that PFA1 Fab binds to AKFRHD (a sequence found in glutamate receptor

Table 1: Alignment of Peptide Sequences Relevant to This Work and K_d Measurements for Peptide Binding to the PFA1 Fab^a

protein/peptide	sequence	K_d (nM) measurements from SPR
A β (1–40) WT ("WT")	DAEFRDHS...	39.0
A β (2–7) WT	AEFRHD	60
pyro-Glu3-A β (3–8)	pEFRHDS	3000
Ror2(518–525)	REEFRHEA	24
Grip1(110–115)	AKFRHD	3400
position 4 mutant	AEIRHD	no binding
position 6 mutant	AEFRSD	no binding

^a Taken from ref 6.

interacting protein 1 (GRIP1)) with an affinity 28 times lower than to AEFRHD (6). Perhaps more significantly, the peptide sequence REEF-RHEA, found in the cytosolic tyrosine kinase domain of human receptor-related neurotrophic tyrosine kinase (Ror2(518–525)), actually binds to PFA1 and PFA2 with approximately twice the affinity of the WT A β (2–7) peptide AEFRHD. Interestingly, Ror2 plays a role in bone formation (23), while Grip1 is responsible for maintaining the plasticity of α -amino-3-hydroxy-5-methyl-4-isoxazolepropionate (AMPA) glutamate receptors by PDZ domain interactions (24, 25).

Here we report the crystal structures of PFA1 complexed with pyro-Glu3-A β (3–8) and Ror2(518–525). These structures provide the molecular details that should make possible the optimization of mAb structure to appropriately adjust its binding affinities for these peptides.

MATERIALS AND METHODS

Protein Production and Purification. The hybridoma production of PFA1 and its purification and Fab generation were described previously(6).

Crystallization, Data Collection, and Processing. Crystal screening was performed with the PEGs Suite (QIAGEN) using the PHOENIX crystallization robot (Art Robbins Instruments). The PFA1-Ror2(518–525) complex and the PFA1-pyro-Glu3-A β (3–8) complex were crystallized using the sitting drop vapor diffusion method. Cocrystals of the PFA1-Ror2(518–525) peptide complex appeared in a solution containing 0.2 M LiCl and 20% w/v PEG3350 without a buffer. The cocrystals of the PFA1-pyro-Glu3-A β (3–8) complex were grown in a solution consisting of 0.1 M Tris-HCl, pH 8.5, and 25% (w/v) PEG2000-MME. For X-ray data collection, all crystals were harvested into a cryoprotectant solution. For the PFA1-Ror2(518–525) cocrystals, this consisted of 0.2 M LiCl, 25% PEG3350, and 20% glycerol. The PFA1-pyro-Glu3-A β (3–8) cocrystals were cryoprotected using 0.1 M Tris-HCl, pH 8.5, 28% PEG2000-MME, and 20% glycerol. Crystals were flash-frozen in liquid nitrogen. All diffraction data were collected at the Advanced Photon Source, beamline 14BMC (BIOCARS-CAT), with a Q315 detector. The oscillation range was 0–250° in 0.5° increments for the PFA1-Ror2(518–525) cocrystals and was 0–190° in 0.5° increments for the PFA1-pyro-Glu3-A β (3–8) cocrystals. Data for both of the cocrystals were indexed, integrated, and scaled using the HKL2000 package (26). Although the data for PFA1 complexed with pyro-Glu is 92% complete at 2.2 Å resolution, the data can be processed to 1.95 Å resolution with greater than 50% completeness in the last shell. The PFA1-Ror2 complex diffracted to 2.7 Å resolution with 85% completeness. The lack

Table 2: Crystallographic Data Collection and Refinement Parameters

	PFA1-Ror2 (518–525)	PFA1-pyro- Glu3-A β (3–8)
	Data Collection	
space group	<i>P</i> 1	<i>P</i> 1
cell dimensions		
<i>a</i> , <i>b</i> , <i>c</i> (Å)	41.6, 42.6, 58.7	41.7, 42.7, 58.6
α , β , γ (deg)	96.1, 93.6, 91.7	96.1, 93.8, 91.7
resolution (Å)	30–2.7 (2.8–2.7)	35–1.9 (2.02–1.95)
(last shell)		
<i>R</i> _{sym}	0.083 (0.222)	0.049 (0.284)
<i>I</i> / <i>I</i> σ	12.8 (2.8)	17.5 (2.2)
completeness (%)	85.0 (49.9)	86.4 (54.2)
redundancy	2.3 (1.6)	1.8 (1.5)
	Refinement	
resolution (Å)	2.7	1.9
no. of reflections	8780	23906
TLS groups	5	0
<i>R</i> _{work} / <i>R</i> _{free} (%)	17.7/24.8	20.4/25.6
no. of atoms		
protein (non-H)	3421	3495
ligand/ion	0	6 (glycerol)
water	23	113
<i>B</i> -factors (Å ²)	32	35
bound peptide	40	52
ligand/ion	none	41 (glycerol)
water	30	40
rmsd bond lengths (Å)	0.010	0.010
rmsd bond angles (deg)	1.2	1.3
Ramachandran	99.5	99.7
allowed region (%)		
PDB code	3EYU	3EYS

of >90% completeness can be attributed to the large blind spot region encountered in the *P*1 space group.

Molecular Replacement and Structure Refinement. The structures were solved by molecular replacement using the constant (CL + CH1) and variable portions of PFA1 (PDB code 1IPU) separately using MOLREP (27). Protein rebuilding, including water picking, was performed using Coot (28). Refinement was conducted using Refmac5 (29), and the structures were validated with SFCHECK (30), PROCHECK (31), and MOLPROBITY (32). As reported by PROCHECK, 99.7% and 99.5% of the residues lie in the allowed regions of the Ramachandran plot for PFA1-pyro-Glu3-A β (3–8) and PFA1-Ror2(518–525) complex structures, respectively. Data collection and refinement statistics appear in Table 2. Calculation of buried surface area was carried out using AREAIMOL (33, 34) with a 1.4 Å probe radius.

RESULTS

To explore the binding mode of pyro-Glu3-F-R-H-D-S (pyro-Glu3-A β (3–8)) peptide to the mAb PFA1, as well as the potential for cross-reactivity with the protein Ror2, we determined the three-dimensional X-ray structures of the two binary cocomplexes PFA1-Fab-pyro-Glu3-A β (3–8) and PFA1-Fab-Ror2(518–525) to 1.9 and 2.7 Å, respectively (see Table 2 for peptide sequences). Both structures have reasonable refinement statistics as shown in Table 2, and they are isomorphous to the *apo* PFA1 structure (6). The $2F_o - F_c$ difference Fourier electron density map clearly defines the first five residues of the pyro-Glu3-A β (3–8) and the EFRHEA sequence of the Ror2(518–525) of the




FIGURE 1: Stereoview of $2F_o - F_c$ maps of PFA1 binding to the pyro-Glu3-A β (3–8) peptide (a) and the Ror2(518–525) (b). The maps were contoured at 1.1σ (within 2 Å of the peptide). Numbering scheme is that of A β (1–8) WT (see Table 1).

two structures (Figure 1 and Table 1), and the electron density for the complementarity determining regions (CDRs) is readily traced. The antigen-binding site, which lies in a cleft between the light- and heavy-chain variable domains, is formed by four of the six CDRs: CDR-L1 (QSIVHSNGNTY), CDR-L3 (FQGS HVPLTF), CDR-H2 (IW-WDDDR), and CDR-H3 (VRRHTTVLGDFWFA). The residues in bold-face type directly interact with the antigen.

Overall, the pyro-Glu3-A β (3–8) peptide adopts a slightly extended coil conformation that is stabilized by hydrogen bonds, van der Waals, and ion pair interactions (see Figure 2A). The crucial mAb-binding epitope is the ³EFRH⁶ at the N-terminus of A β , which we previously showed to be extremely sensitive to mutations. Mutating any single residue in this epitope to alanine essentially abolishes binding (6). The (FR) of the ³EFRH⁶ motif makes interactions with a WWDD motif belonging to CDR-H2, similar to that observed in the A β WT complex.

The Glu to pyro-Glu substitution at position 3 has a drastic effect on mAb binding (6). However, the C α rms deviation for residues 3–8 between the WT A β and pyro-Glu3-A β (3–8) is only 0.24 Å. This is not surprising as the main-chain and side-chain conformations between the two peptides are extremely conserved beyond residue 3 (see Figure 2A). In fact, the main-chain conformation of all the peptide complexes determined by our laboratory is extremely conserved as reflected by their small rms deviations (see Table 3 and Figure 2C). The greatest structural difference between WT A β (1–8) and pyro-Glu3-A β (3–8) is found at position 3. Interestingly, the pyroglutaminyl ring occupies overlapping space with the E3 side chain of WT A β (see Figure 2A). Consequently, this places the pyro-Glu3 amide nitrogen within hydrogen-bonding distance of the carbonyl of




FIGURE 2: Comparison of Aβ and related peptide structures. Stereoview of the overlay of Aβ(1–8) WT peptide (green) with (a) pyropeptide (blue) and (b) Ror2(518–525) (pink). The PFA1 residues are drawn with thinner bonds and similarly color-coded, save that the light chain is shown in a paler shade and the heavy chain is shown in a darker shade. The numbering scheme is that of Aβ(1–8) WT. The WWDDD motif appears in the lower left corner of (a) and (b); D57 points away from the bound peptide and does not bind to it. (c) Superposition of Aβ-related peptide structures demonstrating coupled position changes (“cross-talk”) of residues 3 and 8: Aβ(1–8) (cyan), Grip1(110–115) (orange), pyro-Glu3-Aβ (white), and Ror2(518–525) (purple) bound to PFA1. The side chains are marked P2–P8, where P stands for the peptide position.

S97, an interaction that the WT Aβ cannot make. However, while the carbonyl oxygen of pyro-Glu3 occupies a position merely 1.9 Å away from that of the Oε1 atom of the unmodified Aβ E3 side chain, this is still too far away to form compensatory hydrogen bonds; the nearest hydrogen-bonding donor to the carbonyl oxygen atom of pyro-Glu3 is 4 Å away. The carboxylate side chain of E3 in WT Aβ forms two hydrogen bonds to the amide of the main chain and the hydroxyl group of LC S32. The distance between the amide of LC S32 and the Oε1 of E3 is 2.6 Å, while the distance between the hydroxyl of S32 and the Oε2 of E3 is 2.8 Å. In the PFA1-pyro-Glu3-Aβ(3–8) structure, pyro-Glu3 does not form these two important hydrogen bonds. Instead, the oxygen that is attached to the pyroglutaminyl ring is almost 4.7 Å away from the amide nitrogen of S32. Furthermore, E3 makes some long-range (<5 Å) ion pair interactions with the LC H98 side chain. A complete list of pyro-Glu3 interactions with PFA1 is compared with those of the WT Aβ(1–8) in Supporting Information Table 1A,B.

Although there are sequence differences at four positions between WT Aβ(1–8) and the Ror2(518–525) (Table 1), the Ror2(518–525) binds PFA1 similarly to the WT Aβ(1–8). The rmsd between the Cα atoms is 0.3 Å (Table 3). The greatest differences between the two peptide structures occur at E3, at position 7 where the WT Aβ residue D7 is substituted by E in Ror2(518–525), and at the C-terminus, where S8 of the WT Aβ is replaced by A in Ror2(518–525) (Table 1 and Figures 2B and 3A). The E3 side chain adopts a different conformation in the WT Aβ; the carboxylate side chain hydrogen bonds with the hydroxyl of LC S27E, while in Ror2(518–525) the E3 side chain hydrogen bonds with the imidazole side chain of LC H93. Interestingly, the differing E3 interactions between the two peptides appear to result from the S → A substitution at position 8 in Ror2(518–525) that leads to a chain of events spanning nearly an 8 Å distance. First, the substitution causes a main-chain conformational difference that places the carboxyl terminus of Ror2(518–525) further from LC S27E, abrogating the formation of a water-mediated second sphere hydrogen bond between the COOH and LC S27E observed in the WT Aβ complex (see Figure 3A). Second, the Ror2(518–525)’s COOH group is positioned such that it is unable to stabilize the aforementioned water molecule. Indeed, we do not observe electron density for a water molecule near this position in the PFA1- Ror2(518–525) complex structure. Third, the absence of the water molecule prevents the formation of the hydrogen bond between the water and E3 that is observed in WT Aβ, resulting in side-chain conformational flexibility of Ror2(518–525)’s E3 which finds a new hydrogen bond partner in LC H93. These observations suggest that there is cross-talk between the binding sites 3 and 8 (Figures 2B and 3A). Also, the substitution of D to E at position 7 results in a slight change in the main-chain conformation between WT Aβ and Ror2(518–525) (Figure 2B).

Table 3: Root Mean Square Differences (RMSDs) for the CDRs of Five PFA1 Structures

RMSD	WT					
	APO	mon1	Pyro	Grip1	Ror2	
APO		0.276	0.237	0.301	0.276	CA atoms of CDR
WT mon1	0.37		0.242	0.203	0.26	
Pyro	0.288	0.272		0.32	0.189	
Grip1	0.46	0.257	0.36		0.295	
Ror2	0.355	0.3	0.26	0.387		
All CDR atoms						




FIGURE 3: Superposition of A β and related structures. (a) A β (1–8) (green) and Ror2(518–525) (purple) bound to PFA1 (PFA1 residues are drawn in the same colors corresponding to each peptide). (b) Superposition of PFA1 (CDRs and peptide drawn in green) and WO2 (CDRs in cyan and peptide drawn in red) CDRs. (c) Comparison of PFA1-A β (1–8) (peptide and Fab residues are in green) and WO2-A β (1–16) (peptide drawn in red and the Fab residues are in cyan) binding sites. One hydrogen bond is shown in dashed lines between HC Y52 of WO2 and the amide of R3 of the A β (1–16) to highlight the CDR sequence difference at HC Y52. The Fab residues are labeled according to the PFA1 sequence and numbered according to the Kabat convention (<http://www.biochem.ucl.ac.uk/~martin/abs/GeneralInfo.html>).

This, combined with the larger E side chain of Ror2(518–525), positions its carboxyl group further away from the imidazole hydrogen bond donor of HC H102, resulting in a longer hydrogen bond (3.4 Å for PFA1-Ror2(518–525) versus 2.6 Å for PFA1-A β (1–8)) between D7 of the bound peptide and H-H102 of PFA1. However, compared to the WT A β , the larger E7 side chain of Ror2(518–525) makes a much shorter intramolecular ion pair interaction with the guanidinium group of R5 (4.7 versus 5.7 Å) belonging to the EFRH motif. A complete list of the Ror2(518–525) peptide's interactions with PFA1 is compared with those of the WT A β (1–8) in Supporting Information Table 1B,C. PFA1 binds Ror2(518–525) with an approximately 3-fold improved affinity compared to WT A β .

DISCUSSION

To understand why the pyro-Glu modification lowers affinity to PFA1, it is helpful to review the core binding for the WT A β peptide. The E3 side chain of the A β peptide makes important hydrogen bonds with the antibody, its only charge complementarity is with LC H31's side chain, which has an average ~10% charge at physiological pH, so the mAb does not exploit charge complementarity to significantly stabilize E3 (Figure 2A). The A β peptide residues F4, R5, and H6 make more specific interactions, however. F4 sits in a complementarily shaped hydrophobic pocket made up of heavy chain residues W47, H50, W52, and F100(E) (Kabat numbering) and light chain residues V94 and L96 (Figure 4A). Arginine 5 fits neatly into a shallow cavity consisting of heavy chain residues D54 and D56 for charge complementarity, with residue H97 providing π -overlap for the




FIGURE 4: Electrostatics of binding: (a) the electrostatic potential surface of PFA1 with bound A β (1–8) peptide, (b) with bound pyro-Glu3-peptide, and (c) Ror2(518–525). Blue represents positive charge, red indicates negative charge, and the uncharged portions are shown in white. Each of the peptides is drawn with carbon (yellow), nitrogen (blue), and oxygen (red). The numbering scheme is that of A β (1–8) WT.

guanidinium group and with residue W52 providing a hydrophobic patch near the aliphatic atoms of R5's side chain. Finally, H6 points into a complementary pocket, with π -overlap provided by light chain residue Y32. H6 forms hydrogen bonds to light chain residue G91 and to heavy chain residue D100(C); light chain residue H27(D) also contributes to the shape of this cavity. The full impact of the pyro-Glu3 modification to PFA1 binding can be observed when comparing electrostatic binding surfaces (Figure 4A,B). Binding energy of protein–protein or protein–ligand can be expressed as a linear combination of the differences in polar and apolar solvent-accessible surface areas between those states (35). The pyro-Glu3-A β (3–8) peptide buries 432.9 Å² of the antibody surface when binding to PFA1, which is some 60 Å² less than that of the WT A β . The difference in SA is ultimately attributable to the reduced length of the pyro-Glu3-A β (3–8) peptide (6 residues) compared to WT A β (8 residues). The PFA1-pyro-Glu3-A β (3–8) cocrystal structure reveals that the loss of binding of pyro-Glu3-A β to PFA1 is in part due to the loss of crucial hydrogen bonds and long-range ion pair interactions with E3. It is also due to the loss of buried SA caused by the shorter pyro-Glu3-A β (3–8) peptide compared to WT A β peptide.

As noted before (19), the following factors may contribute to pyro-Glu-A β peptides' acceleration of fibril forming kinetics. First, the pyro-Glu(3–40)-A β lacks a charged N-terminal group, making it more hydrophobic (depicted in Figure 4). Second, it has been noted that the capped N-terminus resulting from cyclization reduces A β 's propensity to form intramolecular antiparallel β -sheets (19). This is particularly important, as fiber diffraction and solid-state NMR data have shown the fibrils are composed of intermolecular, in-register, *parallel* β -sheets formed by hydrogen bonding between A β monomers (36, 37). Moreover, the introduction of a lactam modification that restrains rotation of the peptide bond promotes fibrillogenesis by favoring a conformation prone to aggregation (38). Finally, unlike other N-terminal modifications, which are extremely unstable, the pyro-Glu residue stabilizes the A β peptide against degradation by aminopeptidases (16, 17). Hence, its stability, hydrophobicity, and structure make it a perfect seed for fibrillogenesis and, thus, a good target for mAb therapy.

Recently, two other related structures have been reported (39, 40). The first structure is of A β (17–36) in complex with an affibody adopting a β -conformation (39). However, as the N-terminus of A β is absent in this structure, we are unable to do a comparison with our PFA1-A β (1–8) complex. The second structure is that of A β (1–16) in complex with the Fab fragment of the mAb WO2 (40). Although residues 1–16 of A β were used in the cocrystallization experiment, only residues 1–8 of A β were observed in the electron density maps. Comparison of the PFA1-A β (1–8) with the WO2-Fab-A β (1–16) structure reveals that the greatest conformational differences between the two Fabs occur at the two CDRs H2 and H3 that interact with residues 7 and 8 of the A β peptide (see Figure 3b,c). The A β (1–8) peptide conformations bound to PFA1 and WO2 are extremely well conserved between residues 1–6 (see Figure 3c). The FR residues of the EFRH motif in the WO2Fab-A β (1–16) structure, as in our PFA1-A β (1–8) structure, interact with the CDRH2 residues 52–56; in PFA1-A β (1–8), this segment comprises a WWDDD motif. This motif is conserved in WO2, except that in the WO2 mAb, HC W52 is substituted by Y. The side-chain hydroxyl of HC Y52 in WO2 makes a hydrogen bond with the main-chain amide of R5 of A β , an interaction not observed in the PFA1-A β (1–8) complex. Residues D7 and S8 of the A β (1–8) peptide

show the greatest main-chain conformational difference between the PFA1-A β (1–8) and the WO1Fab-A β (1–16) structure. One reason for this can be attributed to the previously mentioned chemical and conformational difference at CDRH3 between the two structures: we observe D7 of A β (1–8) in the WO1 complex occupying a sterically overlapping position with the imidazole side chain of HC H97 of PFA1 (Figure 3b,c). For these reasons, the Fab contact surface of A β (7–8) differs between the two complexes. In the WO2-Fab-A β (1–16) complex, residues 7 and 8 contact HC Y100B; in PFA1, they interact instead with HC T98 and HC E100C. The E100C does not have a spatially overlapping residue in PFA1 due to the conformational differences in loop CDRH3 between the two mAbs. Other substitutions between the PFA1 and WO2 include HC S58 \rightarrow R and LC H93 \rightarrow L.

The EFRH motif of the A β N-terminus (comprising of residues 3–6) is the dominant binding epitope for PFA1 and for the recently reported IgG, WO2. The examination of the Ror2 (518–525)-PFA1 complex reveals that the Fab's enhanced affinity for Ror2(518–525) when compared to A β is likely due to an increased number of electrostatic interactions as well as more favorable intramolecular interactions, possibly stabilizing the peptide conformation. Although the Ror2(518–525) sequence is found in the (predicted) cytosolic portion of the protein, Tampellini et al. have shown evidence for anti-A β mAb internalization(41, 42). Hence, its enhanced affinity for PFA1 does strikingly illustrate the risk of cross-reactivity that stems from the short EFRH epitope. Based on our four PFA1-peptide complex structures (Table 1), the greatest variation in the peptide conformations occurs at the N-terminus (positions 2 and 3) and at position 8 at the C-terminus, and less so at position 7 (Figure 2C). Structurally, the FRH main-chain and side-chain conformations are extremely conserved. Hence, to ensure binding to N-terminal variants of A β such as pyro-Glu3-A β while improving the specificity of PFA1-derived mAbs toward the N-terminus of A β , mutations to PFA1 will have to concentrate on optimizing interactions at subsites 7 and beyond. Two such possibilities that we have modeled are to mutate HC 98 to K, or to mutate HC 100 (Kabat numbering) from V to R or K, introducing charge complementarity to enhance binding to D7 of A β . To enhance affinity to the N-terminus of the pyro-Glu3-A β peptide, mutating mAb residue LC S27E \rightarrow Q would put that side chain in a position to form a hydrogen bond with the lactam carbonyl oxygen atom of peptide residue pyro-Glu3. While these mutations might not help to distinguish A β from Ror2 as such (Ror2 is E7 at that position), they could nonetheless address a general problem with most anti-A β mAbs: the short length of immunodominant EFRH binding epitope.

ACKNOWLEDGMENT

We thank the members of the BMC beamline at BioCARS at APS for assistance with data collection. We also thank Dr. Brad Bennett and Jay Prendergast for proofreading the manuscript. Use of the Advanced Photon Source was supported by the U.S. Department of Energy, Basic Energy Sciences, Office of Science, under Contract W-31-109-Eng-38. Use of the BioCARS Sector 14 was supported by the National Institutes of Health, National Center for Research Resources, under Grant RR007707.

SUPPORTING INFORMATION AVAILABLE

One table containing hydrogen-bonding data for the PFA1-pyro-Glu3-A β and PFA1-Ror2(518–525) structures. This

material is available free of charge via the Internet at <http://pubs.acs.org>.

REFERENCES

- Haass, C., Koo, E. H., Mellon, A., Hung, A. Y., and Selkoe, D. J. (1992) Targeting of cell-surface beta-amyloid precursor protein to lysosomes: alternative processing into amyloid-bearing fragments. *Nature* 357, 500–503.
- Schenk, D., Barbour, R., Dunn, W., Gordon, G., Grajeda, H., Guido, T., Hu, K., Huang, J., Johnson-Wood, K., Khan, K., Kholodenko, D., Lee, M., Liao, Z., Lieberburg, I., Motter, R., Mutter, L., Soriano, F., Shopp, G., Vasquez, N., Vandeventer, C., Walker, S., Wogulis, M., Yednock, T., Games, D., and Seubert, P. (1999) Immunization with amyloid-beta attenuates Alzheimer-disease-like pathology in the PDAPP mouse. *Nature* 400, 173–177.
- Bard, F., Cannon, C., Barbour, R., Burke, R. L., Games, D., Grajeda, H., Guido, T., Hu, K., Huang, J., Johnson-Wood, K., Khan, K., Kholodenko, D., Lee, M., Lieberburg, I., Motter, R., Nguyen, M., Soriano, F., Vasquez, N., Weiss, K., Welch, B., Seubert, P., Schenk, D., and Yednock, T. (2000) Peripherally administered antibodies against amyloid beta-peptide enter the central nervous system and reduce pathology in a mouse model of Alzheimer disease. *Nat. Med.* 6, 916–919.
- Solomon, B., Koppel, R., Frankel, D., and Hanan-Aharon, E. (1997) Disaggregation of Alzheimer beta-amyloid by site-directed mAb. *Proc. Natl. Acad. Sci. U.S.A.* 94, 4109–4112.
- DeMattos, R. B., Bales, K. R., Cummins, D. J., Dodart, J. C., Paul, S. M., and Holtzman, D. M. (2001) Peripheral anti-A beta antibody alters CNS and plasma A beta clearance and decreases brain A beta burden in a mouse model of Alzheimer's disease. *Proc. Natl. Acad. Sci. U.S.A.* 98, 8850–8855.
- Gardberg, A. S., Dice, L. T., Ou, S., Rich, R. L., Helmbrecht, E., Ko, J., Wetzel, R., Myska, D. G., Patterson, P. H., and Dealwis, C. (2007) Molecular basis for passive immunotherapy of Alzheimer's disease. *Proc. Natl. Acad. Sci. U.S.A.* 104, 15659–15664.
- Kuo, Y. M., Emmerling, M. R., Woods, A. S., Cotter, R. J., and Roher, A. E. (1997) Isolation, chemical characterization, and quantitation of A beta 3-pyroglutamy peptide from neuritic plaques and vascular amyloid deposits. *Biochem. Biophys. Res. Commun.* 237, 188–191.
- Roher, A. E., Lowenson, J. D., Clarke, S., Wolkow, C., Wang, R., Cotter, R. J., Reardon, I. M., Zurcher-Neely, H. A., Heinrichson, R. L., and Ball, M. J.; et al. (1993) Structural alterations in the peptide backbone of beta-amyloid core protein may account for its deposition and stability in Alzheimer's disease. *J. Biol. Chem.* 268, 3072–3083.
- He, W., and Barrow, C. J. (1999) The A beta 3-pyroglutamy and 11-pyroglutamy peptides found in senile plaque have greater beta-sheet forming and aggregation propensities in vitro than full-length A beta. *Biochemistry* 38, 10871–10877.
- Sergeant, N., Bombois, S., Ghestem, A., Drobecq, H., Kostanjevecki, V., Missiaen, C., Watzel, A., David, J. P., Vanmechelen, E., Sergheraert, C., and Delacourte, A. (2003) Truncated beta-amyloid peptide species in pre-clinical Alzheimer's disease as new targets for the vaccination approach. *J. Neurochem.* 85, 1581–1591.
- Saido, T. C., Iwatsubo, T., Mann, D. M., Shimada, H., Ihara, Y., and Kawashima, S. (1995) Dominant and differential deposition of distinct beta-amyloid peptide species, A beta N3(pE), in senile plaques. *Neuron* 14, 457–466.
- Russo, C., Schettini, G., Saido, T. C., Hulette, C., Lippa, C., Lannfelt, L., Ghetti, B., Gambetti, P., Tabaton, M., and Teller, J. K. (2000) Presenilin-1 mutations in Alzheimer's disease. *Nature* 405, 531–532.
- Russo, C., Salis, S., Dolcini, V., Venezia, V., Song, X. H., Teller, J. K., and Schettini, G. (2001) Amino-terminal modification and tyrosine phosphorylation of [corrected] carboxy-terminal fragments of the amyloid precursor protein in Alzheimer's disease and Down's syndrome brain. *Neurobiol. Dis.* 8, 173–180.
- Hosoda, R., Saido, T. C., Otvos, L. Jr., Arai, T., Mann, D. M., Lee, V. M., Trojanowski, J. Q., and Iwatsubo, T. (1998) Quantification of modified amyloid beta peptides in Alzheimer disease and Down syndrome brains. *J. Neuropathol. Exp. Neurol.* 57, 1089–1095.
- Saido, T. C., Yamao-Harigaya, W., Iwatsubo, T., and Kawashima, S. (1996) Amino- and carboxyl-terminal heterogeneity of beta-amyloid peptides deposited in human brain. *Neurosci. Lett.* 215, 173–176.
- Kraus, M., Bienert, M., and Krause, E. (2003) Hydrogen exchange studies on Alzheimer's amyloid-beta peptides by mass spectrometry using matrix-assisted laser desorption/ionization and electrospray ionization. *Rapid Commun. Mass Spectrom.* 17, 222–228.
- Piccini, A., Russo, C., Gliozzi, A., Relini, A., Vitali, A., Borghi, R., Giliberto, L., Armirotti, A., D'Arrigo, C., Bachi, A., Cattaneo, A., Canale, C., Torrasa, S., Saido, T. C., Markesbery, W., Gambetti, P., and Tabaton, M. (2005) beta-amyloid is different in cerebrospinal fluid and in Alzheimer disease. *J. Biol. Chem.* 280, 34186–34192.
- Vanderstichele, H., De Meyer, G., Andreasen, N., Kostanjevecki, V., Wallin, A., Olsson, A., Blennow, K., and Vanmechelen, E. (2005) Amino-truncated beta-amyloid42 peptides in cerebrospinal fluid and prediction of progression of mild cognitive impairment. *Clin. Chem.* 51, 1650–1660.
- Schilling, S., Lauber, T., Schaupp, M., Manhart, S., Scheel, E., Bohm, G., and Demuth, H. U. (2006) On the seeding and oligomerization of pGlu-amyloid peptides (in vitro). *Biochemistry* 45, 12393–12399.
- Tekirian, T. L., Yang, A. Y., Glabe, C., and Geddes, J. W. (1999) Toxicity of pyroglutaminated amyloid beta-peptides 3(pE)-40 and -42 is similar to that of A beta1-40 and -42. *J. Neurochem.* 73, 1584–1589.
- Geddes, J. W., Tekirian, T. L., and Mattson, M. P. (1999) N-terminus truncated beta-amyloid peptides and C-terminus truncated secreted forms of amyloid precursor protein: distinct roles in the pathogenesis of Alzheimer's disease. *Neurobiol. Aging* 20, 75–79 (discussion 87).
- Schilling, S., Zeitschel, U., Hoffmann, T., Heiser, U., Francke, M., Kehlen, A., Holzer, M., Hutter-Paier, B., Prokesh, M., Windisch, M., Jagla, W., Schlenzig, D., Lindner, C., Rudolph, T., Reuter, G., Cynis, H., Montag, D., Demuth, H. U., and Rossner, S. (2008) Glutaminyl cyclase inhibition attenuates pyroglutamate A beta and Alzheimer's disease-like pathology. *Nat. Med.* 14, 1106–1111.
- Liu, Y., Ross, J. F., Bodine, P. V., and Billiard, J. (2007) Homodimerization of Ror2 tyrosine kinase receptor induces 14-3-3{beta} phosphorylation and promotes osteoblast differentiation and bone formation. *Mol. Endocrinol.* 21(12), 305–61.
- Burette, A., Wyszynski, M., Valtschanoff, J. G., Sheng, M., and Weinberg, R. J. (1999) Characterization of glutamate receptor interacting protein-immunopositive neurons in cerebellum and cerebral cortex of the albino rat. *J. Comp. Neurol.* 411, 601–612.
- Wyszynski, M., Valtschanoff, J. G., Naisbitt, S., Dunah, A. W., Kim, E., Standaert, D. G., Weinberg, R., and Sheng, M. (1999) Association of AMPA receptors with a subset of glutamate receptor-interacting protein in vivo. *J. Neurosci.* 19, 6528–6537.
- Otwinowski, Z., and Minor, W. (1997) Processing of X-ray diffraction data collected in oscillation mode, in *Macromolecular Crystallography, Part A* (Carter, C. W., and Sweet, R. M., Eds.) pp 307–326, Academic Press, New York.
- Zhu, X., Heine, A., Monnat, F., Houk, K. N., Janda, K. D., and Wilson, I. A. (2003) Structural basis for antibody catalysis of a cationic cyclization reaction. *J. Mol. Biol.* 329, 69–83.
- Emsley, P., and Cowtan, K. (2004) Coot: model-building tools for molecular graphics. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* 60, 2126–2132.
- Murshudov, G. N., Vagin, A. A., and Dodson, E. J. (1997) *Acta Crystallogr., Sect. D: Biol. Crystallogr.* 53, 240–255.
- Vaguine, A. A., Richelle, J., and Wodak, S. J. (1999) SFCHECK: a unified set of procedures for evaluating the quality of macromolecular structure-factor data and their agreement with the atomic model. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* 55, 191–205.
- Laskowski, R. A., MacArthur, M. W., Moss, D. S., and Thornton, J. M. (1993) PROCHECK: a program to check the stereochemical quality of protein structures. *J. Appl. Crystallogr.* 26, 283–291.
- Davis, I. W., Murray, L. W., Richardson, J. S., and Richardson, D. C. (2004) MOLPROBITY: structure validation and all-atom contact analysis for nucleic acids and their complexes. *Nucleic Acids Res.* 32, W615–619.
- Collaborative Computational Project N. (1994) The CCP4 suite: programs for protein crystallography. *Acta Crystallogr. D* 50, 760–763.
- Lee, B., and Richards, F. M. (1971) The interpretation of protein structures: estimation of static accessibility. *J. Mol. Biol.* 55, 379–400.
- Murphy, K. P., Xie, D., Garcia, K. C., Amzel, L. M., and Freire, E. (1993) Structural energetics of peptide recognition: angiotensin II/antibody binding. *Proteins* 15, 113–120.
- Blake, C. C., Serpell, L. C., Sunde, M., Sandgren, O., and Lundgren, E. (1996) A molecular model of the amyloid fibril. *CIBA Found. Symp.* 199, 6–15 (discussion 15–21, 40–16).
- Petkova, A. T., Leapman, R. D., Guo, Z., Yau, W. M., Mattson, M. P., and Tycko, R. (2005) Self-propagating, molecular-level polymorphism in Alzheimer's beta-amyloid fibrils. *Science* 307, 262–265.

38. Sciarretta, K. L., Gordon, D. J., Petkova, A. T., Tycko, R., and Meredith, S. C. (2005) Abeta40-lactam(D23/K28) models a conformation highly favorable for nucleation of amyloid. *Biochemistry* 44, 6003–6014.
39. Hoyer, W., Gronwall, C., Jonsson, A., Stahl, S., and Hard, T. (2008) Stabilization of a beta-hairpin in monomeric Alzheimer's amyloid-beta peptide inhibits amyloid formation. *Proc. Natl. Acad. Sci. U.S.A.* 105, 5099–5104.
40. Miles, L. A., Wun, K. S., Crespi, G. A., Fodero-Tavoletti, M. T., Galatis, D., Bagley, C. J., Beyreuther, K., Masters, C. L., Cappai, R., McKinstry, W. J., Barnham, K. J., and Parker, M. W. (2008) Amyloid-beta-anti-amyloid-beta complex structure reveals an extended conformation in the immunodominant B-cell epitope. *J. Mol. Biol.* 377, 181–192.
41. Tampellini, D., Magrane, J., Takahashi, R. H., Li, F., Lin, M. T., Almeida, C. G., and Gouras, G. K. (2007) Internalized antibodies to the Abeta domain of APP reduce neuronal Abeta and protect against synaptic alterations. *J. Biol. Chem.* 282, 18895–18906.
42. Arbel, M., and Solomon, B. (2007) Immunotherapy for Alzheimer's disease: attacking amyloid-beta from the inside. *Trends Immunol.* 28, 511–513.