


Effects of Mg^{2+} on the Free Energy Landscape for Folding a Purine Riboswitch RNA

Desirae Leipply[‡] and David E. Draper^{*,§}

[‡]Program in Molecular Biophysics and Department of Biophysics, and [§]Department of Biophysics and Department of Chemistry, Johns Hopkins University Baltimore, Maryland 21218, United States

S Supporting Information

ABSTRACT: There are potentially several ways Mg^{2+} might promote formation of an RNA tertiary structure: by causing a general “collapse” of the unfolded ensemble to more compact conformations, by favoring a reorganization of structure within a domain to a form with specific tertiary contacts, and by enhancing cooperative linkages between different sets of tertiary contacts. To distinguish these different modes of action, we have studied Mg^{2+} interactions with the adenine riboswitch, in which a set of tertiary interactions that forms around a purine-binding pocket is thermodynamically linked to the tertiary “docking” of two hairpin loops in another part of the molecule. Each of four RNA forms with different extents of tertiary structure were characterized by small-angle X-ray scattering. The free energy of interconversion between different conformations in the absence of Mg^{2+} and the free energy of Mg^{2+} interaction with each form have been estimated, yielding a complete picture of the folding energy landscape as a function of Mg^{2+} concentration. At 1 mM Mg^{2+} (50 mM K^+), the overall free energy of stabilization by Mg^{2+} is large, -9.8 kcal/mol, and about equally divided between its effect on RNA collapse to a partially folded structure and on organization of the binding pocket. A strong cooperative linkage between the two sets of tertiary contacts is intrinsic to the RNA. This quantitation of the effects of Mg^{2+} on an RNA with two distinct sets of tertiary interactions suggests ways that Mg^{2+} may work to stabilize larger and more complex RNA structures.



The stabilization of RNA tertiary structure by low concentrations of Mg^{2+} ions was first described in early studies of tRNA folding.^{1–4} Much attention has been focused on those factors which promote strong Mg^{2+} interactions with native structures: a high charge density causes accumulation of Mg^{2+} in an “ion atmosphere”,⁵ and the tertiary fold of an RNA may create special environments with unusually strong ion interactions.^{6,7} To understand how Mg^{2+} stabilizes RNA structures, it is also important to consider partially unfolded forms of the RNA, which also interact strongly with Mg^{2+} .^{8–10} Particularly intriguing is the ability of Mg^{2+} to cause the “collapse” of RNAs from extended forms to compact intermediates that lack fully formed tertiary contacts.^{11–15}

A relatively simple RNA with which to explore the varied effects of Mg^{2+} on the formation of both a native RNA structure and its folding intermediates is the aptamer domain of riboswitches that sense purine ligands.^{16–18} These RNAs have a simple topology with two distinct sets of tertiary hydrogen bonds (Figure 1A): the two hairpin loops dock with each other, and a binding pocket becomes organized around the bound purine ligand. From FRET and NMR studies of several purine riboswitch RNAs, it is known that Mg^{2+} favors the docked conformation in the absence of ligand.^{19–21} Furthermore, docking of the loops strengthens the ligand affinity of the binding pocket.²² The purine riboswitch is thus a particularly simple case in which Mg^{2+} -dependent formation of a compact but not fully native

intermediate is coupled to formation of a core functional tertiary structure.

In this work, we ask how Mg^{2+} alters the folding landscape of an adenine-specific riboswitch (A-riboswitch, Figure 1A) by measuring the free energy of Mg^{2+} interaction with the native structure and each of several partially folded forms that are potentially part of the equilibrium folding pathway. We use a titration method that monitors the thermodynamic activity of Mg^{2+} in the presence of RNA and yields the extent of charge neutralization by Mg^{2+} (the “excess” Mg^{2+}) for a particular RNA conformation. An advantage of this approach for our purposes is that the effect of Mg^{2+} on the free energy of individual RNA forms at different positions in the folding landscape (e.g., extended, collapsed, or native conformations) can be examined over a wide range of Mg^{2+} concentrations.⁸ Together with structural characterization of the RNA forms by small-angle X-ray scattering (SAXS) and hydroxyl radical probing, we have developed a quantitative picture of the folding transitions taking place in the A-riboswitch in the absence of Mg^{2+} (50 mM K^+), and the ways these folding equilibria shift as Mg^{2+} is titrated. Mg^{2+} strongly stabilizes both the docked and binding pocket tertiary structures, and because of a large cooperative linkage between

Received: December 7, 2010

Revised: February 9, 2011

Published: March 01, 2011




Figure 2. Titration of wild type A-riboswitch (open symbols) or C60G RNA (gray circles) with MgCl_2 in buffer containing 50 mM K^+ (20 °C), monitored by absorbance at 260 nm. Mg^{2+} -dependent baselines have been subtracted from the data sets (see Materials and Methods). Wild type A-riboswitch was titrated in the presence of 82 μM adenine (open circles), or no ligand (open squares); the C60G RNA titration included 82 μM DAP (gray circles). Titration of the wild type RNA with ligand has been fit to the Hill equation (midpoint 29.0 μM Mg^{2+} , $n = 1.58$). The other two curves are optimum fits of a Mg^{2+} -dependent two-state equilibrium between E and I_{dock} forms (wild type RNA) or E and I_{LBP} forms (C60G RNA) (see Results for further explanation).

DAP,¹⁰ we found that DAP binds 34-fold more tightly over a wide range of Mg^{2+} concentrations.] Comparing titrations of the wild type RNA (in the presence of 82 μM adenine) and C60G RNA (82 μM DAP), the midpoint of the C60G RNA titration is shifted to higher Mg^{2+} concentration and the hypochromic change is less than half (Figure 2). The extent of folding of the two RNAs can be directly compared at 0.1 mM Mg^{2+} (Figure 2); taking into account the 34 fold tighter affinity of DAP over adenine, C60G RNA has ~ 570 fold weaker affinity for ligand than does wild type RNA at this Mg^{2+} concentration. Similar experiments with G38C RNA failed to observe any reproducible folding transition at Mg^{2+} concentrations up to 1 mM. A trend of increasing stability for G38C, C60G, and wild type RNAs was found in UV melting experiments in the presence of 1 mM Mg^{2+} and ligand, and a very weak affinity of G38C RNA for DAP ($\sim 26 \text{ mM}^{-1}$) was estimated from melting experiments carried out in 10 mM Mg^{2+} (Figure S1 of Supporting Information). Overall, these titration and melting experiments imply that the mutation G38C severely weakens the docked loop conformation while C60G has a substantial but less drastic effect, consistent with ligand affinities reported for analogous mutations in other purine riboswitch sequences.^{19,28} In subsequent sections, we present evidence that C60G RNA bound to ligand is in the desired I_{LBP} form, with native-like tertiary structure formed around the ligand but undocked hairpin loops.

As previously noted in melting experiments with the Figure 1A A-riboswitch sequence,¹⁰ Mg^{2+} may induce tertiary structure formation in the absence of ligand. The Mg^{2+} -dependence of this folding is seen by Mg^{2+} titration of the wild type RNA (Figure 2). We argue below that the RNA conformation adopted at 1 mM Mg^{2+} is the desired I_{dock} form.

RNA Dimensions from Small-Angle X-ray Scattering. We used SAXS to monitor docking of the hairpin loops by changes in the overall solution dimensions of the RNAs. In these experiments, wild-type A-riboswitch in the presence of both ligand and Mg^{2+} has the same radius of gyration (R_g) as calculated from the

crystal structure (PDB 1Y26, 20.3 Å). In the absence of either Mg^{2+} or ligand (DAP), the RNA adopts a much more extended conformation ($R_g = 24.9$ Å). Titration with Mg^{2+} in the absence of ligand gradually brings R_g close to that of the native form (Figure 3A). In contrast to the wild type RNA, C60G RNA dimensions in the absence of ligand are much less sensitive to the presence of Mg^{2+} (Figure 3A); even conditions that saturate C60G RNA with ligand (1 mM Mg^{2+} , 250 μM DAP) leave the RNA in an extended conformation (Figure 3A).

The distance distribution function derived from the scattering data, $P(r)$, contains a more detailed description of macromolecular conformation than reported by the single parameter R_g . Comparing $P(r)$ for wild type and C60G RNAs, it is clear that the extended forms found in the absence of Mg^{2+} and ligand are not identical; the wild type A-riboswitch has a slightly more compact form than C60G RNA (Figure 3B). Addition of 1 mM Mg^{2+} has very little effect on the $P(r)$ function of C60G RNA in the absence of ligand (Figure 3B,C), unlike the strong effect of Mg^{2+} on the wild-type RNA.

Another point of consequence is that $P(r)$ curves for wild type RNA in 1 mM Mg^{2+} are nearly identical whether or not ligand is bound (Figure 3C). Thus, any organization of the binding pocket that accompanies ligand binding does not affect the dimensions of the RNA in a way that is detectable by SAXS; the experiment is primarily sensitive to the disposition of the two loops.

We attempted to calculate low resolution models of the wild type and variant A-riboswitch RNAs that satisfy the constraints of the various scattering data sets (see Materials and Methods). The shape found for the wild type RNA in the presence of Mg^{2+} and ligand corresponds to the overall dimensions of the crystal structure (Figure 3D). Similar shapes were calculated for wild type RNA in 1 or 10 mM Mg^{2+} without ligand (not shown). Reconstructions of C60G and G38C RNAs in the absence of Mg^{2+} gave similar extended structures suggestive of undocked hairpin loops (Figure 3E). It is ambiguous which of the three arms of this T-shaped structure might be the 5'-3' terminal helix. A proposed model of ligand-free purine riboswitch based on FRET measurements²¹ has the 3' 20 nucleotides of helices P1 and P3 (A64–C83) remain coaxially stacked as they are in the native crystal structure,¹⁸ and the P2 helix protruding at right angles. However, the P1–P3 helices span ~ 70 Å, while the extended state RNA reconstruction has a long dimension of ~ 100 Å; it seems unlikely that the sequence in question could be stretched enough to cover this distance. Alternatively, the 29 nucleotides of backbone extending between the P2 and P3 hairpins (A35–U63) could form a stacked structure of about the right dimensions. Although the identity of the three helical segments in the E state structure is not crucial for the present work, the nature of the three helix junction structure that enforces the extended conformation is an interesting question for further investigation.

Calculations with scattering data collected on either the wild type A-riboswitch without ligand or Mg^{2+} , or C60G RNA saturated with ligand at 1 mM Mg^{2+} , did not yield unique structures, a possible indication that an ensemble of disparate structures is present in solution.²⁹

Hydroxyl Radical Probing of Tertiary Structure in C60G RNA. To further delineate conformational differences between wild type and C60G RNAs, we measured the reactivity of the two sequences toward hydroxyl radical, a probe sensitive to ribose exposure to solvent.³⁰ Loop nucleotides G37 and G38 were protected in the native structure of the wild type RNA (relative to




Figure 3. SAXS studies of wild type and variant A-riboswitch RNAs in standard buffer; ligand is 250 μ M DAP, where present. Samples were equilibrated with the indicated bulk Mg^{2+} concentrations (C_{2+}). (A) R_g as a function of bulk Mg^{2+} concentration for wild type RNA with (blue) or without (red) ligand, or C60G RNA with (green) or without (orange) ligand. The solid red line is the best fit of a Mg^{2+} -dependent two-state equilibrium between E and I_{dock} forms of the RNA (see text). (B) Distance distribution function $P(r)$ for RNAs in the absence of Mg^{2+} : wild type RNA with (blue) or without (red) ligand, or C60G RNA without ligand (orange). (C) $P(r)$ for RNAs with 1 mM bulk Mg^{2+} concentration: wild type RNA with (blue) or without (red) ligand, and C60G RNA with (green) or without (orange) ligand. (D) Reconstruction from SAXS data of the wild type A-riboswitch bound to ligand in the presence of Mg^{2+} . The RNA backbone from the corresponding crystal structure (1Y26) has been aligned with the calculated density; the bound adenine ligand is shown as a space-filling model. The two views differ by 90° rotation. (E) Reconstruction from SAXS data of the C60G (green) and G38C (yellow) RNAs in the absence of ligand or Mg^{2+} . The two structures are superimposed to show their similar overall shape.

unfolded RNA in the absence of both Mg^{2+} and ligand), but neither residue changed reactivity in response to high Mg^{2+} concentration and saturating ligand concentration in the C60G RNA (Figure 4). Six different residues within the binding pocket became protected in the wild-type RNA in the presence of ligand; the same residues were protected to about the same extent in C60G RNA bound to ligand (Figure 4). Disruption of the docked loops apparently does not affect the structure of the RNA around the bound ligand, at least at a level detectable by hydroxyl radical reactivity.

Direct Measurement of Excess Mg^{2+} . The excess Mg^{2+} accumulated by different conformations of the wild type and mutant RNAs was measured using a fluorescent chelator that monitors the effect of an RNA on the thermodynamic activity of Mg^{2+} ions.^{8,23} The results are reported in terms of the interaction parameter Γ_{2+} , which is the excess number of Mg^{2+} ions present in compensation for the RNA negative charge at a given level of bulk ion concentration C_{2+} . It is essentially the same as the ions accumulated in a dialysis experiment in response to the presence of an RNA, where C_{2+} is the ion concentration in the solution external to the RNA dialysis solution.³¹ Similar interaction parameters describe the excess monovalent ion (Γ_{+}) and excluded anion (Γ_{-}), though they are not measured here.

The wild type RNA was titrated with Mg^{2+} either in the absence of ligand or in the presence of a high enough concentration of DAP that the RNA was in the native structure without Mg^{2+} present¹⁰ (Figure 5A). As expected, the more compact

native form accumulated more Mg^{2+} than the unfolded form of the RNA. The difference between these two curves, $\Delta\Gamma_{2+}$, is the uptake of Mg^{2+} upon folding. Excess Mg^{2+} for the native RNA structure is 15.5 ± 0.2 ions/RNA at 1 mM Mg^{2+} (Table 2, Supporting Information), which corresponds to 43% of the RNA phosphate charge neutralized by Mg^{2+} . The data shown here extend our previous measurements of Γ_{2+} with this RNA to much higher Mg^{2+} concentrations.¹⁰

Because the C60G RNA tertiary structure does not form in the absence of Mg^{2+} , even at very high ligand concentrations, the Γ_{2+} curves obtained with this RNA in the presence or absence of DAP were initially identical to each other (Figure 5B) and to the titration curve of unfolded wild type RNA (Figure 5A). The two C60G RNA curves diverge significantly when C_{2+} exceeds ~ 50 μ M Mg^{2+} ; based on UV-monitored titrations (Figure 2 and data not shown), the midpoint of ligand binding is expected at ~ 90 μ M Mg^{2+} at the ligand concentration of the experiment. Notably, the ligand-bound form of C60G RNA, which is essentially the only form present above $C_{2+} \approx 0.6$ mM, did not reach the same level of Γ_{2+} as the wild type RNA (at 1 mM Mg^{2+} , Γ_{2+} was 13.4 ± 0.2 ions/RNA for C60G RNA). This observation is consistent with SAXS measurements that showed a more extended structure for ligand-bound C60G RNA (Figure 3A). In the absence of ligand, the wild type and C60G RNA Γ_{2+} curves matched at low Mg^{2+} concentrations but diverged above ~ 0.1 mM (Figure 5A), again implying a different, probably more extended structure for the mutant.




Figure 4. Hydroxyl radical footprinting results for the wild-type and C60G variant A-riboswitch (data analysis was as described in Materials and Methods). Experiments were performed in standard buffer at 0 °C, with the indicated amount of Mg^{2+} and 250 μM DAP present in all but the unfolded control samples. The difference in normalized reactivity between the control unfolded form (RNA in the absence of Mg^{2+} and DAP) and the sample RNA (with DAP and Mg^{2+}) is plotted; a positive number represents reduced reactivity in the sample RNA. Open column, wild type RNA with 5 mM Mg^{2+} ; gray – black columns, C60G RNA with increasing Mg^{2+} concentrations of 0.1, 0.2, 0.5, 1.0, and 5.0 mM. Residue numbering is the same as Figure 1A. Relative reactivities of nucleotides 32–75 are in Supporting Information (Figure S2).

An important point is that the uptake of Mg^{2+} upon ligand binding ($\Delta\Gamma_{2+}$, as calculated from Figure 5 data) is about the same for C60G RNA (2.6 ± 0.3 ions/RNA) as wild type (3.2 ± 0.3 ions) when evaluated at 1 mM Mg^{2+} (Table S2, Supporting Information). However, $\Delta\Gamma_{2+}$ only appears insensitive to the mutation because Mg^{2+} interactions with both the ligand-bound and unbound states of the RNA have been altered to comparable extents, presumably because C60G RNA loops remain undocked whether or not ligand is bound.

Free Energies of Mg^{2+} –RNA Interactions. If no conformational changes take place in an RNA during its titration with Mg^{2+} , the Γ_{2+} curve may be integrated to give the total free energy of Mg^{2+} –RNA interaction ($\Delta G_{\text{RNA-}2+}$) as a function of bulk Mg^{2+} concentration.^{8,31} On the basis of scattering measurements (Figure 3A), neither the native, ligand-bound structure (labeled N in Figure 5A) nor the ligand-free form of C60G RNA (E in Figure 5B) is affected by the presence or absence of Mg^{2+} ; $\Delta G_{\text{N-}2+}$ and $\Delta G_{\text{E-}2+}$ have been calculated from these respective Γ_{2+} curves (Figure 5C). The other two forms of the RNA observed at 1 mM Mg^{2+} , I_{dock} for wild type RNA in the absence of ligand and I_{LBP} for C60G RNA saturated with ligand (Figure 5A,B), were achieved only when Mg^{2+} was present. We calculated $\Delta G_{\text{RNA-}2+}$ for these two RNA forms by noting that the $\Delta G_{\text{N-}2+}$ and $\Delta G_{\text{E-}2+}$ curves differ by a uniform factor (1.95 ± 0.19 fold) independent of Mg^{2+} concentration (Figure S3, Supporting Information). We assume that corresponding curves for the two intermediate forms will also have similar shapes and differ from $\Delta G_{\text{N-}2+}$ or $\Delta G_{\text{E-}2+}$ by a constant factor over the entire range of Mg^{2+} concentrations. The free energies $\Delta G_{\text{Idock-}2+}$ and $\Delta G_{\text{ILBP-}2+}$ (shown in Figure 5C) have therefore been scaled from an average of the $\Delta G_{\text{N-}2+}$ and $\Delta G_{\text{E-}2+}$ curves; the interaction free energies found at 1 mM bulk Mg^{2+} (Table S1, Supporting Information) were used to calculate the scaling factor.




Figure 5. Excess Mg^{2+} (Γ_{2+}) as a function of bulk Mg^{2+} concentration (C_{2+}) for wild type and C60G RNAs in standard conditions (50 mM K^+ , 20 °C), with and without the ligand DAP present (250 μM). Six or seven independent data sets were merged to obtain each titration curve; errors are shown in the plots as lighter color bars (see Materials and Methods). (A) Wild type RNA in the presence (blue) or absence (magenta) of ligand. C60G RNA in the absence of ligand is shown for comparison (orange). (B) C60G RNA in the presence (green) or absence (orange) of ligand. Wild type RNA in the presence of ligand is shown for comparison (blue). (C) Free energies of Mg^{2+} –RNA interaction ($\Delta G_{\text{RNA-}2+}$) calculated from Γ_{2+} curves in panels A and B (see Materials and Methods). Blue, native RNA (N); orange, C60G RNA in the absence of ligand (E); green, the I_{dock} conformation, extrapolated from wild type RNA at 1 mM Mg^{2+} ; magenta, I_{LBP} conformation, extrapolated from C60G RNA bound to ligand at 1 mM Mg^{2+} .

Intrinsic Free Energies of Formation of Intermediates from Extended State RNA. To fully evaluate the effects of Mg^{2+} on the A-riboswitch folding landscape, it would be useful to know the initial free energy differences between the four RNA forms (Figure 1B) before Mg^{2+} is added. Equilibria between the extended form of the RNA and the two intermediates, I_{dock} and I_{LBP} , strongly favor the extended form in the absence of Mg^{2+} and are therefore difficult to measure directly. However, using SAXS or UV absorbance, we have observed these equilibria shift toward the intermediate form as Mg^{2+} is titrated (Figures 2 and 3A). The Mg^{2+} -dependence of the observed equilibrium between


an extended (E) and intermediate (I) form is derived as

$$K_{E-I} = K_{E-I}^{\text{int}} \exp[(\Delta G_{E-2+} - \Delta G_{I-2+})/RT] \quad (1)$$

where the ΔG values are the known free energies of Mg^{2+} –RNA interaction with either RNA form (Figure 5C) and R is the gas constant. K_{E-I} is the Mg^{2+} -dependent, observed equilibrium constant for interconverting the extended and intermediate forms. K_{E-I}^{int} is the intrinsic equilibrium constant in the absence of Mg^{2+} , and the only unknown in the equation. A two-state equilibrium between E and I is assumed; potential limitations imposed by the assumption are considered below. In deriving the equation (see Supporting Information), the reference state is defined as our standard experimental conditions (50 mM K^+) in the absence of Mg^{2+} . The free energy associated with the conformational equilibrium in the absence of Mg^{2+} , K_{E-I}^{int} , is therefore a standard state free energy, ΔG_{E-I}° . The corresponding free energies in the presence of Mg^{2+} are denoted by ΔG_{E-I} , which varies with the Mg^{2+} concentration.

For the docking reaction, $E \rightleftharpoons I_{\text{dock}}$, we compare eq 1 with the Mg^{2+} -dependences of UV absorption (Figure 2) and R_g (Figure 3A) measurements with wild type RNA in the absence of ligand. When $K_{E-I_{\text{dock}}}^{\text{int}}$ was systematically varied, the best fit values were 0.12 ($\Delta G_{E-I_{\text{dock}}}^{\circ} = 1.3$ kcal/mol) or 0.065 ($\Delta G_{E-I_{\text{dock}}}^{\circ} = 1.6$ kcal/mol) for the UV and SAXS data, respectively. The Mg^{2+} -dependence of ligand binding to C60G RNA (Figure 2) corresponds to the equilibrium $E \rightleftharpoons I_{\text{LBP}}$; we obtain the intrinsic equilibrium constant $K_{E-I_{\text{LBP}}}^{\text{int}} = 110 \text{ M}^{-1}$ for the binding of DAP. Given that adenine binds the wild type RNA 34-fold more weakly than DAP, we estimate $\Delta G_{E-I_{\text{LBP}}}^{\circ} = -0.7$ kcal/mol as the standard state affinity of C60G RNA for adenine in the absence of Mg^{2+} .

There is reason to think that the docking reaction, $E \rightleftharpoons I_{\text{dock}}$, might not be strictly two-state: single-molecule FRET experiments with a similar A-riboswitch sequence as used here detected a population of molecules with hairpin loops in an intermediate conformation between docked and extended.¹⁹ We have therefore considered the folding pathway



in which the intermediate I' is assigned an R_g halfway between those of the E and I_{dock} forms, and various Mg^{2+} interaction energies between ΔG_{E-2+} and $\Delta G_{I_{\text{dock}}-2+}$. Although it is possible to find combinations of the intrinsic equilibrium constants for the $E \rightleftharpoons I'$ and $I' \rightleftharpoons I_{\text{dock}}$ reactions that give slightly better fits to the scattering data than the two-state model (Figure 3A), the product of the two equilibrium constants is always within 10% of $K_{E-I_{\text{LBP}}}^{\text{int}}$; the maximum population of the I' state in these simulations was less than 10%. Although intermediate states between the fully extended and docked RNA forms may well exist, we conclude that the overall free energy estimated for the $E \rightleftharpoons I_{\text{dock}}$ reaction is reasonable.

DISCUSSION

Γ_{2+} and R_g Define Four Forms of the A-Riboswitch RNA. To determine the ways Mg^{2+} might affect folding of the A-riboswitch, we used the complementary methods of X-ray scattering and excess Mg^{2+} ion (Γ_{2+}) measurements to characterize the A-riboswitch–ligand complex and its various partially unfolded forms. Four species have been studied, each with a distinct combination of size (R_g) and excess Mg^{2+} . We first summarize essential characteristics of the four RNA forms and their interconversions




Figure 6. Effects of Mg^{2+} on the A-riboswitch folding landscape. (A) Four states of the A-riboswitch, distinguished by their dimensions (R_g) and excess Mg^{2+} (Γ_{2+} , reported at 1 mM Mg^{2+}). See Discussion for definitions of the four states. Arrows connecting the RNA forms define a thermodynamic cycle with three free energies, all of which are potentially Mg^{2+} -dependent. The number of ions taken up ($\Delta\Gamma_{2+}$) in the reaction defined by each arrow is indicated. (B) Stabilities of each of the RNA forms from panel A, relative to the extended form (E) in its standard state (50 mM K^+ buffer without Mg^{2+}). Stabilities of the ligand-bound RNA forms (N and I_{LBP}) are calculated for an adenine concentration of 10 μM . Thick arrows, free energies of the two-state equilibria between E and each of the three other forms; cyan and blue bars, stabilization of each form by 0.1 or 1.0 mM bulk Mg^{2+} concentration, respectively. Calculations of the three ΔG values defined in panel A are illustrated for 1 mM Mg^{2+} (arrows labeled by ΔG). Error bars for the N and E state Mg^{2+} –RNA interaction free energies are based on the reproducibility of the Γ_{2+} data that were integrated to find $\Delta G_{\text{RNA-2+}}$ (Figure 5); errors for the other two forms include an additional estimated uncertainty from the method used to interpolate the free energies.

and in the following section consider the different ways Mg^{2+} stabilizes the RNA tertiary structure and enhances ligand binding affinity.

The form of the RNA with the highest value of Γ_{2+} is the native structure N, the ligand complex formed with the wild type RNA (Figure 6A). The wild type RNA in the absence of ligand adopts a docked-loop conformation at 1 mM Mg^{2+} , as anticipated from FRET and NMR studies of related purine riboswitch sequences.^{19–21} This form of the RNA, I_{dock} , has similar dimensions and shape as the native form (Figure 3C) but accumulates Mg^{2+} to a lesser extent than the native structure (Figure 6A); the smaller Γ_{2+} must be due to structural differences in the binding pocket when ligand is absent.

The C60G riboswitch variant maintains an extended structure with dimensions that are insensitive to Mg^{2+} in the absence of ligand (E in Figure 6A). The convergence of reconstructions on a unique structure with plausible dimensions (Figure 3E) argues that the RNA adopts a single conformation, within the resolution of SAXS experiments. C60G RNA can be saturated with ligand at 1 mM Mg^{2+} , but based on hydroxyl radical probing and scattering measurements (Figures 3C and 4), the loops remain undocked with an average angle between helices P2 and P3 that must be intermediate between the docked and fully extended structures (I_{LBP} in Figure 6A). The binding pocket, however, probably retains the same tertiary structure around the bound ligand as the native RNA (Figure 4).

The free energy differences between an extended E state and each of the other three forms in the absence of Mg^{2+} can be estimated, if two-state equilibria are assumed. (As described in Results, we use ΔG° to refer to our standard buffer conditions, 50 mM K^+ without Mg^{2+} . ΔG° for formation of the two ligand-bound states, N and I_{LBP} , includes the standard free energy of adenine binding.) Two of these ΔG° values were calculated by extrapolating the equilibria observed over a range of Mg^{2+} concentrations to zero Mg^{2+} (see Results). We can also estimate ΔG° for a third transition, $\text{E} \rightarrow \text{N}$. Upon melting of the wild type RNA in the presence of 11 μM DAP without Mg^{2+} , RNA unfolding from the native, ligand-bound conformation to an equilibrium mixture of E and I_{dock} states was observed with a T_m of 20.9 $^\circ\text{C}$.³¹ After correcting for the small amount of I_{dock} calculated to be present and for the 34-fold weaker binding affinity of adenine, ΔG° for $\text{E} \rightarrow \text{N}$ is estimated as -4.7 kcal/mol (20 $^\circ\text{C}$).

We combine our ΔG° estimates for interconversion between different RNA forms with the Mg^{2+} –RNA interaction energies of each form ($\Delta G_{\text{RNA-2+}}$) to construct a free energy landscape for A-riboswitch folding.⁴ The relative free energies of the four RNA forms are shown in Figure 6B; the free energies of the two ligand-bound species have been adjusted to reflect 10 μM adenine rather than the standard state 1 M, as an order-of-magnitude approximation of the in vivo adenine concentration.^{32,33} In the absence of Mg^{2+} , the extended E state is the most stable RNA conformation; as Mg^{2+} is added the relative stabilities change until the ligand-bound N state becomes by far the most stable conformation.

As a caution about applying the data in Figure 6B to physiological conditions, note that the free energies determined in 1 mM Mg^{2+} are probably larger than those attained in vivo. The K^+ activity in our standard 50 mM buffer is about 3-fold lower than the K^+ activity present in bacteria under typical growth conditions.³⁴ K^+ competes with Mg^{2+} for interactions with any RNA; thus, the higher cellular levels of K^+ would shift our Γ_{2+} curves to a higher Mg^{2+} concentration range. An increase in Γ_{2+} by one ion is coupled to a decrease in Γ_+ by ~ 1.8 K^+ ions;^{35,36} therefore, a 3-fold increase in K^+ activity should cause a ~ 6 fold ($3^{1.8}$) increase in the Mg^{2+} concentration needed to attain a given Γ_{2+} value. Estimates of the effective Mg^{2+} concentration in vivo span 0.5 to ~ 2 mM;^{37–39} in our experiments, the same accumulation of excess Mg^{2+} as occurs with an RNA in vivo would be expected in the range 83 μM to 0.33 mM.

How Mg^{2+} Affects the A-Riboswitch Energy Landscape. Having defined four A-riboswitch conformations with different combinations of tertiary interactions, we are now able to quantitatively parse the Mg^{2+} -dependence of RNA folding and ligand binding into several components. We consider all four conformations

in Figure 6A as structures that are accessible to the wild type sequence and capable of interconversion; the diagram then becomes a four-sided thermodynamic cycle that is defined, in the simplest possible model, by three ΔG values. There is a free energy associated with docking of the hairpin loops (ΔG_{dock}), a free energy for ligand-binding pocket formation (ΔG_{LBP} , which includes interactions with the ligand and therefore depends on the concentration of ligand present), and a coupling factor that links loop docking and ligand binding (ΔG_w). Each of these three free energies can be deduced from the relative free energy values of the four RNA forms (Figure 6B). The Mg^{2+} -dependence of each ΔG represents a different potential mechanism for translating Mg^{2+} –RNA interactions into an enhancement of ligand binding affinity; each is considered in turn below.

First, the binding of ligand to an RNA with undocked loops ($\Delta G_{\text{LBP}}^\circ$) is intrinsically very weak but strongly enhanced by Mg^{2+} . It is known from structure mapping (ref 22 and Figure 4), NMR,²⁰ and fluorescent tag⁴⁰ studies that ligand binding induces distinctive structural changes in the binding pocket; presumably the organization of the binding pocket needed to accommodate ligand is energetically costly. Mg^{2+} strongly favors the structural rearrangements that accompany binding; ΔG_{LBP} is enhanced -5.1 kcal/mol by 1 mM Mg^{2+} . Backbone surrounding the bound ligand forms two sharp bends which bring phosphate oxygens as close as 3.6 Å;¹⁸ such regions of high charge density could be responsible for the sensitivity of the binding pocket structure to Mg^{2+} ⁴⁰ and the accumulation of excess Mg^{2+} .

Second, docking of the hairpin loops (ΔG_{dock}) is also an intrinsically unfavorable reaction but strongly favored by Mg^{2+} , as expected from other studies.^{20,21,41} Here we find that 1 mM Mg^{2+} stabilizes the docked conformation by ~ -3.5 kcal/mol. However, the docking reaction will influence ligand binding only to the degree that loop docking and formation of the binding pocket are favorably linked ($\Delta G_w < 0$, Figure 6A). From studies of a purine riboswitch RNA with both hairpins deleted, this linkage has been estimated as -3.3 kcal/mol in the presence of 10 mM Mg^{2+} (125 mM K^+);²² it has not been known whether a favorable ΔG_w depends on Mg^{2+} . From our estimations of $\Delta G_{\text{dock}}^\circ$ and $\Delta G_{\text{LBP}}^\circ$, we deduce $\Delta G_w^\circ \approx -5.3$ kcal/mol (see arrow in Figure 6B). This value becomes more negative by -0.5 kcal/mol in 0.1 mM Mg^{2+} , or -1.2 kcal/mol in 1 mM Mg^{2+} . (These free energies, because they are small differences between large $\Delta G_{\text{RNA-2+}}$, have uncertainties comparable to their size.) The increase in Γ_{2+} that accompanies loop docking also appears somewhat larger when the binding pocket is occupied (2.1 ions/RNA, at 1 mM Mg^{2+}) than unoccupied (1.8 ions/RNA), though the difference of 0.3 ions is comparable to the measurement error. We conclude that a strong linkage between the two tertiary structures is inherent to the RNA structure, but the linkage may be enhanced by Mg^{2+} to a small degree.

The value of ΔG_w° we calculate assumes that the DAP complex with C60G RNA mimics a ligand-bound state that is accessible to the wild type riboswitch sequence. The G38C RNA is more difficult to saturate ($\Delta G_{\text{LBP}}^\circ$ more positive) and would therefore yield a somewhat more negative value of ΔG_w° if used as a mimic of the ligand-bound intermediate. The conclusion that there is strong coupling between the two sets of tertiary structure in the absence of Mg^{2+} nevertheless remains.

Overall, we conclude that Mg^{2+} enhances the A-riboswitch affinity for adenine principally by two mechanisms of nearly equal effectiveness: stabilization of the docked loop structure, and stabilization of the ligand binding pocket (cf. $\Delta G_{\text{Idock-E2+}}$

and ΔG_{ILBP} , Figure 5C). The Mg^{2+} interaction free energies are large compared to the intrinsic stability differences between the various RNA folds, and above ~ 0.1 mM Mg^{2+} the relative stabilities of the RNA forms reflect, to a large degree, differences in the strengths of their interactions with Mg^{2+} .

Parallels to Mg^{2+} -Induced Collapse in Larger RNAs. Mg^{2+} -induced “collapse” of an RNA to compact, non-native structures has been documented in several larger sequences (~ 200 – 400 nt), with implications for the kinetic and equilibrium pathways by which tertiary structures form in response to Mg^{2+} .^{11,15,42,43} There is a structural parallel between these larger RNAs and the A-riboswitch RNA, in that large RNAs tend to use pseudoknot, tetraloop receptor,⁴⁴ and A-minor⁴⁵ motifs to link peripheral elements together structurally. Such tertiary cross-links presumably restrain the RNA dimensions and the organization of the central core of the structure [see, for instance, the structures of RNase P⁴⁶ and *Azoarcus* group I intron⁴⁷]. In a similar way, a peripheral cross-link in the A-riboswitch (the docked loops) is linked to formation of a core tertiary structure (the ligand binding site). Are there also parallels between the Mg^{2+} -induced collapse phenomenon and the response of the A-riboswitch RNA to Mg^{2+} ?

Although specific characteristics of the collapse phenomenon vary between RNAs, usually two distinct types of collapse can be distinguished. Partial collapse takes place at lower Mg^{2+} concentrations, does not depend on tertiary contacts,^{11,15,48} and is promoted by a variety of ions, presumably because it relies on simple screening of the electrostatic repulsion between phosphates.^{48,49} At higher Mg^{2+} concentrations, further collapse takes place to a form with nearly the same dimensions as the native structure. This second collapse phase depends on the potential of the RNA to form specific tertiary contacts.^{11,15,48} However, chemical probing of the near-native collapsed form does not uncover any fully formed tertiary contacts.^{15,48,50} Among several tertiary interactions mutated in one study of a group I intron, only disruption of a peripheral tetraloop-receptor contact destabilized the compact intermediate in addition to the native structure.¹⁵ These near-native intermediates apparently sample peripheral contact(s) without fully forming either peripheral or central core tertiary structures. The thermodynamic linkages between peripheral contacts and the central core are therefore important parameters that determine the cooperativity of the folding pathway in response to titration with Mg^{2+} .

The Mg^{2+} -dependent equilibrium folding pathway of the A-riboswitch is similar to that of larger RNAs in that it is predicted to proceed through a compact intermediate with docked loops. On the basis of calculations with the free energy landscape, the I_{dock} RNA form may reach 30% of the total RNA at some ligand and Mg^{2+} concentrations. In the case of the A-riboswitch, we were able to estimate the cooperative free energy that links the peripheral and core structures; it is large (-5.3 kcal/mol) and only weakly dependent on Mg^{2+} . In the A-riboswitch, at least, the core and peripheral structures take advantage of Mg^{2+} for stability, while the linkage between them is largely built into the RNA structure.

In contrast to the two collapse phases seen with larger RNAs, we do not observe the first, electrostatic collapse that is independent of peripheral tertiary contacts; the extended E state of the A-riboswitch appears completely insensitive to Mg^{2+} . Screening of the RNA negative charge by ions must not be able to alter the intrinsic stiffness of the unliganded junction in this RNA. A similar finding has been made in a study of the S domain of RNase P,

which concluded that a Mg^{2+} -independent, extended conformation of a folding intermediate is maintained because of a specific, non-native structure within a three-helix junction, and that electrostatic repulsion plays a minor role in enforcing the extended conformation.⁵¹ Modulation of the stiffness vs. flexibility of folding intermediates is another way RNA sequence may adjust the stability of native structures in the presence Mg^{2+} .

Concluding Remarks. Γ_{2+} measurements provide a useful perspective for considering the effects of Mg^{2+} on RNA folding. As illustrated with the A-riboswitch RNA in this work, a knowledge of Γ_{2+} as a function of Mg^{2+} concentration can be used to extract the intrinsic (Mg^{2+} -independent) free energy of folding reactions which might not be directly observable. Γ_{2+} measurements also reveal the strong Mg^{2+} -dependences of ion uptake ($\Delta\Gamma_{2+}$) and Mg^{2+} –RNA interaction free energies ($\Delta G_{\text{RNA-2+}}$), which are usually not apparent from folding reaction analyses based on the Hill equation.¹⁰ Lastly, Γ_{2+} provides benchmarks for computational efforts to quantitatively model Mg^{2+} interactions with known RNA conformations.⁹

■ ASSOCIATED CONTENT

S Supporting Information. Three figures showing melting curves of wild type, C60G, and G38C RNAs, the hydroxyl radical reactivity of C60G RNA, and the correspondence of normalized Mg^{2+} –RNA interaction free energies for the native A-riboswitch and extended C60G RNAs; two tables list R_g values for the three RNAs under various conditions and Γ_{2+} and $\Delta G_{\text{RNA-2+}}$ values evaluated at 1 mM Mg^{2+} for wild type and C60G RNAs; a complete derivation of eq 1. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Tel 410 516 7448. Fax 410 516 8420. E-mail draper@jhu.edu.

Funding Sources

This work was supported by National Institutes of Health Grants GM58545 (to D.E.D.) and T32 GM008403 (Program in Molecular Biophysics).

■ ACKNOWLEDGMENT

We thank Drs. Xiaobing Zuo and Sonke Seifert for help with collection and analysis of the scattering data, and Robert Trachman for providing Mg^{2+} titration data for the A-riboswitch in the absence of ligand.

■ ABBREVIATIONS USED

A-riboswitch, the adenine-binding domain of an adenine riboswitch RNA; DAP, 2,6-diaminopurine; HQS, 8-hydroxyquinoline-5-sulfonic acid; SAXS, small-angle X-ray scattering; R_g , radius of gyration

■ ADDITIONAL NOTE

^aWe assume that the extended states of wild type and C60G RNAs have the same standard state chemical potential, set to zero in Figure 6B, so that free energy changes measured with each RNA may be compared. This assumption implies that the two RNA E forms have similar structures and interaction with ions. Although the C60G mutation may extend the P2 helix by one

base pair, we found no significant difference in Γ_{2+} between the two RNAs up to $\sim 30 \mu\text{M}$ Mg^{2+} in the absence of ligand (Figure 5A) and suggest that the assumption is a reasonable approximation within the errors of the Figure 6B free energy landscape.

REFERENCES

- (1) Lynch, D. C., and Schimmel, P. R. (1974) Cooperative binding of magnesium to transfer ribonucleic acid studied by a fluorescent probe. *Biochemistry* 13, 1841–1852.
- (2) Römer, R., and Hach, R. (1975) tRNA conformation and magnesium binding. A study of a yeast phenylalanine-specific tRNA by a fluorescent indicator and differential melting curves. *Eur. J. Biochem.* 55, 271–284.
- (3) Stein, A., and Crothers, D. M. (1976) Conformational changes of transfer RNA. The role of magnesium(II). *Biochemistry* 15, 160–167.
- (4) Leroy, J. L., Guéron, M., Thomas, G., and Favre, A. (1977) Role of divalent ions in folding of tRNA. *Eur. J. Biochem.* 74, 567–574.
- (5) Misra, V. K., and Draper, D. E. (2000) $\text{Mg}(2+)$ binding to tRNA revisited: the nonlinear Poisson-Boltzmann model. *J. Mol. Biol.* 299, 813–825.
- (6) Cate, J. H., Hanna, R. L., and Doudna, J. A. (1997) A magnesium ion core at the heart of a ribozyme domain. *Nat. Struct. Biol.* 4, 553–558.
- (7) Misra, V. K., and Draper, D. E. (2001) A thermodynamic framework for Mg^{2+} binding to RNA. *Proc. Natl. Acad. Sci. U. S. A.* 98, 12456–12461.
- (8) Grilley, D., Soto, A. M., and Draper, D. E. (2006) Mg^{2+} -RNA interaction free energies and their relationship to the folding of RNA tertiary structures. *Proc. Natl. Acad. Sci. U. S. A.* 103, 14003–14008.
- (9) Soto, A. M., Misra, V., and Draper, D. E. (2007) Tertiary structure of an RNA pseudoknot is stabilized by “diffuse” $\text{Mg}(2+)$ ions. *Biochemistry* 46, 2973–2983.
- (10) Leipply, D., and Draper, D. E. (2010) The dependence of RNA tertiary structure stability on Mg^{2+} concentration: interpretation of the Hill equation and coefficient. *Biochemistry* 49, 1843–1853.
- (11) Buchmueller, K. L., Webb, A. E., Richardson, D. A., and Weeks, K. M. (2000) A collapsed non-native RNA folding state. *Nat. Struct. Biol.* 7, 362–366.
- (12) Russell, R., Millett, I. S., Doniach, S., and Herschlag, D. (2000) Small angle X-ray scattering reveals a compact intermediate in RNA folding. *Nat. Struct. Biol.* 7, 367–370.
- (13) Rangan, P., Masquida, B., Westhof, E., and Woodson, S. A. (2003) Assembly of core helices and rapid tertiary folding of a small bacterial group I ribozyme. *Proc. Natl. Acad. Sci. U. S. A.* 100, 1574–1579.
- (14) Perez-Salas, U. A., Rangan, P., Krueger, S., Briber, R. M., Thirumalai, D., and Woodson, S. A. (2004) Compaction of a bacterial group I ribozyme coincides with the assembly of core helices. *Biochemistry* 43, 1746–1753.
- (15) Chauhan, S., Caliskan, G., Briber, R. M., Perez-Salas, U., Rangan, P., Thirumalai, D., and Woodson, S. A. (2005) RNA tertiary interactions mediate native collapse of a bacterial group I ribozyme. *J. Mol. Biol.* 353, 1199–1209.
- (16) Mandal, M., Boese, B., Barrick, J. E., Winkler, W. C., and Breaker, R. R. (2003) Riboswitches control fundamental biochemical pathways in *Bacillus subtilis* and other bacteria. *Cell* 113, 577–586.
- (17) Mandal, M., and Breaker, R. R. (2004) Adenine riboswitches and gene activation by disruption of a transcription terminator. *Nat. Struct. Mol. Biol.* 11, 29–35.
- (18) Serganov, A., Yuan, Y. R., Pikovskaya, O., Polonskaia, A., Malinina, L., Phan, A. T., Hobartner, C., Micura, R., Breaker, R. R., and Patel, D. J. (2004) Structural basis for discriminative regulation of gene expression by adenine- and guanine-sensing mRNAs. *Chem. Biol.* 11, 1729–1741.
- (19) Lemay, J. F., Penedo, J. C., Tremblay, R., Lilley, D. M., and Lafontaine, D. A. (2006) Folding of the adenine riboswitch. *Chem. Biol.* 13, 857–868.
- (20) Noeske, J., Schwalbe, H., and Wohnert, J. (2007) Metal-ion binding and metal-ion induced folding of the adenine-sensing riboswitch aptamer domain. *Nucleic Acids Res.* 35, 5262–5273.
- (21) Brenner, M. D., Scanlan, M. S., Nahas, M. K., Ha, T., and Silverman, S. K. (2010) Multivector fluorescence analysis of the xpt guanine riboswitch aptamer domain and the conformational role of guanine. *Biochemistry* 49, 1596–1605.
- (22) Stoddard, C. D., Gilbert, S. D., and Batey, R. T. (2008) Ligand-dependent folding of the three-way junction in the purine riboswitch. *RNA* 14, 675–684.
- (23) Grilley, D., Soto, A. M., and Draper, D. E. (2009) Direct quantitation of Mg^{2+} -RNA interactions by use of a fluorescent dye. *Methods Enzymol.* 455, 71–94.
- (24) Lambert, D., Leipply, D., and Draper, D. E. (2010) The osmolyte TMAO stabilizes native RNA tertiary structures in the absence of $\text{Mg}(2+)$: Evidence for a large barrier to folding from phosphate dehydration. *J. Mol. Biol.* 404, 138–157.
- (25) Petoukhov, M. V., Konarev, P. V., Kikhney, A. G., and Svergun, D. I. (2007) ATSAS 2.1 – towards automated and web-supported small-angle scattering data analysis. *J. Appl. Crystallogr.* 40, s223–s228.
- (26) Kozin, M. B., and Svergun, D. I. (2001) Automated matching of high- and low-resolution structural models. *J. Appl. Crystallogr.* 34, 33–41.
- (27) Das, R., Laederach, A., Pearlman, S. M., Herschlag, D., and Altman, R. B. (2005) SAFA: semi-automated footprinting analysis software for high-throughput quantification of nucleic acid footprinting experiments. *RNA* 11, 344–354.
- (28) Gilbert, S. D., Love, C. E., Edwards, A. L., and Batey, R. T. (2007) Mutational analysis of the purine riboswitch aptamer domain. *Biochemistry* 46, 13297–13309.
- (29) Rambo, R. P., and Tainer, J. A. (2010) Improving small-angle X-ray scattering data for structural analyses of the RNA world. *RNA* 16, 638–66.
- (30) Latham, J. A., and Cech, T. R. (1989) Defining the inside and outside of a catalytic RNA molecule. *Science* 245, 276–245.
- (31) Leipply, D., Lambert, D., and Draper, D. E. (2009) Ion–RNA interactions: Thermodynamic analysis of the effects of mono- and divalent ions on RNA conformational equilibria. *Methods Enzymol.* 469, 433–463.
- (32) Burton, K. (1994) Adenine transport in *Escherichia coli*. *Proc. Biol. Sci.* 255, 153–157.
- (33) Nygaard, P., and Saxild, H. H. (2005) The purine efflux pump PbuE in *Bacillus subtilis* modulates expression of the PurR and G-box (XptR) regulons by adjusting the purine base pool size. *J. Bacteriol.* 187, 791–794.
- (34) Cayley, S., Lewis, B. A., Guttman, H. J., and Record, M. T., Jr. (1991) Characterization of the cytoplasm of *Escherichia coli* K-12 as a function of external osmolarity. Implications for protein–DNA interactions in vivo. *J. Mol. Biol.* 222, 281–300.
- (35) Record, M. T., Jr., deHaseth, P. L., and Lohman, T. M. (1977) Interpretation of monovalent and divalent cation effects on the lac repressor-operator interaction. *Biochemistry* 16, 4791–4796.
- (36) Misra, V. K., and Draper, D. E. (1999) The interpretation of Mg^{2+} binding isotherms for nucleic acids using Poisson-Boltzmann theory. *J. Mol. Biol.* 294, 1135–47.
- (37) Alatossava, T., Jutte, H., Kuhn, A., and Kellenberger, E. (1985) Manipulation of intracellular magnesium content in polymyxin B nonapeptide-sensitized *Escherichia coli* by ionophore A23187. *J. Bacteriol.* 162, 413–419.
- (38) London, R. E. (1991) Methods for measurement of intracellular magnesium: NMR and fluorescence. *Annu. Rev. Physiol.* 53, 241–258.
- (39) Froschauer, E. M., Kolisek, M., Dieterich, F., Schweigel, M., and Schweyen, R. J. (2004) Fluorescence measurements of free $[\text{Mg}^{2+}]$ by use of mag-fura 2 in *Salmonella enterica*. *FEMS Microbiol. Lett.* 237, 49–55.
- (40) Rieder, R., Lang, K., Graber, D., and Micura, R. (2007) Ligand-induced folding of the adenosine deaminase A-riboswitch and implications on riboswitch translational control. *ChemBiochem* 8, 896–902.
- (41) Lemay, J. F., and Lafontaine, D. A. (2007) Core requirements of the adenine riboswitch aptamer for ligand binding. *RNA* 13, 339–350.
- (42) Fang, X., Littrell, K., Yang, X. J., Henderson, S. J., Siefert, S., Thiagarajan, P., Pan, T., and Sosnick, T. R. (2000) Mg^{2+} -dependent compaction and folding of yeast tRNAPhe and the catalytic domain of

the B. subtilis RNase P RNA determined by small-angle X-ray scattering. *Biochemistry* 39, 11107–11113.

(43) Russell, R., Millett, I. S., Tate, M. W., Kwok, L. W., Nakatani, B., Gruner, S. M., Mochrie, S. G., Pande, V., Doniach, S., Herschlag, D., and Pollack, L. (2002) Rapid compaction during RNA folding. *Proc. Natl. Acad. Sci. U. S. A.* 99, 4266–4271.

(44) Costa, M., and Michel, F. (1995) Frequent use of the same tertiary motif by self-folding RNAs. *EMBO J.* 14, 1276–1285.

(45) Nissen, P., Ippolito, J. A., Ban, N., Moore, P. B., and Steitz, T. A. (2001) RNA tertiary interactions in the large ribosomal subunit: the A-minor motif. *Proc. Natl. Acad. Sci. U. S. A.* 98, 4899–4903.

(46) Kazantsev, A. V., Krivenko, A. A., Harrington, D. J., Holbrook, S. R., Adams, P. D., and Pace, N. R. (2005) Crystal structure of a bacterial ribonuclease P RNA. *Proc. Natl. Acad. Sci. U. S. A.* 102, 13392–13397.

(47) Adams, P. L., Stahley, M. R., Kosek, A. B., Wang, J., and Strobel, S. A. (2004) Crystal structure of a self-splicing group I intron with both exons. *Nature* 430, 45–50.

(48) Das, R., Kwok, L. W., Millett, I. S., Bai, Y., Mills, T. T., Jacob, J., Maskel, G. S., Seifert, S., Mochrie, S. G., Thiyagarajan, P., Doniach, S., Pollack, L., and Herschlag, D. (2003) The fastest global events in RNA folding: electrostatic relaxation and tertiary collapse of the Tetrahymena ribozyme. *J. Mol. Biol.* 332, 311–319.

(49) Moghaddam, S., Caliskan, G., Chauhan, S., Hyeon, C., Briber, R. M., Thirumalai, D., and Woodson, S. A. (2009) Metal ion dependence of cooperative collapse transitions in RNA. *J. Mol. Biol.* 393, 753–764.

(50) Buchmueller, K. L., and Weeks, K. M. (2003) Near native structure in an RNA collapsed state. *Biochemistry* 42, 13869–13878.

(51) Baird, N. J., Gong, H., Zaheer, S. S., Freed, K. F., Pan, T., and Sosnick, T. R. (2010) Extended structures in RNA folding intermediates are due to nonnative interactions rather than electrostatic repulsion. *J. Mol. Biol.* 397, 1298–1306.

(52) Lambert, D., and Draper, D. E. (2007) Effects of osmolytes on RNA secondary and tertiary structure stabilities and RNA–Mg²⁺ interactions. *J. Mol. Biol.* 370, 993–1005.