

A Switch I Mutant of Cdc42 Exhibits Less Conformational Freedom


Reena Chandrashekar,[†] Omar Salem,[†] Hana Krizova,[‡] Robert McFeeters,[‡] and Paul D. Adams^{*,†}

[†]Department of Chemistry and Biochemistry, University of Arkansas, Fayetteville, Arkansas 72701, United States

[‡]Department of Chemistry, University of Alabama, Huntsville, Alabama 35899, United States

S Supporting Information

ABSTRACT: Cdc42 is a Ras-related small G-protein and functions as a molecular switch in signal transduction pathways linked with cell growth and differentiation. It is controlled by cycling between GTP-bound (active) and GDP-bound (inactive) forms. Nucleotide binding and hydrolysis are modulated by interactions with effectors and/or regulatory proteins. These interactions are centralized in two relatively flexible “Switch” regions as characterized by internal dynamics on multiple time scales [Loh, A. P., et al. (2001) *Biochemistry* 40, 4590–4600], and this flexibility may be essential for protein interactions. In the Switch I region, Thr³⁵ seems to be critical for function, as it is completely invariant in Ras-related proteins. To investigate the importance of conformational flexibility in Switch I of Cdc42, we mutated threonine to alanine, determined the solution structure, and characterized the backbone dynamics of the single-point mutant protein, Cdc42(T35A). Backbone dynamics data suggest that the mutation changes the time scale of the internal motions of several residues, with several resonances not being discernible in wild-type Cdc42 [Adams, P. D., and Oswald, R. E. (2007) *Biomol. NMR Assignments* 1, 225–227]. The mutation does not appear to affect the thermal stability of Cdc42, and chymotrypsin digestion data further suggest that changes in the conformational flexibility of Switch I slow proteolytic cleavage relative to that of the wild type. In vitro binding assays show less binding of Cdc42(T35A), relative to that of wild type, to a GTPase binding protein that inhibits GTP hydrolysis in Cdc42. These results suggest that the mutation of T³⁵ leads to the loss of conformational freedom in Switch I that could affect effector–regulatory protein interactions.



The success of target-based approaches to combat Ras-related signal transduction events leading to diseases such as cancer will depend upon understanding the correlations among structural biology, protein–protein interactions, and protein function. Indeed, binding events and the activation of protein receptors may play a role in abnormal cell signaling activities. Understanding such activities will require approaches that elucidate changes in both protein conformation and protein dynamics that lead to particular molecular events and behaviors. Cdc42 is a member of the Rho subfamily of Ras proteins, and they are closely related to small GTPases. These proteins are ubiquitously expressed, evolutionarily conserved, and among the first human oncogenic proteins to have been identified.^{1,2} Mutations or abnormal expression patterns of Ras proteins have been found in up to 30% of cancer cell types.^{1,2} Ras proteins act as molecular switches that couple extracellular signals to a variety of cellular responses. Ras protein activities are controlled via cycling between a biologically active GTP form and an inactive GDP-bound form.^{1,3} The function of Cdc42, as well as other Ras proteins, is largely mediated through protein–protein interactions. Therefore, understanding the specificity governing these interactions and associated conformational changes is vital for understanding molecular mechanisms underlying the onset of

diseases such as cancer. Different signals can activate guanine nucleotide exchange factors (GEFs) that induce the small G-protein to switch from the inactive, GDP-bound state to the active, GTP-bound state.^{4,5} In the active state, the GTPase binds the downstream effector and the effector in turn can mediate diverse biological responses. These different signals are terminated when Cdc42 is deactivated, either through its own ability to hydrolyze GTP or by means of a reaction catalyzed by GTPase-activating proteins (GAPs).⁶ Alternatively, interactions with regulatory proteins that block GDP dissociation, such as GDP dissociation inhibitors (GDIs), stabilize the GDP-bound inactive state and prevent effective GTP–GDP cycling.⁷ Mechanisms by which Cdc42 and most Ras proteins participate in cellular proliferation and/or transformation include mutations that alter the GTPase activity^{1,8–11} and mutations that alter the interactions with effectors or regulators.¹² The best understood mutations have been those that result in cellular transformation by inhibiting GTPase activity.^{1,8} In addition to this, Cdc42 mutations have also been shown to facilitate cell transformation by

Received: March 22, 2011

Revised: June 10, 2011

Published: June 13, 2011

increasing the rate of cycling between active and inactive forms of the protein.¹³ Structural studies of the mutant Cdc42(F28L) provided evidence that local environmental changes affect the stability of the nucleotide-binding site leading to aberrant cell signaling behavior.¹⁴

The binding of effector molecules to Cdc42 essentially involves two regions, Switch I (residues 28–40) and Switch II (residues 57–74), which can undergo rapid conformational changes. In fact, it has been reported that the Switch I region in H-Ras possesses two distinct conformational states, bound to effectors and unbound.^{15,16} Indeed, Switch I is the principal interacting surface for Cdc42 with downstream effectors.^{11,17} For Cdc42 in solution, several resonances in the Switch regions cannot be observed because of increased flexibility in these regions.¹⁸ The binding of effectors to Cdc42 results in the loss of backbone mobility of the Switch I region, suggesting that the inherent dynamics may allow Cdc42 to bind to different effector proteins.¹⁹ It has been widely thought that the conformational freedom of the Switch I region may provide the basis for nucleotide-dependent signal propagation of Cdc42, and that the conformational flexibility of this region is central to the engagement of downstream effectors.^{17,20,21} This premise agrees with biological studies.²² Structural studies of Ras have implicated Thr³⁵ in the Switch I region as being critical for the conformational changes as it is completely invariant in Ras-related proteins.³¹ ³¹P nuclear magnetic resonance (NMR) studies with H-Ras(T35A) and Cdc42(T35A) suggested that this mutation causes a signaling defective state.^{11,16} The ³¹P NMR spectra of both proteins in their GTP-bound state revealed only small changes, suggesting that the nucleotide-binding site of the mutant was similar to that of the wild type. However, binding peptide effectors that target the Switch I region produced significant shifts in ³¹P resonances in the wild-type Ras–peptide complexes but not in the T35A–effector complexes, suggesting a weaker interaction between effector peptide derivatives and the mutant protein.^{11,16}

To investigate the importance of the conformational flexibility in Switch I of Cdc42, we have determined the solution structure, examined the backbone dynamics, and characterized the biochemical activity of Cdc42(T35A) [Protein Data Bank (PDB) entry 2kb0]. Our results suggest that the mutation of Thr³⁵ leads to the loss of conformational freedom in Switch I relative to the rest of the protein. Our results furthermore provide molecular details about why the conformational flexibility of this region is inherently important for the recognition of the Ras-related effectors. Recent studies have implied that innovative strategies for targeting cancer treatment in the future should target specific mutant Ras proteins and directly block Ras protein interactions with oncogenic potential, for example, using antibody fragments.²³ Taken together, the structural and dynamics results for Cdc42(T35A) suggest that, in addition to strategies for blocking Ras–protein interactions, strategies for restricting the conformational flexibility of important binding regions of Ras leading to the inhibition of effector interactions that facilitate aberrant signaling activity may also have therapeutic potential.

MATERIALS AND METHODS

Protein Expression and Purification. Cdc42(T35A) was cloned, expressed, and purified as described previously for Cdc42.^{14,24} Briefly, the Cdc42(T35A) construct was cloned into a pET-15b vector and subsequently overexpressed in *Escherichia*

coli. Cultures (5 mL) were initially grown to saturation and used to seed 1 or 2 L cultures grown in LB broth for unlabeled samples, and in minimal medium containing ¹⁵NH₄Cl for ¹⁵N-labeled samples and ¹⁵NH₄Cl and [¹³C]glucose for ¹⁵N- and ¹³C-labeled samples. Cell cultures were grown at 37 °C to an OD₅₆₀ reading of 0.4–0.6, and expression was induced by addition of 1 mM IPTG (isopropyl β-D-thiogalactopyranoside) for 7–12 h. (His)₆-Cdc42(T35A) and the wild type were expressed and purified by column chromatography. Matrix-assisted laser desorption ionization mass spectroscopy was used to verify that the molecular weight of the protein was correct, and gel electrophoresis was used to validate the purity of the protein samples.

NMR Sample Conditions, Data Collection, and Processing.

All protein samples were prepared in an NMR sample buffer solution containing 25 mM NaCl, 10 mM NaH₂PO₄, 5 mM MgCl₂, and 1 mM NaN₃ with 8% D₂O at pH 5.5 (not adjusted for isotope effect) at Cdc42(T35A) concentrations of 0.2–0.5 mM in volumes of 500–550 mL for structural studies. The concentrations of wild-type Cdc42 and T35A used for relaxation data collection were 1 mM. NMR spectra for structural studies were recorded using Varian Inova 500, 600, and 800 MHz spectrometers. All spectrometers were equipped with triple-resonance pulsed-field gradient probes. All NMR spectra were recorded at 25 °C in States-TPPI mode for quadrature detection.²⁵ Carrier frequencies for ¹H and ¹⁵N were 4.77 and 115.9 ppm, respectively. Two-dimensional ¹H–¹⁵N HSQC spectra were recorded on Cdc42(T35A) homogeneously labeled with ¹⁵N, and resonance assignments were made and deposited (BMRB entry 15424). Hydrogen–deuterium exchange was assessed by lyophilizing wild-type Cdc42 or T35A and resuspending each in 100% D₂O. A series of ¹H–¹⁵N HSQC experiments were performed, and the decrease in the volume and eventual disappearance of ¹H–¹⁵N cross-peaks were used to determine the extent of exchange of deuterons for protons. Three-dimensional ¹H–¹⁵N TOCSY-HSQC experiments were conducted with saturating delays of 50 and 70 ms. Three-dimensional ¹H–¹⁵N NOESY-HSQC spectra were recorded using a mixing time of 125 ms. The pulse sequences used included a water suppression by a selective pulse on the water resonance followed by dephasing using a gradient in the mixing period, as well as by a gradient applied while the magnetization was spin-ordered during the final INEPT transfer step. NMR spectra were processed using NMRPipe, version 1.6.²⁶ The spectra were zero-filled and then apodized using a Gaussian window function prior to Fourier transformation. After Fourier transformation, a baseline correction was applied. Spectra were visualized and analyzed using SPARKY.²⁷

Structure Calculation. The structure of Cdc42(T35A) was modeled and refined using SPARKY²⁷ and CNS.²⁸ A combination of ¹H–¹⁵N NOESY-HSQC and ¹H–¹³C NOESY-HSQC experiments was used to derive distance constraints. In the spectra, the peak volumes of several well-dispersed and isolated sequential *d*_{αN} and *d*_{αN}(*i*,*i*+1) NOE peaks within and between parallel β-strands, respectively, were compared to the distances expected for ideal β-sheets (2.2 and 3 Å, respectively). Approximately 1057 NOESY constraints were classified according to their relative strength in the NOESY spectrum (<2.4, <3.4, <4.0, and <5.5 Å). Also, 71 hydrogen bond constraints were also input into the structure calculations on the basis of the rate of deuterium exchange, the chemical shift index, and expected NOE patterns indicative of hydrogen bonding. For each hydrogen bond, the constraint for the amide proton–carbonyl oxygen

Table 1. Structural Statistics and Restraint Data for the Structure Calculation of Cdc42(T35A)^a

	ensemble of 12 structures	most representative structure
rmsd from experimental constraints		
NOE distances (Å)	0.018 ± 0.004	0.018
dihedral angles (deg)	0.37 ± 0.05	0.38
rmsd from secondary structure		
bonds (Å)	0.0015 ± 0.0001	0.0015
angles (deg)	0.33 ± 0.02	0.33
impropers (deg)	0.35 ± 0.02	0.34
Restraint Data (no.)		
long-range NOEs ($ i - j > 4$)		174
medium-range NOEs ($4 > i - j > 1$)		883
total NOEs		1057
intraresidue		457
sequential		475
medium-range		169
long-range		246
hydrogen bond		142
ψ		101
φ		100
NH backbone		74
RDC restraints		

^a From the average structure for backbone non-hydrogen atoms: 0.859 ± 0.01.

distance for the residue was held between 1.8 and 2.3 Å, and that for the amide nitrogen–carbonyl oxygen distance was between 2.5 and 3.3 Å.¹⁸ Dihedral restraints (total numbers of 101 for φ and 100 for ψ) were included for residues in α -helices and β -sheets, and the constraints were held to favorable regions of the φ – ψ space (for α -helix, $\varphi = -80 \pm 50^\circ$ and $\psi = -20 \pm 50^\circ$; for β -sheet, $\varphi = -105 \pm 65^\circ$ and $\psi = 145 \pm 45^\circ$) using methods developed by Wishart, Sykes, and Richardson^{29,30} as done in the solution structure determination of other Cdc42 constructs.^{14,18} Residual dipolar coupling restraints were generated using a sample of Cdc42(T35A)-GDP that was partially aligned with a nonionic liquid crystalline medium composed of 4% C12E6 [*n*-dodecyl hexa-(ethylene glycol)] with hexanol as described previously.¹⁴ RDCs were measured from partially aligned and unaligned samples using TROSY experiments with the phase cycling modified to observe the nitrogen components individually. RDC restraints were developed using PALES,³¹ with optimal values of D_a and R being -9.8 and 0.64 , respectively. All restraints were used to calculate an ensemble of structures for the protein using CNS. Twelve “best” structures were selected on the basis of their overall low energies and minimal constraint violations, and these structures have been deposited in the RCSB Protein Data Bank (entry 2kb0). Table 1 gives the experimental statistics of the structure calculations. The structures were visualized using Swiss PDBviewer (version 3.7)³² and Chimera³³ and analyzed using AQUA,³⁴ PROCHECK-NMR,³⁵ and the NMR ensemble program, OLDERADO.³⁶

Dynamics Data Analysis. For relaxation studies, T_1 , T_2 , and ^1H – ^{15}N NOE experiments were conducted with wild-type and T35A samples at 500 and 800 MHz.^{37,38} The relaxation delays for T_1 experiments were 0, 150, 300, 450, 600, 750, 900, and 1050 ms with triplicate experiments recorded with 0 and 450 ms delays at 500 MHz. Delays of 0, 150, 500, 750, 1000, 1250, 1500, and 1750 ms with triplicate experiments with 0 and 1000 ms delays were collected at 800 MHz. T_2 experiments used 5, 15, 25, 40, 55, 70, 85, and 100 ms delays in triplicate at 5 and 40 ms at 500 MHz. Experiments conducted at 800 MHz used 5, 15, 25, 40, 55, 70, and 85 ms delays with 5 and 40 ms delay experiments repeated. Heteronuclear NOE experiments utilized a 5 s saturation period and a 5 s recycle delay. Spectra recorded with a 10 s recycle delay were used as references. All NOE experiments were conducted in triplicate. T_1 and T_2 values were obtained by fitting the peak intensities to a single exponential using NMRPipe.²⁶ Heteronuclear NOEs were measured as the ratio of peak intensity from the spectra collected with saturation divided by the volume of the same peak in the reference spectra. The relaxation data sets were subjected to model-free analysis using TENSOR³⁹ and Model-free4.⁴⁰ The diffusion tensors of wild-type Cdc42 and T35A were determined using the three-dimensional coordinate files of PDB entries 1aje and 2kb0, respectively.

In Vitro Binding Assays. For in vitro pull down assays, a peptide derivative of p21-activated serine/threonine kinase (PAK) that binds to wild-type Cdc42, PBD46 (H₂N-GSKER-PEISLPDSFEHTIHVGFDVAVTGFTGIPEQWARLLQTS-NIT-COO[−]), was expressed and purified,¹⁷ as was GST (control). Aliquots (5 mg) of GST-PBD46 (or GST alone) were mixed with 5 mg aliquots of partner proteins, wild-type Cdc42 or Cdc42(T35A), according to quantification, and 0.5 mM GMPPCP was added to the mixtures. The total mixture volume was increased to 2 mL via addition of PBS buffer. After incubation of the mixture for 4 h at 4 °C, 50 μL of Glutathione Sepharose beads (GE Healthcare) was added to precipitate GST-PBD46 (or GST) after incubation for 1 h; then the supernatant was discarded, and the beads were washed with PBS buffer at least three times. Fifty milliliters of 10 mM glutathione in 25 mM Tris (pH 8) was used to elute protein on the Sepharose beads. Ten microliters of each elution sample was run on 15% sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–PAGE), and band intensities were measured to calculate the relative amount of complex formed. In all gels, there was an equal density representing a doublet of GST-PBD46 indicating that equal amounts were pulled down by glutathione beads in wild-type Cdc42/PBD46 and Cdc42(T35A)/PBD46 mixtures, which has been seen in other in vitro binding assays using PBD with Cdc42 bound to various other nucleotides.¹¹ The corresponding intensity of the 20 kDa band on a 15% SDS–PAGE gel was measured using a UVP Biospectrum Imaging System. As a negative control, wild-type Cdc42 and Cdc42(T35A) were incubated with GST alone, as mentioned above, and the incubated mixture did not show a band for the wild type or T35A in the eluted sample, indicating that there was specific binding with PBD46 (data not shown).

Proteolytic Digestion Assay. Chymotrypsin (Sigma-Aldrich), trichloroacetic acid (TCA), NaCl (BDH), sodium phosphate, and urea (J. T. Baker) were of analytical or ultrapure grade. The proteolytic experiments with wild-type Cdc42 and Cdc42(T35A) were conducted at 25 °C using chymotrypsin in a buffer containing 100 mM NaCl and 10 mM NaH₂PO₄ (pH 8.0). Proteolytic




Figure 1. Structure of Cdc42(T35A)-GDP. The 12 lowest-energy structures were superimposed relative to the average structure generated using Xplor-NIH. They were visualized using Chimera.³³ The GDP nucleotide was superimposed on the structure(s) using coordinates from the X-ray crystal structure of wild-type Cdc42 (PDB entry 1AN0). (A) All 12 twelve structures overlaid with each other and (B) the most representative structure, as defined by OLDERADO.³⁶

digestions were performed in triplicate at an enzyme:substrate (wild type or T35A) ratio of 10:1 (enzyme:protein). The protease activity was stopped after incremental time intervals (20, 40, 50, 60, 70 80, 90, 100, and 110 min) by the addition of 100% TCA. After TCA precipitation, 8 M urea was added to the protein sample, and the degree of proteolytic cleavage was measured from the intensity of the 20 kDa band on a 15% SDS–PAGE gel corresponding to the uncleaved protein (time zero) (wild type or T35A), using UN-SCAN-IT gel version 6.1 (Silk Scientific). The control band was taken to be 100% to calculate the percentage of cleavage that occurred.

Differential Scanning Calorimetry. Temperature-induced unfolding of wild-type Cdc42 versus Cdc42(T35A) was performed using a Nano-DSC III differential scanning calorimeter (Calorimetry Sciences Corp.). A 1 mg/mL solution of Cdc42 (wild type and T35A) was dialyzed extensively against 1 × PBS with 0.5 mM Mg²⁺ at pH 7.5 and 4 °C. After dialysis, each protein sample was degassed before it was heated from 20 to 80 °C using a scan rate of 1.0 °C/min. The cells containing each protein sample were equilibrated for approximately 10 min before each run, and buffer samples without protein were run prior to each experiment to obtain a baseline and subtracted from the protein scan. Each protein sample was scanned in triplicate. CpCalc (Calorimetry Sciences Corp.) was used to construct the melting curve and to determine the T_m value.

RESULTS AND DISCUSSION

Overall Structure and Backbone Dynamics of Cdc42(T35A).

The solution structure of Cdc42(T35A) was characterized to examine conformational differences imparted by the mutation in the Switch I region when the side chain OH group of residue 35 is removed. An ensemble of the 12 lowest-energy structures of Cdc42(T35A) (PDB entry 2kb0) calculated from NMR-determined distance, hydrogen bonding, dihedral, and residual dipolar coupling restraints is shown (Figure 1A). The core of Cdc42(T35A) is comprised of residues 3–11, 15–24, 28, 29, 39–46, 48–58, 75–103, 109–117, 123–133, 136–150, and 152–177. The core region is similar to wild-type Cdc42¹⁸ and to other Cdc42 mutants, such as Cdc42(F28L).¹⁴ There were no changes observed for the secondary structure regions in Cdc42(T35A)

relative to those of the wild type, as Cdc42(T35A) consists of a six-stranded β -sheet structure around its C-terminal α -helix just as in the wild type (Figure 1B). As seen with the wild type, and other mutants of Cdc42,^{14,18} the Switch I and II regions and, to a lesser extent, the Insert region (α I) show structural disorder relative to the rest of the protein. The nucleotide-binding site seemed to be unaffected by the mutation, as there were no noticeable differences observed for residues that comprise the nucleotide-binding pocket of Cdc42(T35A) relative to that of the wild type in ¹⁵N HSQC spectra.²⁴ This is in agreement with ³¹P NMR studies of H-Ras-(T35A) and Cdc42(T35A) that showed very small, if any, changes in the ³¹P NMR spectra.¹¹ Therefore, the mutation in Cdc42-(T35A) does not seem to alter the overall folding of Cdc42, and the globular structure of Cdc42(T35A) is largely similar to that of wild-type Cdc42.

The backbone dynamics for Cdc42(T35A) were characterized from ¹⁵N relaxation parameters. Plots of the longitudinal (T_1) and transverse relaxation (T_2) rates and steady state heteronuclear NOE values are shown in Figure 2. The primary structure of Cdc42(T35A) consists of 166 non-proline residues, and ~72% of the backbone amide resonances for both the wild type (119 residues) and T35A (120 residues) were observable. One notable difference was that resonances for residues 40 and 41 were broadened and unobservable in the wild-type spectra. This broadening is less significant in the Cdc42(T35A) spectra, allowing analysis of NH bond vector internal dynamics for these residues. Because of resonance overlap in both the wild type and Cdc42(T35A), there were a number of resonances excluded from the analysis, including residues 35–39 in the Switch I region. However, in contrast to the wild-type spectra, the Cdc42(T35A) spectra contained resonances from residues 38, 40, and 41 in Switch I and residues 57, 59, 61, 63, 71, and 74 in Switch II for which internal dynamics parameters could be obtained. The NH resonances from residues 35 and 36 were also observed for Cdc42(T35A); however, they suffered from resonance overlap, making confident analysis of relaxation data impossible. Overall, relaxation data from regions with well-defined secondary structure of both Cdc42 constructs correspond very well to the previously published data for wild-type Cdc42 and Cdc42(F28L).^{19,41} Because the Switch I (residues 28–41) and Switch II (residues 57–74) regions participate in the interaction of Cdc42 with effector proteins, the internal motions of these regions are of particular interest. However, direct comparisons for these regions are not possible because of the resonance broadening and signal overlap in the NMR spectra of wild-type Cdc42.

Relaxation data were subjected to a model-free formalism,⁴² to extract parameters describing the internal motion of the individual NH bond vectors. We compared the relaxation data collected at both 500 and 800 MHz for both wild-type Cdc42 and Cdc42(T35A) and found that the trends in the plots of the data as a function of residue number agreed very well for both proteins. However, to minimize effects that can arise in the amplification of the CSA term at high field strengths, only the data collected at 500 MHz were used in the dynamics analysis. The τ_c for wild-type Cdc42 was 10.9 ± 0.3 ns and that for Cdc42(T35A) 11.0 ± 0.2 ns. An axially symmetric diffusion tensor fit best, with a $D_{\text{par}}/D_{\text{per}}$ for wild-type Cdc42 equal to 0.84 ± 0.02 and that for Cdc42(T35A) equal to 0.71 ± 0.02 . Most of the core internal dynamics could be described by models 1 and 2 (S^2 or S^2 and τ_e) (Figure S1 of the Supporting Information). S^2 order parameter values, which are used to




Figure 2. Relaxation data for wild-type Cdc42 (black circles) and Cdc42(T35A) (red circles) measured at 800 MHz. The blue shaded bars indicate residues 28–41 (Switch II), and the pink shaded bars highlight residues 57–74 (Switch 2): (A) T_1 , (B) T_2 , and (C) NOE.

characterize the mobility of backbone N–H bonds, calculated from both TENSOR and Modelfree4 were in agreement for both wild-type Cdc42 and Cdc42(T35A). The average per residue magnitude of the difference in S^2 between TENSOR and Modelfree4 was 0.031 for the wild type and 0.039 for Cdc42(T35A), both being on the same order of magnitude as the error of S^2 determination. S^2 values were consistently larger in Cdc42(T35A) relative to the corresponding S^2 values determined for the wild type (Figure 3). Thus, it appeared that some regions of Cdc42(T35A) were more conformationally constrained than the wild type, with the largest differences occurring in the Switch regions. As listed in Table 2, the overall average S^2 for both proteins was ~ 0.89 . However, the range of S^2 values for the Switch I region was 0.78–0.86 for Cdc42(T35A) compared to 0.60–0.79 for the wild type. These results indicated that the Switch I region of T35A showed some differences in the amplitudes of motion, and/or less conformational freedom, relative to the corresponding region in the wild type. NMR studies of Cdc42 in complex with PBD46 (a peptide derivative of p21-activated serine/threonine kinase) also revealed chemical shift changes that were largest within Switch I and Switch II, and the P-loop (phosphate-binding loop).¹⁷ These results may account for the inhibition of GTP hydrolysis by PBD46.¹⁷

Structure and Backbone Dynamics of Switch Regions. *Switch I.* The observation of additional backbone NH resonances suggests that the Switch I region in Cdc42(T35A), while still poorly structured, behaves differently than the corresponding region in




Figure 3. Residue-specific S^2 values extracted for wild-type Cdc42 (black circles) and Cdc42(T35A) (red circles) from the extended Lipari–Szabo Modelfree formalism. The Switch I and II regions are highlighted with a light green block around the regions. The data shown are from an analysis of the relaxation data using TENSOR.

the wild type (Figure 4). Thr³⁵ in Ras proteins is crucial for binding to effector proteins; however, it may not be directly involved in effector interactions.¹⁶ This implies that other residues in the Switch I region may be affected by the T35A mutation, as observed

in the NMR spectra. Shima et al.⁴³ have shown via X-ray structures that two conformational states for the Switch I region of H-Ras may account for its ability to interact freely with effector proteins. We were able to observe approximately 1.5 times as many backbone NH NOE restraints in the Switch I region for Cdc42(T35A) (58 total, 23 inter-residue, 8 long-range) relative to the number in the wild type (39 total, 20 inter-residue, 3 long-range). Figure 5A depicts only the backbone of the ensemble of structures from wild-type Cdc42 (PDB entry 1aje) and Cdc42(T35A) (PDB entry 2kb0). A comparison of the backbone rmsd values revealed that Switch I BBRMSD Cdc42(T35A) = 1.48 ± 0.065 , whereas Switch I BBRMSD wild-type Cdc42 = 2.95 ± 0.115 . As can be seen, the backbone of Switch I from the ensemble of structures, while still random coil in both proteins, highlights a reduced level of conformational freedom, particularly at the C-terminal end of Switch I (residues 38–41) for Cdc42(T35A). In addition, we observed new medium to weak NOEs between residues near the C-terminal end of Switch I and residues in and just adjacent to the Switch II region in T35A; for example, one observes with Cdc42-(T35A), but not with the wild type, medium and weak backbone

Table 2. Average Values of Overall Order Parameters (S^2) and Deviations from the Average Overall S^2 for the Switch I and II Regions for Wild-Type Cdc42 and Cdc42(T35A)^a

	wild-type Cdc42	Cdc42(T35A)
average S^2 protein	0.89 ± 0.04	0.89 ± 0.04
average Switch I S^2 (28–41)	0.73 ± 0.10	0.81 ± 0.03
average Switch II S^2 (57–74)	0.85 ± 0.091	0.88 ± 0.05
% ΔS^2 (Switch I) ^b	18.5%	8.6%
% ΔS^2 (Switch II) ^b	4.6%	1.5%

^aBoth proteins were in the GDP-bound form. Residue ranges of each region are given in parentheses. ^b% ΔS^2 = $|(^{avg}S^2_{protein} - ^{avg}S^2_{region}) / ^{avg}S^2_{protein}| \times 100$.

NOEs involving 39NH and 59HB, and 40HN and 55NH, and a side chain NOE between 39HD and 56HZ (aromatic proton). Mutagenesis studies and NMR measurements with Cdc42 and PAK (p21-activated serine/threonine kinase) furthermore have shown that residues 38 and 40 of Switch I are vital to the binding and maximal interaction of the two proteins.^{7,44} Both the Switch I




Figure 5. (A) Stereoview of the backbone ensemble of the Switch I region of wild-type Cdc42 and Cdc42(T35A). (B) Backbone of the Switch I region colored on the basis of best-fit dynamics model assignments from dynamics calculations. Model assignments are as follows: white for unassigned, blue for model 1 (S^2 only), green for model 2 (S^2 and τ_e), orange for model 3 (S^2 and R_{ex}), red for model 4 (S^2 , R_{ex} , and τ_e), and yellow for model 5 (S^2 , S_s^2 , and τ_e).




Figure 4. Two-dimensional ^{15}N HSQC spectrum of wild-type Cdc42 (black) overlaid with that of Cdc42(T35A) (red). The inset (top right) highlights a region of the spectrum containing resonances from three residues on the C-terminal portion of the Switch I region of Cdc42T35A (residues 38, 40, and 41). These resonances are well-defined in the Cdc42T35A spectrum but could not be detected in the HSQC spectrum of wild-type Cdc42 because of excessive line broadening.¹⁸

region and the $\beta 2$ region (residues 40–46) following Switch I have been implicated as the interface for effector protein binding with Cdc42.¹⁷ As residues in these regions cannot be observed on the NMR time scale in wild-type Cdc42, it seems plausible that the fast time scales of motions may result from multiple, interconverting conformations of Switch I made possible by the presence of the OH group of Thr³⁵ in Switch I, as has been examined previously.²⁴ This could, in turn, lead to the notion that interactions with effectors, such as PAK, are made possible by conformational selectivity⁴⁵ for a particular conformation, facilitating the strong binding interactions that occur between effector proteins and wild-type Cdc42.¹⁷ Nonetheless, our data highlight an apparent reduced flexibility that may reduce the number of possible conformations for the Switch I region in Cdc42(T35A) when the OH group of Thr³⁵ is not present. A study of the molecular details of the complex of Cdc42(T35A) and effectors, such as PAK or GTPase-binding derivatives of PAK, would be useful to confirm these possibilities.

Spoerner et al.¹⁶ suggested from kinetics and ³¹P NMR studies there are two dominant conformational states of Ras, one in the absence and another in the presence of effectors; however, in Ras(T35A), this was not the case. The arguments for two conformational states of the Switch I region were strengthened by X-ray studies of M- and H-Ras.⁴³ The increased number of observable resonances in Switch I in Cdc42(T35A) indicates that the conformational flexibility of Switch I in T35A may be more restricted than in wild-type Cdc42. This result could be due to a change in the transition rate and/or populations of the possible conformational states that have been suggested for H- and M-Ras by Shima et al.⁴³

The “Switch” binding regions in wild-type Cdc42 are unstructured and flexible, yet crucial for interactions of Cdc42 with effectors that inhibit important regulatory control mechanisms, such as GTP hydrolysis,¹⁷ a property seen also with other Ras proteins.^{46,47} Cdc42 behaves like H-, K-, and M-Ras and Rap in that, in these Ras proteins, millisecond time scale motions are seen for the Switch I region.^{15,48} We characterized the backbone dynamics of Cdc42(T35A) to discover which regions of the protein might be most influenced by the T35A mutation. Figure 5B depicts the models of motion, extracted for the observable residues of Switch I for the wild type relative to Cdc42(T35A). Even though the average overall order parameter, S^2 , is 0.89 for both proteins (Table 2), a residue-by-residue comparison leads to an average S^2 difference of -0.06 ± 0.04 between the two forms [Cdc42(T35A) vs wild-type Cdc42]. There were, furthermore, regions in which backbone order parameters differed significantly. As seen in Figure 3, the S^2 values for residues that can be observed in both wild-type Cdc42 and Cdc42(T35A) in the Switch regions are consistently larger in the mutant. For Cdc42(T35A), a number of observable residues in Switch I have an S^2 that is near the average value for the entire protein. For wild-type Cdc42, some values of S^2 are significantly lower than the average S^2 value calculated for the protein, particularly N–H bond vectors of residues 31 and 32 (32 not measured for T35A), which have S^2 values of <0.65 (Figure 3), with the internal dynamics of the N–H bond of residue 32 being characterized by model 5 (S_s^2 , S_f^2 , and τ_e). This suggests that the internal dynamics of N–H bond vectors of residues following Y³² were probably complicated and contributed to resonance broadening in the NMR spectra of the wild type. The Switch I region shows no residues with an S^2 value of <0.7 in T35A.




Figure 6. Representative in vitro binding assay comparing the binding affinities between GST-tagged Cdc42wt (left) and GST-tagged Cdc42(T35A) (right) for PBD46, a minimal GTPase-binding domain peptide for an important Ras regulatory protein, mPAK-3. PBD46 is shown as a doublet as has been shown previously.¹¹ Fractions of purified GST alone were used as nonspecific background binding control (data not shown) and did not show any appreciable binding to Cdc42 or Cdc42(T35A); however, any nonspecific binding was subtracted out by the equation

$$\Delta I = \frac{(\text{Cdc42wt} - \text{PBD} - \text{GST}) - (\text{Cdc42wt} - \text{GST})}{[\text{Cdc42(T35A)} - \text{PBD} - \text{GST}] - [\text{Cdc42(T35A)} - \text{GST}]}$$

To understand the extent of the changes in the backbone dynamics, we calculated $\% \Delta S^2$ for the observable residues of Switch I relative to the average S^2 for each entire protein (Table 2). We observed that S^2 was 18.5% lower for the Switch I residues in the wild type but only 8.6% lower for those in T35A. Previous studies have implied that Thr mutants in the Switch I region of Ras proteins exhibit a reduced affinity for effector proteins, yet Thr³⁵ is not directly involved in protein interactions in this region.¹⁶ Our results clearly show that the T35A mutation affects the backbone mobility of the Switch I region of Cdc42. As previous studies suggested, when Ser is present at position 35, two conformational states and a two-state binding model have been implied for wild-type Ras and Ras(T35S), but not seen in Ras(T35A).¹⁶ Therefore, the backbone flexibility in Switch I seems likely to be important for effector interactions for Cdc42 and other Ras proteins and is lost in Cdc42(T35A).

Single-point mutations in Switch I involving residues 35, 37, and 40 markedly reduce the strength of the association of the Ras protein with effector proteins.⁴⁹ To test whether the binding of a GTP hydrolysis inhibitory protein is altered for Cdc42(T35A) as postulated from other studies, we performed an in vitro binding assay of PBD46 with wild-type Cdc42 (Cdc42wt) and Cdc42(T35A) by measuring the intensity change of the pull-down band of Cdc42 relative to that of Cdc42(T35A) in the presence of equimolar amounts of PBD46 and compared the change in the intensity (ΔI) (Figure 6). The ΔI of the wild-type Cdc42–PBD46 complex compared to that of the Cdc42(T35A)–PBD46 complex was 4.2 ± 0.59 , consistent with previous observations of Cdc42(T35A) showing weakened binding to effector proteins.⁵⁰ We also examined the overall stability of wild-type Cdc42 relative to the Switch I mutant protein using differential scanning calorimetry (Figure 7). The apparent T_m is seemingly unaffected by the mutation as $^{Cdc42(wt)}T_m = 63.5 \pm 0.06$ °C and $^{Cdc42(T35A)}T_m = 64.2 \pm 0.11$ °C, suggesting no change in the overall conformational




Figure 7. Representative differential scanning calorimetry thermograms showing the unfolding of Cdc42wt (top) and Cdc42(T35A) (bottom). The T_m values are calculated from experiments conducted in triplicate for both proteins. The protein concentration for each protein was 1.0 mg/mL in buffer as described in Materials and Methods. Both thermograms were corrected for background noise, and each of the experiments (conducted in triplicate) were performed at a scan rate of 1.0 °C/min.

stability or folding of the constructs. The finding of similar overall folding and stability, yet reduced conformational flexibility, for Switch I in Cdc42(T35A) may help to explain some of the biological consequences of mutations in the Switch I region.⁴⁹

We also examined whether the Switch I region of Cdc42(T35A) would exhibit altered sensitivity to chymotrypsin as four of the 13 residues in Switch I are aromatic. We observed that ~90% of the parent band for Cdc42wt, but only ~38% of Cdc42(T35A), is cleaved during a 110 min exposure to chymotrypsin (Figure 8), which suggests that the Cdc42(T35A) construct is somewhat stabilized against chymotrypsin cleavage. This finding is consistent with the structural and dynamics results that showed that Switch I is more conformationally restricted in Cdc42(T35A). In summary, our results point to reduced conformational flexibility for Cdc42(T35A). The reduced flexibility in turn may lead to less efficient interactions with effector proteins.

Switch II. Switch II as well as Switch I shows increased conformational flexibility in wild-type Cdc42.¹⁹ The Switch II region in Cdc42 is also believed to serve as a part of the binding pocket for effector proteins, such as the Ser/Thr PAK kinase that inhibits GTP hydrolysis. Indeed, chemical shifts could be observed for some residues of Switch II upon binding of effector proteins to Cdc42.¹⁷ The structure and the backbone dynamics of this region are also changed by the T35A mutation in Cdc42. Indeed, several NH backbone resonances can be

identified relative to the Switch II region in the wild type. In Cdc42(T35A), only one non-proline residue (R^{68}) is unobserved, whereas the wild type is missing resonances for residues 59, 60, 66–68, 71, 72, and 74. The NOE data for Switch II resulted in more than 3 times as many observed backbone NH NOEs in Cdc42(T35A) (76 total, 45 inter-residue, 15 long-range) relative to the wild type (24 total, 13 inter-residue, 2 long-range). The region is random coil; however, unlike Switch I in T35A, the backbone residues of Switch II display similar conformational freedom like the region in the wild type, as most of the NOEs are sequential and there are no NOEs seen in Switch II of Cdc42(T35A) that are more than two residues apart in the primary sequence.

The dynamics of Switch II of Cdc42(T35A) (see Figure 3) reveal comparisons of S^2 values for the wild type and Cdc42(T35A). Thirteen residues in Switch II of the wild type have either broad, barely visible, or unassigned peaks, compared to only nine in T35A. As was true for Switch I, many Switch II residues have S^2 values that are higher in Cdc42(T35A) than in the wild type. In Switch II, many of the N–H bond vectors are best described by model 3 (S^2 and R_{ex}), with contributions of motions on the micro- to millisecond time scale. The S^2 values for observable residues in Switch II for both proteins (Table 2) show very little change relative to the overall S^2 values, as there is an only 4.6% change in the wild type and a 1.5% change in Cdc42(T35A). The small changes suggest that, while important for regulatory and effector interactions in Ras-related proteins, Switch II may serve a more complementary role in effector binding, whereas the functionally significant binding region is likely Switch I.^{16,17,43} Solution structure studies have shown that when the PAK derivative, PBD46, binds to Cdc42, the Switch II region shows some chemical shift changes, but there are no NOE constraints between the effector peptide and Switch II.^{17,21} X-ray studies have suggested that the primary importance of Switch II is direct interaction with the phosphate-binding or P-loop, as residues 59–61 show contacts with the γ - PO_4 group in Ras and Cdc42.^{10,51} These studies also argued that GTP hydrolysis results in the loss of stabilizing interactions with residues 59–61 when the γ - PO_4 group leaves, suggesting that Switch II may provide stability to the nucleotide cavity.^{51,52} In addition, structural studies using other transforming mutants have implicated a series of congruent hydrophobic interactions between Switch II and the nucleotide.⁵³ Recent ^{31}P NMR studies showed significant chemical shift changes upon PBD46 binding for the wild type (γ - ^{31}P shift of 7.473 for free vs 6.740 for bound), whereas for Cdc42(T35A), the chemical shift difference was negligible (γ - ^{31}P shift of 7.808 for free vs 7.844 for bound).¹¹ Our data suggest that, for Cdc42(T35A), the dynamics and conformational changes observed for Switch I might facilitate the weakened binding of PBD46, and thus, the Switch II– γ - PO_4 interaction in Cdc42(T35A) is minimally disturbed.

X-ray studies of transforming Switch II mutants (such as Q^{61}) of Ras have suggested a Switch II region more ordered than that in wild-type Ras.^{53,54} The studies also suggested that, in each Switch II mutant studied, Switch I and Switch II come closer together forming a more hydrophobic pocket around the nucleotide. In vitro binding studies using the Q^{61} mutant of Cdc42 with two Ras effector proteins (IQGAP1 and -2) suggested that both proteins could bind at various concentrations to this Cdc42 mutant, but that at least two Switch I mutations (Y32K and T35A) lead to no appreciable binding.⁵⁰ Moreover, structural studies of Cdc42 with Cdc42 GAPs have highlighted several




Figure 8. (A) Chymotrypsin digestion products of Cdc42wt and Cdc42(T35A). Intensity of the parent Cdc42 protein bands in lane 3 in both gels that were not treated with chymotrypsin (zero incubation time with enzyme) and assumed to be 100% protein for each gel. Initial concentrations used for each protein were the same. In each gel, corresponding Cdc42wt or Cdc42(T35A) bands were normalized to the parent band (lane 3) at ~20 kDa (denoted with an asterisk). Lanes 4–12 in each gel represent the chymotrypsin digestion products obtained after incubation with either Cdc42wt or Cdc42(T35A) for 20, 40, 50, 60, 70, 80, 90, 100, and 110 min, respectively. In each gel, lane 1 contains the MW marker and lane 2 contains chymotrypsin alone to show corresponding bands from the autolysis of chymotrypsin. (B) Graphical depiction of the percentage of undigested Cdc42wt (●) or Cdc42(T35A) (■) incubated with chymotrypsin as a function of time. Error bars represent the standard deviation of triplicate measurements.

Switch II residues (residues 61–64) that are oriented toward Switch I residues and also make direct contacts with GAPs.⁵⁵ Therefore, while Switch II may not be critical for the optimal binding of all Ras-related effectors, Switch II is nevertheless likely to be important in facilitating interactions with Switch I that indirectly support the binding of some effectors. The conformational freedom exhibited by both of these regions likely helps in effector identification for the wild-type protein. We are not able to compare through-space distances between Switch I and Switch II in wild-type Cdc42 relative to Cdc42(T35A), primarily because of the lack of observable resonances for some Switch II residues in the wild type. However, it seems plausible, in part because of the presence of long-range NOEs between portions of the two regions, that the slowed dynamics caused by the T35A mutation in Switch I facilitates interactions between portions of Switch I and Switch II that do not allow optimal interaction with effector proteins, in part because of some lost entropic component in the mutant construct.

The backbone dynamics in the Switch II region in Cdc42wt and Cdc42(T35A) are similar to that of Cdc42 as a whole. However, our studies imply that the formation of the binding pocket involving the backbone residues of Switch I and Switch II

is altered, and perhaps in a more packed environment in T35A than in the wild type. The major change again is the reduced conformational flexibility of Switch I in this mutant. The minimal changes seen in the backbone dynamics of Switch II suggest a minimal or indirect role for this region in terms of effector interactions. A more systematic investigation of Switch II, via site-directed mutagenesis, in Cdc42 with PBD46, should help confirm this hypothesis.

Clearly, the Switch regions have important features necessary for proper effector binding, and Switch I has been shown to be largely responsible for the regulation of effector binding that could potentially influence cell signaling.⁹ Cdc42 mutations that alter the intrinsic GTPase activity,⁹ facilitate anti-apoptotic events,⁵⁶ or alter interactions with regulatory proteins¹² are all involved in aberrant signaling activity. These findings have their foundation in the impact of the Switch regions on the effector interactions of Ras proteins. Therefore, Cdc42 has been and will continue to be an excellent model for probing new aspects of structure–function relationships with regard to cell signaling processes. Our characterization of the solution structure and dynamics of Cdc42(T35A) has outlined molecular details of the Switch regions, particularly Switch I, that likely account for the

importance of this region in facilitating interactions with a number of Ras-related effectors.

The Ras GTPases are promising drug targets for anticancer therapy, presumably because of the high abundance of abnormal cell signaling activities produced by mutation. Progress is evident in the development of approaches to target aberrantly functioning Ras proteins. For example, small molecule inhibitors have shown promise in impacting Ras-related cell death by inhibiting transformation events by preventing Ras proteins from attaching to the cell membrane; however, these drugs have also shown differences in selectivity for individual Ras proteins.^{57,58} Also, progress has been made in the design of small molecule inhibitors to block interactions of Rho GTPases with GEFs via screening of large databases.^{59,60} Recently, it was shown that Ras signaling in transgenic mouse models could be altered by hindering RAS–protein interactions using an antibody fragment, and that this blocked Ras protein interaction resulted in the regression of tumor growth.⁶¹ Characterization of human papilloma virus protein complexes also has suggested that alternative mechanisms for blocking HPV-related infections should include the development of ways to inhibit protein interactions.⁶² These findings highlight the importance of understanding Ras-related protein interactions that contribute to abnormal cell signaling activities that lead to proliferation and transformation. New approaches to inhibiting the transformation capability of Ras proteins should involve considerations of dynamics as well as structure. The results of our study of the solution structure and backbone dynamics of Cdc42(T35A) suggest that, in addition to strategies for blocking Ras–protein interactions with small molecules, the development of strategies to restrict the conformational flexibility of regions of Ras that interact with effectors may be a beneficial way to target Ras-related abnormal cell signaling activity. One such approach could involve the use of molecules to catalyze the formation of covalent bonds to “tie down” important interacting regions. Efforts are underway in our laboratory to develop approaches to restrict the conformational freedom of Switch I of Cdc42 without altering its overall globular conformation. This study foreshadows the importance of understanding how aberrant Ras-related protein interactions might be controlled via the restriction of conformational flexibility of the Switch region(s).

■ ASSOCIATED CONTENT

S Supporting Information. Best-fit dynamics model assignment as a function of residue for Cdc42wt and Cdc42-(T35A) (Figure S1). This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: pxa001@uark.edu. Phone: (479) 575-5621. Fax: (479) 575-4049.

Funding Sources

We are grateful for National Institutes of Health Grants 1K-01-CA113753 and ARRA/3K01CA113753 to P.D.A., National Center of Research Resources COBRE Grant 1P30RR031154-01, and the Arkansas Biosciences Institute for financial support. O.S. was supported by an Undergraduate Research Fellowship from the State of Arkansas.

■ ACKNOWLEDGMENT

We thank Drs. Roger Koeppe, II, Colin Heyes, and Robert Oswald for critical review of the manuscript.

■ ABBREVIATIONS

Cdc42, cell division cycle 42; T35A, threonine 35 to alanine; C12E6, *n*-dodecyl hexa(ethylene glycol); D₂O, deuterium oxide; PAK, p21-activated kinase; PBD46, PAK binding domain 46; IPTG, isopropyl β -D-thiogalactopyranoside; BMRB, Biological Magnetic Resonance Data Bank; HSQC, heteronuclear single-quantum coherence; TOCSY, total correlation spectroscopy; NOE, nuclear Overhauser effect; INEPT, insensitive nuclei enhanced by polarization; RDCs, residual dipolar couplings; TROSY, transverse relaxation optimized spectroscopy; GMPPCP, guanosine 5'-[(α,β)-methylene]triphosphate; rmsd, root-mean-square deviation.

■ REFERENCES

- (1) Barbacid, M. (1987) *ras* Genes. *Annu. Rev. Biochem.* 56, 779–827.
- (2) White, M. A., Nicolette, C., Minden, A., Polverino, A., Van Aelst, L., Karin, M., and Wigler, M. H. (1995) Multiple Ras functions can contribute to mammalian cell transformation. *Cell* 80, 533–541.
- (3) Bourne, H. R., Sanders, D. A., and McCormick, F. (1991) The GTPase superfamily: Conserved Structure and Molecular Mechanism. *Nature* 117–127.
- (4) Hart, M., Eva, A., Evans, T., Aaronson, S., and Cerione, R. (1991) Catalysis of Guanine Nucleotide Exchange on the Cdc42Hs Protein by the *dbl* Oncogene Product. *Nature* 354, 311–314.
- (5) Horii, Y., Beeler, J. F., Sakaguchi, K., Tachibana, M., and Miki, T. (1994) A novel oncogene, *ost*, encodes a guanine nucleotide exchange factor that potentially links Rho and Rac signaling pathways. *EMBO J.* 13, 4776–4786.
- (6) Hart, M., Shinjo, K., Hall, A., Evans, T., and Cerione, R. A. (1991) Identification of the Human Platelet GTPase Activating Protein for the CDC42Hs Protein. *J. Biol. Chem.* 266, 20840–20848.
- (7) Leonard, D., Hart, M. J., Platko, J. V., Eva, A., Henzel, W., Evans, T., and Cerione, R. A. (1992) The identification and characterization of a GDP-dissociation inhibitor (GDI) for the CDC42Hs protein. *J. Biol. Chem.* 267, 22860–22868.
- (8) Bos, J. L., Toksoz, D., Marshall, C. J., Verlaan-de Vries, M., Veeneman, G. H., van der Eb, A. J., van Boom, J. H., Janssen, J. W., and Steenvoorden, A. C. (1985) Amino-acid substitutions at codon 13 of the N-ras oncogene in human acute myeloid leukaemia. *Nature* 315, 726–730.
- (9) Miller, A.-F., Halkides, C., and Redfield, A. (1993) An NMR Comparison of the Changes Produced by Different Guanosine 5'-Triphosphate Analogs in Wildtype and Oncogenic Mutant p21^{ras}. *Biochemistry* 32, 7367–7376.
- (10) Pai, E. F., Krengel, U., Petsko, G. A., Goody, R. S., Kabsch, W., and Wittinghofer, A. (1990) Refined crystal structure of the triphosphate conformation of h-ras p21 at 1.35 Å resolution: Implication for the mechanism of GTP hydrolysis. *EMBO J.* 9, 2351–2359.
- (11) Phillips, M. J., Calero, G., Chan, B., Ramachandran, S., and Cerione, R. A. (2008) Effector proteins exert an important influence on the signaling-active state of the small GTPase Cdc42. *J. Biol. Chem.* 283, 14153–14164.
- (12) McCormick, F. (1989) ras GTPase activating protein: Signal transmitter and signal terminator. *Cell* 56, 5–8.
- (13) Lin, R., Bagrodia, S., Cerione, R., and Manor, D. (1997) A novel Cdc42Hs mutant induces cellular transformation. *Curr. Biol.* 7, 794–797.
- (14) Adams, P. D., and Oswald, R. E. (2006) Solution structure of an oncogenic mutant of Cdc42Hs. *Biochemistry* 45, 2577–2583.
- (15) Geyer, M., Schweins, T., Herrmann, C., Prisner, T., Wittinghofer, A., and Kalbitzer, H. R. (1996) Conformational transitions in p21^{ras} and in its complexes with the effector protein Raf-RBD and the GTPase activating protein GAP. *Biochemistry* 35, 10308–10320.

- (16) Spoerner, M., Herrmann, C., Vetter, I. R., Kalbitzer, H. R., and Wittinghofer, A. (2001) Dynamic properties of the Ras switch I region and its importance for binding to effectors. *Proc. Natl. Acad. Sci. U.S.A.* 98, 4944–4949.
- (17) Guo, W., Sutcliffe, M. J., Cerione, R. A., and Oswald, R. E. (1998) Identification of the binding surface on Cdc42Hs for p21-activated kinase. *Biochemistry* 37, 14030–14037.
- (18) Feltham, J. L., Dotsch, V., Raza, S., Manor, D., Cerione, R. A., Sutcliffe, M. J., Wagner, G., and Oswald, R. E. (1997) Definition of the switch surface in the solution structure of Cdc42Hs. *Biochemistry* 36, 8755–8766.
- (19) Loh, A. P., Guo, W., Nicholson, L. K., and Oswald, R. E. (1999) Backbone dynamics of inactive, active, and effector-bound Cdc42Hs from measurements of ¹⁵N relaxation parameters at multiple field strengths. *Biochemistry* 38, 12547–12557.
- (20) Mott, H. R., Owen, D., Nietlispach, D., Lowe, P. N., Manser, E., Lim, L., and Laue, E. D. (1999) Structure of the small G protein Cdc42 bound to the GTPase-binding domain of ACK. *Nature* 399, 384–388.
- (21) Rittinger, K., Walker, P. A., Eccleston, J. F., Nurmahomed, K., Owen, D., Laue, E., Gamblin, S. J., and Smerdon, S. J. (1997) Crystal structure of a small G protein in complex with the GTPase-activating protein rhoGAP. *Nature* 388, 693–697.
- (22) Gladfelter, A. S., Moskow, J. J., Zyla, T. R., and Lew, D. J. (2001) Isolation and characterization of effector-loop mutants of CDC42 in yeast. *Mol. Biol. Cell* 12, 1239–1255.
- (23) Tanaka, T., and Rabbitts, T. H. (2010) Interfering with RAS-effector protein interactions prevent RAS-dependent tumour initiation and causes stop-start control of cancer growth. *Oncogene* 29, 6064–6070.
- (24) Adams, P. D., and Oswald, R. E. (2007) NMR assignment of Cdc42(T35A), an active Switch I mutant of Cdc42. *Biomol. NMR Assignments* 1, 225–227.
- (25) States, D., Haberkorn, R., and Ruben, D. (1982) A two-dimensional nuclear Overhauser experiment with pure absorption phase in four quadrants. *J. Magn. Reson.* 48, 282–292.
- (26) Delaglio, F., Grzesiek, S., Vuister, G. W., Zhu, G., Pfeifer, J., and Bax, A. (1995) NMRPipe: A multidimensional spectral processing system based on UNIX pipes. *J. Biomol. NMR* 6, 277–293.
- (27) Goddard, T. D., and Kneller, D. G. (2007) SPARKY 3, University of California, San Francisco.
- (28) Brunger, A. T., Adams, P. D., Clore, G. M., DeLano, W. L., Gros, P., Grosse-Dunstleve, R. W., Jiang, J. S., Kuszewski, J., Nilges, M., Pannu, N. S., Read, R. J., Rice, L. M., Simonson, T., and Warren, G. L. (1998) Crystallography and NMR system: A New Software suite for Macromolecular Structure Determination. *Acta Crystallogr. D* 54, 905–921.
- (29) Wishart, D. S., Sykes, B. D., and Richards, F. M. (1991) Relationship between nuclear magnetic resonance chemical shift and protein secondary structure. *J. Mol. Biol.* 222, 311–333.
- (30) Wishart, D. S., Sykes, B. D., and Richards, F. M. (1992) The chemical shift index: A fast and simple method for the assignment of protein secondary structure through NMR spectroscopy. *Biochemistry* 31, 1647–1651.
- (31) Zweckstetter, M., and Bax, A. (2000) Prediction of Sterically Induced Alignment in a Dilute Liquid Crystalline Phase: Aid to Protein Structure Determination by NMR. *J. Am. Chem. Soc.* 122, 3791–3792.
- (32) Guex, N., and Peitsch, M. C. (1997) SWISS-MODEL and the Swiss-PdbViewer: An environment for comparative protein modeling. *Electrophoresis* 18, 2714–2723.
- (33) Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C., and Ferrin, T. E. (2004) UCSF Chimera: A visualization system for exploratory research and analysis. *J. Comput. Chem.* 25, 1605–1612.
- (34) Laskowski, R. A., Rullmann, J. A., MacArthur, M. W., Kaptein, R., and Thornton, J. M. (1996) AQUA and PROCHECK-NMR: Programs for checking the quality of protein structures solved by NMR. *J. Biomol. NMR* 8, 477–486.
- (35) Laskowski, R. A., MacArthur, M. W., Moss, D. S., and Thornton, J. M. (1993) PROCHECK: A Program to Check the Stereochemical Quality of Protein Structures. *J. Appl. Crystallogr.* 26, 283–291.
- (36) Kelley, L. A., and Sutcliffe, M. J. (1997) OLDERADO: On-Line Database of Ensemble Representatives And DOfains. *Protein Sci.* 6, 2628–2630.
- (37) Kay, L., Nicholson, L., Delaglio, F., Bax, A., and Torchia, D. (1992) Pulse sequences for removal of the effects of cross-correlation between dipolar and chemical-shift anisotropy relaxation mechanism on the measurement of heteronuclear T1 and T2 values in proteins. *J. Magn. Reson.* 97, 359–375.
- (38) Kay, L., Torchia, D., and Bax, A. (1989) Backbone Dynamics of Proteins as Studied by ¹⁵N Inverse Detected Heteronuclear NMR Spectroscopy: Application to Staphylococcal Nuclease. *Biochemistry* 28, 8972–8979.
- (39) Dossset, P., Hus, J. C., Blackledge, M., and Marion, D. (2000) Efficient analysis of macromolecular rotational diffusion from heteronuclear relaxation data. *J. Biomol. NMR* 16, 23–28.
- (40) Palmer, A. (1998) ModelFree, version 4.0, <http://cpmcnet.columbia.edu/dept/gsas/biochem/labs/palmer>.
- (41) Adams, P. D., Loh, A. P., and Oswald, R. E. (2004) Backbone dynamics of an oncogenic mutant of Cdc42Hs shows increased flexibility at the nucleotide-binding site. *Biochemistry* 43, 9968–9977.
- (42) Lipari, G., and Szabo, A. (1982) Model-free Approach to the Interpretation of Nuclear Magnetic Resonance Relaxation in Macromolecules. 1. Theory and Range of Validity. *J. Am. Chem. Soc.* 104, 4546–4559.
- (43) Shima, F., Ijiri, Y., Muraoka, S., Liao, J., Ye, M., Araki, M., Matsumoto, K., Yamamoto, N., Sugimoto, T., Yoshikawa, Y., Kumasaka, T., Yamamoto, M., Tamura, A., and Kataoka, T. (2010) Structural basis for conformational dynamics of GTP-bound Ras protein. *J. Biol. Chem.* 285, 22696–22705.
- (44) Lamarche, N., Tapon, N., Stowers, L., Burbelo, P. D., Aspenstroem, P., Bridges, T., Chant, J., and Hall, A. (1996) Rac and Cdc42 induce action polymerization and G1 cell cycle progression independently of p65 and the JNK/SAPK MAP kinase cascade. *Cell* 87, 519–529.
- (45) Hammes, G. G., Chang, Y. C., and Oas, T. G. (2009) Conformational selection or induced fit: A flux description of reaction mechanism. *Proc. Natl. Acad. Sci. U.S.A.* 106, 13737–13741.
- (46) Abdul-Manan, N., Aghazadeh, B., Liu, G. A., Majumdar, A., Ouerfelli, O., Siminovich, K. A., and Rosen, M. K. (1999) Structure of Cdc42 in complex with the GTPase-binding domain of the 'Wiskott-Aldrich syndrome' protein. *Nature* 399, 379–383.
- (47) Gizachew, D., Guo, W., Chohan, K. K., Sutcliffe, M. J., and Oswald, R. E. (2000) Structure of the complex of Cdc42Hs with a peptide derived from P-21 activated kinase. *Biochemistry* 39, 3963–3971.
- (48) Liao, J., Shima, F., Araki, M., Ye, M., Muraoka, S., Sugimoto, T., Kawamura, M., Yamamoto, N., Tamura, A., and Kataoka, T. (2008) Two conformational states of Ras GTPase exhibit differential GTP-binding kinetics. *Biochem. Biophys. Res. Commun.* 369, 327–332.
- (49) Fiordalisi, J. J., Holly, S. P., Johnson, R. L., Il, Parise, L. V., and Cox, A. D. (2002) A distinct class of dominant negative Ras mutants: Cytosolic GTP-bound Ras effector domain mutants that inhibit Ras signaling and transformation and enhance cell adhesion. *J. Biol. Chem.* 277, 10813–10823.
- (50) McCallum, S. J., Wu, W. J., and Cerione, R. A. (1996) Identification of a putative effector for Cdc42Hs with high sequence similarity to the RasGAP-related protein IQGAP1 and a Cdc42Hs binding partner with similarity to IQGAP2. *J. Biol. Chem.* 271, 21732–21737.
- (51) Pai, E. F., Kabsch, W., Krengel, U., Holmes, K. C., John, J., and Wittinghofer, A. (1989) Structure of the guanine nucleotide binding domain of the Ha-Ras oncogene product p21 in the triphosphate conformation. *Nature* 341, 209–214.
- (52) Stouten, P. F., Sander, C., Wittinghofer, A., and Valencia, A. (1993) How does the switch II region of G-domains work? *FEBS Lett.* 320, 1–6.
- (53) Buhrman, G., Wink, G., and Mattos, C. (2007) Transformation efficiency of RasQ61 mutants linked to structural features of the switch regions in the presence of Raf. *Structure* 15, 1618–1629.

- (54) Krengel, U., Schlichting, L., Scherer, A., Schumann, R., Frech, M., John, J., Kabsch, W., Pai, E., and Wittinghofer, A. (1990) Three-Dimensional Structures of H-Ras p21 Mutants: Molecular Basis for their Inability to Function as Signal Switch Molecules. *Cell* 62, 539–548.
- (55) Nassar, N., Hoffman, G. R., Manor, D., Clardy, J. C., and Cerione, R. A. (1998) Structures of Cdc42 bound to the active and catalytically compromised forms of Cdc42GAP. *Nat. Struct. Biol.* 5, 1047–1052.
- (56) Tu, S. S., Wu, W. J., Yang, W., Nolbant, P., Hahn, K., and Cerione, R. A. (2002) Antiapoptotic Cdc42 mutants are potent activators of cellular transformation. *Biochemistry* 41, 12350–12358.
- (57) Lebowitz, P. F., and Prendergast, G. C. (1998) Non-Ras targets of farnesyltransferase inhibitors: Focus on Rho. *Oncogene* 17, 1439–1445.
- (58) Schmidmaier, R., Baumann, P., Simsek, M., Dayyani, F., Emmerich, B., and Meinhardt, G. (2004) The HMG-CoA reductase inhibitor simvastatin overcomes cell adhesion-mediated drug resistance in multiple myeloma by geranylgeranylation of Rho protein and activation of Rho kinase. *Blood* 104, 1825–1832.
- (59) Gao, Y., Dickerson, J. B., Guo, F., Zheng, J., and Zheng, Y. (2004) Rational design and characterization of a Rac GTPase-specific small molecule inhibitor. *Proc. Natl. Acad. Sci. U.S.A.* 101, 7618–7623.
- (60) Nassar, N., Cancelas, J., Zheng, J., Williams, D. A., and Zheng, Y. (2006) Structure-function based design of small molecule inhibitors targeting Rho family GTPases. *Curr. Top. Med. Chem.* 6, 1109–1116.
- (61) Yagi, R., Tanaka, M., Sasaki, K., Kamata, R., Nakanishi, Y., Kanai, Y., and Sakai, R. (2010) ARAP3 inhibits peritoneal dissemination of scirrhous gastric carcinoma cells by regulating cell adhesion and invasion. *Oncogene* 30, 1413–1421.
- (62) Chi, C. N., Bach, A., Engstrom, A., Stromgaard, K., Lundstrom, P., Ferguson, N., and Jemth, P. (2011) Biophysical Characterization of the Complex between Human Papillomavirus E6 Protein and Synapse-associated Protein 97. *J. Biol. Chem.* 286, 3597–3606.
- (63) Loh, A. P., Pawley, N., Nicholson, L. K., and Oswald, R. E. (2001) An Increase in Side Chain Entropy Facilitates Effector Binding: NMR Characterization of the Side Chain Methyl Group Dynamics in Cdc42Hs. *Biochemistry* 40, 4590–4600.