

Two Functional S100A4 Monomers Are Necessary for Regulating Nonmuscle Myosin-IIA and HCT116 Cell Invasion


Reniqua P. House,[†] Maria Pozzuto,[†] Purvi Patel,[†] Natalya G. Dulyaninova,[†] Zhong-Hua Li,[†] Wendy D. Zencheck,[†] Michele I. Vitolo,[‡] David J. Weber,[‡] and Anne R. Bresnick^{*,†}

[†]Department of Biochemistry, Albert Einstein College of Medicine, 1300 Morris Park Avenue, Bronx, New York 10461, United States

[‡]Department of Biochemistry and Molecular Biology, University of Maryland School of Medicine, 108 North Greene Street, Baltimore, Maryland 21201, United States

Supporting Information

ABSTRACT: S100A4, a member of the Ca^{2+} -activated S100 protein family, regulates the motility and invasiveness of cancer cells. Moreover, high S100A4 expression levels correlate with poor patient survival in several cancers. Although biochemical, biophysical, and structural data indicate that S100A4 is a noncovalent dimer, it is unknown if two functional S100A4 monomers are required for the productive recognition of protein targets and the promotion of cell invasion. To address this question, we created covalently linked S100A4 dimers using a glycine rich flexible linker. The single-chain S100A4 (sc-S100A4) proteins exhibited wild-type affinities for calcium and nonmuscle myosin-IIA, retained the ability to regulate nonmuscle myosin-IIA assembly, and promoted tumor cell invasion when expressed in S100A4-deficient colon carcinoma cells. Mutation of the two calcium-binding EF-hands in one monomer, while leaving the other monomer intact, caused a 30–60-fold reduction in binding affinity for nonmuscle myosin-IIA concomitant with a weakened ability to regulate the monomer–polymer equilibrium of nonmuscle myosin-IIA. Moreover, sc-S100A4 proteins with one monomer deficient in calcium responsiveness did not support S100A4-mediated colon carcinoma cell invasion. Cross-linking and titration data indicate that the S100A4 dimer binds a single myosin-IIA target peptide. These data are consistent with a model in which a single peptide forms interactions in the vicinity of the canonical target binding cleft of each monomer in such a manner that both target binding sites are required for the efficient interaction with myosin-IIA.



S100A4 is a member of the S100 Ca^{2+} -binding family of proteins. The S100 designation is derived from the observation that these proteins are soluble in 100% saturated ammonium sulfate at neutral pH.¹ At present, there are 21 identified human S100 family members,^{2,3} which make up a subclass of the larger EF-hand-containing family of proteins represented by calmodulin and troponin C.^{4,5} The S100 proteins are 25–65% identical with one another in terms of amino acid sequence and are expressed solely in vertebrates in a cell and tissue specific manner.^{2,6,7} This family of proteins is involved in diverse cellular processes and contributes to various disease states such as neurodegeneration, inflammation, and cancer.^{2,8,9}

The S100 proteins are primarily α -helical, and most family members form noncovalent antiparallel dimers.^{10–15} The dimerization interface is composed of an X-type four-helix bundle that is comprised of helices 1 and 4 from each monomer.^{10–15} Each S100 monomer contains two helix–loop–helix (EF-hand) Ca^{2+} -binding motifs connected by a hinge region: a 14-residue N-terminal pseudo-EF-hand and a 12-residue C-terminal canonical EF-hand.^{10–15} The canonical EF-hand (EF2) uses several side chain oxygen atoms, an oxygen

from a water molecule, and two oxygen atoms from a glutamate side chain to coordinate the Ca^{2+} ion, whereas the N-terminal EF-hand (EF1) coordinates Ca^{2+} via several backbone carbonyl oxygens, an oxygen from water, and two oxygen atoms from a glutamate side chain.¹⁶ Consistent with these structural data, the reported Ca^{2+} equilibrium dissociation constants (K_D) for S100A4 indicate that EF1 has a weak affinity for calcium ($K_D > 500 \mu\text{M}$), whereas EF2 has a relatively high affinity for Ca^{2+} ($K_D = 2.6 \pm 1.0 \mu\text{M}$).¹⁷ Most S100 proteins, including S100A4, undergo a significant conformational rearrangement following binding of Ca^{2+} to the C-terminal EF-hand. This conformational rearrangement increases the interhelical angle between helices 3 and 4 and exposes a hydrophobic target binding cleft comprised of residues from helix 3, helix 4, and the hinge region.

The biological function of S100A4 has been characterized most extensively in the context of cancer. S100A4 overexpression in benign rat mammary tumor cells confers a

Received: April 4, 2011

Revised: June 30, 2011

Published: July 1, 2011



metastatic phenotype with no apparent effect on tumor proliferation.¹⁸ In addition, overexpression of S100A4 in the mammary epithelium of MMTV-neu mice, which normally have a low incidence of metastasis, increased the incidence of metastasis by 72%.¹⁹ Conversely, the loss of S100A4 expression in highly metastatic MMTV-PyMT mice resulted in a 60% decrease in the number of metastatic lesions in the lungs.²⁰ Consistent with these animal studies, S100A4 expression in breast cancer patients correlates with patient death.²¹

Intracellular S100A4 targets include the p53 homologues p63 and p73, nonmuscle tropomyosin, liprin β 1, and nonmuscle myosin-IIA.^{22–25} For the S100A4–myosin-IIA interaction, S100A4 binding disrupts the monomer–polymer equilibrium of myosin-IIA by promoting the disassembly of preexisting filaments or by inhibiting the assembly of monomers into filaments.^{26–28} Moreover, S100A4 modulates the polarization and motility of mammary tumor cells through a direct interaction with myosin-IIA.²⁹ Given the importance of myosin-II in modulating cell migration, the regulation of myosin-IIA assembly by S100A4 likely contributes to the ability of S100A4 to promote tumor invasion and metastasis.³⁰

Although there has been some examination of the molecular determinants involved in S100A4 dimerization,^{31–33} it is not known whether both monomers of the S100A4 homodimer are required for the recognition of its protein targets and the promotion of cell invasion. Using covalently linked S100A4 monomers or single-chain S100A4 proteins (sc-S100A4), we demonstrate that two functional S100A4 monomers are necessary for the efficient interaction of S100A4 with myosin-IIA and for the promotion of tumor cell invasion.

EXPERIMENTAL PROCEDURES

Creation of sc-S100A4 Constructs. To create the sc-S100A4 constructs, the S100A4 cDNA was amplified using a primer to the 3' end that contained the DNA sequence for the linker and a BamHI restriction site (monomer 1). The S100A4 polymerase chain reaction (PCR) product containing the linker sequence was cloned into pET23a using the NdeI and BamHI sites, and monomer 2 was cloned into the BamHI and HindIII sites. The resulting sc-S100A4 constructs were subcloned into pET26b using the NdeI and HindIII sites. The final linker sequence between the two S100A4 monomers was (GGGGS)_nGS.

The EF-hand substitutions were created in the wild-type S100A4 pET23a construct using the QuikChange site-directed mutagenesis kit (Stratagene). Reverse primers contained either the E33A or E74A substitution. Single-EF-hand substitutions were introduced first, and the E33A/E74A double-EF-hand substitutions were introduced by subcloning the E33A mutant into the E74A mutant S100A4 construct using the NdeI and XmaI sites. To incorporate the EF-hand substitutions into the sc-S100A4 constructs, we amplified E33A/E74A S100A4 with primer sets containing the appropriate linker addition and subcloned it as described above.

The sc-S100A4–GFP constructs were created by ligation-independent cloning.^{34,35} The pEGFP-N3 vector and S100A4 primers were as follows: vector forward primer, 5'TCTGTGCTGTCGTCGGTACCGCGGGCCC3'; vector reverse primer, 5'TGGTGCTGCTTCCGTCCGGTAGCGC-TAGCGGAT3'; S100A4 forward primer, 5'ACCGGACGGAAG-CAGCACCATGGCGTGCCTCTG3'; and S100A4 reverse primer, 5'ACCGACGACAGCAACAGATTTCTTCCTGGG3'.

pEGFP-N3 was amplified using KOD polymerase (Novagen), and S100A4 was amplified using a standard PCR protocol. The resulting PCR products were treated with T4 polymerase in separate reaction mixtures containing either 2.5 mM dATP (pEGFP-N3) or 2.5 mM dTTP (S100A4). For ligation-independent cloning, the reaction mixture contained equimolar amounts of T4-digested S100A4 and pEGFP-N3 and the reaction was performed at 22 °C for 25 min. The reaction was quenched by the addition of 10 mM EDTA (pH 8.0) and incubation at 22 °C for 5 min.

Cloning of the S100A4–GFP Fusion Protein. GFP was fused to the C-terminus of S100A4 using primer pairs that removed the S100A4 stop codon and added a 5' HindIII site and a 3' SacII site. The resulting PCR product was subcloned into the pEGFP-N3 (Clontech) vector using these sites.

Peptides. Myosin-IIA peptides [FITC-Ahx-DAMNR-EVSSLKNKLRR-CONH₂ (FITC-MIIA^{1908–1923}) and FITC-Ahx-TETADAMNREVSSLKNKLRRGDLP-CONH₂ (FITC-MIIA^{1904–1927})] were synthesized by Biosynthesis Inc. (Lewisville, TX). The molecular weight and purity (>95%) of all peptides were verified using mass spectrometry and high-performance liquid chromatography (HPLC), respectively. The concentration of all peptide solutions was determined using the FITC moiety.

Protein Purification. MIIA^{1851–1960} and MIIA^{1338–1960} were expressed and purified as described previously.^{16,27} The wild-type sc-S100A4 constructs and S100A4–GFP fusion protein were expressed and purified as described for the native S100A4 dimer.¹⁴ The masses of the sc-S100A4 proteins and S100A4–GFP fusion protein were confirmed by mass spectrometry.

For purification of the EF-hand substitutions, BL21(DE3)* cells were transformed and grown in autoinduction TB medium for 20 h at 37 °C.³⁶ The cells were pelleted and resuspended in 10 mL of lysis buffer [50 mM Tris (pH 7.5), 10% glycerol, 300 mM KCl, 2 mM DTT, 1 mM EDTA, 1 mM PMSF, and chymostatin, leupeptin, and pepstatin (5 μ g/mL each)] per gram of cell pellet. Resuspended cell pellets were sonicated and then centrifuged at 16000g for 30 min. Ammonium sulfate was added to the supernatant to 30% saturation, and the sample was centrifuged at 16000g for 10 min. CaCl₂ (2 mM) was added to the supernatant, and the sample was applied to a Phenyl-Sepharose (GE Healthcare) column equilibrated with buffer 1 [20 mM Tris (pH 7.5), 2 mM CaCl₂, 300 mM KCl, 1 mM DTT, 1 mM EDTA, 16.4 g/100 mL ammonium sulfate, and 0.02% NaN₃]. The column was washed with 3 column volumes of buffer 1 and washed with 3 column volumes of buffer 2 (buffer 1 without ammonium sulfate). sc-S100A4 EF-hand mutant proteins were eluted using buffer 3 [20 mM Tris (pH 7.5), 5 mM EGTA, 1 mM DTT, 1 mM EDTA, and 0.02% NaN₃]. Collected protein fractions were dialyzed against 20 mM PIPES (pH 6.8), 20 mM KCl, 0.5 mM DTT, and 0.02% NaN₃, centrifuged at 16000g for 15 min, and applied to a Mono S column (GE Healthcare) equilibrated with dialysis buffer. The column was washed with 10 column volumes of dialysis buffer, and sc-S100A4 EF-hand mutant proteins were eluted using a 0 to 0.5 M KCl gradient. Pooled protein fractions were dialyzed against 20 mM Tris (pH 7.5), 20 mM KCl, 1.5 mM DTT, and 0.02% NaN₃. The dialyzed protein was centrifuged for 15 min at 16000g and concentrated using a Vivaspin 3 kDa cutoff concentrator (Sartorius). Concentrated sc-S100A4 EF-hand mutant proteins were applied to a Superdex 75 gel

filtration column (GE Healthcare) equilibrated with dialysis buffer. On the basis of sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–PAGE), pure protein fractions were pooled and concentrated as described above. All protein concentrations were determined using the Bio-Rad Protein Assay (Bio-Rad) and a S100A4 standard at a known concentration.

Analytical Size Exclusion Chromatography. Wild-type S100A4 (0.7–15 μ M), 15 μ M sc-S100A4 proteins, or 1 mg/mL protein standards [BSA (67 kDa), ovalbumin (43 kDa), chymotrypsin (27 kDa), and ribonuclease A (13 kDa)] were injected onto a Superdex 75HR 10/30 column (GE Healthcare) equilibrated with 20 mM Tris (pH 7.5), 150 mM KCl, 2 mM CaCl_2 , 1 mM DTT, and 0.02% NaN_3 . For studies examining the elution of apo-S100A4, the calcium was replaced with 2 mM EGTA. Absorbance at 280 nm was used to monitor protein elution.

Chemical Denaturation. Guanidine hydrochloride (GdnHCl) stocks (7 M) were made using EGTA buffer [20 mM Tris (pH 7.5), 20 mM KCl, 2 mM EGTA, and 0.02% NaN_3] or calcium buffer in which the EGTA was replaced with 2 mM CaCl_2 . S100A4 proteins were dialyzed into either EGTA or calcium buffer, and 5 μ M S100A4 dimer was used per 200 μ L reaction mixture with fresh DTT (1 mM) added the day of the experiment. Reaction mixtures contained GdnHCl concentrations ranging from 0 to 6.5 M. Individual reaction mixtures were incubated for 22 h at 22 °C. The intrinsic fluorescence of the two S100A4 tyrosine residues was monitored using a Fluoromax-3 spectrofluorometer (Jobin Yvon Inc.). The data were fit to a sigmoidal dose–response curve using GraphPad Prism version 5.0a (GraphPad Software Inc.), and the fraction unfolded and $D_{1/2}$ were calculated using formulas described previously.³⁷

Circular Dichroism. S100A4 proteins were dialyzed into 20 mM Tris (pH 7.5), 20 mM KCl, 1 mM DTT, and 0.02% NaN_3 . Either 0.3 mM CaCl_2 or 2 mM EGTA was added to each reaction mixture containing 6.8 μ M S100A4 dimer and the mixture incubated for 1 h prior to the acquisition of spectra. Samples were placed into a 0.1 cm cuvette, and spectra from 190 to 250 nm were obtained at 22 °C using a Jasco (Easton, MD) J-18 spectrometer. Molar ellipticity values were calculated using the formula $\theta_{\text{molar}} = (\theta \times \text{MW} \times 100) / (lc)$, where θ is the measured ellipticity in degrees, MW is the molecular weight of the S100A4 dimer, 100 is the conversion of molar concentration to decimoles per cubic centimeter, l is the path length of the cuvette in centimeters, and c is the protein concentration in milligrams per milliliter.³⁸

Calcium Binding. Calcium affinities were determined as described previously.^{17,39} Briefly, S100A4 and MIIA^{1851–1960} were dialyzed into calcium free buffer [20 mM Tris (pH 7.5), 150 mM KCl, 1 mM DTT, and 0.02% NaN_3] that had been pretreated with Chelex 100 resin. Calcium was titrated into reaction mixtures containing 12.5 μ M S100A4 dimer and 25 μ M 5,5'-Br₂-BAPTA alone or in the presence 100 μ M MIIA^{1851–1960} monomer, and the absorbance was monitored at 263 nm. Using Caligator, macroscopic binding affinities were obtained by fitting the resulting titration curves to two Ca^{2+} -binding sites using a stepwise macroscopic binding equation in the presence of chelator.⁴⁰

Myosin-IIA Binding Assay. Binding affinities of S100A4 for myosin-IIA were determined using a fluorescence anisotropy assay as described previously.¹⁷ FITC-MIIA^{1908–1923}, which comprises the minimal S100A4 binding

site, was used to monitor the interaction of the wild-type and sc-S100A4 proteins and myosin-IIA. Reaction mixtures (200 μ L) contained 100 nM FITC-MIIA^{1908–1923} and 0–20 μ M wild-type or sc-S100A4 dimer, or 0–500 μ M EF-hand mutant sc-S100A4 in 20 mM Tris (pH 7.5), 150 mM KCl, 1 mM DTT, 0.02% NaN_3 , and 0.5 mM CaCl_2 . Anisotropy was measured with excitation at 494 nm and emission at 516 nm. Dissociation constants were obtained by fitting to a single-site saturation binding equation with a floating Y_{min} value using GraphPad Prism.

For fluorescence anisotropy titrations, 1 μ M FITC-MIIA^{1904–1927} was titrated with 0–8 μ M S100A4 dimer in 20 mM Tris (pH 7.5), 150 mM KCl, 1 mM DTT, 0.02% NaN_3 , and 0.5 mM CaCl_2 . Linear regression was used to determine the intersecting points of the titration curve.

Promotion of Disassembly. The ability of wild-type S100A4, sc-S100A4–5, sc-S100A4–10, and sc-S100A4 EF-hand mutant proteins to regulate myosin-IIA disassembly was assessed as described previously.²⁷ Briefly, reaction mixtures containing 1.5 μ M MIIA^{1338–1960} dimer and 0–15 μ M S100A4 dimer in 20 mM Tris (pH 7.5), 300 μ M CaCl_2 , 1.5 mM MgCl_2 , 150 mM NaCl, 1 mM DTT, and 0.02% NaN_3 were incubated for 1 h at 22 °C and centrifuged at 80000 rpm for 15 min using an Optima TLX ultracentrifuge (Beckman). A sample of the mixture was obtained by removing a 10 μ L aliquot after incubation for 10 min. The 5 \times sample buffer was added to aliquots of the mix and the supernatant, and they were loaded onto a 12% Tris-Tricine gel. The amount of myosin-IIA in the supernatant was quantified by densitometric analysis of the Coomassie-stained protein bands.

Cross-Linking Experiments. Wild-type S100A4 and MIIA^{1851–1960} were dialyzed into 20 mM HEPES (pH 7.5), 150 mM KCl, 1 mM DTT, and 0.02% NaN_3 . Reaction mixtures containing equimolar concentrations of S100A4 monomer and MIIA^{1851–1960} monomer (either 25 or 50 μ M of each) were incubated with either 0.5 mM CaCl_2 or 4 mM EGTA for 60 min at 22 °C. Disuccinimidyl tartrate (DST) was added to a final concentration of 1.2 mM, and the reaction mixture was incubated for an additional 60 min. Cross-linking reactions with S100A4 alone or MIIA^{1851–1960} alone were performed in the presence of 0.5 mM CaCl_2 . Reactions were quenched by the addition of 5 \times Laemmli gel loading dye or 50 mM Tris-HCl (pH 7.5). Samples were run on a 12% Tris-Tricine gel and stained with Coomassie brilliant blue. For immunoblot analysis, the 12% Tris-Tricine gel was transferred to a polyvinylidene difluoride (PVDF) membrane (Millipore) and reacted with antibodies against S100A4 or the myosin-IIA C-terminus.^{41,42}

Cross-linked S100A4–myosin-IIA complexes were evaluated by LC-ESI MS. Seventeen micrograms of the protein complexes was separated on a 1.0 mm \times 50 mm C3 column (MicroTech Scientific) attached to an LC Packings Ultimate Plus HPLC system. The column was equilibrated in buffer A (2% acetonitrile and 0.1% formic acid). After a 5 min desalt, protein complexes were separated using a 15 min gradient from 15 to 60% buffer B (90% acetonitrile and 0.1% formic acid) with a flow rate of 75 μ L/min. The effluent was directly delivered into an LTQ Linear Ion Trap (Thermo) for mass analysis, which has an error of ± 5 Da for samples in this mass range.

Cell Culture. HCT116 parental and S100A4^{−/−} cells, which will be described elsewhere, were maintained in McCoy's 5A medium (Cellgro) supplemented with 10% fetal bovine

serum and 1% penicillin/streptomycin. BAC1.2F5 macrophages were maintained in α MEM medium supplemented with L-asparagine, L-glutamine, 10% FBS, and 3000 units/mL human recombinant CSF-1.

Transfection. HCT116 cells were plated in a 10 cm dish 24 h prior to transfection at a cell density that yielded approximately 40% cell confluence the next day. Cells were transfected with S100A4 plasmid DNA using the HeLaMON-STER reagent (Mirus). Maximal expression was observed 48 h post-transfection.

Immunoblots. Whole cell lysates were prepared from HCT116 cell lines at a cell density of 70–80% confluence. Cells were lysed in RIPA buffer [50 mM Tris, 150 mM NaCl, 0.5% deoxycholate, 0.1% SDS, and leupeptin, chymostatin, and pepstatin (5 μ g/mL each)], and total protein was quantified using the Dc Bio-Rad protein assay. Whole cell lysates (20 μ g) were separated on a 12% Tris-Tricine gel, transferred to a PVDF membrane, and immunoblotted for β -actin and S100A4. Immunoreactive proteins were imaged using the West Pico chemiluminescence substrate (Pierce).

3D Invasion Assays. Invasion assays were conducted as described previously.⁴³ Briefly, 80000 HCT116 cells labeled with Cell Tracker Green CMFDA and 200000 BAC1.2F5 cells labeled with CellTracker Red CMPTX were plated on 35 mm glass-bottom dishes in macrophage medium. After 18–24 h, the cells were overlaid with a \sim 1100 μ m thick layer of 5.8 mg/mL collagen I (BD Biosciences). After 24 h, the cells were fixed with 4% formaldehyde for 30 min and analyzed by confocal microscopy on a Bio-Rad Radiance 2000 Laser Scanning Confocal Microscope. Optical z-sections were taken every 5 μ m, beginning at the base of the dish to 70 μ m into the collagen gel. The percent cell invasion was calculated by dividing the fluorescence from z-sections 20–70 μ m by the sum of the fluorescence from all sections. Experiments were performed in duplicate, and three different fields from each dish were imaged. For S100A4–GFP or sc-S100A4–GFP protein rescue experiments, transfected HCT116 cells were used 48 h post-transfection in the 3D invasion assay. The GFP fluorescence was used to monitor the invasion of the collagen gel by tumor cells.

RESULTS

Development of the Single-Chain S100A4 Proteins.

With the exception of S100G (also known as calbindin D9k), the S100 proteins form noncovalent dimers.⁴⁴ Because each monomer is comprised of two EF-hands and a target binding site, the presumed functional unit of these proteins consists of four EF-hands and two target binding sites.^{2,10,14–16} To investigate whether both monomers in the S100A4 dimer are necessary for biochemical and biological function, we created an obligate S100A4 dimer. The two S100A4 chains were connected with a flexible linker comprised of four glycines and one serine, thus allowing S100A4 to fold properly even though it is encoded by a single polypeptide chain. Glycine residues were used to promote a flexible linker between the two S100A4 chains, and a serine residue was added to aid in the solubility and stability of the resulting single-chain S100A4 protein (sc-S100A4) (Figure 1).^{45,46} The linker sequence (GGGGS) was repeated one, two, and three times, and the proteins were designated sc-S100A4–5, sc-S100A4–10, and sc-S100A4–15, respectively. Both sc-S100A4–5 and sc-S100A4–10 expressed well in bacteria, whereas expression of sc-S100A4–15 was not detected in whole cell lysates (data not shown). On SDS–PAGE, the purified native S100A4 (wt-S100A4) migrated at



Figure 1. Cartoon of sc-S100A4 proteins. The C-terminus of monomer 1 was fused to the N-terminus of monomer 2 using the linker sequence (GGGGS)_n, where $n = 1, 2$, or 3 .

approximately 11 kDa, consistent with a noncovalent dimer. Consistent with the mass spectrometry data, purified sc-S100A4–5 and sc-S100A4–10 migrated above 21 kDa as expected for two native S100A4 chains covalently attached via a linker sequence (Figure 2A).

Size-Exclusion Chromatography of the sc-S100A4 Proteins. To confirm that the proteins were properly folded in solution, they were examined by analytical size-exclusion chromatography. In the presence of EGTA or Ca^{2+} , the elution




Figure 2. Analytical gel filtration of sc-S100A4 proteins. (A) SDS–PAGE of purified wt-S100A4, sc-S100A4–5, and sc-S100A4–10 on a 12% Tris-Tricine gel. Molecular mass markers are indicated on the left. (B) Elution profile of wt-S100A4 (—) and protein standards (---) on a Superdex 75HR column in the presence of EGTA. The inset is a plot of protein standard molecular masses vs elution volume. (C and D) Elution profiles of sc-S100A4–5 and sc-S100A4–10 in the presence of EGTA: wt-S100A4 (—) and sc-S100A4–5 and sc-S100A4–10 (---).

profile of wt-S100A4 showed one major peak, which was coincident with the 27 kDa marker (Figure 2B). The apparent molecular mass of the native S100A4 dimer agreed well with the calculated mass (Table 1). Notably, at loading concentrations of 0.7 μ M, we observed a single peak for wt-S100A4, demonstrating that the native S100A4 forms a dimer at submicromolar concentrations. Both sc-S100A4–5 and sc-S100A4–10 had elution profiles comparable to that of wt-S100A4, indicating that in solution both sc-S100A4 constructs adopted an overall organization similar to that of the noncovalent dimer of the wild-type protein (Figure 2C,D). In the presence of calcium, all three proteins eluted as more compact proteins (Table 1), consistent with other Ca^{2+} -bound S100 proteins.⁴⁷

Table 1. Apparent Molecular Masses and Stokes Radii of S100A4 Proteins

protein	condition	elution volume (mL) ^a	Stokes radius (Å) ^a	apparent molecular mass (×10 ³ Da) ^a
wt-S100A4	EGTA	11.4 ± 0.1	22.4 ± 0.4	24.2 ± 1.2
wt-S100A4	Ca ²⁺	12.0 ± 0.2	19.2 ± 1.1	19.4 ± 1.9
sc-S100A4–5	EGTA	11.3 ± 0.0	22.4 ± 0.0	24.0 ± 0.0
sc-S100A4–5	Ca ²⁺	12.0 ± 0.2	19.2 ± 1.1	19.4 ± 1.9
sc-S100A4–10	EGTA	11.2 ± 0.2	23.1 ± 1.0	25.7 ± 2.5
sc-S100A4–10	Ca ²⁺	12.0 ± 0.2	19.2 ± 1.1	19.4 ± 1.9

^aMean ± standard deviation of two or three gel filtration runs in the presence of 2 mM EGTA or 2 mM CaCl₂. Molecular masses and Stokes radii were calculated on the basis of the molecular masses and Stokes radii of the protein standards, respectively.

Chemical Denaturation of S100A4 Proteins. Previous studies demonstrated that some S100 family members are more stable to denaturation in the presence of calcium.^{48,49} To determine if S100A4 shares this characteristic and to establish if the flexible linker affected the stability of the single-chain proteins, we monitored unfolding in the presence of guanidine hydrochloride. In the absence of calcium, wt-S100A4, sc-S100A4–5, and sc-S100A4–10 had similar midpoints of denaturation, 1.71 ± 0.03, 1.84 ± 0.05, and 1.81 ± 0.06 M guanidine hydrochloride, respectively (Figure 3A). Consistent with other S100 proteins, the wt-S100A4, sc-S100A4–5, and sc-S100A4–10 proteins were more resistant to denaturant in the presence of calcium.^{48,50} Interestingly, the sc-S100A4 proteins were more stable than

S100A4 (Figure 3B). These observations suggest that the addition of the flexible linker between the two S100A4 monomers has a stabilizing effect on the protein.

Secondary Structural Analysis of S100A4 Proteins. Structural studies of wt-S100A4 showed that it is primarily composed of α-helices.^{14,16,51,52} Circular dichroism spectroscopy of sc-S100A4–5 and sc-S100A4–10 demonstrated that these proteins are mostly α-helical (Figure 4B,C). As for wt-S100A4, we observed a small increase in the helical content of sc-S100A4–5 and sc-S100A4–10 in the presence of calcium




Figure 3. Guanidine hydrochloride denaturation of sc-S100A4 proteins. (A) Unfolding of 5 μM wild-type S100A4 dimer or sc-S100A4 proteins in the presence of 2 mM EGTA. wt-S100A4, sc-S100A4–5, and sc-S100A4–10 exhibited similar midpoints of denaturation. (B) Unfolding in the presence of 2 mM CaCl₂. The sc-S100A4 proteins were more resistant to denaturant-induced unfolding than wt-S100A4. wt-S100A4 (black), sc-S100A4–5 (red), and sc-S100A4–10 (blue). Values represent the mean ± standard error of the mean from three or four independent experiments.

the wt-S100A4 protein with midpoints of denaturation of 3.95 ± 0.02 and 3.76 ± 0.06 M for sc-S100A4–5 and sc-S100A4–10, respectively, compared to 3.10 ± 0.04 M for wt-




Figure 4. Circular dichroism spectroscopy of S100A4 proteins. Far-UV spectra of 6.8 μM S100A4 protein dimers of (A) wt-S100A4, (B) sc-S100A4–5, and (C) sc-S100A4–10. Solid and dotted lines correspond to spectra in the presence of 2 mM EGTA and 0.3 mM CaCl₂, respectively. The α-helical structure of S100A4 is not altered by the addition of the flexible linker.

(Figure 4A). These data confirm that the addition of the linker has not grossly perturbed the secondary structure of S100A4 in either the apo or Ca²⁺-bound state.

Biochemical Evaluation of sc-S100A4 Proteins. The ability of the sc-S100A4 proteins to bind myosin-IIA was assessed in an anisotropy assay using FITC-labeled MIIA^{1908–1923}, which comprises the minimal binding site for S100A4.^{16,17} The measured dissociation constant for wt-S100A4 (1.8 ± 0.2 μM) was

comparable to previously reported values.^{16,17} Both sc-S100A4–5 ($1.0 \pm 0.1 \mu\text{M}$) and sc-S100A4–10 ($1.3 \pm 0.2 \mu\text{M}$) demonstrated binding affinities for FITC-MIIA^{1908–1923} comparable to that of wt-S100A4 (Figure 5A). Moreover, binding was not observed in the presence of EGTA, demonstrating that the sc-S100A4 proteins exhibit Ca^{2+} -dependent binding to myosin-IIA.




Figure 5. Single-chain S100A4 proteins bind and regulate myosin-IIA assembly. (A) Affinities of S100A4 proteins for myosin-IIA were measured using a fluorescence anisotropy assay. wt-S100A4 (□), sc-S100A4–5 (○), and sc-S100A4–10 (◇) exhibit similar affinities for myosin-IIA. (B) Promotion of myosin-IIA disassembly. wt-S100A4 (black bars), sc-S100A4–5 (white bars), and sc-S100A4–10 (gray bars) all promote maximal filament disassembly at a 1:1 molar ratio of S100A4 dimer to myosin-IIA dimer. Myosin-IIA disassembly in the absence of S100A4 (hatched bar). Values represent the mean \pm standard error of the mean from three independent experiments.

Biochemical studies have shown that S100A4 regulates the monomer–polymer equilibrium of myosin-IIA by promoting the monomeric state.^{26,27} Using an assembly competent myosin-IIA construct, MIIA^{1338–1960}, we monitored the ability of the sc-S100A4 proteins to promote the disassembly of preexisting myosin-IIA filaments. Similar to that of wt-S100A4, maximal promotion of myosin-IIA disassembly occurred at a ratio of 1 mol of sc-S100A4 protein dimer to 1 mol of MIIA^{1338–1960} dimer with disassembly of approximately 90% of the myosin-IIA filaments (Figure 5B).

Dissociation constants for Ca^{2+} binding were determined using a competition assay with the chromophoric calcium chelator 5,5'-Br₂-BAPTA.^{17,39} In agreement with previous reports,^{16,17} wt-S100A4 exhibited relatively strong affinities for calcium in the presence of a 4-fold molar excess of myosin-IIA ($^{\text{EF1}}K_D = 4.26 \pm 0.04 \mu\text{M}$; $^{\text{EF2}}K_D = 0.44 \pm 0.04 \mu\text{M}$) (Table 2 and Figure 6A). The Ca^{2+} binding affinities for sc-S100A4–5 ($^{\text{EF1}}K_D = 1.65 \pm 0.05 \mu\text{M}$; $^{\text{EF2}}K_D = 0.74 \pm 0.11 \mu\text{M}$) and sc-

S100A4–10 ($^{\text{EF1}}K_D = 3.20 \pm 0.30 \mu\text{M}$; $^{\text{EF2}}K_D = 1.10 \pm 0.18 \mu\text{M}$) in the presence of myosin-IIA were comparable to that of wt-S100A4 (Table 2 and Figure 6B,C).

Table 2. Dissociation Constants^a for Binding of Ca^{2+} to S100A4

protein	IIIA ^{1851–1960}	EF1 (μM)	EF2 (μM)
wt-S100A4 ^b	+	4.26 ± 0.04	0.44 ± 0.04
wt-S100A4 ^c	–	>500	7.6 ± 0.4
sc-S100A4–5 ^b	+	1.65 ± 0.05	0.74 ± 0.11
sc-S100A4–10 ^b	+	3.20 ± 0.30	1.10 ± 0.18
sc-S100A4–5M1 ^b	+	169 ± 23	0.51 ± 0.05
sc-S100A4–5M1 ^c	–	366 ± 93	1.83 ± 0.34
sc-S100A4–5M2 ^b	+	205 ± 41	0.24 ± 0.03
sc-S100A4–5M2 ^c	–	>500	0.75 ± 0.21
sc-S100A4–10M1 ^b	+	244 ± 46	0.43 ± 0.06
sc-S100A4–10M1 ^c	–	223 ± 26	1.46 ± 0.31
sc-S100A4–10M2 ^b	+	>500	0.92 ± 0.30
sc-S100A4–10M2 ^c	–	>500	1.26 ± 0.12

^aValues represent the average from two or three independent experiments. ^bMacroscopic Ca^{2+} binding constants determined for 12.5 μM S100A4 dimer in the presence of 100 μM MIIA^{1851–1960} monomer. ^cMacroscopic Ca^{2+} binding constants determined for 12.5 μM S100A4 dimer in the absence of target.

Development and Biochemical Evaluation of sc-S100A4 EF-Hand Mutant Proteins. To test whether both monomers in the S100A4 homodimer are required for function, we incorporated single-amino acid substitutions in both the pseudo (EF1) and canonical (EF2) EF-hands in either monomer 1 (M1) or monomer 2 (M2) of the sc-S100A4 constructs (Figure 7A). Previous studies with S100B demonstrated that substitution of E31 and E72 with alanine was sufficient to disrupt binding of Ca^{2+} to EF1 and EF2; therefore, we made the analogous substitutions in sc-S100A4 (E33A/E74A). Circular dichroism spectroscopy of the M1 and M2 proteins showed that the alanine substitutions did not disrupt the overall secondary structure of the single-chain proteins in the apo and Ca^{2+} -bound states (Figure 1 of the Supporting Information).

Ca^{2+} binding affinities were measured for “wild-type” monomer 2 in sc-S100A4–5M1 and sc-S100A4–10M1 proteins and the wild-type monomer 1 in sc-S100A4–5M2 and sc-S100A4–10M2 proteins. In the absence of target, we obtained the following values: $^{\text{EF1}}K_D = 366 \pm 93 \mu\text{M}$ and $^{\text{EF2}}K_D = 1.83 \pm 0.34 \mu\text{M}$ for sc-S100A4–5M1, $^{\text{EF1}}K_D = 223 \pm 26 \mu\text{M}$ and $^{\text{EF2}}K_D = 1.46 \pm 0.31 \mu\text{M}$ for sc-S100A4–10M1, $^{\text{EF1}}K_D > 500 \mu\text{M}$ and $^{\text{EF2}}K_D = 0.75 \pm 0.21 \mu\text{M}$ for sc-S100A4–5M2, and $^{\text{EF1}}K_D > 500 \mu\text{M}$ and $^{\text{EF2}}K_D = 1.26 \pm 0.12 \mu\text{M}$ for sc-S100A4–10M2 (Table 2). Interestingly, in the presence of a 4-fold molar excess of myosin-IIA, the Ca^{2+} binding affinities for sc-S100A4–5M1 ($^{\text{EF1}}K_D = 169 \pm 23 \mu\text{M}$; $^{\text{EF2}}K_D = 0.51 \pm 0.05 \mu\text{M}$), sc-S100A4–10M1 ($^{\text{EF1}}K_D = 244 \pm 46 \mu\text{M}$; $^{\text{EF2}}K_D = 0.43 \pm 0.06 \mu\text{M}$), sc-S100A4–5M2 ($^{\text{EF1}}K_D = 205 \pm 41 \mu\text{M}$; $^{\text{EF2}}K_D = 0.24 \pm 0.03 \mu\text{M}$), and sc-S100A4–10M2 ($^{\text{EF1}}K_D > 500 \mu\text{M}$; $^{\text{EF2}}K_D = 0.92 \pm 0.30 \mu\text{M}$) were comparable to the K_D values measured in the absence of target (Table 2 and Figure 7B–E). These observations indicate that mutation of either monomer 1 or monomer 2 in both sc-S100A4–5 and sc-S100A4–10 proteins affects binding of Ca^{2+} to the wild-type monomer for both proteins. Binding of Ca^{2+} to EF1 was weak both in the




Figure 6. Single-chain proteins and wt-S100A4 exhibit similar Ca^{2+} binding affinities in the presence of myosin-IIA. Ca^{2+} binding affinities were measured using a chromophoric calcium chelator 5,5'-Br₂-BAPTA competition assay. The absorbance at 263 nm was monitored as a function of the Ca^{2+} concentration: (A) wt-S100A4, (B) sc-S100A4-S, and (C) sc-S100A4-10. The data represent the average from three independent experiments. The insets show the saturation curve representation for the best fit in Caligator.

absence and in the presence of myosin-IIA, whereas EF2 exhibited a high affinity for Ca^{2+} irrespective of the presence of target.

An examination of binding of sc-S100A4-SM1 and -M2 and sc-S100A4-10M1 and -M2 to FITC-MIIA^{1908–1923} in the anisotropy assay demonstrated that these proteins had 30–60-fold reduced affinities for myosin-IIA with dissociation constants of 93.4 ± 9.4 , 125.1 ± 17.0 , 69.3 ± 17.4 , and 68.9 ± 9.7 μM , respectively (Figure 8A). These data demonstrate that neither the sc-S100A4-M1 nor -M2 proteins retained wild-type binding affinities for myosin-IIA.

To confirm that the combined E33A and E74A substitutions in EF1 and EF2 disrupt the Ca^{2+} -dependent interactions of S100A4 with myosin-IIA, we examined the ability of a noncovalent S100A4 dimer that contained alanine substitution

in all four EF hands to promote myosin-IIA disassembly. As expected, even at a molar ratio of 2.5:1 (S100A4 dimer:MIIA^{1338–1960} dimer), myosin-IIA disassembly was not observed (Figure 8B). Although the wild-type sc-S100A4 proteins promoted maximal disassembly of myosin-IIA at a 1:1 sc-S100A4 dimer:MIIA^{1338–1960} dimer molar ratio (Figure 6B), the sc-S100A4-M1 and -M2 proteins poorly regulated myosin-IIA disassembly (Figure 8B). We observed modest myosin-IIA disassembly at molar ratios of 2.5:1 (sc-S100A4-M1 or -M2 dimer to MIIA^{1338–1960} dimer). At ratios of 10:1, approximately 50% of the myosin-IIA had shifted to the supernatant fraction. These findings are consistent with the reduced affinities that the sc-S100A4-M1 and M2 proteins exhibit for myosin-IIA and suggest that two functional monomers are needed for binding and regulation of myosin-IIA.

Biological Characterization of sc-S100A4 Proteins.

S100A4 overexpression promotes tumor cell migration and invasion.^{18,53} To evaluate the requirement for two functional S100A4 monomers to promote invasion, we used an in vitro assay that reconstitutes macrophage-dependent invasion of tumor cells into a 3D collagen gel⁴³ using colorectal HCT116 carcinoma cells that are homozygous null for S100A4 (Figure 9A). When cultured with macrophages, 26% of the parental S100A4-expressing HCT116 cells invaded at least 20 μm into the collagen gel, whereas only 9% of parental HCT116 cells invaded the matrix when cultured alone (Figure 9C). The S100A4^{-/-} HCT116 cell lines, L5 and N3, did not exhibit macrophage-dependent invasion (Figure 9C).

To establish that HCT116 invasion required S100A4, we monitored the invasive capabilities of N3 S100A4^{-/-} cells transfected with GFP alone or with wild-type S100A4–GFP fusion protein. To demonstrate that attachment of GFP to the native S100A4 did not disrupt protein function, we evaluated the ability of bacterially expressed and purified S100A4–GFP protein to promote myosin-IIA filament disassembly. Notably, identical purification protocols were used for S100A4–GFP protein and for the untagged wild-type S100A4, indicating that the addition of the GFP tag does not affect the biochemical properties of S100A4. At a 1:1 molar ratio of S100A4–GFP dimer to MIIA^{1338–1960} dimer in the presence of calcium, we observed filament disassembly comparable to that elicited by wild-type S100A4 (Figure 2A of the Supporting Information). These observations demonstrate that S100A4–GFP protein is functionally equivalent to untagged wild-type S100A4. To compare the localization of the endogenous S100A4 and S100A4–GFP proteins, parental HCT116 cells were transfected with the S100A4–GFP construct. A comparison of the localization of S100A4–GFP protein with total S100A4 showed that the endogenous and the majority of the exogenously expressed S100A4 colocalize (Figure 2B of the Supporting Information). Moreover, the overall distribution of the S100A4–GFP protein is similar to the distribution of endogenous S100A4.

Transfection of N3 S100A4^{-/-} cells with GFP alone did not rescue macrophage-dependent invasion; however, transfection with S100A4–GFP protein rescued the invasion of HCT116 cells with 32% of tumor cells invading more than 20 μm into the collagen gel (Figure 9C). The enhanced invasion observed for S100A4–GFP protein-expressing N3 cells likely results from an increased level of S100A4 expression (Figure 9B). Immunoblot analysis demonstrated that total S100A4 levels in the transfected N3 cells are comparable to levels of expression




Figure 7. S100A4 single-chain EF-hand mutant proteins exhibit altered Ca^{2+} affinities. (A) Cartoon of sc-S100A4 EF-hand mutant constructs. Both E33 (EF1) and E74 (EF2) residues in either monomer 1 or monomer 2 were substituted with alanine. Calcium titrations of (B) sc-S100A4–5M1, (C) sc-S100A4–10M1, (D) sc-S100A4–5M2, and (E) sc-S100A4–10M2 in the presence of myosin-IIA and 5,5'-Br₂-BAPTA. The data represent the average from two or three independent experiments. The insets show the saturation curve representation for the best fit in Caligator.

in parental HCT116 cells; however, the transfection efficiency was approximately 40–50% for N3 cells, and thus, the invading cells express higher S100A4 levels. The expression of sc-S100A4–5–GFP protein in L5 and N3 S100A4^{−/−} cells, which had a transfection efficiency of 40–50%, also rescued the macrophage-dependent invasion of these null lines to varying extents (Figure 10). N3 cells expressing sc-S100A4–5M1–GFP protein did not exhibit macrophage-dependent invasion, whereas L5 cells exhibited a slight increase in the level of invasion in the presence of macrophages that was significantly reduced compared to that of parental HCT116 cells. The transfection efficiency for sc-S100A4–5M1–GFP protein was consistently 10–20% lower than that observed for sc-S100A4–5–GFP protein (Figure 9B). Taking into account the differences in transfection efficiency, we found the transfected cells express comparable levels of sc-S100A4–5–GFP and sc-S100A4–5M1–GFP proteins. Thus, it is unlikely that the inability of sc-S100A4–5M1–GFP protein to rescue HCT116 invasion results from a decreased level of expression of this construct.

Chemical Cross-Linking of S100A4–Myosin-IIA Complexes. To experimentally evaluate the composition and stoichiometry of the S100A4–myosin-IIA complex, the native S100A4 dimer and MIIA^{1851–1960} were cross-linked in the presence of EGTA or calcium using disuccinimidyl tartrate (DST). To maximize binding, these studies were performed at S100A4 and myosin-IIA concentrations approximately 50-fold

above the equilibrium dissociation constant for the S100A4–MIIA^{1851–1960} interaction.¹⁶ For both S100A4 and myosin-IIA alone, we observed both monomeric and dimeric species in the cross-linked sample (Figure 11A). Cross-linked S100A4 or MIIA^{1851–1960} dimers were not detected by mass spectrometry, which is likely due to the low abundance of these protein species in the sample. However, multiple monomeric species comprised of unmodified S100A4 or MIIA^{1851–1960} monomers (S100A4, 11599.4 Da; MIIA, 12399.5 Da) and monomers containing multiple intramolecular and hanging cross-links (S100A4, 11844.2 and 11977.2 Da; MIIA, 12530.4 and 12662.3 Da) were identified. Moreover, the S100A4 monomer was more extensively modified in the absence of MIIA^{1851–1960} than in the presence of MIIA^{1851–1960}. The weak reactivity of the S100A4 and MIIA^{1851–1960} monomers on immunoblots is likely a consequence of modification by the DST cross-linker (Figure 11B).

In samples containing equimolar S100A4 and myosin-IIA, we observed two Ca^{2+} -dependent species (Figure 11A, asterisks). Immunoblot analysis indicated that the upper band was comprised of both S100A4 and myosin-IIA, whereas the lower band contained myosin-IIA and exhibited very weak cross-reactivity with the S100A4 antibody (Figure 11B). Mass spectrometry showed that the upper band was comprised of two protein species with molecular masses of 36264 and 36130 Da; however, we did not identify any species associated with the lower Ca^{2+} -dependent band. In the absence of any DST cross-links, the calculated mass of a S100A4 dimer bound to a




Figure 8. S100A4 single-chain EF-hand mutant proteins only weakly promote myosin-IIA disassembly. (A) Fluorescence anisotropy measurements of sc-S100A4-5M1 (solid gray line), sc-S100A4-10M1 (solid black line), sc-S100A4-5M2 (dashed gray line), and sc-S100A4-10M2 (dashed black line) demonstrated that sc-S100A4-M1 and -M2 EF-hand mutants have a 30–60-fold reduced affinity for FITC-MIIA^{1908–1923}. Values represent the mean \pm standard deviation from two or three independent experiments. (B) Myosin-IIA disassembly assays. Myosin-IIA disassembly in the absence of any added S100A4 (light gray bar). Ca²⁺-dependent myosin-IIA disassembly in the presence of wt-S100A4 (black bars) and EF-hand mutant proteins sc-S100A4-5M1 (white bars), sc-S100A4-5M2 (white-hatched bars), sc-S100A4-10M1 (dark gray bars), sc-S100A4-10M2 (dark gray-hatched bars), and wt-S100A4-E33A/E74A (white-lined bars). Values represent the mean \pm standard error of the mean for three independent experiments.

single myosin-IIA polypeptide is 35592 Da, whereas the calculated mass of a MIIA^{1851–1960} dimer complexed to a S100A4 monomer is 36392 Da. The observed 36264 and 36130 Da species are thus consistent with a complex comprised of an S100A4 dimer and one MIIA^{1851–1960} polypeptide with several cross-links. Although we were not able to identify the lower Ca²⁺-dependent band by mass spectrometry, its migration via SDS-PAGE is consistent with a complex of a S100A4 monomer bound to a single myosin-IIA polypeptide (calculated mass of 23995 Da) with several cross-links.

Because the cross-linking data suggested that one S100A4 dimer binds a single myosin-IIA polypeptide chain, we performed anisotropy titration measurements to directly measure the stoichiometry of binding. The measured dissociation constant for the binding of FITC-MIIA^{1904–1927} to wild-type S100A4 was $0.26 \pm 0.03 \mu\text{M}$, which is approximately 7-fold tighter than the affinity observed for FITC-MIIA^{1908–1923} ($1.8 \pm 0.2 \mu\text{M}$) (Figure 12A). Consistent with the cross-linking data, titration of $1 \mu\text{M}$ FITC-MIIA^{1904–1927} with wild-type S100A4 revealed a stoichiometry of one S100A4 dimer per myosin-IIA peptide (Figure 12B).




Figure 9. HCT116 cells require S100A4 for macrophage-dependent invasion. (A) Immunoblot of S100A4 expression in parental and L5 and N3 S100A4^{-/-} HCT116 cells. β -Actin was used as a loading control. (B) Immunoblot of S100A4 protein expression in L5 and N3 S100A4^{-/-} HCT116 cells transfected with wild-type S100A4-GFP, sc-S100A4-5-GFP, or sc-S100A4-5M1-GFP proteins. β -Actin was used as a loading control. (C) Invasion of parental HCT116 cells, L5 and N3 S100A4^{-/-} cells, and N3 S100A4^{-/-} cells transfected with GFP or S100A4-GFP fusion protein in the absence (white bars) or presence (black bars) of macrophages. Values represent the mean \pm standard error of the mean for three independent experiments.




Figure 10. Single-chain S100A4-5 rescues HCT116 cell invasion. Invasion of parental HCT116 cells and L5 and N3 S100A4^{-/-} HCT116 cells transfected with sc-S100A4-5-GFP or sc-S100A4-5M1-GFP fusion proteins in the absence (white bars) or presence (black bars) of macrophages. Values represent the mean \pm standard deviation from two independent experiments.

DISCUSSION

Here we show that S100A4, a known metastasis factor, requires both monomers to effectively interact with nonmuscle myosin-IIA. Our single-chain S100A4 proteins, sc-S100A4, allowed us to disrupt Ca²⁺ binding in one monomer while leaving the other intact. Similar approaches involving the linking of subunits to evaluate or enhance protein function have been




Figure 11. Composition of S100A4–myosin-IIA complexes. (A) Coomassie-stained SDS–PAGE of 12.5 μ M wt-S100A4 dimer and 25 μ M MIIA monomer in the presence of DST, a 6.4 Å cross-linker. Asterisks denote complexes observed in the presence of calcium. (B) Western blots of cross-linked samples. Duplicate protein samples were separated by SDS–PAGE and transferred to a PVDF membrane, which was cut in half and probed with antibodies to either S100A4 or myosin-IIA.

used previously to increase the blood circulation time of CuZn superoxide dismutase,⁵⁴ to create bispecific antibodies,⁵⁵ and to assess the DNA binding affinity for lambda Cro repressor.⁵⁶

The biophysical and biochemical characterization of the wild-type sc-S100A4 proteins revealed that the sc-S100A4–5 and sc-S100A4–10 proteins are true mimics of the wt-S100A4 noncovalent dimer. One exception is that the Ca²⁺-bound sc-S100A4 proteins were more resistant to chemical denaturants than the wild-type noncovalent S100A4 dimer, suggesting that the flexible linker stabilizes the S100A4 dimer against chemical denaturation. The precise mechanism responsible for the enhanced stability of the sc-S100A4 proteins is likely to be complex, though it is likely to benefit at least in part from entropic contributions. For example, the condensation of two distinct chains into a dimeric species results in the loss of six degrees of freedom (three rotational and three translational), while in the case of the sc-S100A4 proteins, there is no analogous entropic penalty. The absence of this entropic penalty in the folding of the sc-S100A4 proteins may contribute to the observed enhancement in stability relative to that of the wild-type noncovalent dimer.

Increased resistance to denaturation has been documented for other single-chain proteins, including thermal denaturation of the single-chain variant of monellin and urea denaturation of the Arc-L1-Arc single-chain protein.^{57–59} This stabilizing effect is not due to gross structural changes as circular dichroism spectroscopy of the sc-S100A4 proteins showed that these proteins remain mostly α -helical with spectra similar to those of wt-S100A4. Similar to other S100 proteins, this single transition suggests that the native S100A4 dimer unfolds via a two-state mechanism, where unfolding of the dimer proceeds to the




Figure 12. Stoichiometry of binding of FITC-MIIA^{1904–1927} to wt-S100A4. (A) Fluorescence anisotropy measurements of wt-S100A4 binding to MIIA^{1904–1927} demonstrated that wt-S100A4 has an approximately 7-fold increased affinity for FITC-MIIA^{1904–1927} as compared to FITC-MIIA^{1909–1923}. Values represent the mean \pm standard deviation from two or three independent titrations. (B) Stoichiometry of binding of wt-S100A4 to FITC-MIIA^{1904–1927}. wt-S100A4 binds to FITC-MIIA^{1904–1927} at a 1:1 molar ratio (wt-S100A4 dimer:peptide). Gray dotted lines represent the linear regression of the rise and plateau of the titration curve. Values represent the mean \pm standard deviation from two or three independent titrations.

denatured state without an intermediate.^{48,49} Although folding intermediates have been detected for S100B, S100A11, and S100A12, our data do not support the existence of a folding intermediate.^{48,60,61} While it remains possible that weakly populated intermediates, which are beyond the detection limits of this study, contribute to the folding of S100A4, additional studies will be required to evaluate the mechanism of S100A4 unfolding.

The selection of the EF1 E33 and EF2 E74 residues for mutation was based on analogous Ca²⁺-disrupting substitutions examined in S100B.⁶² Other studies assessing binding of Ca²⁺ to S100A4 used the EF-hand substitutions EF1 E33Q and EF2 D63N, which also disrupt Ca²⁺ binding.^{63,64} An examination of the canonical EF-hand sequence for all S100 proteins, as well as other EF-hand-containing proteins, shows that both D63 and E74 are invariant, indicating that these positions are necessary for Ca²⁺ ligation. This is also supported by the X-ray structure of the Ca²⁺-bound S100A4, which shows that D63 and E74 are critical for Ca²⁺ ion coordination in EF2.^{16,51,52} For the native S100A4 dimer, Ca²⁺ binding affinities increase in the presence of myosin-IIA 10- and 100-fold for EF2 and EF1, respectively.¹⁶ For the sc-S100A4-M1 and -M2 proteins, inactivation of monomer 1 or monomer 2 yielded S100A4 proteins whose Ca²⁺ affinities were the same in the absence and presence myosin-IIA. It is not surprising that EF1 did not exhibit an increased affinity for Ca²⁺ in the presence of myosin-IIA as the

sc-S100A4-M1 and -M2 proteins have 30–60-fold reduced affinities for myosin-IIA and the calcium measurements were taken at concentrations below the K_D for myosin-IIA. However, EF2 in sc-S100A4-M1 and M2 proteins exhibited affinities for Ca^{2+} in the absence of target that are typically only observed for the wild-type protein in the presence of myosin-IIA. These data suggest that there is communication between the two S100A4 monomers in which the inactive conformation of one monomer effects binding of Ca^{2+} to the second wild-type monomer.

Mutagenesis studies demonstrated that binding of Ca^{2+} to EF2 alone is sufficient for binding of S100A4 to myosin-IIA.⁶³ On the basis of this finding, the wild-type monomer in the sc-S100A4-M1 and -M2 mutant proteins should retain myosin-IIA binding activity; however, these proteins exhibited significantly decreased myosin-IIA binding activity and only weakly promoted myosin-IIA filament assembly. The inability of the sc-S100A4 mutant proteins to bind myosin-IIA is reminiscent of studies with the single-chain tetracycline repressor (sc-TetR).⁶⁵ TetR binds with high affinity to the *tet* operator as a dimer. Tetracycline binding induces a conformational rearrangement in the dimerization domain that is transmitted to the DNA binding domain and reduces the affinity of TetR for DNA.⁶⁵ Covalent tethering of the two TetR monomers revealed that a functional tetracycline binding site in each TetR monomer was necessary to promote the conformational rearrangements required for DNA release.⁶⁵ S100A4 may function in a similar manner, where the open, active conformation of both monomers is required for interactions with myosin-IIA.

This idea is supported by our cross-linking data that suggest that the S100A4 dimer binds a single MIIA^{1851–1960} polypeptide. Moreover, these data are consistent with our observation that S100A4 binds a 24mer myosin-IIA peptide with a stoichiometry of one peptide per S100A4 dimer and a recent report that a 32mer myosin-IIA peptide binds to S100A4 with a stoichiometry of one peptide per S100A4 dimer.⁶⁴ Given the length of the MIIA^{1851–1960} construct, it may be surprising that only one polypeptide chain is cross-linked to the S100A4 dimer; however, recent NMR studies of residues 1850–1960 of myosin-IIA demonstrated that the 17 N-terminal amino acids and the 35 C-terminal residues are unstructured.⁶⁴ We proposed previously that S100A4 binding promotes myosin-IIA filament disassembly by inducing local unwinding of the myosin-IIA coiled coil.²⁷ Because a significant portion of the MIIA^{1851–1960} construct is unstructured, S100A4-mediated conformational rearrangements of the coiled coil could result in dissociation of the two myosin-IIA polypeptides, resulting in the cross-linking of a single MIIA^{1851–1960} polypeptide chain. Structural studies indicate that most S100 protein dimers bind the target peptide in the hydrophobic cleft between helices 3 and 4, which is exposed upon Ca^{2+} binding, thus allowing two target peptides to bind per S100 dimer.^{66–69} However, the recent structure of S100A6 bound to the C-terminal domain of the Siah-1 interacting protein showed that although the S100A6 dimer binds two target peptides, each peptide binds across the S100A6 dimer interface and forms interactions with the canonical target binding cleft.⁷⁰ On the basis of these observations, one may envision that a target peptide, which binds across the dimer interface and is of sufficient length, could make contacts with both canonical target binding clefts. Such a configuration would result in a complex of one target peptide bound per S100 dimer.

On the basis of our biochemical analysis of the sc-S100A4 proteins with myosin-IIA, our cross-linking data, and the

determination of S100A4:myosin-IIA stoichiometries, we propose a model in which the myosin-IIA heavy chain wraps around the S100A4 dimer to form interactions in the vicinity of the canonical target binding cleft of each monomer in such a manner that both target binding sites are required for the efficient interaction with myosin-IIA.

■ ASSOCIATED CONTENT

§ Supporting Information

Circular dichroism spectra of sc-S100A4–5M1 and -M2 EF-hand mutant proteins, Ca^{2+} -dependent myosin-IIA promotion of disassembly by S100A4–GFP protein, and immunofluorescent localization of S100A4 in untransfected parental HCT116 cells and parental HCT116 cells transfected with S100A4–GFP protein. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Telephone: (718) 430-2741. Fax: (718) 430-8565. E-mail: anne.bresnick@einstein.yu.edu.

Funding

This work was supported by National Institutes of Health Grants CA129598 (A.R.B.) and GM58888 (D.J.W.).

■ ACKNOWLEDGMENTS

We thank Dr. Michael Brenowitz and Dr. Steven Almo for helpful discussions and Ms. Janine Stanford and Mr. Christopher Pudney for technical assistance.

■ ABBREVIATIONS

MMTV, mouse mammary tumor virus; PyMT, polyoma middle T; sc-S100A4, single-chain S100A4; wt-S100A4, native S100A4 dimer; GdnHCl, guanidine hydrochloride; BAPTA, 1,2-bis(*o*-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid; MIIA, non-muscle myosin-IIA; FITC, fluorescein isothiocyanate; M1, monomer 1; M2, monomer 2; 3D, three-dimensional; DST, disuccinimidyl tartrate; MS, mass spectrometry.

■ REFERENCES

- (1) Moore, B. W. (1965) A soluble protein characteristic of the nervous system. *Biochem. Biophys. Res. Commun.* 19, 739–744.
- (2) Marenholz, I., Heizmann, C. W., and Fritz, G. (2004) S100 proteins in mouse and man: From evolution to function and pathology (including an update of the nomenclature). *Biochem. Biophys. Res. Commun.* 322, 1111–1122.
- (3) Seal, R. L., Gordon, S. M., Lush, M. J., Wright, M. W., and Bruford, E. A. (2011) *genenames.org*: The HGNC resources in 2011. *Nucleic Acids Res.* 39, D514–D519.
- (4) Persechini, A., Moncrief, N. D., and Kretsinger, R. H. (1989) The EF-hand family of calcium-modulated proteins. *Trends Neurosci.* 12, 462–467.
- (5) Yap, K. L., Ames, J. B., Swindells, M. B., and Ikura, M. (1999) Diversity of conformational states and changes within the EF-hand protein superfamily. *Proteins* 37, 499–507.
- (6) Ravasi, T., Hsu, K., Goyette, J., Schroder, K., Yang, Z., Rahimi, F., Miranda, L. P., Alewood, P. F., Hume, D. A., and Geczy, C. (2004) Probing the S100 protein family through genomic and functional analysis. *Genomics* 84, 10–22.
- (7) Santamaria-Kisiel, L., Rintala-Dempsey, A. C., and Shaw, G. S. (2006) Calcium-dependent and -independent interactions of the S100 protein family. *Biochem. J.* 396, 201–214.

- (8) Schafer, B. W., and Heizmann, C. W. (1996) The S100 family of EF-hand calcium-binding proteins: Functions and pathology. *Trends Biochem. Sci.* 21, 134–140.
- (9) Zimmer, D. B., Cornwall, E. H., Landar, A., and Song, W. (1995) The S100 protein family: History, function, and expression. *Brain Res. Bull.* 37, 417–429.
- (10) Donato, R. (2003) Intracellular and extracellular roles of S100 proteins. *Microsc. Res. Tech.* 60, 540–551.
- (11) Drohat, A. C., Baldisseri, D. M., Rustandi, R. R., and Weber, D. J. (1998) Solution structure of calcium-bound rat S100B($\beta\beta$) as determined by nuclear magnetic resonance spectroscopy. *Biochemistry* 37, 2729–2740.
- (12) Koch, M., Diez, J., and Fritz, G. (2008) Crystal structure of Ca^{2+} -free S100A2 at 1.6-Å resolution. *J. Mol. Biol.* 378, 933–942.
- (13) Otterbein, L. R., Kordowska, J., Witte-Hoffmann, C., Wang, C. L., and Dominguez, R. (2002) Crystal structures of S100A6 in the Ca^{2+} -free and Ca^{2+} -bound states: The calcium sensor mechanism of S100 proteins revealed at atomic resolution. *Structure* 10, 557–567.
- (14) Vallely, K. M., Rustandi, R. R., Ellis, K. C., Varlamova, O., Bresnick, A. R., and Weber, D. J. (2002) Solution structure of human Mts1 (S100A4) as determined by NMR spectroscopy. *Biochemistry* 41, 12670–12680.
- (15) Wright, N.T., Cannon, B. R., Zimmer, D. B., and Weber, D. J. (2009) S100A1: Structure, Function, and Therapeutic Potential. *Curr. Chem. Biol.* 3, 138–145.
- (16) Malashkevich, V. N., Varney, K. M., Garrett, S. C., Wilder, P. T., Knight, D., Charpentier, T. H., Ramagopal, U. A., Almo, S. C., Weber, D. J., and Bresnick, A. R. (2008) Structure of Ca^{2+} -bound S100A4 and its interaction with peptides derived from nonmuscle myosin-IIA. *Biochemistry* 47, 5111–5126.
- (17) Garrett, S., Hodgson, L., Rybin, A., Touthkine, A., Hahn, K. M., Lawrence, D. S., and Bresnick, A. R. (2008) A Biosensor of S100A4 Metastasis Factor Activation: Inhibitor Screening and Cellular Activation Dynamics. *Biochemistry* 47, 986–996.
- (18) Davies, B. R., Davies, M. P., Gibbs, F. E., Barraclough, R., and Rudland, P. S. (1993) Induction of the metastatic phenotype by transfection of a benign rat mammary epithelial cell line with the gene for p9Ka, a rat calcium-binding protein, but not with the oncogene EJ-ras-1. *Oncogene* 8, 999–1008.
- (19) Davies, M. P., Rudland, P. S., Robertson, L., Parry, E. W., Jolicoeur, P., and Barraclough, R. (1996) Expression of the calcium-binding protein S100A4 (p9Ka) in MMTV-neu transgenic mice induces metastasis of mammary tumours. *Oncogene* 13, 1631–1637.
- (20) Xue, C., Plieth, D., Venkov, C., Xu, C., and Neilson, E. G. (2003) The gatekeeper effect of epithelial-mesenchymal transition regulates the frequency of breast cancer metastasis. *Cancer Res.* 63, 3386–3394.
- (21) Rudland, P. S., Platt-Higgins, A., Renshaw, C., West, C. R., Winstanley, J. H., Robertson, L., and Barraclough, R. (2000) Prognostic significance of the metastasis-inducing protein S100A4 (p9Ka) in human breast cancer. *Cancer Res.* 60, 1595–1603.
- (22) van Dieck, J., Brandt, T., Teufel, D. P., Veprintsev, D. B., Joerger, A. C., and Fersht, A. R. (2010) Molecular basis of S100 proteins interacting with the p53 homologs p63 and p73. *Oncogene* 29, 2024–2035.
- (23) Takenaga, K., Nakamura, Y., Sakiyama, S., Hasegawa, Y., Sato, K., and Endo, H. (1994) Binding of pEL98 protein, an S100-related calcium-binding protein, to nonmuscle tropomyosin. *J. Cell Biol.* 124, 757–768.
- (24) Kriajevska, M., Fischer-Larsen, M., Moertz, E., Vorm, O., Tulchinsky, E., Grigorian, M., Ambartsumian, N., and Lukanidin, E. (2002) Liprin β 1, a member of the family of LAR transmembrane tyrosine phosphatase-interacting proteins, is a new target for the metastasis-associated protein S100A4 (Mts1). *J. Biol. Chem.* 277, 5229–5235.
- (25) Kriajevska, M. V., Cardenas, M. N., Grigorian, M. S., Ambartsumian, N. S., Georgiev, G. P., and Lukanidin, E. M. (1994) Non-muscle myosin heavy chain as a possible target for protein encoded by metastasis-related mts-1 gene. *J. Biol. Chem.* 269, 19679–19682.
- (26) Murakami, N., Kotula, L., and Hwang, Y. W. (2000) Two distinct mechanisms for regulation of nonmuscle myosin assembly via the heavy chain: Phosphorylation for MIB and mts 1 binding for MIIA. *Biochemistry* 39, 11441–11451.
- (27) Li, Z. H., Spektor, A., Varlamova, O., and Bresnick, A. R. (2003) Mts1 regulates the assembly of nonmuscle myosin-IIA. *Biochemistry* 42, 14258–14266.
- (28) Mitsuhashi, M., Sakata, H., Kinjo, M., Yazawa, M., and Takahashi, M. (2011) Dynamic assembly properties of nonmuscle myosin II isoforms revealed by combination of fluorescence correlation spectroscopy and fluorescence cross-correlation spectroscopy. *J. Biochem.* 149, 253–263.
- (29) Betapudi, V., Licate, L. S., and Egelhoff, T. T. (2006) Distinct roles of nonmuscle myosin II isoforms in the regulation of MDA-MB-231 breast cancer cell spreading and migration. *Cancer Res.* 66, 4725–4733.
- (30) Garrett, S.C., Varney, K. M., Weber, D. J., and Bresnick, A. R. (2006) S100A4, a mediator of metastasis. *J. Biol. Chem.* 281, 677–680.
- (31) Tarabykina, S., Scott, D. J., Herzyk, P., Hill, T. J., Tame, J. R., Kriajevska, M., Lafitte, D., Derrick, P. J., Dodson, G. G., Maitland, N. J., Lukanidin, E. M., and Bronstein, I. B. (2001) The dimerization interface of the metastasis-associated protein S100A4 (Mts1): In vivo and in vitro studies. *J. Biol. Chem.* 276, 24212–24222.
- (32) Helfman, D. M., Kim, E. J., Lukanidin, E., and Grigorian, M. (2005) The metastasis associated protein S100A4: Role in tumour progression and metastasis. *Br. J. Cancer* 92, 1955–1958.
- (33) Ismail, T. M., Zhang, S., Fernig, D. G., Gross, S., Martin-Fernandez, M. L., See, V., Tozawa, K., Tynan, C. J., Wang, G., Wilkinson, M. C., Rudland, P. S., and Barraclough, R. (2010) Self-association of calcium-binding protein S100A4 and metastasis. *J. Biol. Chem.* 285, 914–922.
- (34) Rashtchian, A. (1995) Novel methods for cloning and engineering genes using the polymerase chain reaction. *Curr. Opin. Biotechnol.* 6, 30–36.
- (35) Aslanidis, C., and de Jong, P. J. (1990) Ligation-independent cloning of PCR products (LIC-PCR). *Nucleic Acids Res.* 18, 6069–6074.
- (36) Studier, F. W. (2005) Protein production by auto-induction in high density shaking cultures. *Protein Expression Purif.* 41, 207–234.
- (37) Pace, C. N., Shirley, B. A., and Thomson, J. A. (1997) Measuring the conformational stability of a protein. In *Protein Structure: A Practical Approach* (Creighton, T., Ed.) pp 299–321, IRL Press at Oxford University Press, New York.
- (38) Schmid, F. (1997) Optical spectroscopy to characterize protein conformation and conformational change. In *Protein Structure: A Practical Approach* (Creighton, T., Ed.) pp 289–297, IRC Press at Oxford University Press, New York.
- (39) Linse, S. (2002) Calcium binding to proteins studied via competition with chromophoric chelators. In *Calcium Binding Protein Protocols* (Vogel, H. J., Ed.) pp 15–24, Humana Press, Totowa, NJ.
- (40) Andre, I., and Linse, S. (2002) Measurement of Ca^{2+} -binding constants of proteins and presentation of the CaLigator software. *Anal. Biochem.* 305, 195–205.
- (41) Dulyaninova, N. G., House, R. P., Betapudi, V., and Bresnick, A. R. (2007) Myosin-IIA heavy-chain phosphorylation regulates the motility of MDA-MB-231 carcinoma cells. *Mol. Biol. Cell* 18, 3144–3155.
- (42) Li, Z. H., and Bresnick, A. R. (2006) The S100A4 metastasis factor regulates cellular motility via a direct interaction with myosin-IIA. *Cancer Res.* 66, 5173–5180.

- (43) Goswami, S., Sahai, E., Wyckoff, J. B., Cammer, M., Cox, D., Pixley, F. J., Stanley, E. R., Segall, J. E., and Condeelis, J. S. (2005) Macrophages promote the invasion of breast carcinoma cells via a colony-stimulating factor-1/epidermal growth factor paracrine loop. *Cancer Res.* 65, 5278–5283.
- (44) Akke, M., Drakenberg, T., and Chazin, W. J. (1992) Three-dimensional solution structure of Ca^{2+} -loaded porcine calbindin D9k determined by nuclear magnetic resonance spectroscopy. *Biochemistry* 31, 1011–1020.
- (45) Bird, R. E., Hardman, K. D., Jacobson, J. W., Johnson, S., Kaufman, B. M., Lee, S. M., Lee, T., Pope, S. H., Riordan, G. S., and Whitlow, M. (1988) Single-chain antigen-binding proteins. *Science* 242, 423–426.
- (46) Hagemeyer, C. E., von Zur Muhlen, C., von Elverfeldt, D., and Peter, K. (2009) Single-chain antibodies as diagnostic tools and therapeutic agents. *Thromb. Haemostasis* 101, 1012–1019.
- (47) Drohat, A. C., Nenortas, E., Beckett, D., and Weber, D. J. (1997) Oligomerization state of S100B at nanomolar concentration determined by large-zone analytical gel filtration chromatography. *Protein Sci.* 6, 1577–1582.
- (48) Garcia, A.F., Garcia, W., Nonato, M. C., and Araujo, A. P. (2008) Structural stability and reversible unfolding of recombinant porcine S100A12. *Biophys. Chem.* 134, 246–253.
- (49) Botelho, H. M., Koch, M., Fritz, G., and Gomes, C. M. (2009) Metal ions modulate the folding and stability of the tumor suppressor protein S100A2. *FEBS J.* 276, 1776–1786.
- (50) Ferguson, P. L., and Shaw, G. S. (2002) Role of the N-terminal helix I for dimerization and stability of the calcium-binding protein S100B. *Biochemistry* 41, 3637–3646.
- (51) Pathuri, P., Vogeley, L., and Luecke, H. (2008) Crystal structure of metastasis-associated protein S100A4 in the active calcium-bound form. *J. Mol. Biol.* 383, 62–77.
- (52) Gingras, A. R., Basran, J., Prescott, A., Kriajevska, M., Bagshaw, C. R., and Barsukov, I. L. (2008) Crystal structure of the Ca^{2+} -form and Ca^{2+} -binding kinetics of metastasis-associated protein, S100A4. *FEBS Lett.* 582, 1651–1656.
- (53) Bjornland, K., Winberg, J. O., Odegaard, O. T., Hovig, E., Loennechen, T., Aasen, A. O., Fodstad, O., and Maelandsmo, G. M. (1999) S100A4 involvement in metastasis: Deregulation of matrix metalloproteinases and tissue inhibitors of matrix metalloproteinases in osteosarcoma cells transfected with an anti-S100A4 ribozyme. *Cancer Res.* 59, 4702–4708.
- (54) Hallewell, R. A., Laria, I., Tabrizi, A., Carlin, G., Getzoff, E. D., Tainer, J. A., Cousens, L. S., and Mullenbach, G. T. (1989) Genetically engineered polymers of human CuZn superoxide dismutase. Biochemistry and serum half-lives. *J. Biol. Chem.* 264, 5260–5268.
- (55) Mallender, W. D., and Voss, E. W. Jr. (1994) Construction, expression, and activity of a bivalent bispecific single-chain antibody. *J. Biol. Chem.* 269, 199–206.
- (56) Jana, R., Hazbun, T. R., Fields, J. D., and Mossing, M. C. (1998) Single-chain lambda Cro repressors confirm high intrinsic dimer-DNA affinity. *Biochemistry* 37, 6446–6455.
- (57) Kim, S. H., Kang, C. H., Kim, R., Cho, J. M., Lee, Y. B., and Lee, T. K. (1989) Redesigning a sweet protein: Increased stability and renaturability. *Protein Eng.* 2, 571–575.
- (58) Robinson, C.R., and Sauer, R. T. (1996) Equilibrium stability and sub-millisecond refolding of a designed single-chain Arc repressor. *Biochemistry* 35, 13878–13884.
- (59) Aghera, N., Earanna, N., and Udgaonkar, J. B. (2011) Equilibrium Unfolding Studies of Monellin: The Double-Chain Variant Appears To Be More Stable Than the Single-Chain Variant. *Biochemistry* 50, 2434–2444.
- (60) Shaw, G. S., Marlatt, N. M., Ferguson, P. L., Barber, K. R., and Bottomley, S. P. (2008) Identification of a dimeric intermediate in the unfolding pathway for the calcium-binding protein S100B. *J. Mol. Biol.* 382, 1075–1088.
- (61) Pan, J., Rintala-Dempsey, A. C., Li, Y., Shaw, G. S., and Konermann, L. (2006) Folding kinetics of the S100A11 protein dimer studied by time-resolved electrospray mass spectrometry and pulsed hydrogen-deuterium exchange. *Biochemistry* 45, 3005–3013.
- (62) Markowitz, J., Rustandi, R. R., Varney, K. M., Wilder, P. T., Udan, R., Wu, S. L., Horrocks, W. D., and Weber, D. J. (2005) Calcium-binding properties of wild-type and EF-hand mutants of S100B in the presence and absence of a peptide derived from the C-terminal negative regulatory domain of p53. *Biochemistry* 44, 7305–7314.
- (63) Kim, E. J., and Helfman, D. M. (2003) Characterization of the metastasis-associated protein, S100A4. Roles of calcium binding and dimerization in cellular localization and interaction with myosin. *J. Biol. Chem.* 278, 30063–30073.
- (64) Badyal, S. K., Basran, J., Bhanji, N., Kim, J. H., Chavda, A. P., Jung, H. S., Craig, R., Elliott, P. R., Irvine, A. F., Barsukov, I. L., Kriajevska, M., and Bagshaw, C. R. (2011) Mechanism of the Ca^{2+} -dependent interaction between S100A4 and tail fragments of nonmuscle myosin heavy chain IIA. *J. Mol. Biol.* 405, 1004–1026.
- (65) Kamionka, A., Majewski, M., Roth, K., Bertram, R., Kraft, C., and Hillen, W. (2006) Induction of single chain tetracycline repressor requires the binding of two inducers. *Nucleic Acids Res.* 34, 3834–3841.
- (66) Rustandi, R. R., Baldisseri, D. M., and Weber, D. J. (2000) Structure of the negative regulatory domain of p53 bound to S100B($\beta\beta$). *Nat. Struct. Biol.* 7, 570–574.
- (67) Bhattacharya, S., Large, E., Heizmann, C. W., Hemmings, B., and Chazin, W. J. (2003) Structure of the Ca^{2+} /S100B/NDR kinase peptide complex: Insights into S100 target specificity and activation of the kinase. *Biochemistry* 42, 14416–14426.
- (68) Wright, N. T., Prosser, B. L., Varney, K. M., Zimmer, D. B., Schneider, M. F., and Weber, D. J. (2008) S100A1 and calmodulin compete for the same binding site on ryanodine receptor. *J. Biol. Chem.* 283, 26676–26683.
- (69) Wright, N. T., Cannon, B. R., Wilder, P. T., Morgan, M. T., Varney, K. M., Zimmer, D. B., and Weber, D. J. (2009) Solution structure of S100A1 bound to the CapZ peptide (TRTK12). *J. Mol. Biol.* 386, 1265–1277.
- (70) Lee, Y. T., Dimitrova, Y. N., Schneider, G., Ridenour, W. B., Bhattacharya, S., Soss, S. E., Caprioli, R. M., Filipek, A., and Chazin, W. J. (2008) Structure of the S100A6 complex with a fragment from the C-terminal domain of Siah-1 interacting protein: A novel mode for S100 protein target recognition. *Biochemistry* 47, 10921–10932.