


# What Determines the Activity of Antimicrobial and Cytolytic Peptides in Model Membranes

Kim S. Clark, James Svetlovics, Alesia N. McKeown, Laura Huskins, and Paulo F. Almeida\*

Department of Chemistry and Biochemistry, University of North Carolina, Wilmington, North Carolina 28403, United States

**ABSTRACT:** We previously proposed three hypotheses relating the mechanism of antimicrobial and cytolytic peptides in model membranes to the Gibbs free energies of binding and insertion into the membrane [Almeida, P. F., and Pokorny, A. (2009) *Biochemistry* 48, 8083–8093]. Two sets of peptides were designed to test those hypotheses, by mutating of the sequences of  $\delta$ -lysin, cecropin A, and magainin 2. Peptide binding and activity were measured on phosphatidylcholine membranes. In the first set, the peptide charge was changed by mutating basic to acidic residues or vice versa, but the amino acid sequence was not altered much otherwise. The type of dye release changed from graded to all-or-none according to prediction. However, location of charged residues in the sequence with the correct spacing to form salt bridges failed to improve binding. In the second set, the charged and other key residues were kept in the same positions, whereas most of the sequence was significantly but conservatively simplified, maintaining the same hydrophobicity and amphipathicity. This set behaved completely different from predicted. The type of release, which was expected to be maintained, changed dramatically from all-or-none to graded in the mutants of cecropin and magainin. Finally, contrary to the hypotheses, the results indicate that the Gibbs energy of binding to the membrane, not the Gibbs energy of insertion, is the primary determinant of peptide activity.



*Irrefutability is not a virtue of a theory (as people often think) but a vice. Every genuine test of a theory is an attempt to falsify it, or to refute it.*

— Karl Popper, 1963, *Conjectures and Refutations*

Our understanding of the relation between the sequence of antimicrobial peptides and their mechanism remains poor. Perhaps the main consensual conclusions are that positive charge on antimicrobial peptides improves binding and, to some extent, determines the specificity of these peptides to anionic bacterial membranes, and that sufficient hydrophobicity renders the peptide hemolytic, because binding to the outer monolayer of eukaryotic membranes, which contains almost exclusively zwitterionic and neutral lipids, improves.<sup>1–3</sup> Beyond that, we have learned remarkably little about the importance of amino acid sequence in spite of a vast number of studies. It remains impossible to predict mechanism and function from sequence alone.

The cell membrane provides a tightly controlled barrier between cell interior and environment. Protein export involves a secretory pathway from the endoplasmic reticulum, through the Golgi apparatus, to the extracellular matrix. A number of proteins and peptides, however, circumvent the regular transport machinery of the cell and appear to cross the lipid bilayer directly.<sup>4</sup> Cell-penetrating peptides (CPPs),<sup>1</sup> for example, appear to accomplish this goal.<sup>5</sup> Amphipathic CPPs are similar to antimicrobial peptides, and their mechanisms may have more in common than often appreciated.<sup>6,7</sup>

We have proposed three hypotheses to understand the mechanism of membrane-active peptides in model membranes,

including antimicrobial, cytolytic, and amphipathic cell-penetrating peptides, on the basis of sequence.<sup>6</sup> By mechanism, we mean the series of steps in the kinetics leading to membrane disruption, including the process of vesicle leakage. We do not mean the molecular models commonly discussed for the structural arrangements of the peptides in the membrane, such as the barrel-stave,<sup>8</sup> carpet,<sup>9–11</sup> toroidal pore,<sup>12,13</sup> or sinking-raft models.<sup>14,15</sup> The central idea of the first hypothesis we proposed is that the mechanism of membrane-active peptides is determined by the Gibbs free energy of insertion into the bilayer from the membrane-bound state.<sup>6</sup> More precisely, if that energy is smaller than a certain threshold, the peptides are predicted to translocate across the lipid bilayer. This seems to be the case for amphipathic CPPs.<sup>16</sup> If the Gibbs free energy of insertion is larger than the threshold, the peptides cannot cross the membrane and accumulate instead on its surface. A point may be reached when the membrane yields, forming a pore. This distinction is important because peptides that translocate can be used as carriers for delivery of drugs or nucleic acids into cells, whereas those that cause major membrane disruption should be primarily antimicrobial or cytolytic, depending on specificity. According to this hypothesis, the peptide sequence affects the mechanism primarily through its effect on the Gibbs free energy of insertion.


This concept is described more precisely with reference to Figure 1. In water, an equilibrium exists between helical and unfolded conformations of the peptide, which favors the unfolded

**Received:** June 6, 2011

**Revised:** August 12, 2011

**Published:** August 26, 2011





**Figure 1.** Thermodynamic cycle for peptide binding to the membrane interface and insertion into the bilayer. The folding equilibrium in water lies toward the unstructured state and is determined by  $\Delta G_f^\circ$ , which is typically small in comparison with the other terms. The Gibbs energy of binding to the interface ( $\Delta G_{if}^\circ$ ) includes contributions from the hydrophobic effect and secondary structure formation. Transfer, as an  $\alpha$ -helix, from water to the bilayer interior is approximated by transfer to octanol ( $\Delta G_{oct}^\circ$ ). The Gibbs energy of transfer from the surface to the interior of the bilayer is approximately  $\Delta G_{oct}^\circ - \Delta G_{if}^\circ = \Delta G_{oct-if}^\circ$ . Modified with permission from ref 38. Copyright 2007 Elsevier.

state. Upon binding to the bilayer–water interface, the peptide folds to an  $\alpha$ -helix. The Gibbs energy of binding to the interface is given by  $\Delta G_{if}^\circ$ . If we ignore the free energy associated with folding in water, which is typically small compared to the other terms,<sup>6</sup> the Gibbs energy of insertion from the surface-bound state should be approximately given by  $\Delta G_{oct}^\circ - \Delta G_{if}^\circ = \Delta G_{oct-if}^\circ$ , where  $\Delta G_{oct}^\circ$ , the Gibbs energy of transfer from water to octanol, is used to estimate transfer to the bilayer hydrophobic core.<sup>17</sup> The idea is that  $\Delta G_{oct-if}^\circ$  provides a tool for predicting the behavior of the peptides. The Gibbs energy of binding measured experimentally is designated here by  $\Delta G_{bind}^\circ$ . It can also be calculated using the Wimley–White interfacial hydrophobicity scale<sup>18–21</sup> and is then designated by  $\Delta G_{if}^\circ$ .  $\Delta G_{oct-if}^\circ$  is calculated with the Wimley–White octanol hydrophobicity scale.<sup>18,22</sup>

Previously, we examined the kinetics and thermodynamics of interaction of a set of amphipathic,  $\alpha$ -helical peptides with model membranes.<sup>6</sup> This original set consisted of  $\delta$ -lysin, a hemolytic peptide from *Staphylococcus aureus*;<sup>23</sup> cecropin A, an antimicrobial peptide from *Hyalophora cecropia*;<sup>24</sup> magainin 2, an antimicrobial peptide from *Xenopus laevis*;<sup>25</sup> and transportan 10 (TP10), an amphipathic CPP.<sup>26,27</sup> For most peptides,  $\Delta G_{bind}^\circ$  was in very good agreement with  $\Delta G_{if}^\circ$ .<sup>6,16</sup> However, for two peptides ( $\delta$ -lysin and cecropin A), the calculated and experimental binding free energies did not agree. Because these were the only peptides of the set originally examined in which intramolecular salt bridges between side chains could be established, we proposed salt bridge formation on the membrane was the reason for the disagreement in those two cases.<sup>6</sup> Formation of an intramolecular salt bridge between acidic and basic groups of the peptide contributes  $-4$  kcal/mol to transfer from water to octanol.<sup>28</sup> We suggested that formation of a salt bridge may contribute approximately  $-2$  kcal/mol to transfer from water to the POPC membrane interface. Using this value, and postulating two intramolecular salt bridges in

$\delta$ -lysin and cecropin A, we found  $\Delta G_{if}^\circ$  could be brought into agreement with  $\Delta G_{bind}^\circ$ .<sup>6</sup>

For the sake of clarity and completeness, we now restate the three hypotheses proposed.<sup>6</sup> Note also that when  $\Delta G_{bind}^\circ$  is known, this experimental value is used in the relationship  $\Delta G_{oct-if}^\circ = \Delta G_{oct}^\circ - \Delta G_{bind}^\circ$  instead of  $\Delta G_{if}^\circ$ .

(1) If  $\Delta G_{oct-if}^\circ \leq 20$  kcal/mol, the peptides can translocate across the bilayer. This should result in graded dye release because translocation provides a means of dissipating the mass imbalance across the bilayer, created by peptide binding. However, if  $\Delta G_{oct-if}^\circ > 23$  kcal/mol, the energy barrier for translocation is prohibitively large; the peptides accumulate on the membrane surface until, in a stochastic manner, the membrane yields, releasing the contents of the vesicle in an all-or-none manner. To be clear, hypothesis 1 contains in fact the following logical structure:

$$\begin{aligned} \Delta G_{oct-if}^\circ \leq 20 \text{ kcal/mol} &\Rightarrow \text{translocation} \\ &\Rightarrow \text{graded release} \end{aligned} \quad (1)$$

$$\begin{aligned} \Delta G_{oct-if}^\circ \geq 23 \text{ kcal/mol} &\Rightarrow \text{no translocation} \\ &\Rightarrow \text{all-or-none release} \end{aligned} \quad (2)$$

A “gray zone” may exist for  $\Delta G_{oct-if}^\circ$  between  $\sim 20$  and  $\sim 23$  kcal/mol, in which either mechanism may prevail.

(2) Formation of intramolecular salt bridges (hydrogen-bonded ion pairs) by the residue side chains can lower  $\Delta G_{if}^\circ$  and  $\Delta G_{oct-if}^\circ$ , allowing the peptides to bind more strongly and translocate across the bilayer.

(3) The sequence of antimicrobial and other membrane-active peptides can be simplified without altering the peptide mechanism and the type of dye release as long as the Gibbs energies of binding and insertion into the membrane remain approximately constant. The idea is that the thermodynamics of membrane binding and insertion determine the mechanism. Therefore, it should be possible to reduce the diversity of amino acids in the sequence, simplifying it considerably, while preserving the amino acid types to ensure that the peptide remains amphipathic, provided those sequence changes do not alter the thermodynamics of binding and insertion. If this is correct, mutations may occur naturally to impart variation to the sequences, to circumvent acquired bacterial resistance.

TP10 variants were the subject of our previous study, which showed that the changes to the sequence resulted in peptide mechanisms consistent with these hypotheses.<sup>16</sup> The results for designed variants of other membrane-active peptide families are presented here.

The peptide sequences are listed in Table 1, and their helical wheel projections are shown in Figure 2. They are divided into two sets and named according to the original peptides from which they are derived:  $\delta$ -lysin (DL), cecropin A (CE), and magainin 2 (MG). Thus, DL-1 and DL-2 variants are based on  $\delta$ -lysin, CE-1 and CE-2 on cecropin A, and MG-1 and MG-2 on magainin 2. Set 1 consists of DL-1, CE-1, and MG-1; here, charge and putative salt bridges were changed relative to those of the parent peptides. During design, we tried to keep membrane binding affinity within a measurable range, assuming that the helicity would not change much relative to those of the parent peptides, while retaining the general amphipathicity of the peptide. Set 1 was designed to test hypothesis 1, that the

**Table 1. Parent Set of Peptides and Sets 1 and 2 of Mutants<sup>a</sup>**

peptide	charge (pH 7.5)	length (no. of residues)	$\mu_H^b$	sequence
Set of Parent Peptides				
$\delta$ -lysin	0	26	7.8	formyl-MAQDIISTIGDLVKWIIDTVNKFTKK
cecropin A	+7	37	4.9	KWKLFKKIEKVGQNIRDGIKAGPAVAVVGQATQIAK-amide
magainin 2 F12W	+3	23	6.9	GIGKFLHSAKKWGKAFVGEIMNS
Set 1 of Mutant Peptides				
DL-1	+6	26	7.2	formyl-MAQKIISTIGKLVKWIITVNVKFTKK
CE-1	+1	37	7.3	EWKLFKEIEKLGQNILDGIKLGPLLALLGQLTQIAL-amide
MG-1	0	23	9.1	GILKFLESAKKWLEAFLAEIMNS
Set 2 of Mutant Peptides				
DL-2a	0	26	7.7	formyl-LAADLLAALGDLAKWLLDALAKAAKK
DL-2b	0	26	8.8	formyl-LAADLLAALGDLKWLDDALAKLAKK
CE-2	+7	37	5.2	KWKLLKKLEKAGAAKEGLLKAGPALALLGAAAAALAK-amide
MG-2	+3	23	7.0	GLGKLLHAAKKLGKAWLGELLAA

<sup>a</sup>Unless indicated, N- and C-termini are free. <sup>b</sup>Hydrophobic moment of the complete helix calculated with MPEx using the Wimley–White interfacial scale.<sup>18</sup>

peptide mechanism depends on the difficulty of peptide insertion. Namely, the release type should be graded or all-or-none depending on whether  $\Delta G_{\text{oct-if}}^\circ$  is below or above the threshold. We sought to change peptides between those that can translocate across the membrane (equated with causing graded release) and those that cannot (equated with causing all-or-none release) by engineering mutations that change  $\Delta G_{\text{oct-if}}^\circ$  in a predictable way. Furthermore, at least if the mechanism of the peptides does not change with the mutations, the easier the insertion, the faster dye release should be. To alter  $\Delta G_{\text{oct-if}}^\circ$ , basic residues were mutated to acidic and vice versa, but with minimal other changes to the sequence. This resulted in changes to the net charge of the peptide and also eliminated or allowed new salt bridges. In this way, hypothesis 2 was also tested.

Set 2 consists of DL-2a, DL-2b, CE-2, and MG-2; here, the sequence was simplified, but the charge, hydrophobicity, and amphipathicity (hydrophobic moment) were maintained. Set 2 was designed to test hypothesis 3, that is, the importance of amino acid diversity in the sequence. The sequences of the original peptides were simplified to contain a minimal number of amino acid types, while keeping key residues, such as those charged, in place. In practice, this amounted to changing hydrophobic residues to Leu, small polar ones to Ala, and charged residues to Lys or Glu. The peptides of set 2 are experimental analogues of the HP (hydrophobic/polar) model<sup>29</sup> that has been extensively studied in protein folding in water. Can these experimental models capture the essence of the activity of antimicrobial peptides?

## MATERIALS AND METHODS

**Chemicals.** DL-1 (93% pure) was purchased from Bachem (Torrance, CA). DL-2a (95%), CE-2 (95%), and MG-2 (88%) were from New England Peptide (Gardner, MA). DL2b (99%), CE-1 (98%), and MG-1 (95%) were from Genscript (Scotch Plains, NJ). Their identity was ascertained by mass spectrometry, and the purity was determined by high-performance liquid chromatography, with instruments provided by the manufacturer.  $\delta$ -Lysine was a gift from H. Birkbeck (University of Glasgow, Glasgow, Scotland); cecropin A and magainin 2 were purchased from American Peptide Co. (Sunnyvale, CA). and magainin 2 F12W was a gift from R. Biltonen (University of Virginia, Charlottesville, VA). Stock solutions were prepared by


dissolving lyophilized peptide in deionized water or a 1:1 (v/v) water/ethyl alcohol mixture (AAPER Alcohol and Chemical, Shelbyville, KY). Stock peptide solutions were stored at  $-80^\circ\text{C}$  and kept on ice during experiments. 1-Palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC), 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphatidylethanolamine (POPE), and 1-palmitoyl-2-oleoyl-*sn*-glycero-3-[phospho-*rac*-(1-glycerol)] (POPG), in a chloroform solution, were purchased from Avanti Polar Lipids (Alabaster, AL). 7-Methoxycoumarin-3-carboxylic acid (7MC) succinimidyl ester, 8-aminonaphthalene-1,3,6-trisulfonic acid (ANTS) disodium salt, *p*-xylene bis-pyridinium bromide (DPX), and carboxyfluorescein (CF) were purchased from Molecular Probes/Invitrogen (Carlsbad, CA). Organic solvents (high-performance liquid chromatography/American Chemical Society grade) were purchased from Burdick & Jackson (Muskegon, MI). Lipids and fluorophores were tested by thin layer chromatography (TLC) and used without further purification. Peptide concentrations were determined by Trp absorbance at 280 nm.

**Synthesis of Fluorescent Probes.** The syntheses of fluorescent probes, using POPE and a fluorophore attached through an amide bond to the amino group of the ethanolamine, were performed as previously described in detail,<sup>30–33</sup> following the method of Vaz and Hallmann.<sup>34</sup>

**Preparation of Large Unilamellar Vesicles.** Large unilamellar vesicles (LUV) were prepared by extrusion through 0.1  $\mu\text{m}$  pore size filters (Nuclepore, Whatman, Florham, NJ), as previously described,<sup>15,16,31,32</sup> in 20 mM MOPS buffer (pH 7.5), 0.1 mM EGTA, 0.02%  $\text{NaN}_3$ , and 100 mM KCl or appropriately modified, as indicated below. Lipid concentrations were assayed by the Bartlett phosphate method,<sup>35</sup> modified as previously described.<sup>14</sup>

**Circular Dichroism.** Peptide secondary structure was assessed by circular dichroism (CD) on a Chirascan CD spectrometer (Applied Photophysics, Leatherhead, Surrey, U.K.), as previously described.<sup>16</sup> CD spectra were recorded in aqueous solution, with 1–5  $\mu\text{M}$  peptide, in 10 mM phosphate buffer (pH 7.5), and in the presence of POPC vesicles, at low concentrations (5  $\mu\text{M}$  peptide and 300  $\mu\text{M}$  POPC LUV) and high concentrations (20  $\mu\text{M}$  peptide and 5 mM POPC LUV). CD measurements of amphipathic peptides in LUV at high lipid concentrations (3–7 mM) yield results equivalent to those obtained with small unilamellar vesicles (SUV), with several advantages.<sup>36</sup> To obtain the helicity of the bound peptide at low





**Figure 2.** Helical wheel projection of the original and variant peptides studied. The colors reflect the hydrophobicities according to the Wimley–White interfacial scale:<sup>18</sup> white for hydrophilic or neutral, light gray for hydrophobic, magenta for aromatic, red for negatively charged, and blue for positively charged (at pH 7.5). The only difference between DL-2a and DL-2b is the replacement of Ala 13 and Ala 23 with Leu in DL-2b.


lipid concentrations, it is necessary to know the dissociation constant ( $K_D$ ) and the helicity of the peptides in solution in equilibrium with the vesicles. The helicities in solution are difficult to obtain for several of these peptides because of aggregation or adsorption to the cuvette walls in the absence of vesicles. For the membrane-associated peptides, the use of high LUV concentrations ( $\geq 10K_D$ ) has the additional advantage that larger peptide concentrations can be used, while maintaining a low peptide:lipid ratio ( $< 1:200$ ). We relied on determinations at high concentrations for the calculation of peptide helicity on the membrane. Nevertheless, determinations at high and low concentrations yielded helical contents on the membrane that differed by only 5–10%. The fractional helicity on the membrane ( $f_H$ ) was calculated from the average of at least two independent samples, according to the method of Luo and Baldwin,<sup>37</sup> as previously described.<sup>16</sup>

**Membrane Binding Kinetics.** The kinetics of peptide binding to lipid LUV were measured on a stopped-flow

fluorimeter (SX.18MV, Applied Photophysics). The fluorescence signal recorded was the emission of 7MC-POPE (maximum at 396 nm) incorporated into the bilayer, upon Förster resonance energy transfer (FRET) from a Trp residue on the peptide, with excitation at 280 nm, as previously described.<sup>16,31–33</sup> After mixing in the stopped-flow system, the concentration of peptide was 0.5–1  $\mu\text{M}$  and that of the lipid varied between 25 and 400  $\mu\text{M}$ . The kinetics of peptide binding to membranes were analyzed as previously described in detail.<sup>31–33</sup> Briefly, each kinetic trace (Figure 3), or the average of several traces from the same sample, was fit with a single-exponential rising function of the form

$$F(t) = a_0[1 - \exp(-k_{\text{app}}t)] + a_1 \quad (3)$$

where  $a_0$  and  $a_1$  are constants,  $t$  is time, and  $k_{\text{app}}$  is the apparent rate constant. (In the cases where a very slow process was also apparent, which was not included in  $k_{\text{app}}$ , it could be well



**Figure 3.** Example of a curve of binding kinetics, acquired for DL-1 binding to 100  $\mu\text{M}$  POPC LUV. The data correspond to fluorescence emission from the lipid fluorophore 7MC-POPE incorporated in the membrane, through FRET, upon excitation of the Trp residue on the peptide. The line is a single-exponential fit to the data.

approximated by a linear ramp.) This rate constant contains contributions from the on and off rate constants

$$k_{\text{app}} = k_{\text{on}}[L] + k_{\text{off}} \quad (4)$$

where  $[L]$  is the lipid concentration, which was varied in a series of different experiments. A plot of  $k_{\text{app}}$  versus  $[L]$  yields  $k_{\text{on}}$  from the slope and  $k_{\text{off}}$  from the  $y$ -intercept.

**ANTS/DPX Quenching Assay.** Steady state fluorescence measurements were performed in a spectrofluorimeter (8100 SLM-Aminco, Urbana, IL) upgraded by ISS (Champaign, IL), as previously done for the original peptides.<sup>15,31,32,38,39</sup> In the ANTS/DPX assay,<sup>40–42</sup> excitation was at 365 nm (8 nm slit width) and emission at 515 nm (16 nm slit width). The solution encapsulated in the LUV contained 5 mM ANTS, 10 mM DPX, 20 mM MOPS (pH 7.5), 0.1 mM EGTA, 0.02%  $\text{NaN}_3$ , and 70 mM KCl. The titrating solution contained 45 mM DPX, 20 mM MOPS (pH 7.5), 0.1 mM EGTA, 0.02%  $\text{NaN}_3$ , and 30 mM KCl. Following extrusion, the LUV with encapsulated ANTS and DPX were passed through a Sephadex-G25 column to separate the dye in the external buffer from the vesicles. Typical concentrations were 0.1–2  $\mu\text{M}$  peptide and  $\sim 600$   $\mu\text{M}$  lipid. The data were analyzed as described in detail by Ladokhin et al.<sup>42</sup> The curve for graded release is described by<sup>41,42</sup>

$$Q_{\text{in}} = \frac{F_i}{F_i^{\text{max}}} = 1 / \{ [1 + K_{\text{dyn}}[\text{DPX}]_0(1 - f_{\text{out}})]^\alpha \times [1 + K_{\text{sta}}[\text{DPX}]_0(1 - f_{\text{out}})]^\alpha \} \quad (5)$$

where  $F_i$  and  $F_i^{\text{max}}$  are the fluorescence intensities from the vesicle interior with and without quencher (DPX), respectively,  $[\text{DPX}]_0$  is the initial concentration of DPX encapsulated,  $f_{\text{out}}$  is the ANTS fraction outside the vesicles,  $K_{\text{dyn}}$  is the dynamic quenching constant, fixed at 50  $\text{M}^{-1}$  in the fits,<sup>41</sup>  $K_{\text{sta}}$  is the static quenching constant, and  $\alpha$  is the ratio of the rate of release of DPX to that of ANTS.

**Carboxyfluorescein Release Kinetics.** Carboxyfluorescein (CF) release kinetics were measured as described previously in detail.<sup>15,31–33,38</sup> Briefly, LUV 0.1  $\mu\text{m}$  in diameter, containing


50 mM CF, were prepared by extrusion in 20 mM MOPS buffer (pH 7.5), 0.1 mM EGTA, and 0.02%  $\text{NaN}_3$ . Removal of external CF was accomplished by size-exclusion chromatography (Sephadex-G25). The kinetics of CF release, measured by the relief of self-quenching of CF fluorescence, were recorded in a stopped-flow fluorimeter (SX.18MV, Applied Photophysics). The peptide concentration was 1  $\mu\text{M}$  after mixing, in all experiments. The fraction of CF released was determined by comparison of the fluorescence with that obtained upon addition of 1% Triton X-100 (maximal release). The efficiency of dye release was characterized quantitatively, in a model-free way, by the average time constant of dye release ( $\tau$ )<sup>43,44</sup>

$$\tau = \frac{\int t f(t) dt}{\int f(t) dt} \quad (6)$$

where  $f(t) [=dF(t)/dt]$  is the time derivative of the fractional release  $[F(t)]$  as a function of time. This derivative behaves as a probability density function.<sup>45,46</sup>

## RESULTS

**Peptide Secondary Structure.** The secondary structure of the peptides in aqueous solution and on the membrane was assessed by CD (Figure 4), in POPC or 1:1 POPC/POPG LUV (if binding to POPC was weak). The percent helicity of




**Figure 4.** Representative CD spectra of the mutant peptides (20  $\mu\text{M}$ ) in 5 mM POPC LUV, except for CE-2 and MG-2, which were in 1:1 POPC/POPG LUV: (A) DL-1, (B) CE-1, (C) MG-1, (D) DL-2a, (E) CE-2, and (F) MG-2.

the membrane-associated peptides was obtained from the ellipticity at 222 nm, at a lipid concentration (5 mM) much larger than the dissociation constants ( $K_D$ ), ensuring that the peptides were fully bound. All peptides were helical on the membrane, usually to a degree similar to that of the corresponding original peptides (Table 3). The helicity on the membrane is necessary for calculation of  $\Delta G_{\text{if}}^\circ$  with the Wimley–White interfacial scale, using Membrane Protein Explorer, MPEx.<sup>47</sup> In aqueous solution, however, the helicities were difficult to obtain for several of these peptides. If the helicity in solution is below  $\approx 30\%$ , the experimental error is

large, in part because of the low peptide concentrations used ( $\sim 2 \mu\text{M}$ ) to minimize aggregation, and the measurements may not reflect the structure that exists in solution in equilibrium with the membrane. Thus, the helicities in solution, which are not used in any calculation, are approximate, and several were rounded to the nearest 5 or 10%, reflecting our estimate of the error. The values calculated using AGADIR<sup>48–53</sup> are included in Table 3 for comparison. (Note that the helicities on the membrane were determined for fully bound peptides and do not rely on the measurements in solution.)

**Binding to POPC Membranes.** The equilibrium binding constants for binding of the peptides to POPC membranes were determined from the kinetics of binding, measured by stopped-flow fluorescence, using the change in FRET from a Trp residue on the peptide to a lipid fluorophore (7MC-POPE) incorporated into the bilayer (Figure 3). Typically, these measurements were performed on a time scale of  $\sim 1$  s, to capture only the binding event and avoid contributions from changes imparted to the vesicles by the peptides, which may occur concomitant with deeper insertion. Figure 3 is very representative in showing that the binding process is complete in less than  $\sim 1$  s. A fit of eq 3 to the data (Figure 3) yields the apparent rate constant for binding,  $k_{\text{app}}$ .<sup>31,32,38,39,58,59</sup> To obtain the on and off rate constants from the equation  $k_{\text{app}} = k_{\text{on}}[\text{L}] + k_{\text{off}}$ , this experiment was performed as a function of lipid concentration  $[\text{L}]$ . The results are shown in Figure 5 for DL-1 (A), CE-1 (B), MG-1 (C), and DL-2a (D). (Note that, in all figures,



**Figure 5.** Kinetics of peptide binding to POPC LUV (DL-1, CE-1, MG-1, and DL-2a) or 1:1 POPC/POPG LUV (CE-2 and MG-2). The apparent rate constant ( $k_{\text{app}}$ ), obtained from a single-exponential fit to the traces of binding kinetics, is plotted vs lipid concentration: (A) DL-1, (B) CE-1, (C) MG-1, and (D) DL-2a. The error bars are standard deviations from two to four independent experiments, and the lines are linear regressions, which yield  $k_{\text{on}}$  from the slope and  $k_{\text{off}}$  from the y-intercept. In the case of MG-1, the data are averages from 50:50, 70:30, 80:20, and 90:10 POPC/POPG LUV, because there was no detectable dependence on membrane composition. The values of  $k_{\text{on}}$ ,  $k_{\text{off}}$ , and  $K_D$  obtained are listed in Table 2.

triangles refer to data for the DL family, circles to data for the CE family, and squares to data for the MG family.) The equilibrium dissociation constants were calculated by

$$K_D = \frac{k_{\text{off}}}{k_{\text{on}}} \quad (7)$$

and are listed in Table 2, together with those of the original peptides, for comparison.

**Table 2. Kinetic and Equilibrium Constants for Binding to POPC Bilayers at Room Temperature<sup>a</sup>**

peptide	$K_D$ ( $\mu\text{M}$ )	$k_{\text{on}}$ ( $\text{M}^{-1} \text{s}^{-1}$ )	$k_{\text{off}}$ ( $\text{s}^{-1}$ )
$\delta$ -lysin <sup>b</sup>	30	$2 \times 10^3$	0.06
DL-1	400	$2.5 \times 10^4$	10
DL-2a	200	$5.5 \times 10^2$	0.11
DL-2b <sup>c</sup>	3400	$2.5 \times 10^3$	8.6
cecropin A	1000	$2.8 \times 10^5$	300
CE-1	400	$1.9 \times 10^4$	7.5
CE-2	4700	$3.0 \times 10^4$	140
magainin 2 <sup>d</sup>	2000	$2.8 \times 10^5$	550
MG-1	200	$6.1 \times 10^4$	13
MG-2	1100	$2.1 \times 10^4$	24


<sup>a</sup>The data for the original peptides were from ref 6 and references cited therein. The estimated relative error in the rate and equilibrium constants is  $\sim 30\%$ , except for  $\delta$ -lysin, for which it is larger (see footnote b).

<sup>b</sup>Measuring binding for  $\delta$ -lysin has been especially difficult, for reasons we do not fully understand. Data were collected previously, for POPC and other unsaturated phosphatidylcholines to which binding is similar.<sup>6,59</sup> On the basis of all those data, the best previous estimate of  $K_D$  was  $60 \mu\text{M}$ .<sup>6</sup> We have now re-examined the earlier data and supplemented it with new data (not shown). The numbers listed here are our improved estimates at this point. However,  $k_{\text{on}}$  is strikingly small. From our previous estimates,<sup>6,15</sup> the monomer should be the dominant species in solution at a concentration of  $0.5 \mu\text{M}$ , which was used in the binding kinetics. Nevertheless, the small  $k_{\text{on}}$  suggests that either the monomer conformation in solution is such that binding is impaired or some peptide oligomerization, which is known to occur in solution above  $1 \mu\text{M}$ ,<sup>14,54–57</sup> persists at  $0.5 \mu\text{M}$ . If the oligomerization equilibrium contributes to the measured on rates, the true  $K_D$  would be even smaller. <sup>c</sup>Data from only one experiment. <sup>d</sup>The binding data were obtained for the F12W variant of magainin 2, which, like the F16W variant, behaves very much like the original magainin.<sup>71</sup> The  $K_D$  value of  $5 \text{ mM}$ <sup>32</sup> was revised. The new estimate of  $K_D$  ( $2 \text{ mM}$ ) was obtained by direct extrapolation of the  $K_D$  in POPC/POPG mixtures, as a function of POPG content in the membrane, to pure POPC. Previously, we had extrapolated  $k_{\text{on}}$  and  $k_{\text{off}}$  and obtained  $K_D$  from their ratio,<sup>32</sup> but  $k_{\text{on}}$  had been overestimated. The new estimates of the rate constants for magainin 2 (F12W) in POPC are listed here.

CE-2 and MG-2 bind very weakly to POPC, with  $K_D$  values in the millimolar range, like those of the original peptides, cecropin A and magainin 2. To obtain those dissociation constants,  $k_{\text{on}}$  and  $k_{\text{off}}$  were determined in POPC/POPG mixtures, varying the POPC content between 50 and 100%, and the results were extrapolated to pure POPC (Figure 6), as previously done for the original peptides.<sup>31,32</sup> The extrapolation using the data from all the lipid mixtures provides a much better estimate of the rate and equilibrium constants for POPC than the direct determination with POPC only. This is because if binding to POPC is weak, the corresponding data are noisier than those acquired in the mixtures containing POPG.

In the case of MG-1, we found that binding to POPC/POPG mixtures is independent of POPG content. This is consistent with MG-1 having no net charge at pH 7.5. However, the FRET signal was better in the presence of POPG, perhaps





**Figure 6.** Kinetics of binding of CE-2 (left) and MG-2 (right) to LUV as a function of membrane composition, from 1:1 POPC/POPG to pure POPC LUV. The data show the dependence of  $k_{on}$  (A and E),  $k_{off}$  (B and F),  $K_D$  (C and D), and  $\ln K_D$  on lipid composition. Extrapolation to pure POPC, to which binding is weak, provides the best estimates of the rate and equilibrium constants for CE-2 and MG-2 in POPC (listed in Table 2).

because of the different position of the Trp or the fluorescent moiety of 7MC-POPE in the anionic membranes. Therefore, the data for MG-1 binding to 50:50, 70:30, 80:20, and 90:10

POPC/POPG LUV were all averaged, at each lipid concentration, which is shown in Figure 5C.

The Gibbs free energies of binding to POPC were obtained from the experimental value of  $K_D$  by

$$\Delta G_{bind}^{\circ} = RT \ln K_D - RT \ln [W] \quad (8)$$


where  $[W]$  (55.5 M) is the molar concentration of water. The term  $RT \ln 55.5 = 2.4$  kcal/mol (at room temperature) converts  $K_D$  to a partition coefficient of the peptide between lipid and water, with concentrations expressed in mole fraction units. To be able to compare the values of the Gibbs energy of binding derived from experiment ( $\Delta G_{bind}^{\circ}$ ) with those calculated ( $\Delta G_{if}^{\circ}$ ) with the Wimley–White interfacial scale,<sup>18–20</sup> mole fraction units must be used. MPEX<sup>47</sup> was employed to calculate  $\Delta G_{if}^{\circ}$ , assuming a free energy  $\Delta G_{hb}^{\circ}$  of  $-0.4$  kcal/mol for hydrogen bond formation at the interface, in a helical conformation.<sup>21</sup> The values of  $\Delta G_{bind}^{\circ}$  and  $\Delta G_{if}^{\circ}$  obtained are listed in Table 3.

**Kinetics of Dye Release.** Peptide-induced release of carboxyfluorescein (CF) from lipid vesicles was measured in POPC. Examples of release curves are shown by the solid lines in Figure 7 for set 1 (left panels) and set 2 (right panels). The corresponding curves for the original peptides are shown by the dashed lines for comparison. The overall release rates of DL-1 and  $\delta$ -lysine are similar, but the shape of the curves is completely different (Figure 7A). CE-1, on the other hand, is dramatically more efficient than cecropin A, which hardly releases dye from POPC vesicles (Figure 7B). The same is true of MG-1 in comparison with magainin 2 (Figure 7C). DL-2a induces significantly slower release than DL-1 and  $\delta$ -lysine (Figure 7D); DL-2b induces even slower release (Figure 7D, dotted line). The solid lines in panels E and F of Figure 7 show release induced by CE-2 and MG-2 from POPC, respectively, which occur on approximately the same time scale as for the original peptides, cecropin A and magainin 2 (dashed lines). The efficiency of dye release is expressed quantitatively by the overall apparent rate constant, which is the inverse of the

**Table 3.** Thermodynamic and Kinetic Parameters for the Mutant Peptides Binding and Insertion into POPC Bilayers at Room Temperature

peptide	$\Delta G_{bind}^{\circ}$ (kcal/mol)		helix (%)			$\Delta G_{oct}^{\circ}$ (kcal/mol)		$\Delta G_{oct-if}^{\circ}$ (kcal/mol)		dye release	
	exptl	calcd	membrane	water		$\Delta G_{oct}^{\circ}$ (kcal/mol)	$\Delta G_{oct-if}^{\circ}$ (kcal/mol)	type	$\tau$ (s) <sup>d</sup>	ANTS/ DPX	CF
				exptl	calcd <sup>e</sup>						
$\delta$ -lysine	−8.5	−5.8	100	53	1	21.9	30.4	graded	2.2		
DL-1	−7.0	−1.5	52	~20	1	19.3	26.3	all-or-none	4.9		
DL-2a	−7.4	−2.3	54	54	35	23.2	30.6	graded	14.8		
DL-2b	−5.7	−6.2	77	59	59	19.7	25.4	—	650		
cecropin A	−6.4	−1.7	63	~10	1	31.1	37.5	all-or-none	~15000		
CE-1	−7.0	−6.1	55	~30	3	17.3	24.3	all-or-none	35		
CE-2	−5.6	−5.7	70	~30	4	29.6	36.2	graded	11000		
magainin 2 <sup>f</sup>	−6.1	−3.8 (−6.2) <sup>g</sup>	57 (83)	~5	0	20.1	26.2	all-or-none	~20000		
MG-1	−7.4	−3.6	69	46	16	18.2	25.6	all-or-none	13.5		
MG-2	−6.4	−2.7 (−6.4)	40 (80)	~10	2	19.1	25.5	graded	20000		

<sup>a</sup>Calculated from the experimentally determined dissociation constant by the relationship  $\Delta G_{bind}^{\circ} = \ln K_D - 2.4$  kcal/mol. <sup>b</sup>Calculations of  $\Delta G_{if}^{\circ}$  and  $\Delta G_{oct}^{\circ}$  were performed with MPEX.<sup>47</sup> <sup>c</sup>Calculated using the experimental value of  $\Delta G_{bind}^{\circ}$  for the free energy of binding to the interface. <sup>d</sup>Measured in 50  $\mu$ M POPC LUV. <sup>e</sup>Helicity in aqueous solution calculated with AGADIR. <sup>f</sup>Binding was assessed with the F12W variant of magainin 2. The calculations assume His is neutral in magainin 2 and MG-2, because the experiments were performed at pH 7.5. The value of  $\Delta G_{bind}^{\circ}$  for magainin 2 (F12W) on POPC, previously reported as  $-5.5$  kcal/mol,<sup>6</sup> which was based on a  $K_D$  of 5 mM,<sup>32</sup> was revised. The  $\Delta G_{bind}^{\circ}$  value of  $-6.1$  kcal/mol listed here is based on a better estimate of  $K_D$  of 2 mM (see Table 2). <sup>g</sup>Binding of magainin 2 to POPC is too weak to measure helicity accurately. The value of  $-3.8$  kcal/mol is calculated using the helicity (57%) that we measured in 1:1 POPC/POPG LUV. The value of  $-6.2$  is calculated using the helicity (83%) determined from NMR in DPC micelles.<sup>64</sup>




**Figure 7.** Kinetics of CF release induced by the mutant peptides (—) from 50  $\mu$ M POPC LUV. The curves for the original peptides are shown for comparison (---) under the same conditions: (A) DL-1 (—) and  $\delta$ -lysin (---) from POPC, (B) CE-1 (—) and cecropin A (---), (C) MG-1 (—) and magainin 2 (---), (D) DL-2a (—) and DL-2b (---) from POPC, (E) CE-2 (—) and cecropin A (---), and (F) MG-2 (—) and magainin 2 (---). Some curves were acquired for longer times but are shown in the time frames that allow the best comparison. In panel D, only the beginning of the curve for DL-2b is shown; the curve has the same shape as that for DL-2a, but on a time scale 1 order of magnitude larger.

characteristic average time constant of the process,  $\tau$  (eq 6). The values of  $\tau$ , obtained under similar experimental conditions, namely at an  $\approx 1:50$  peptide:lipid molar ratio, with peptide and lipid concentrations of 0.5–1 and 30–50  $\mu$ M, respectively, are listed in Table 3.

**Type of Dye Release: Graded or All-or-None.** To determine if graded or all-or-none release occurred, the conventional ANTS/DPX requenching assay<sup>40–42</sup> was performed with the mutant peptides, as previously done with the original set. Briefly, if release is all-or-none, the fluorescence inside the vesicles remains constant as ANTS (fluorophore) and DPX (quencher) leak out, resulting in a horizontal line, but if release is graded, the degree of quenching inside the vesicles decreases as DPX leaks out, resulting in a rising curve.<sup>40–42</sup> The results of the ANTS/DPX assay for all mutant peptides are shown in Figure 8A–F. The solid lines represent the best fits to the equation for graded release (eq 5), and the dashed lines represent all-or-none behavior.

DL-1 exhibits all-or-none behavior in POPC (Figure 8A), whereas  $\delta$ -lysin was weakly graded.<sup>15</sup> CE-1 and MG-1 also exhibit all-or-none behavior in POPC (Figure 8B,C), which is the same as for the original peptides, cecropin A<sup>31,60</sup> and magainin 2.<sup>32,61,62</sup> DL-2a exhibits graded release (Figure 8D), as the original  $\delta$ -lysin, with similar values of the ratio of DPX/ANTS release rates,  $\alpha$  (eq 5). For DL-2a and  $\delta$ -lysin,  $\alpha = 0.28$  and 0.22, respectively. Finally, CE-2 and MG-2 also exhibit graded release (Figure 8E,F), in striking contrast with cecropin A and magainin 2, which released in an all-or-none manner



**Figure 8.** ANTS/DPX requenching assay for the mutant peptides: (A) DL-1 in POPC, (B) CE-1 in POPC, (C) MG-1 in POPC, (D) DL-2a in POPC, (E) CE-2 in 1:1 POPC/POPG, and (F) MG-2 in 1:1 POPC/POPG. The solid lines in panels D–F (DL-2a, CE-2, and MG-2) represent the best fits to the equation for graded release (eq 5). The corresponding fit parameters are as follows: for DL-2a,  $\alpha = 0.28$  and  $K_{sta} = 80 \text{ M}^{-1}$ ; for CE-2,  $\alpha = 8.5$  and  $K_{sta} = 140 \text{ M}^{-1}$ ; for MG-2,  $\alpha = 3.0$  and  $K_{sta} = 140 \text{ M}^{-1}$ . For comparison with DL-2a, for  $\delta$ -lysin in POPC, the parameters were as follows:  $\alpha = 0.22$  and  $K_{sta} = 220 \text{ M}^{-1}$ .<sup>15</sup> The dashed horizontal lines represent the behavior expected for all-or-none release.

from 1:1 POPC/POPG LUV. The extreme, graded shape of the curve for CE-2 and MG-2 indicates, furthermore, that DPX is released much faster than ANTS by these peptides, which is reflected in  $\alpha$  values of 8.5 and 3.0 for CE-2 and MG-2, respectively.

## DISCUSSION

Two sets of peptide variants were designed to alter the properties of the original peptides ( $\delta$ -lysin, cecropin A, and magainin 2) in a way predicted by the hypotheses we had proposed. DL-1 and DL-2 variants were based on  $\delta$ -lysin, CE-1 and CE-2 on cecropin A, and MG-1 and MG-2 on magainin 2 (Table 1 and Figure 2). Set 1 (DL-1, CE-1, and MG-1) was designed to test hypotheses 1 and 2, that the peptide mechanism depends on the difficulty of peptide insertion, which can be altered by intramolecular salt bridges. The dye release should be graded or all-or-none depending on whether  $\Delta G_{\text{oct-if}}^{\circ}$  is below or above the threshold. Set 2 (DL-2a, DL-2b, CE-2, and MG-2) was designed to test hypothesis 3, which states that amino acid diversity in the sequence is not important for the mechanism as long as the thermodynamics of binding and insertion remain the same. Here, the sequences of the original peptides were simplified to contain a minimal number of amino acid types (hydrophobic, polar, and charged) in the same positions, but drastically reducing their diversity. In the



TP10 family of CPPs that we examined previously, the agreement between the Gibbs energies of binding measured experimentally ( $\Delta G_{\text{bind}}^{\circ}$ ) and calculated with the Wimley–White interfacial scale ( $\Delta G_{\text{if}}^{\circ}$ ) was excellent.<sup>16</sup> This is not true of all the peptides studied here.

In the next two sections, we compare the experimental value of the Gibbs energy of binding to POPC ( $\Delta G_{\text{bind}}^{\circ}$ ) with the value calculated with the Wimley–White scale ( $\Delta G_{\text{if}}^{\circ}$ ), which are listed in Table 3. Originally, we reasoned that considering salt bridge formation was justified if  $\Delta G_{\text{bind}}^{\circ}$  was more negative than  $\Delta G_{\text{if}}^{\circ}$  by approximately  $\geq 2$  kcal/mol.<sup>6</sup> Putative contributions from salt bridge formation are not included in the calculations of  $\Delta G_{\text{if}}^{\circ}$  and  $\Delta G_{\text{oct}}^{\circ}$  listed in Table 3, but we comment on their possible effects.

### Binding to POPC Membranes: Test of Hypothesis 2.

First, let us consider the peptides of set 1. In DL-1, the three acidic residues (Asp) of  $\delta$ -lysin were converted to basic ones (Lys), imparting on DL-1 a net charge of +6 at pH 7.5, instead of zero as for  $\delta$ -lysin. Because of the high charge, weak binding of DL-1 to POPC was predicted by the Wimley–White scale ( $\Delta G_{\text{if}}^{\circ} = -1.5$  kcal/mol), but we found that  $\Delta G_{\text{bind}}^{\circ} = -7.0$  kcal/mol (Table 3). This difference of 5.5 kcal/mol cannot be explained by salt bridge formation because DL-1 does not have any acidic residues. One possible salt bridge between a Lys side chain and the C-terminal carboxylate group would be insufficient to bring  $\Delta G_{\text{if}}^{\circ}$  and  $\Delta G_{\text{bind}}^{\circ}$  into agreement. CE-1 was designed to form one more intramolecular salt bridge than cecropin A, between the N-terminal amino group and Glu2. Peptide binding to POPC was therefore expected to improve relative to that of cecropin A. Indeed, CE-1 binds better than cecropin A to POPC, but for CE-1,  $\Delta G_{\text{bind}}^{\circ} \approx \Delta G_{\text{if}}^{\circ}$ . Thus, the reason for the improved binding is simply the better partitioning to the membrane interface of the amino acid side chains of CE-1, not salt bridge contributions. MG-1 was designed to form one intramolecular salt bridge (E7/K10,K11 or K10,K11/E13), the EK type being more likely than the KE type.<sup>63</sup> Binding of MG-1 to POPC should be similar to that of magainin 2 if no salt bridge formed, but better if it did. We found that  $\Delta G_{\text{bind}}^{\circ}$  ( $-7.4$  kcal/mol) for MG-1 is significantly more favorable than the calculated value ( $\Delta G_{\text{if}}^{\circ} = -3.6$  kcal/mol), but one salt bridge (the maximum possible) is insufficient to account for the difference.

The peptides of set 2 have the same distribution of amino acid types as the corresponding original peptides, and it was expected that their calculated  $\Delta G_{\text{if}}^{\circ}$  would be similar to the original ones. In DL-2b, the experimental values of  $\Delta G_{\text{bind}}^{\circ}$  are fairly close to those calculated. In DL-2a, however,  $\Delta G_{\text{if}}^{\circ} = -2.3$  kcal/mol but  $\Delta G_{\text{bind}}^{\circ} = -7.4$  kcal/mol. Three salt bridges would be sufficient to account for this difference, but it seems unlikely that this would happen only in DL-2a. CE-2 and MG-2 have the same charged amino acid residues as cecropin A and magainin 2, respectively, with net charges of +7 and +3, respectively, at pH 7.5. Like the parent peptides, they bind poorly to POPC. Nevertheless, reliable values of  $K_{\text{D}}$  in POPC were obtained, by extrapolating  $K_{\text{D}}$  in POPC/POPG mixtures, as a function of POPG content, to pure POPC. For CE-2,  $\Delta G_{\text{bind}}^{\circ} = -5.6$  kcal/mol, in very good agreement with the calculated  $\Delta G_{\text{if}}^{\circ}$  of  $-5.7$  kcal/mol. However, for MG-2, the values are different ( $\Delta G_{\text{bind}}^{\circ} = -6.4$ , and  $\Delta G_{\text{if}}^{\circ} = -2.7$  kcal/mol), and the difference cannot be explained by salt bridge formation.

The calculated  $\Delta G_{\text{if}}^{\circ}$  is small because of the low helical content of the peptide on the membrane (40%).

In general, the  $\Delta G_{\text{if}}^{\circ}$  value cannot be made to match  $\Delta G_{\text{bind}}^{\circ}$  by invoking salt bridge contributions to binding. Even though formation of those salt bridges cannot be directly assessed by the methods used, but only suggested by discrepancies in those two measures of the Gibbs energy of binding, it is clear that salt bridge formation fails to explain all the results reported here. Thus, either intramolecular salt bridges do not form on the membrane interface, even though partner residues were positioned with the correct spacing in the sequence, or they form but provide no significant contribution to  $\Delta G_{\text{bind}}^{\circ}$ . This is clearly shown by DL-1 compared to  $\delta$ -lysin, in which elimination of the putative salt bridges failed to abolish the discrepancy between  $\Delta G_{\text{if}}^{\circ}$  and  $\Delta G_{\text{bind}}^{\circ}$ . This is corroborated by CE-1 or MG-1, in which the possible salt bridges introduced failed to significantly improve binding. Thus, as a general mechanism for enhancing binding, hypothesis 2 is most probably wrong.

### Sources of Error in the Gibbs Energies of Binding.

What is the source of these discrepancies? Is there a significant error in the experimental values of the Gibbs free energy of binding ( $\Delta G_{\text{bind}}^{\circ} = RT \ln K_{\text{D}} - 2.4$  kcal/mol) or in the calculated ones ( $\Delta G_{\text{if}}^{\circ}$ ) that can explain the differences? In general, no. Let us consider the error in those thermodynamic parameters. Typically, the relative error in  $K_{\text{D}}$  arising from variability in experimental measurements is on the order of 30%, which corresponds to only  $\approx 0.2$  kcal/mol in  $\Delta G_{\text{bind}}^{\circ}$ . A larger source of error arises from the uncertainty regarding peptide translocation and therefore whether all the lipid or only the outer monolayer of each vesicle should be considered in calculating  $K_{\text{D}}$ . This corresponds to a factor of 2 in  $K_{\text{D}}$ , which translates into 0.4 kcal/mol in  $\Delta G_{\text{bind}}^{\circ}$ . Thus, in the worst case scenario, the cumulative error from those sources is  $\sim 0.6$  kcal/mol; this is  $<10\%$  of a typical value of  $\Delta G_{\text{bind}}^{\circ}$  ( $-7.0$  kcal/mol).

For  $\delta$ -lysin, DL-2a, DL-2b, CE-1, and MG-1, however, the measurements of  $K_{\text{D}}$  ( $=k_{\text{off}}/k_{\text{on}}$ ) proved to be difficult. The signal amplitudes in the stopped-flow fluorescence measurements of binding were small, which suggests that a significant fraction of the peptides were in a state that does not associate with membranes in the short time frame of the binding kinetics. The most likely explanation is that peptide oligomers exist in aqueous solution, which dissociate slowly. It is the monomer that binds to the membrane, and if its concentration is low, the signal is small. These five peptides have low or no net charge at pH 7.5, consistent with a tendency to oligomerize. This is known for  $\delta$ -lysin above  $1 \mu\text{M}$ .<sup>14,54–57</sup> This idea is corroborated by high helicities in aqueous buffer, considerably higher than those expected from AGADIR.<sup>53</sup> Oligomerization may distort the measured on and off rates, which are used to calculate  $K_{\text{D}}$ . Thus,  $K_{\text{D}}$  values for these five peptides may have an error larger than those for the others. However, as discussed below, there is a very good correlation between the experimental values of  $\Delta G_{\text{bind}}^{\circ}$  and the rate of induced dye release, suggesting that the estimate of binding affinities is nearly correct.

Consider now the error in  $\Delta G_{\text{if}}^{\circ}$ , calculated with MPEx using the Wimley–White interfacial scale, which originates from uncertainty in the helicity of the peptide on the membrane. Typically, the relative error in peptide helicity on the membrane arising from variability in experimental measurements is on the order of 5%. Using the actual standard deviations obtained in

the CD determinations, this yields errors of 0.1–0.6 kcal/mol in  $\Delta G_{if}^\circ$ . In DL-1, the error in  $\Delta G_{if}^\circ$  is  $\sim 0.2$  kcal/mol, clearly insufficient to bridge the difference of 5.5 kcal/mol relative to  $\Delta G_{bind}^\circ$ . The helicity would have to be increased from the experimental value of 52% to 95% to make  $\Delta G_{if}^\circ$  equal to  $\Delta G_{bind}^\circ$ . In cecropin A, the error in  $\Delta G_{if}^\circ$  is 0.5 kcal/mol, which is much smaller than the 3.7 kcal/mol difference between  $\Delta G_{if}^\circ$  and  $\Delta G_{bind}^\circ$ ; in fact, a helicity of 95% on the membrane would be necessary to bring the two values into agreement. In MG-1,  $\Delta G_{if}^\circ = -3.6$  kcal/mol with an error of  $\approx 0.6$  kcal/mol, which is insufficient to bring it even close to the  $\Delta G_{bind}^\circ$  of  $-7.4$  kcal/mol. The helicity of MG-1 is 69% in POPC, which is a typical average value for antimicrobial peptides, and even increasing it to 100% would only make  $\Delta G_{if}^\circ$  equal  $-6.4$  kcal/mol. As a counterexample, in CE-1 the formal error in  $\Delta G_{if}^\circ$  is 0.1 kcal/mol, but a small increase in helicity, from 55 to 62%, is sufficient to render  $\Delta G_{if}^\circ$  equal to  $\Delta G_{bind}^\circ$ ; it is reasonable that CD may underestimate the helicity by 7%.

In the case of magainin 2, the helicity is 57% in 1:1 POPC/POPG LUV; using this value,  $\Delta G_{if}^\circ = -3.8$  kcal/mol, whereas  $\Delta G_{bind}^\circ = -6.1$  kcal/mol. However, the helicity in pure POPC may be higher; in dodecylphosphocholine (DPC) micelles, the helicity is 83%,<sup>64</sup> which yields a  $\Delta G_{if}^\circ$  of  $-6.2$  kcal/mol, in agreement with the experimental value. A similar situation occurs with MG-2. The CD measurement had to be made in 1:1 POPC/POPG LUV because MG-2 does not bind well to POPC. If the helicity was 80% in pure POPC, the calculated and experimental values would agree. The cases of magainin 2 and MG-2 can thus be explained.

Another factor may be at play in DL-2a, and perhaps to a lesser extent in MG-1. Their helicities in aqueous solution are large, much greater than those expected from AGADIR, suggesting that the peptides oligomerize in water. When they are mixed with POPC to measure the CD, if only a small fraction dissociates from the oligomers and binds to the membrane, the measured value of the helicity may be heavily weighted by oligomers remaining in solution. However, we could not see a change in the CD spectrum of DL-2a added to 5 mM POPC even after 1 day. It may be a coincidence that the helicities in solution and on the membrane are the same, but it could also be that the amount of peptide bound to lipid is small. In that case, the value of the helicity measured “on the membrane” would be incorrect. If DL-2a were in fact as helical on the membrane as  $\delta$ -lysin (close to 100%), we would obtain a  $\Delta G_{if}^\circ$  of  $-7$  kcal/mol, in agreement with the experimental value.

In conclusion, in many cases, the experimental  $\Delta G_{bind}^\circ$  values do not agree with those calculated from the Wimley–White interfacial scale ( $\Delta G_{if}^\circ$ ). Whereas these differences can perhaps be attributed to errors in helicity in the cases of magainin 2, MG-2, and DL-2a, the discrepancies observed in cecropin A,  $\delta$ -lysin, DL-1, and MG-1 are real. There appear to be contributions to binding that are nonadditive and are therefore not captured by the simple sum of the Wimley–White parameters that  $\Delta G_{if}^\circ$  represents. We had previously explained the discrepancies by formation of salt bridges by the side chains of Lys and Arg with Asp and Glu. Now, that explanation is untenable in general. However, the inter-residue spacing that was required for formation of salt bridges remains in DL-1, because three Asp residues were changed to Lys, but their positions in the sequence were maintained. These residues are

close to each other in space in the folded, helical peptide on the membrane surface. It has recently been shown by simulation that transfer of spatially close Arg residues to the bilayer interior is nonadditive.<sup>65</sup> That is, once a first Arg is transferred to the bilayer and creates a water defect, additional Arg residues can transfer at essentially no energetic cost to the same defect. The presence of a water defect that hydrates the first charge is costly, but additional charges can use that defect for free. Could a similar situation occur with the transfer of the six Lys residues of DL-1 to the membrane interface? If the unfavorable contributions of all but one Lys were considered, a Gibbs energy of transfer ( $\Delta G_{if}^\circ$ ) of  $-7.5$  kcal/mol would be obtained, which is close to the experimental  $\Delta G_{bind}^\circ$  value of  $-7.0$  kcal/mol.


**Effect of Mutations on Dye Release Kinetics.** We now compare the effects of mutations on the rate of dye release. The peptide activity toward model membranes was characterized by measuring the kinetics of release of CF from LUV, under similar experimental conditions. The release kinetics were quantified by calculating the average characteristic time constants  $\tau$ , which are listed in Table 3.

The release curves of  $\delta$ -lysin characteristically exhibit a lag time (Figure 7A). This may be due to oligomerization on the membrane<sup>14,15</sup> or to an apparent slow on rate. The observed  $k_{on}$  is indeed small, which could be due to a rate-limiting dissociation of oligomers in solution. On the other hand, dye release by DL-1 has no lag time. Unlike  $\delta$ -lysin, which has zero net charge at pH 7.5, DL-1 has a charge of +6. This suggests that repulsive electrostatic interactions in DL-1 minimize oligomerization in solution and corroborates the idea that oligomerization of  $\delta$ -lysin in solution causes its apparent slow rate of binding. DL-1 was expected to be less efficient than the original  $\delta$ -lysin, because of a less favorable Gibbs energy of binding. However, the difference in the experimental Gibbs energies of binding is only  $\approx 1$  kcal/mol. Accordingly, dye release induced by DL-1 is only a factor of  $\approx 2$  slower than by  $\delta$ -lysin (Figure 7A and Table 3).

In the cases of CE-1 and MG-1, the rate of dye release was predicted to increase significantly, compared to those of cecropin A and magainin 2, and it did. This expectation was based on the improved binding observed ( $\Delta G_{bind}^\circ$  changed from  $-6.4$  to  $-7.0$  kcal/mol in CE-1 and from  $-6.1$  to  $-7.4$  kcal/mol in MG-1), but also on a lower  $\Delta G_{oct-if}^\circ$  in the case of CE-1, which decreased by 13 kcal/mol relative to that of cecropin A. As predicted, dye release became dramatically faster for both mutants compared with the original peptides (Figure 7B,C).

DL-2a and DL-2b were expected to exhibit dye release kinetics similar to those of  $\delta$ -lysin. Indeed, their release curves (Figure 7D) show the sigmoidal shape characteristic of  $\delta$ -lysin (Figure 7A). The much lower efficiency of DL-2a and DL-2b in dye release, with  $\tau$  values of 14.8 and 650 s compared to a  $\tau$  of 2.2 s for  $\delta$ -lysin, is probably a consequence of a greater extent of oligomerization in solution. As expected, CE-2 and MG-2 are approximately as inefficient as cecropin A and magainin 2 in causing the release of dye from POPC LUV, which occurs on time scales on the order of 10000 s for these four peptides (Table 3).

To understand these results, let us consider the Gibbs energies of peptide binding and insertion. The caveats regarding the experimental  $\Delta G_{bind}^\circ$  for some of the peptides notwithstanding, there is a clear correlation between the binding affinity for the membrane and the characteristic time  $\tau$  of peptide-induced dye release (Figure 9A). Previously, we argued that the mean time



**Figure 9.** (A) Gibbs energy of binding to the membrane interface determined experimentally ( $\Delta G_{\text{bind}}^{\circ}$ ) as a function of the mean characteristic time  $\tau$  for release of CF from POPC LUV. Each gray symbol corresponds to a peptide examined here. Data for  $\delta$ -lysin, DL-1, DL-2a, and DL-2b are shown with gray triangles, data for cecropin A, CE-1, and CE-2 with gray circles, and data for magainin 2, MG-1, and MG-2 with gray squares. The data for TP10 variants previously published<sup>16</sup> are shown here for comparison (diamonds): TP10W, TPW-1, TPW-2, and TPW-3; TP10, TP10-COO<sup>-</sup>, TP10W-COO<sup>-</sup>, and TP10-7MC.<sup>38,58</sup> The black symbols correspond to experimental data; the white symbols correspond to data for TP10 and TP10-COO<sup>-</sup>, for which  $\tau$  is experimental but the binding affinity is calculated with the Wimley–White scale (neither contains Trp). The straight line is a fit with a slope of 0.5. The dashed line represents qualitatively the expected behavior limited by diffusion or bilayer response. (B) Gibbs energy of transfer to octanol calculated with the Wimley–White scale ( $\Delta G_{\text{oct}}^{\circ}$ ) as a function of mean characteristic time  $\tau$  for release of CF from POPC LUV. The points correspond to the same peptides as in panel A, and the same symbols were used. Again, the TP10 variant data are from our previous paper.<sup>16</sup> The white circle corresponds to TP10-7MC for which  $\Delta G_{\text{oct}}^{\circ}$  was estimated assuming Tyr for the Lys-MC residue.<sup>16</sup> The lines are fits with slopes of 1.

constant for dye release ( $\tau$ ) should depend on the Gibbs energy of binding ( $\Delta G_{\text{bind}}^{\circ}$ ) and on the activation free energy for insertion into the bilayer ( $\Delta G^{\ddagger}$ ) through an equation of the Arrhenius type<sup>16</sup>

$$1/\tau = A_0 e^{-\Delta G_{\text{bind}}^{\circ}/RT} e^{-\Delta G^{\ddagger}/RT} \quad (9)$$

The pre-exponential factor  $A_0$  includes effects of lipid and peptide concentrations as well as molecular details specific for each peptide family but should remain fairly constant within a given family under similar experimental conditions. This is the case for the TP10 family.<sup>16</sup> According to eq 9, if the activation free energy for insertion  $\Delta G^{\ddagger} \approx \Delta G_{\text{oct-if}}^{\circ}$ ,<sup>66</sup> and recalling that  $\Delta G_{\text{oct-if}}^{\circ}$  was calculated using the experimental value of  $\Delta G_{\text{bind}}^{\circ}$  ( $\Delta G_{\text{oct-if}}^{\circ} = \Delta G_{\text{oct}}^{\circ} - \Delta G_{\text{bind}}^{\circ}$ ), we obtain

$$1/\tau = A_0 e^{-\Delta G_{\text{oct}}^{\circ}/RT} \quad (10)$$

which, after taking logarithms and multiplying by  $-RT$ , yields

$$\Delta G_{\text{oct}}^{\circ} = RT \ln \tau + \text{constant} \quad (11)$$

Hence, a linear relation with a slope of 1 is expected between  $\Delta G_{\text{oct}}^{\circ}$  and  $RT \ln \tau$ , which was observed for the family of the

cell-penetrating peptide TP10 (Figure 9B, diamonds). When the data for the other three peptide families now examined are compared (Figure 9B, gray symbols), we notice that the dispersion is much larger, though the average behavior is still captured by a straight line with a slope of 1 (dashed line). Within each family, the data are too sparse to allow any trends to be gleaned. As a whole, however, it is clear that the mutants of antimicrobial and cytolytic peptides behaved differently from the TP10 variants, in that the Gibbs energy of transfer into the nonpolar interior of the membrane is much more unfavorable. A small  $\Delta G_{\text{oct-if}}^{\circ}$ , which is a consequence of a small  $\Delta G_{\text{oct}}^{\circ}$ , would allow the CPPs, but not the antimicrobial and cytolytic peptides, to translocate across the membrane.

It is also clear from Figure 9 that  $\Delta G_{\text{bind}}^{\circ}$  is a much better overall predictor of the peptide activity (measured by  $\tau$ ) than  $\Delta G_{\text{oct}}^{\circ}$ . Previously, on the basis of only the TP10 variants, we thought that the Gibbs energy of insertion was the most important parameter, which led to eq 11.<sup>16</sup> Now, with a larger data set, it is apparent that the only point that deviates significantly from the line in Figure 9A is that corresponding to TPW-2, on the left of the plot. A deviation from linearity at very low  $\tau$  is, however, unavoidable. No matter how good binding is,  $\tau$  eventually approaches a minimum value ( $\tau_{\text{min}}$ ) that is determined by the diffusion limit for peptide binding to the membrane and by the time of bilayer response. The diffusion limit for binding to LUV ( $k_{\text{lim}}$ ) is  $\approx 10^6 \text{ M}^{-1} \text{ s}^{-1}$  expressed in terms of lipids (1 LUV  $\approx 10^5$  lipids). With a lipid concentration ( $c$ ) of  $50 \mu\text{M}$ , this means that the maximum rate of binding is  $k_{\text{lim}}c = 1/\tau_{\text{min}} = 50 \text{ s}^{-1}$ . Thus,  $\tau \geq 0.02 \text{ s}$ , and  $RT \ln \tau \geq -2.3 \text{ kcal/mol}$ . This limit can be only asymptotically approached as  $\Delta G_{\text{bind}}^{\circ} \rightarrow -\infty$ . The dashed line in Figure 9A depicts qualitatively the expected behavior limited by diffusion or bilayer response. What these results indicate is that binding is the primary step that determines peptide activity.

**Gibbs Energy of Insertion: Test of Hypotheses 1 and 3.** We have determined the type of dye release using the ANTS/DPX requenching assay and confronted the predictions of hypotheses 1 and 3 with the results. The Gibbs energy of insertion was estimated by  $\Delta G_{\text{oct-if}}^{\circ}$ , where the experimental value of  $\Delta G_{\text{bind}}^{\circ}$  was used for the Gibbs energy of binding to the interface. These values are listed in Table 3, which also includes a summary of the ANTS/DPX results. According to hypothesis 1, if the type of release provides an indication of peptide translocation, a  $\Delta G_{\text{oct-if}}^{\circ}$  of  $<20 \text{ kcal/mol}$  corresponds to graded release and a  $\Delta G_{\text{oct-if}}^{\circ}$  of  $>23 \text{ kcal/mol}$  to all-or-none release. These predictions were corroborated by experiments for DL-1, CE-1, and MG-1. On the basis of our original calculations (which included salt bridges),  $\delta$ -lysin was in a gray zone between 20 and 23 kcal/mol; without salt bridges, it should be all-or-none, but experimentally, it is weakly graded.<sup>15</sup> In DL-1, a change to all-or-none release was sought and obtained (Figure 8A). In DL-2a, the type of release of  $\delta$ -lysin was expected to be maintained, and it was (Figure 8D), although without including salt bridges both should be all-or-none. In CE-1 and MG-1, dye release should have remained all-or-none, as in cecropin A<sup>31</sup> and magainin 2,<sup>32,61,62</sup> and it did (Figure 8B,C).

The most interesting results, however, were obtained for CE-2 and MG-2. Both peptides had been designed to remain all-or-none, like cecropin A and magainin 2; the sequences were



simplified, but the mutations were such that the thermodynamics should not have changed. Indeed, the measured  $\Delta G_{\text{bind}}^{\circ}$  values were very close to prediction, and  $\Delta G_{\text{oct-if}}^{\circ}$  values of >25 kcal/mol put these two peptides squarely in the all-or-none class. However, dye release is clearly graded (Figure 8E,F). This demonstrates that the simplification of the sequence can dramatically change the type of release, and hypothesis 3 is wrong. The question addressed by hypothesis 3 was whether sequence diversity is important. In mutants CE-2 and MG-2, the same amino acid types were kept in the same positions. Namely, polar charged residues were conserved, and hydrophobic residues were conserved; but sequence diversity was reduced to a minimum number of amino acids. The situation is similar in the folding problem of globular proteins in water, when the diversity of amino acids is reduced to only hydrophobic (H) and polar (P) types, which is embodied in the HP model.<sup>29</sup> Whereas the HP model captures many features of the protein folding problem, it does not generate unique native conformations.<sup>87</sup> Replacing the HP, two-letter alphabet with a multiletter one, that is, increasing amino acid diversity in the sequence, reduces the degeneracy of the folded state.<sup>68</sup> Furthermore, two- and three-letter alphabet models fold much less cooperatively than 20-letter ones.<sup>69</sup> Experimentally as well, simplified sequences of real proteins fail to fold to unique structures. A three-letter alphabet model of protein G folds to a fluidlike molten globule, but this folded structure is more flexible than the original protein.<sup>70</sup> The low-diversity code produces conformations with abundant secondary structure but is not sufficiently specific to yield a unique tertiary structure. The 20-letter alphabet is thought to eliminate unwanted amino acid contacts and prevent aggregation. Clearly, amino acid diversity in the sequence is also fundamental for the activity of these antimicrobial peptide variants.

Hypothesis 1 is not disproved by these results because its most fundamental statement is that translocation requires  $\Delta G_{\text{oct-if}}^{\circ}$  to be <20 kcal/mol. We had observed that all the peptides whose kinetic mechanism required translocation caused graded release.<sup>6</sup> Now, however, it is clear that either the correlation between translocation and graded release breaks down or hypothesis 1 is wrong. Previously, we proposed that graded release occurs because peptides translocate and dissipate the mass imbalance across the membrane before all the dye has time to escape.<sup>15</sup> Another possibility, however, is that the peptides constantly but slowly perturb the membrane, without ever translocating or causing a major disruption. In that case, release should be complete; indeed, CE-2 induces complete but graded dye release, consistent with this idea.

Thus, either the first or the second implication in each of the statements (eqs 1 and 2) that constitute hypothesis 1 is wrong, but we do not yet know which one. The results obtained with CE-2 and MG-2 show that the type of dye release cannot be predicted on the basis of the Gibbs energy of insertion, and even very conservative changes to the peptide sequence can result in complete switches from all-or-none to graded release. Clearly, it is necessary to determine directly whether the correlation between translocation and graded release holds. If the correlation proves correct for CE-2 and MG-2, hypothesis 1 will be disproved. This will be investigated in future work.

## CONCLUSION

We have attempted to disprove our hypotheses by examining the binding and activity of variants of antimicrobial and

cytolytic peptides in model membranes. First, the results indicate that binding to the membrane, not insertion, is the primary determinant of peptide activity. This is contrary to the concept underlying the proposed hypotheses, in which the Gibbs energy of insertion, estimated by  $\Delta G_{\text{oct-if}}^{\circ}$ , was central. Nevertheless,  $\Delta G_{\text{oct-if}}^{\circ}$  is much more unfavorable for the families of antimicrobial and cytolytic peptides than for variants of the amphipathic cell-penetrating peptide TP10 previously examined.<sup>16</sup> A smaller  $\Delta G_{\text{oct-if}}^{\circ}$  would allow the CPPs, but not the antimicrobial and cytolytic peptides, to translocate across the membrane, as predicted. However, either graded release is not a reliable indicator of peptide translocation, or the first hypothesis is wrong. Second, placing acidic and basic residues along the peptide sequence, with the correct spacing for the establishment of salt bridges by their side chains, does not result in the consistent improvement of binding. Either these salt bridges do not form at the membrane interface, or they form but do not decrease  $\Delta G_{\text{bind}}^{\circ}$ . Third, simplification of the sequence to a few amino acid types, even through very conservative mutations, dramatically changes the mechanism of dye release (though not the rate). Thus, amino acid diversity in the sequence seems to be essential for the function of even these simple polypeptides on the membrane, as it is essential for the folding of globular proteins in water.

## AUTHOR INFORMATION

### Corresponding Author

\*Telephone: (910) 962-7300. Fax: (910) 962-3013. E-mail: almeidap@uncw.edu.

### Funding

This work was supported by National Institutes of Health Grant GM072507, including an ARRA supplement.

## ACKNOWLEDGMENTS

We thank Alex Kreutzberger for performing additional experiments with  $\delta$ -lysin and Antje Pokorny for many discussions.

## ABBREVIATIONS

$\Delta G_{\text{if}}^{\circ}$ , Gibbs energy of peptide binding to the membrane interface, as a helix, calculated with the Wimley–White interfacial scale;  $\Delta G_{\text{oct}}^{\circ}$ , Gibbs energy of transfer of the peptide from water to octanol;  $\Delta G_{\text{oct-if}}^{\circ}$ ,  $\Delta G_{\text{oct}}^{\circ} - \Delta G_{\text{if}}^{\circ}$ ;  $\Delta G_{\text{ins}}^{\circ}$ , Gibbs energy of insertion from the surface into the membrane;  $\Delta G_{\text{bind}}^{\circ}$ , Gibbs energy of binding derived from experiment;  $\Delta G_{\text{f}}^{\circ}$ , Gibbs energy of folding to an  $\alpha$ -helix in water;  $\Delta G^{\ddagger}$ , Gibbs energy of the transition state;  $k_{\text{on}}$ , on rate constant;  $k_{\text{off}}$ , off rate constant;  $K_{\text{D}}$ , equilibrium dissociation constant; SUV, small unilamellar vesicles; LUV, large unilamellar vesicles; Tp10, transportan 10; POPC, 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine; POPG, 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphoglycerol; POPs, 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphoserine; DPC, dodecylphosphocholine; 7MC, 7-methoxycoumarin-3-carboxylic acid; ANTS, 8-amino-naphthalene-1,3,6-trisulfonic acid; DPX, *p*-xylene bis-pyridinium bromide; FRET, Förster (fluorescence) resonance energy transfer; NMR, nuclear magnetic resonance; CD, circular dichroism; MPEx, Membrane Protein Explorer.

# REFERENCES

- (1) Zasloff, M. (2002) Antimicrobial peptides of multicellular organisms. *Nature* 415, 389–395.
- (2) Giangaspero, A., Sandri, L., and Tossi, A. (2001) Amphipathic  $\alpha$ -helical antimicrobial peptides. *Eur. J. Biochem.* 268, 5589–5600.
- (3) Tossi, A., Sandri, L., and Giangaspero, A. (2000) Amphipathic,  $\alpha$ -helical antimicrobial peptides. *Biopolymers* 55, 4–30.
- (4) Nickel, W. (2003) The mystery of nonclassical protein secretion. A current view on cargo proteins and potential export routes. *Eur. J. Biochem.* 270, 2109–2119.
- (5) Lindgren, M., and Langel, U. (2011) Clases and prediction of cell-penetrating peptides. *Methods Mol. Biol.* 683, 3–19.
- (6) Almeida, P. F., and Pokorny, A. (2009) Mechanism of antimicrobial, cytolytic, and cell-penetrating peptides: From kinetics to thermodynamics. *Biochemistry* 48, 8083–8093.
- (7) Splith, K., and Neundorff, I. (2011) Antimicrobial peptides with cell-penetrating peptide properties and vice versa. *Eur. Biophys. J.* 40, 387–397.
- (8) Ehrenstein, G., and Lehar, H. (1977) Electrically gated ionic channels in lipid bilayers. *Q. Rev. Biophys.* 10, 1–34.
- (9) Pouny, Y., Rapaport, D., Mor, A., Nicolas, P., and Shai, Y. (1992) Interaction of antimicrobial dermaseptin and its fluorescently labeled analogues with phospholipid membranes. *Biochemistry* 31, 12416–12423.
- (10) Shai, Y. (2002) Mode of action of membrane active antimicrobial peptides. *Biopolymers* 66, 236–248.
- (11) Papo, N., and Shai, Y. (2003) Can we predict biological activity of antimicrobial peptides from their interactions with model phospholipid membranes? *Peptides* 24, 1693–1703.
- (12) Ludtke, S. J., He, K., Heller, W. T., Harroun, T. A., Yang, L., and Huang, H. W. (1996) Membrane pores induced by magainin. *Biochemistry* 35, 13723–13728.
- (13) Matsuzaki, K., Murase, O., Fujii, N., and Miyajima, K. (1996) An antimicrobial peptide, magainin 2, induced rapid flip-flop of phospholipids coupled with pore formation and peptide translocation. *Biochemistry* 35, 11361–11368.
- (14) Pokorny, A., Birkbeck, T. H., and Almeida, P. F. F. (2002) Mechanism and kinetics of  $\delta$ -lysin interaction with phospholipid vesicles. *Biochemistry* 41, 11044–11056.
- (15) Pokorny, A., and Almeida, P. F. F. (2004) Kinetics of dye efflux and lipid flip-flop induced by  $\delta$ -lysin in phosphatidylcholine vesicles and the mechanism of graded release by amphipathic,  $\alpha$ -helical peptides. *Biochemistry* 43, 8846–8857.
- (16) McKeown, A. N., Naro, J. L., Huskins, L. J., and Almeida, P. F. (2011) A thermodynamic approach to the mechanism of cell-penetrating peptides in model membranes. *Biochemistry* 50, 654–662.
- (17) Jayasinghe, S., Hristova, K., and White, S. H. (2001) Energetics, stability, and prediction of transmembrane helices. *J. Mol. Biol.* 312, 927–934.
- (18) White, S. H., and Wimley, W. C. (1999) Membrane protein folding and stability: Physical principles. *Annu. Rev. Biophys. Biomol. Struct.* 28, 319–365.
- (19) Wimley, W. C., and White, S. H. (1996) Experimentally determined hydrophobicity scale of proteins at membrane interfaces. *Nat. Struct. Biol.* 3, 842–848.
- (20) Hristova, K., and White, S. H. (2005) An experiment-based algorithm for predicting the partitioning of unfolded peptides into phosphatidylcholine bilayer interfaces. *Biochemistry* 44, 12614–12619.
- (21) Ladokhin, A. S., and White, S. H. (1999) Folding of amphipathic  $\alpha$ -helices on membranes: Energetics of helix formation by melittin. *J. Mol. Biol.* 285, 1363–1369.
- (22) Wimley, W. C., Creamer, T. P., and White, S. H. (1996) Solvation energies of amino acid side chains and backbone in a family of host-guest pentapeptides. *Biochemistry* 35, 5109–5124.
- (23) Kreger, A. S., Kim, K.-S., Zaboretzky, F., and Bernheimer, A. W. (1971) Purification and properties of staphylococcal delta hemolysin. *Infect. Immun.* 3, 449–465.
- (24) Hultmark, D., Steiner, H., Rasmuson, T., and Boman, H. G. (1980) Insect immunity. Purification and properties of three inducible bactericidal proteins from hemolymph of immunized pupae of *Hyalophora cecropia*. *Eur. J. Biochem.* 106, 7–16.
- (25) Zasloff, M. (1987) Magainins, a class of antimicrobial peptides from *Xenopus* skin: Isolation, characterization of two active forms, and partial cDNA sequence of a precursor. *Proc. Natl. Acad. Sci. U.S.A.* 84, 5449–5453.
- (26) Soomets, U., Lindgren, M., Gallet, X., Hallbrink, M., Elmquist, A., Balaspiri, L., Zorko, M., Pooga, M., Brasseur, R., and Langel, U. (2000) Deletion analogues of transportan. *Biochim. Biophys. Acta* 1467, 165–176.
- (27) Hallbrink, M., Floren, A., Elmquist, A., Pooga, M., Bartfai, T., and Langel, U. (2001) Cargo delivery kinetics of cell-penetrating peptides. *Biochim. Biophys. Acta* 1515, 101–109.
- (28) Wimley, W. C., Gawrisch, K., Creamer, T. P., and White, S. H. (1996) Direct measurement of salt-bridge solvation energies using a peptide model system: Implications for protein stability. *Proc. Natl. Acad. Sci. U.S.A.* 93, 2985–2990.
- (29) Lau, K. F., and Dill, K. A. (1989) A lattice statistical-mechanics model of the conformational and sequence spaces of proteins. *Macromolecules* 22, 3986–3997.
- (30) Frazier, M. L., Wright, J. R., Pokorny, A., and Almeida, P. F. F. (2007) Investigation of domain formation in sphingomyelin/cholesterol/POPC mixtures by fluorescence resonance energy transfer and Monte Carlo simulations. *Biophys. J.* 92, 2422–2433.
- (31) Gregory, S. M., Cavanaugh, A. C., Journigan, V., Pokorny, A., and Almeida, P. F. F. (2008) A quantitative model for the all-or-none permeabilization of phospholipid vesicles by the antimicrobial peptide cecropin A. *Biophys. J.* 94, 1667–1680.
- (32) Gregory, S. M., Pokorny, A., and Almeida, P. F. F. (2009) Magainin 2 revisited: A test of the quantitative model for the all-or-none permeabilization of phospholipid vesicles. *Biophys. J.* 96, 116–131.
- (33) Almeida, P. F., and Pokorny, A. (2010) Binding and Permeabilization of Model Membranes by Amphipathic Peptides. *Methods Mol. Biol.* 618, 155–169.
- (34) Vaz, W. L. C., and Hallmann, D. (1983) Experimental evidence against the applicability of the Saffman-Delbrück model to the translational diffusion of lipids in phosphatidylcholine bilayer membranes. *FEBS Lett.* 152, 287–290.
- (35) Bartlett, G. R. (1959) Phosphorous assay in column chromatography. *J. Biol. Chem.* 234, 466–468.
- (36) Ladokhin, A. S., Fernandez-Vidal, M., and White, S. H. (2010) CD spectroscopy of peptides and proteins bound to large unilamellar vesicles. *J. Membr. Biol.* 236, 247–253.
- (37) Luo, P., and Baldwin, R. L. (1997) Mechanism of helix induction by trifluoroethanol: A framework for extrapolating the helix-forming properties of peptides from trifluoroethanol/water mixtures back to water. *Biochemistry* 36, 8413–8421.
- (38) Yandek, L. E., Pokorny, A., Floren, A., Knoelke, K., Langel, U., and Almeida, P. F. F. (2007) Mechanism of the cell-penetrating peptide Tp10 permeation of lipid bilayers. *Biophys. J.* 92, 2434–2444.
- (39) Yandek, L. E., Pokorny, A., and Almeida, P. F. F. (2009) Wasp mastoparans follow the same mechanism as the cell-penetrating peptide transportan 10. *Biochemistry* 48, 7342–7351.
- (40) Wimley, W. C., Selsted, M. E., and White, S. H. (1994) Interactions between human defensins and lipid bilayers: Evidence for formation of multimeric pores. *Protein Sci.* 3, 1362–1373.
- (41) Ladokhin, A. S., Wimley, W. C., and White, S. H. (1995) Leakage of membrane vesicle contents: Determination of mechanism using fluorescence quenching. *Biophys. J.* 69, 1964–1971.

- (42) Ladokhin, A. S., Wimley, W. C., Hristova, K., and White, S. H. (1997) Mechanism of leakage of contents of membrane vesicles determined by fluorescence reequenching. *Methods Enzymol.* 278, 474–486.
- (43) Pokorny, A., and Almeida, P. F. F. (2005) Permeabilization of raft-containing lipid vesicles by  $\delta$ -lysin: A mechanism for cell sensitivity to cytotoxic peptides. *Biochemistry* 44, 9538–9544.
- (44) Pokorny, A., Yandek, L. E., Elegbede, A. I., Hinderliter, A., and Almeida, P. F. F. (2006) Temperature and composition dependence of the interaction of  $\delta$ -lysin with ternary mixtures of sphingomyelin/cholesterol/POPC. *Biophys. J.* 91, 2184–2197.
- (45) Colquhoun, D. (1971) *Lectures on Biostatistics*, Clarendon Press, Oxford, U.K.
- (46) Colquhoun, D., and Hawkes, A. G. (1987) The interpretation of single channel recordings. In *Microelectrode Techniques. The Plymouth Workshop Handbook* (Ogden, D., Ed.) 2nd ed., pp 141–188, The Company of Biologists Ltd., Cambridge, U.K.
- (47) Jaysinghe, S., Hristova, K., Wimley, W., Snider, C., and White, S. H. (2010) *Membrane Protein Explorer*. <http://blanco.biomol.uci.edu/MPEX>.
- (48) Muñoz, V., and Serrano, L. (1994) Elucidating the folding problem of helical peptides using empirical parameters. *Nat. Struct. Biol.* 1, 399–409.
- (49) Muñoz, V., and Serrano, L. (1994) Elucidating the folding problem of  $\alpha$ -helical peptides using empirical parameters, II. Helix macrodipole effects and rational modification of the helical content of natural peptides. *J. Mol. Biol.* 245, 275–296.
- (50) Muñoz, V., and Serrano, L. (1994) Elucidating the folding problem of  $\alpha$ -helical peptides using empirical parameters. III: Temperature and pH dependence. *J. Mol. Biol.* 245, 297–308.
- (51) Muñoz, V., and Serrano, L. (1997) Development of the multiple sequence approximation within the Agadir model of  $\alpha$ -helix formation. Comparison with Zimm-Bragg and Lifson-Roig formalisms. *Biopolymers* 41, 495–509.
- (52) Lacroix, E., Viguera, A. R., and Serrano, L. (1998) Elucidating the folding problem of  $\alpha$ -helices: Local motifs, long-range electrostatics, ionic strength dependence and prediction of NMR parameters. *J. Mol. Biol.* 284, 173–191.
- (53) AGADIR, an algorithm to predict the helical content of peptides, is available on-line at <http://agadir.crg.es/>.
- (54) Fitton, J. E., Dell, A., and Shaw, W. V. (1980) The amino acid sequence of the delta haemolysin of *Staphylococcus aureus*. *FEBS Lett.* 115, 209–212.
- (55) Fitton, J. E. (1981) Physicochemical studies on delta haemolysin, a staphylococcal cytolytic polypeptide. *FEBS Lett.* 130, 257–260.
- (56) Thiaudière, E., Siffert, O., Talbot, J. C., Bolard, J., Alouf, J. E., and Dufourcq, J. (1991) The amphiphilic  $\alpha$ -helix concept. Consequences on the structure of staphylococcal  $\delta$ -toxin in solution and bound to lipids. *Eur. J. Biochem.* 195, 203–213.
- (57) Talbot, J. C., Thiaudière, E., Vincent, M., Gallay, J., Siffert, O., and Dufourcq, J. (2001) Dynamics and orientation of amphipathic peptides in solution and bound to membranes: A steady-state and time-resolved fluorescence study of staphylococcal  $\delta$ -toxin and its synthetic analogues. *Eur. Biophys. J.* 30, 147–161.
- (58) Yandek, L. E., Pokorny, A., and Almeida, P. F. F. (2008) Small changes in the primary structure of transportan 10 alter the thermodynamics and kinetics of its interaction with phospholipid vesicles. *Biochemistry* 47, 3051–3060.
- (59) Pokorny, A., Kilelee, E. M., Wu, D., and Almeida, P. F. F. (2008) The activity of the amphipathic peptide  $\delta$ -lysin correlates with phospholipid acyl chain structure and bilayer elastic properties. *Biophys. J.* 95, 4748–4755.
- (60) Silvestro, L., Gupta, K., Weiser, J. N., and Axelsen, P. H. (1997) The concentration-dependent membrane activity of cecropin A. *Biochemistry* 36, 11452–11460.
- (61) Tamba, Y., and Yamazaki, M. (2005) Single giant unilamellar vesicle method reveals effect of antimicrobial peptide magainin 2 on membrane permeability. *Biochemistry* 44, 15823–15833.
- (62) Tamba, Y., and Yamazaki, M. (2009) Magainin 2-induced pore formation in the lipid membranes depends on its concentration in the membrane interface. *J. Phys. Chem. B* 113, 4846–4852.
- (63) Marqusee, S., and Baldwin, R. L. (1987) Helix stabilization by Glu<sup>−</sup>...Lys<sup>+</sup> salt bridges in short peptides of *de novo* design. *Proc. Natl. Acad. Sci. U.S.A.* 84, 8898–8902.
- (64) Gesell, J., Zasloff, M., and Opella, S. J. (1997) Two-dimensional <sup>1</sup>H-NMR experiments show that the 23-residue magainin antibiotic peptide is an  $\alpha$ -helix in dodecylphosphocholine micelles, sodium dodecylsulfate micelles, and trifluoroethanol/water solution. *J. Biomol. NMR* 9, 127–135.
- (65) MacCallum, J. L., Bennett, W. F. D., and Tieleman, D. P. (2011) Transfer of arginine into lipid bilayers is nonadditive. *Biophys. J.* 101, 110–117.
- (66) Hammond, G. S. (1955) A correlation of reaction rates. *J. Am. Chem. Soc.* 77, 334–338.
- (67) Shakhnovich, E. (2006) Protein folding thermodynamics and dynamics: Where physics, chemistry, and biology meet. *Chem. Rev.* 106, 1559–1588.
- (68) Yue, K., Fiebig, K. M., Thomas, P. D., Chan, H. S., Shakhnovich, E. I., and Dill, K. A. (1995) *Proc. Natl. Acad. Sci. U.S.A.* 92, 325–329.
- (69) Kaya, H., and Chan, H. S. (2000) Energetic components of cooperative protein folding. *Phys. Rev. Lett.* 85, 4823–4826.
- (70) Guarnera, E., Pellarin, R., and Caflisch, A. (2009) How does a simplified-sequence protein fold? *Biophys. J.* 97, 1737–1746.
- (71) Matsuzaki, K., Murase, O., Tokuda, H., Funakoshi, S., Fujii, N., and Miyajima, K. (1994) Orientational and aggregational states in magainin 2 in phospholipid bilayers. *Biochemistry* 33, 3342–3349.