

- Engle, R. (1977) *Chem. Rev.* 77, 349.
- Gerlt, J. A., & Whitman, G. J. R. (1975) *J. Biol. Chem.* 250, 5053.
- Gray, M. W. (1976) *Can. J. Biochem.* 54, 413.
- Hanes, L. S., & Isherwood, F. A. (1949) *Nature (London)* 164, 1107.
- Kelly, S. J., & Butler, L. G. (1975) *Biochem. Biophys. Res. Commun.* 66, 316.
- Kelly, S. J., & Butler, L. G. (1977) *Biochemistry* 16, 1102.
- Kelly, S. J., Dardinger, D. E., & Butler, L. G. (1975) *Biochemistry* 14, 4983.
- Kezdy, F. J., & Bender, M. L. (1962) *Biochemistry* 1, 1097.
- Landt, M., & Butler, L. G. (1978) *Biochemistry* 17, 4130.
- Landt, M., Boltz, S. C., & Butler, L. G. (1978) *Biochemistry* 17, 915.
- Laskowski, M., Sr. (1971) *Enzymes*, 3rd Ed. 4, 313.
- Letsinger, R. L., Wilkes, J. S., & Dumas, L. B. (1976) *Biochemistry* 15, 2810.
- Miller, P. S., Fang, K. N., Kondo, N. S., & T'so, P. O. P. (1971) *J. Am. Chem. Soc.* 93, 6657.
- Nichol, A. W., Nomura, A., & Hampton, A. (1967) *Biochemistry* 6, 1008.
- Ogilvie, K. K., & Hruska, F. H. (1976) *Biochem. Biophys. Res. Commun.* 68, 375.
- Osterman, H. L., & Walz, F. G., Jr. (1978) *Biochemistry* 17, 4124.
- Smith, M., Brummond, G. I., & Khorana, H. G. (1961) *J. Am. Chem. Soc.* 83, 698.
- Tolkmitz, M. C. (1958) *J. Org. Chem.* 23, 1685.
- Tsou, K. C., & Yip, K. F. (1973) *Life Sci.* 13, 1505.
- Uesugi, S., Yano, J., Yano, E., & Idehara, M. (1977) *J. Am. Chem. Soc.* 99, 2313.
- Webb, J. L. (1963) *Enzyme and Metabolic Inhibitors*, Vol. 1, Academic Press, New York.
- Yarbrough, L. E. (1978) *Biochem. Biophys. Res. Commun.* 81, 35.


## Adenosine Cyclic 3',5'-Monophosphate Dependent Protein Kinase: Active Site Directed Inhibition by Cibacron Blue F3GA<sup>†</sup>

Jonathan J. Witt and Robert Roskoski, Jr.\*

**ABSTRACT:** Cibacron Blue F3GA, the polycyclic blue chromophore of Blue Dextran 2000, inhibits the catalytic subunit of bovine brain protein kinase. The rate of inactivation exhibits a hyperbolic dependence on the dye concentration. This suggests that an enzyme-dye complex forms prior to inactivation. Protein and peptide substrates or MgATP protects the enzyme against dye inactivation. Kinetic measurements show that the dissociation constant is  $\sim 100 \mu\text{M}$  and the maximal rate of inactivation is  $0.13 \text{ min}^{-1}$  at  $22^\circ\text{C}$ . Inactivation is temperature and time dependent. Exhaustive dialysis, gel filtration, or the addition of substrate fails to reactivate

inhibited enzyme. The failure to reverse the inhibition suggests that the dye forms a covalent complex with the enzyme. Denaturation by sodium dodecyl sulfate also fails to dissociate the dye from enzyme. The hyperbolic kinetics, moreover, suggest that the dye functions as an active site directed reagent. The holoenzyme is resistant to Cibacron Blue inactivation. Addition of cAMP converts the enzyme to a form susceptible to inhibition. In agreement with our previous studies, these results also suggest that the regulatory subunit shields, either physically or functionally, the active site of the catalytic subunit.

Adenosine cyclic 3',5'-monophosphate (cAMP)<sup>1</sup> dependent protein kinase consists of two dissimilar subunits (Gill & Garen, 1970; Tao et al., 1970; Kumon et al., 1970; Reimann et al., 1971). The activation of this enzyme by cAMP is summarized by



in which  $\text{R}_2\text{C}_2$  is the holoenzyme and R and C are the regulatory and catalytic subunits, respectively (Rosen & Erlichman, 1975; Hofmann et al., 1975). The free C subunit is active and cAMP independent. Combining C subunit and the  $(\text{R-cAMP})_2$  complex results in the release of cAMP and the regeneration of the cAMP-dependent holoenzyme.

We reported that the free C subunit binds to Blue Dextran-Sephacrose affinity resin. The blue chromophore of Blue

Dextran, Cibacron Blue F3GA, is a nucleotide analogue (Thompson et al., 1975). These investigators proposed that this nucleotide analogue binds to those proteins which contain the super secondary structure termed the dinucleotide fold. The binding of free C subunit to Blue Dextran-Sephacrose and the failure of the holoenzyme ( $\text{R}_2\text{C}_2$ ) to bind support the notion that the active site is inaccessible in the holoenzyme form.

We present experiments which show that the free chromophore of Blue Dextran, which contains a triazine ring chloride, inhibits the free C subunit in an apparently irreversible manner and exhibits the characteristics of an active site directed reagent. Furthermore, RI, RII, and the protein kinase inhibitor protein (PKI) also protect against blue dye inactivation. These results support the idea that each of these proteins physically or functionally shields the active site of the

<sup>†</sup> From the Department of Biochemistry, The University of Iowa, Iowa City, Iowa 52242. Received November 29, 1978. This work was supported by Grant NS-11310 from the U.S. Public Health Service.

\* Address correspondence to this author at the Department of Biochemistry, Louisiana State University Medical Center, New Orleans, LA 70112.

<sup>1</sup> Abbreviations used: C, catalytic subunit; RI and RII, regulatory subunits I and II, respectively; cAMP, adenosine cyclic 3',5'-monophosphate; PKI, protein kinase inhibitor protein; Mops, 3-(N-morpholino)propanesulfonic acid.

C subunit and thereby inhibits it. Some of these experiments have been reported in preliminary form (Witt & Roskoski, 1978).

#### Methods and Materials

**Protein Kinase Preparation and Assay.** Bovine brain isoenzyme II was isolated by DE-52 chromatography as previously described (Witt & Roskoski, 1975b). The C subunit was prepared to apparent homogeneity as judged by sodium dodecyl sulfate-polyacrylamide gel electrophoresis using the procedure of Beavo et al. (1974). The catalytic subunit had a specific activity of 1.0–1.5  $\mu\text{mol}/(\text{min mg of protein})$ , using histone as substrate. Some of the variability encountered in specific activity was related to the use of different lots of histone IIa (Sigma Chemical Co.).

With histone as substrate, the assays were performed as previously described (Witt & Roskoski, 1975b). With the synthetic heptapeptide corresponding to the phosphorylatable sequence of pig liver pyruvate kinase (Leu-Arg-Arg-Ala-Ser-Leu-Gly from Peninsula Laboratories, Inc.), the assay mixture contained 200  $\mu\text{M}$  peptide substrate and the activity was measured as previously described (Witt & Roskoski, 1975b) except that 30% acetic acid (v/v) was substituted for water to elute  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  and metabolites from the phosphocellulose strips (Glass et al., 1978). Values for enzyme activity presented in the tables and figures represent means of triplicate assays from at least three different experiments. The standard errors of the mean are less than 6% of the values given.

**Catalytic Subunit Inactivation by Cibacron Blue F3GA.** Purified Cibacron Blue F3GA (a generous gift of Dr. H. Bosshard of Ciba Geigy, Basal, Switzerland) was examined by silica gel thin-layer chromatography and found to migrate as a single component with  $R_f$  values at 0.50, 0.62, and 0.58, respectively, using (a) methanol, (b) butanol-acetone-acetic acid-5%  $\text{NH}_4\text{OH}$ -water (7:5:3:3:2 v/v), and (c) tetrahydrofuran-water (48:7 v/v). Dye concentration was determined spectrophotometrically at 610 nm by using an extinction coefficient of  $13600 \text{ M}^{-1} \text{ cm}^{-1}$  (Thompson & Stellwagen, 1976). To catalytic subunit (6  $\mu\text{g}$  in 90  $\mu\text{L}$  of 220 mM potassium phosphate, pH 7.0) was added 10  $\mu\text{L}$  of dye solution to give the final specified concentration. Incubations were performed in 1.5-mL polyethylene (not glass) tubes at 22 °C. Portions (5  $\mu\text{L}$ ) were removed at 6-min intervals and assayed with the synthetic heptapeptide as specified in the previous section to follow the rate of dye inactivation. The high ionic strength and polyethylene tubes were chosen as conditions which stabilize dilute C subunit. This concentration of phosphate, moreover, does not inhibit catalytic activity nor alter the rate of blue dye inactivation compared with the conditions during assay.

The sources of materials and other methods were previously described (Witt & Roskoski, 1975a,b).

#### Results

**Specific Elution of the Catalytic Subunit from Blue Dextran-Sephadex.** Previous studies with the nucleotide analogue, the blue chromophore of Blue Dextran, showed that the bound C subunit is eluted with the substrate MgATP. Guanethidine, at 1 and 5 mM concentrations, also elutes 21 and 76% of the C subunit, respectively, from the nucleotide affinity resin (Table I). Guanethidine dissociates the protein kinase inhibitor from bovine heart C subunit (Demaille et al., 1977) and is proposed to act like the arginyl determinants of protein substrates (Kemp et al., 1976; Demaille et al., 1977). It is not, however, a nucleotide analogue. We propose, therefore,

Table I: Catalytic Subunit Elution from Blue Dextran-Sephadex<sup>a</sup>

eluant	C subunit act. (% applied)
ATP (1 mM)	10
ATP (5 mM)	74
guanethidine (1 mM)	21
guanethidine (5 mM)	76
ATP (1 mM), guanethidine (1 mM)	71
KCl (0.2 M)	97

<sup>a</sup> C subunit was prepared as described under Methods and Materials. It was applied to a Blue Dextran-Sephadex column ( $0.8 \times 1.5 \text{ cm}$ ), 250  $\mu\text{g}$  in 500  $\mu\text{L}$  of buffer containing 4 mM Mops, 0.5 mM dithiothreitol, 5 mM  $\text{MgCl}_2$ , and 5% glycerol (v/v) all at pH 6.7, and washed twice with 2 mL of the same buffer. Then buffer ( $2 \times 2 \text{ mL}$ ) containing the specified compound was used to elute the enzyme. Recovered enzyme was measured as previously described (Witt & Roskoski, 1975b).




FIGURE 1: Time course of Cibacron Blue F3GA inhibition of protein kinase. C subunit was prepared as described under Methods and Materials, and pig heart ( $\text{H}_4$ ) lactate dehydrogenase was purchased from Sigma Chemical Co. C subunit was treated with 100  $\mu\text{M}$  Cibacron Blue F3GA, and enzyme activity was measured in 5- $\mu\text{L}$  portions as described under Methods and Materials. To 90  $\mu\text{L}$  of lactate dehydrogenase ( $1.3 \times 10^{-8} \text{ M}$ ) in 10 mM Tris (10 mM, pH 7.5) containing mercaptoethanol (0.5 mM) was added 10  $\mu\text{L}$  of 1 mM blue dye. At the specified time, portions (5  $\mu\text{L}$ ) were mixed with 500  $\mu\text{L}$  of 0.5 mM sodium pyruvate, 200  $\mu\text{M}$  NADH, and 0.5 mM mercaptoethanol in 10 mM Tris (pH 7.5). Enzyme activity was measured by following the decline in absorbance at 340 nm in a Beckman 25 spectrophotometer.

that the blue chromophore interacts with the portion of the active site complementary to both the MgATP and protein substrate. Thompson & Stellwagen (1976) have previously proposed that the blue chromophore interacts with both the NADH and pyruvate domains of lactate dehydrogenase. Low concentrations of a combination of MgATP and guanethidine possess a greater than additive effect in elution (Table I), supporting the idea that the ligand interacts with both active-site domains. A similar synergistic action of substrates has been suggested in the elution of lactate dehydrogenase and phosphoglycerate kinase (Thompson & Stellwagen, 1976).

**Catalytic Subunit Inhibition by Cibacron Blue F3GA.** Since Cibacron Blue F3GA is a competitive inhibitor of lactate dehydrogenase with respect to its substrate NADH (Thompson & Stellwagen, 1976; Wilson, 1976), similar steady-state kinetic studies were attempted with the bovine brain catalytic subunit. Complicated kinetic patterns, however, resulted. Further experiments indicated that the complexity was due to a time-dependent decrease in enzyme activity which occurs when C subunit is incubated with the blue chromophore (Figure 1). Preincubation with the blue dye, on the other hand, fails to produce a similar time-dependent loss of lactate dehydrogenase (Figure 1) or phosphoglycerate kinase activity (Thompson & Stellwagen, 1976). The steady-state kinetic inhibition is overcome by increasing the NADH concentration. Catalytic





FIGURE 2: MgATP protects C subunit against inactivation by Cibacron Blue F3GA. The C subunit (6  $\mu$ g) was incubated with the indicated concentrations of blue dye without addition (●), with 310  $\mu$ M ATP alone (Δ), or with 310  $\mu$ M ATP and 5 mM  $\text{MgCl}_2$  (▲). The rate of inactivation was measured as described under Methods and Materials, and the values for  $K_{app}$  were graphed according to eq 1.

subunit activity, however, is not restored by addition of large concentrations of MgATP or protein substrate during the assay. These experiments demonstrate that there is a difference in the mechanism by which this chromophore interacts with the protein kinase C subunit when compared with other enzymes.

Specific elution of C subunit from Blue Dextran–Sephacryl affinity resins by MgATP and guanethidine is consistent with the hypothesis that the chromophore binds at the active site of the enzyme. The inactivation of the C subunit by the free chromophore may result, therefore, from complex formation and reaction of the blue dye at the active site. Active site directed enzyme inactivation has kinetic characteristics which are distinguishable from simple bimolecular reactions. Following the presentation of Kitz & Wilson (1962), the reaction between an enzyme with an active site directed agent (similar to that with substrate) can be formulated by



where E and I are free enzyme and inhibitor,  $E \cdot I$  is an enzyme–inhibitor Michaelis complex, EI is the inhibited enzyme, and  $k_3$  is the rate-limiting step. A steady-state treatment of the process yields the equation (Kitz & Wilson, 1962)

$$\frac{1}{K_{app}} = \frac{1}{k_3} + \frac{k_1}{k_3} \frac{1}{[I]} \quad (2)$$

where  $K_{app}$  is the observed rate of inactivation for a given concentration of inhibitor,  $k_3$  is the maximum rate of inactivation, and  $k_1$  is the dissociation constant of inhibitor for the enzyme ( $k_2/k_1$ ).

The inactivation of the catalytic subunit by Cibacron Blue F3GA at pH 7.0, 22 °C, follows pseudo-first-order kinetics.  $K_{app}$  values for varying inhibitor concentrations were measured and plotted in double-reciprocal form (eq 2). A straight line and positive ordinate intercept result (Figure 2), thereby indicating saturating kinetics (and not simple second-order kinetics) for the inactivation process. The maximum rate of inactivation ( $k_3$ ) is 0.13  $\text{min}^{-1}$ , and the  $k_1$  is 100  $\mu$ M, as determined from the ordinate and abscissa intercepts, respectively. These results are consistent with the contention that inactivation of C by Cibacron Blue F3GA occurs after the formation of a reversible  $E \cdot I$  complex.

Substrates and competitive inhibitors characteristically reduce the rate of enzyme inactivation by active site directed reagents. The rate equation is an extension of eq 2, which may be written as (Kitz & Wilson, 1962)

$$1/K_{app} = 1/k_3 + k_1/[k_3(1 + [A]/k_a)/[I]] \quad (3)$$

where A is a substrate or competitive inhibitor and  $k_a$  is its dissociation constant. For substantiation of the contention that Cibacron Blue F3GA is an active site directed reagent, C subunit was incubated with the blue dye in the presence of substrates and competitive enzyme inhibitors.  $K_{app}$  values were measured and graphed according to eq 2. ATP alone fails to alter the rate of dye inactivation (Figure 2). MgATP, however, reduces enzyme inactivation by the blue chromophore (Figure 2). The lines representing inactivation in the presence or absence of MgATP have a common ordinate intercept, thereby indicating that the maximum rate of inactivation ( $k_3$ ) is unchanged. Analogous to competitive inhibition patterns obtained from double-reciprocal plots (Cleland, 1967), this result indicates that the dye and MgATP compete for the same site. From eq 3, the  $k_a$  of MgATP is 620  $\mu$ M. This differs from the  $K_m$  of  $\sim 10$   $\mu$ M and provides another example that the dissociation constant is not necessarily equal to the  $K_m$  (Jencks, 1969).

Since MgATP and guanethidine elute the C subunit from Blue Dextran–Sephacryl synergistically (Table I), we propose that its blue chromophore occupies portions of both the nucleotide and protein substrate domains of the active site. Supporting this proposal, the synthetic heptapeptide substrate (Leu-Arg-Arg-Ala-Ser-Leu-Gly) and guanethidine (a protein substrate competitive inhibitor) also retard the rate of inactivation at a given concentration of blue dye. By use of the procedure outlined in Figure 2, the maximum rate of inactivation ( $k_3$ ), determined from the ordinate intercept, is unchanged. From eq 3, the  $k_a$  for the synthetic peptide and guanethidine is 0.3 and 1.6 mM, respectively. Benzamidine, a feeble competitive inhibitor of C subunit with respect to protein substrate, has no effect on the inactivation process with concentrations to 20 mM. Nucleotide and protein substrate protection at a given dye concentration support the notion that the blue dye is an active site directed reagent.

During the course of these studies, the rate of inactivation declined over a period of several days. Dye decomposition in aqueous solution failed to explain this observation since freshly prepared dye solutions gave  $K_{app}$  values similar to solutions prepared 2 months previously. On the other hand, if  $K_{app}$  values obtained from fresh and aged (3 weeks) enzyme are graphed according to eq 1, the corresponding lines share a common intercept but exhibit a different slope. These results show that the maximum rate of inactivation ( $k_3$ ) is unchanged; enzyme aging, however, is associated with a decreased affinity of the dye for the enzyme. The  $k_1$  for fresh and aged (3 weeks) enzyme was 100 and 210  $\mu$ M, respectively. Some older enzyme preparations (6 weeks), moreover, were resistant to dye inactivation. However, they retained their original specific activity.

Preincubation of aged (3 weeks) C subunit with  $\text{MgCl}_2$  (30 min, 4 °C), however, restores sensitivity to dye inactivation. The  $\text{MgCl}_2$  effect might be related to a direct effect on the enzyme protein or perhaps to the formation of a Mg–dye complex analogous to MgATP. A fivefold excess of EDTA was added to the  $\text{MgCl}_2$ -treated enzyme prior to addition of the dye to distinguish between these possibilities. The  $K_{app}$  for the  $\text{MgCl}_2$ -treated enzyme (minus EDTA) was equivalent to that following EDTA treatment; both, however, were greater than that for the  $\text{MgCl}_2$ -untreated enzyme (Table II). These

Table II:  $\text{MgCl}_2$  Treatment Enhancement of Catalytic Subunit Inactivation by Cibacron Blue F3GA<sup>a</sup>

pretreatment	EDTA (mM)	$K_{\text{app}}$ ( $\text{min}^{-1}$ )
none	0	27.4
none	10	29.3
$\text{MgCl}_2$ (10 mM)	0	15.8
$\text{MgCl}_2$ (10 mM)	10	18.4

<sup>a</sup> Aged C subunit (3 weeks, 30  $\mu\text{g}$  of protein in 200  $\mu\text{L}$ ) was incubated for 30 min at 4 °C in the presence or absence of  $\text{MgCl}_2$ . Portions (20  $\mu\text{L}$ ) were then incubated with 50  $\mu\text{M}$  blue dye in 200 mM potassium phosphate (pH 7.0) at 22 °C with EDTA as specified in a final volume of 100  $\mu\text{L}$ . Enzyme activity and  $K_{\text{app}}$  values were determined as described under Methods and Materials.

results are consistent with the notion that  $\text{MgCl}_2$  has a direct effect on the aged enzyme protein.

**General Characteristics of Cibacron Blue F3GA Inactivation of the Catalytic Subunit.** In addition to time dependence, blue dye inactivation is also temperature dependent. The rate of inactivation at 0 °C is ~3% that at 30 °C. The rate of inactivation was insensitive to pH (6.0–8.0). Since many dyes inhibit enzymes by photooxidation, experiments were performed under various degrees of illumination to test for blue dye mediated photoinactivation. In these experiments, C subunit was incubated for 7 min at 22 °C with 50  $\mu\text{M}$  Cibacron Blue F3GA. Incubations were performed in the dark, under moderate (room fluorescent lighting) or high (375-W bulb, 23-cm distance) intensity lighting. Since no differences in activation were observed, enzyme inactivation is unrelated to photoinactivation.

Several attempts were made to reverse the blue dye inactivation. For example, low concentrations of substrate nucleotides or high NaCl concentrations (200 mM) displace Cibacron Blue F3GA from lactate dehydrogenase and phosphoglycerate kinase as measured spectrophotometrically (Thompson & Stellwagen, 1976). In the present study, if 5 mM MgATP or guanethidine is added alone or in combination to an enzyme solution containing the blue dye, then enzyme inactivation is terminated. This treatment, however, fails to restore lost enzyme activity. Unlike lactate dehydrogenase or phosphoglycerate kinase, the addition of NaCl (to 500 mM) also fails to reactivate the enzyme. Enzyme inactivation is also not reversed by Sephadex G-25 gel filtration or by exhaustive dialysis. Since Thompson et al. (1976) demonstrated that Blue Dextran–Sephadex quantitatively displaces  $\text{NAD}^+$  ( $k_d = 10^{-6}$  M) from glyceraldehyde-3-phosphate dehydrogenase (which can be recovered by salt elution), a similar approach was attempted. If Cibacron Blue F3GA binds noncovalently to the active site of the C subunit, then the Blue Dextran–Sephadex resin may displace it and reverse the inhibition. After treatment of C subunit (6  $\mu\text{g}$ ) with Cibacron Blue F3GA, inhibited catalytic subunit was applied to a small Blue Dextran–Sephadex column and treated with low ionic strength and then high ionic strength buffer to elute specifically adsorbed enzyme, as described in Table I. No activity was detected in any of the elution fractions, indicating that the blue dye was not displaced by the affinity resin. Protein analysis (Bradford, 1976) also shows that the dye-inactivated enzyme fails to bind to the affinity resin.

Chemical attachment of the dye to the enzyme would most likely involve displacement of the chloride group of the triazine ring by a nucleophilic side-chain residue. Blue Dextran–Sephadex, in which sugar alcohols replace the dye chloride, interacts with C in a reversible manner (Table I; Witt & Roskoski, 1975a). For a further test of this hypothesis, the

Table III: Methoxy Cibacron Blue F3GA Reversible Inhibition of C Subunit Activity<sup>a</sup>

treatment	activity [ $\mu\text{mol}/(\text{min mg of protein})$ ]	
	initial assay	gel filtration
none	1.35	1.27
100 $\mu\text{M}$ Cibacron Blue F3GA	0.27	0.30
100 $\mu\text{M}$ methoxy blue dye	0.46	1.30

<sup>a</sup> Catalytic subunit (18  $\mu\text{g}/300 \mu\text{L}$ ) was treated with the specified compound as described under Methods and Materials. After 30 min, samples (5  $\mu\text{L}$ ) were assayed for enzyme activity in the presence of 100  $\mu\text{M}$  Cibacron Blue F3GA or its derivative. The remainder was applied to Sephadex G-50 columns (1  $\times$  15 cm) equilibrated with 0.1 M potassium phosphate (pH 6.8). The protein-containing fractions were combined, and enzyme activity was measured as previously described (Witt & Roskoski, 1975b). Protein was measured by the procedure of Bradford (1976). For replacement of the triazine chloride, the methoxy derivative of Cibacron Blue F3GA was prepared according to methodologies outlined by Dudley et al. (1951) for the synthesis of 2,4-bis-(phenylamino)-6-methoxytriazine. The reaction was initiated by the addition of Cibacron Blue F3GA (125 mg,  $1.5 \times 10^{-4}$  mol) to 50 mL of methanol-dissolved sodium hydroxide (7 mg,  $1.7 \times 10^{-4}$  mol) at 50 °C. After the mixture was refluxed for 2.5 h, the methanol was evaporated. The product was purified by dissolving it in 5 mL of water and then adding 1 g of NaCl. After the mixture was stirred for 2 h at room temperature, the resulting precipitate was collected by suction filtration and washed with 20% NaCl (10 mL). The air-dried precipitate was dissolved in 1 mL of water prior to use.

Table IV: Stoichiometry of Cibacron Blue F3GA Binding to Catalytic Subunit<sup>a</sup>

treatment	dye bound (mol/mol)	act. [ $\mu\text{mol}/(\text{min mg})$ ]
Cibacron Blue F3GA	$1.21 \pm 11$	$0.07 \pm 0.01$
Cibacron Blue F3GA, ATP (5 mM), $\text{MgCl}_2$ (10 mM)	$0.12 \pm 0.04$	$1.31 \pm 0.11$
no addition	0.00	$1.42 \pm 0.12$

<sup>a</sup> Catalytic subunit (150  $\mu\text{g}/250 \mu\text{L}$ ) was treated with 50  $\mu\text{M}$  blue dye for 30 min at 22 °C as specified under Methods and Materials. After Sephadex G-50 medium gel filtration (4  $\times$  1 cm) using 220 mM sodium phosphate (pH 7.0) as eluant, fractions (0.4 mL) were collected and taken for protein and enzyme activity determinations. The content of blue dye was determined spectrophotometrically at 610 nm by using an extinction coefficient of  $13\,600 \text{ M}^{-1} \text{ cm}^{-1}$  (Thompson & Stellwagen, 1976). These experiments represent the mean plus or minus the standard error of the mean of triplicate determinations.

methoxy derivative of the blue dye was synthesized. This compound is a reversible inhibitor of C subunit activity (Table III).

Because of the inability to reactivate the enzyme under nondenaturing conditions, attempts were made to dissociate the blue dye by denaturation. After incubation of 50  $\mu\text{M}$  blue dye for 30 min, the enzyme was subjected to sodium dodecyl sulfate–polyacrylamide gel electrophoresis. The blue dye and Coomassie blue staining protein comigrated as determined by inspection. Inclusion of 1 mM ATP (10 mM  $\text{MgCl}_2$ ) during the period of incubation, however, substantially decreased the amount of dye comigrating with protein.

For a determination of the stoichiometry of chromophore binding, the dye–enzyme complex was resolved from free dye by Sephadex gel filtration. Since the free dye binds tenaciously to the gel, this is an effective means to effect resolution. The amount of dye was measured spectrophotometrically, and the protein was measured by the procedure of Lowry et al. (1951). Under these conditions, the molar ratio of dye to C subunit

Table V: Effect of Cibacron Blue F3GA on Holoenzymes I and II<sup>a</sup>

	act. (pmol/min)	
	isozyme I	isozyme II
control	125	110
Cibacron Blue F3GA (50 $\mu$ M)	120	94
cAMP (10 $\mu$ M)	123	107
Cibacron Blue F3GA (50 $\mu$ M), cAMP (10 $\mu$ M)	88	69

<sup>a</sup> Bovine muscle protein kinases I and II were prepared by using the methodology of Corbin et al. (1975). In a final volume of 100  $\mu$ L (200 mM potassium phosphate, pH 7.0), fractions containing protein kinases I and II were incubated for 20 min at 22 °C with 50  $\mu$ M Cibacron Blue F3GA and cAMP, as specified. Enzyme activity (10  $\mu$ L) was measured as previously described (Witt & Roskoski, 1975b).

was  $1.21 \pm 0.11$  (Table IV). Inclusion of MgATP substantially decreased the amount of bound dye. Similar results were obtained when the incubation was terminated by the addition of 6 M guanidinium hydrochloride, followed by gel filtration (when the column is equilibrated with the salt, protein determinations cannot be performed, but blue dye is associated with protein in the void volume).

**Differential Effects of Cibacron Blue F3GA on C Subunits and Holoenzymes I and II.** In addition to isozyme II of bovine brain, experiments were performed with bovine skeletal muscle isozymes I and II and blue dye. In the absence of cAMP, holoenzyme I is insensitive to blue chromophore treatment and holoenzyme II is inhibited 15% (Table V). This inhibition corresponds to the amount of cAMP-independent activity of this preparation (not shown). After the addition of cAMP, however, the enzymes become susceptible to inactivation by the Cibacron Blue F3GA. For example, isozymes I and II are inhibited 30 and 37%, respectively (Table V). Like the brain C subunit, MgATP protects the enzymes from inactivation. The same results were obtained with reconstituted enzyme. For example, the addition of bovine skeletal muscle RI or RII to brain C converted the latter from a susceptible to a resistant enzyme form. The ability of RI to prevent dye inactivation suggests that it, as well as RII (Witt & Roskoski, 1975a), inhibits holoenzyme activity by binding at or shielding the active site.

Experiments were also performed to determine whether the dye-treated C subunit recombines with the R subunit. Hofmann et al. (1975) have shown that the addition of C subunit to (R-cAMP)<sub>2</sub> complex produces a cAMP-dependent holoenzyme with the release of cAMP. Unlike untreated C, the addition of dye-inactivated C subunit to bovine skeletal muscle RI or RII containing bound [<sup>3</sup>H]cAMP failed to release protein-bound [<sup>3</sup>H]cAMP (Table VI).

## Discussion

Cibacron Blue F3GA, the chromophore of Blue Dextran, inhibits the C subunit of bovine brain and skeletal muscle cAMP-dependent protein kinase in a time- and temperature-dependent fashion. The hyperbolic dependence of inhibition on dye concentration is characteristic of an active site directed process. The protection against inhibitor afforded by MgATP, a synthetic heptapeptide substrate, and guanethidine (a substrate analogue) argues that they and Cibacron Blue F3GA interact with the same enzyme domain, namely, the active site.

That Cibacron Blue F3GA interacts with the active site of the C subunit is expected since the dye is a structural analogue of ATP. The apparently irreversible nature of inhibition,

Table VI: Reassociation of Cibacron Blue F3GA Treated Catalytic Subunit with Regulatory Subunits<sup>a</sup>

catalytic subunit treatment	[ <sup>3</sup> H]cAMP released (pmol)	
	RI	RII
control	30	25
Cibacron Blue F3GA	1	1
Cibacron Blue F3GA (4 °C)	25	20
Cibacron Blue F3GA, ATP (5 mM), MgCl <sub>2</sub> (10 mM)	24	22

<sup>a</sup> RI and RII containing bound [<sup>3</sup>H]cAMP were isolated from rat heart and bovine skeletal muscle, respectively, by the methodology described by Sugden & Corbin (1976). Bovine brain C subunit was treated with 100  $\mu$ M Cibacron Blue F3GA under the specified conditions for 30 min at 30 °C as described under Methods and Materials. Portions (50  $\mu$ L) of treated enzyme were mixed with 250  $\mu$ L of either RI or RII in buffer (5 mM potassium phosphate, pH 7.0, 0.5 mM dithiothreitol, 1 mM EDTA, 1 mM ATP, and 5 mM MgCl<sub>2</sub>). After incubation for 2 h at 4 °C, the mixture was applied to a Sephadex G-25 column (0.9  $\times$  53 cm) and eluted with this buffer. One-mL fractions were collected. Portions (500  $\mu$ L) from those fractions containing protein were transferred to scintillation vials containing 10 mL of Aquasol (purchased from New England Nuclear), and radioactivity was measured by liquid scintillation spectrometry.

however, was unanticipated since its inhibition of other enzymes is reversible (Thompson & Stellwagen, 1976). Weber et al. (1979), on the other hand, reported that commercial sources of Cibacron Blue F3GA inactivate *Escherichia coli* isoleucine-tRNA synthetase and yeast phosphoglycerate kinase by covalent modification. They find, however, that covalent modification and inactivation are associated with contaminants in the blue dye which can be resolved by thin-layer chromatography. In contrast to the heterogeneity observed in commercial sources of the dye, which we confirmed, the dye obtained from Ciba Geigy was homogeneous. The same sample of dye, moreover, reversibly inhibits phosphoglycerate kinase (Thompson & Stellwagen, 1976); this is consistent with the result of Weber et al. (1979), who showed that the dye purified by silica gel chromatography does not irreversibly inhibit this enzyme; they find, however, that a contaminant in the commercial dye irreversibly inhibits the enzyme. The dye used in the present experiments lacks the contaminant based on thin-layer chromatographic analysis and on its failure to irreversibly inhibit phosphoglycerate kinase. Inactivation of the catalytic subunit, therefore, is associated with Cibacron Blue F3GA per se and not with a contaminant.

The  $k_i$  of blue dye for lactate dehydrogenase and C subunit is 0.49 and 100  $\mu$ M, respectively (Thompson & Stellwagen, 1976; this study). The  $K_m$  for NADH and ATP are 7 and 10  $\mu$ M, respectively (Thompson & Stellwagen, 1976; this study). Thompson & Stellwagen (1976) suggest that the blue dye's affinity for the dehydrogenase may be enhanced because it resembles the NAD-pyruvate adduct and may mimic the transition state. In the case of protein kinase, a portion of the dye may span a portion of the protein substrate domain of the active site but does not closely resemble the peptide substrate.

Several previous observations indicate that protein kinase undergoes changes during storage. For example, aged enzyme becomes less dependent on cAMP for expression of activity (Brostrom et al., 1970) and is less capable of releasing cAMP upon regeneration of holoenzyme (Sugden & Corbin, 1976). In the present study, MgCl<sub>2</sub> restores the sensitivity of aged enzyme to dye inactivation. This seems to involve a direct effect on the C subunit per se and not interaction of MgCl<sub>2</sub> with the blue dye. It may be that the C subunit, like *E. coli* glutamine synthetase (Kingdon et al., 1968; Denton &

Ginsburg, 1969), has different conformations which are altered by divalent cations.

The nucleotide and peptide substrates protect the C subunit from blue dye inactivation. The finding that protection occurs over a wide range of substrate and competitive inhibitor concentrations supports the idea that the blue dye interacts with the active site. It seems possible, but less likely, that ligands with such different affinities for the active site would protect against inactivation at a remote site. The ability of the regulatory subunit to protect against inactivation may be due to interaction with the active site of the catalytic subunit. Alternatively, functional shielding of the active site from the dye might involve a conformational change mediated by allosteric interaction of the regulatory with the catalytic subunit. Additional studies are required to test these and other possibilities for the mechanism of inhibition of phosphotransferase activity by the regulatory subunits.

#### References

- Beavo, J. A., Bechtel, P. J., & Krebs, E. G. (1974) *Methods Enzymol.* 38, 299.
- Bradford, M. M. (1976) *Anal. Biochem.* 72, 248.
- Brostrom, M. A., Reimann, E. M., Walsh, D. A., & Krebs, E. G. (1970) *Adv. Enzyme Regul.* 8, 191.
- Cleland, W. W. (1967) *Adv. Enzymol. Relat. Areas Mol. Biol.* 29, 1.
- Corbin, J. D., Keely, S. L., & Park, C. R. (1975) *J. Biol. Chem.* 250, 218.
- Demaille, J. G., Peters, K. A., & Fischer, E. H. (1977) *Biochemistry* 16, 3080.
- Denton, M. D., & Ginsburg, A. (1969) *Biochemistry* 8, 1714.
- Dudley, J. R., Thurston, J. T., Schaefer, F. C., Holm-Hansen, D., Hull, C. J., & Adams, P. (1951) *J. Am. Chem. Soc.* 73, 2986.
- Gill, G. N., & Garren, L. D. (1970) *Biochem. Biophys. Res. Commun.* 39, 335.
- Glass, D. B., Masaracchia, R. A., Feramisco, J. R., & Kemp, B. E. (1978) *Anal. Biochem.* 87, 566.
- Hofmann, F., Beavo, J. A., Bechtel, P. J., & Krebs, E. G. (1975) *J. Biol. Chem.* 250, 7795.
- Jencks, W. P. (1969) *Catalysis in Chemistry and Enzymology*, p 52, McGraw-Hill, New York.
- Kemp, B. E., Benjamini, E., & Krebs, E. G. (1976) *Proc. Natl. Acad. Sci. U.S.A.* 73, 1038.
- Kingdon, H. S., Hubbard, J. S., & Stadtman, E. R. (1968) *Biochemistry* 7, 2136-2142.
- Kitz, R., & Wilson, I. B. (1962) *J. Biol. Chem.* 237, 3245.
- Kumon, A., Yamamura, H., & Nishizuka, Y. (1970) *Biochem. Biophys. Res. Commun.* 41, 1290.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L., & Randall, R. J. (1951) *J. Biol. Chem.* 193, 265-275.
- Reimann, E. M., Brostrom, C. O., Corbin, J. D., King, C. A., & Krebs, E. G. (1971) *Biochem. Biophys. Res. Commun.* 42, 187.
- Rosen, O. M., & Erlichman, J. (1975) *J. Biol. Chem.* 250, 7788.
- Sugden, P. H., & Corbin, J. D. (1976) *Biochem. J.* 159, 423.
- Tao, M., Salas, M. L., & Lipmann, F. (1970) *Proc. Natl. Acad. Sci. U.S.A.* 67, 408.
- Thompson, S. T., & Stellwagen, E. (1976) *Proc. Natl. Acad. Sci. U.S.A.* 73, 361.
- Thompson, S. T., Cass, K. H., & Stellwagen, E. (1975) *Proc. Natl. Acad. Sci. U.S.A.* 72, 669.
- Thompson, S. T., Cass, R., & Stellwagen, E. (1976) *Anal. Biochem.* 72, 293.
- Weber, B. H., Willeford, K., Moe, J. G., & Piskiewicz, D. (1979) *Biochem. Biophys. Res. Commun.* 86, 252-258.
- Wilson, J. E. (1976) *Biochem. Biophys. Res. Commun.* 72, 560.
- Witt, J. J., & Roskoski, R., Jr. (1975a) *Biochemistry* 14, 4503.
- Witt, J. J., & Roskoski, R., Jr. (1975b) *Anal. Biochem.* 66, 253.
- Witt, J. J., & Roskoski, R., Jr. (1978) *Fed. Proc., Fed. Am. Soc. Exp. Biol.* 37, 1330.