

# Comparison of Rate Constants for (PO<sub>3</sub><sup>-</sup>) Transfer by the Mg(II), Cd(II), and Li(I) Forms of Phosphoglucomutase<sup>†</sup>

William J. Ray, Jr.,\* Carol Beth Post, and Joseph M. Puvathingal

Department of Biological Sciences, Purdue University, West Lafayette, Indiana 47907

Received April 29, 1988; Revised Manuscript Received August 16, 1988

**ABSTRACT:** Net rate constants that define the steady-state rate through a sequence of steps and the corresponding effective energy barriers for two (PO<sub>3</sub><sup>-</sup>)-transfer steps in the phosphoglucomutase reaction were compared as a function of metal ion, M, where M = Mg<sup>2+</sup> and Cd<sup>2+</sup>. These steps involve the reaction of either the 1-phosphate or the 6-phosphate of glucose 1,6-bisphosphate (Glc-P<sub>2</sub>) bound to the dephosphoenzyme (E<sub>D</sub>) to produce the phosphoenzyme (E<sub>P</sub>) and the free monophosphates, glucose 1-phosphate (Glc-1-P) or glucose 6-phosphate (Glc-6-P): E<sub>P</sub>·M + Glc-1-P ← E<sub>P</sub>·M·Glc-P<sub>2</sub> → E<sub>P</sub>·M·Glc-6-P<sub>6</sub>. Before this comparison was made, net rate constants for the Cd<sup>2+</sup> enzyme, obtained at high enzyme concentration via <sup>31</sup>P NMR saturation-transfer studies [Post, C. B., Ray, W. J., Jr., & Gorenstein, D. G. (1989) *Biochemistry* (preceding paper in this issue)], were appropriately scaled by using the observed constants to calculate both the expected isotope-transfer rate at equilibrium and the steady-state rate under initial velocity conditions and comparing the calculated values with those measured in dilute solution. For the Mg<sup>2+</sup> enzyme, narrow limits on possible values of the corresponding net rate constants were imposed on the basis of initial velocity rate constants for the forward and reverse directions plus values for the equilibrium distribution of central complexes, since direct measurement is not feasible. The effective energy barriers for both the Mg<sup>2+</sup> and Cd<sup>2+</sup> enzymes, calculated from the respective net rate constants, together with previously published values for the equilibrium distribution of complexes in both enzymic systems [Ray, W. J., Jr., & Long, J. W. (1976) *Biochemistry* 15, 4018-4025], show that the 100-fold decrease in the *k*<sub>cat</sub> for the Cd<sup>2+</sup> relative to the Mg<sup>2+</sup> enzyme is caused by two factors: the increased stability of the intermediate bisphosphate complex and the decreased ability to cope with the phosphate ester involving the 1-hydroxyl group of the glucose ring. In fact, it is unlikely that the efficiency of (PO<sub>3</sub><sup>-</sup>) transfer to the 6-hydroxyl group of bound Glc-1-P (thermodynamically favorable direction) is reduced by more than an order of magnitude in the Cd<sup>2+</sup> enzyme. By contrast, the efficiency of the Li<sup>+</sup> enzyme in the same (PO<sub>3</sub><sup>-</sup>)-transfer step is less than 4 × 10<sup>-8</sup> that of the Mg<sup>2+</sup> enzyme. The large differences in properties of Mg<sup>2+</sup> and Cd<sup>2+</sup> plus the small differences in the above rate of (PO<sub>3</sub><sup>-</sup>) transfer, coupled with the similarity in properties of Mg<sup>2+</sup> and Li<sup>+</sup> plus the enormous differences in the corresponding rate of (PO<sub>3</sub><sup>-</sup>) transfer, suggest that the primary role of the metal ion in the transfer process may involve an electrostatic effect and that the metal ion may be positioned close to an atom that becomes substantially more negative in the transition state. Whether that atom might be the bridging oxygen of the phosphate groups, as in a predominantly dissociative mechanism, or is a peripheral phosphate oxygen or oxygens, as in a predominantly associative mechanism, is considered.

The <sup>31</sup>P saturation-transfer studies described in the accompanying paper (Post et al., 1989) show that one of the (PO<sub>3</sub><sup>-</sup>)-transfer steps<sup>1</sup> facilitated by the Cd<sup>2+</sup> form of phosphoglucomutase, viz., the transfer of the enzymic (PO<sub>3</sub><sup>-</sup>) group to the 6-position of bound Glc-1-P,<sup>2</sup> is surprisingly efficient (*k*<sub>11</sub><sup>+</sup> ≥ 350 s<sup>-1</sup>), given the relative values of *k*<sub>cat</sub> for the Cd<sup>2+</sup> and Mg<sup>2+</sup> enzymes (1/100). The present studies were undertaken in an attempt to rationalize the relatively low value of *k*<sub>cat</sub> for the Cd<sup>2+</sup> enzyme, in view of its efficiency in the above transfer step, since understanding metal-specific differences in (PO<sub>3</sub><sup>-</sup>)-transfer efficiency may provide clues to the role of the metal ion in the bond-breaking/bond-making process (see accompanying paper).

## EXPERIMENTAL PROCEDURES

**Materials.** The preparation, storage, and demetalation of phosphoglucomutase are described in the accompanying paper (Post et al., 1989). Gelatin (Knox) was demetalated as described by Magneson et al. (1987). <sup>31</sup>P phosphoric acid in

dilute HCl (New England Nuclear) was evaporated to dryness, treated with dilute ammonia, and reevaporated before use. A 2 M stock solution of high-purity lithium chloride (Alfa) was passed through a column of Chelex (Bio-Rad), Li<sup>+</sup> form, before use. Sucrose phosphorylase was isolated from *Leuconostoc mesenteroides* (Doudoroff, 1955). DEAE-Spectra/Gel M was from Spectrum. Triethylamine was distilled under vacuum and either stored in the dark at -20 °C or converted to the hydrochloride in aqueous solution. Before use in a HCO<sub>3</sub><sup>-</sup> buffer system, sufficient amine to produce a 1 M solution was suspended in water at 0 °C and bubbled with CO<sub>2</sub> (vigorous stirring) until the pH of a 1:10 dilution fell to about 7.5. ACS scintillation cocktail was from Amersham.

<sup>1</sup> "(PO<sub>3</sub><sup>-</sup>) transfer" is used herein to designate the process R<sub>1</sub>OPO<sub>3</sub><sup>2-</sup> + R<sub>2</sub>OH → R<sub>1</sub>OH + R<sub>2</sub>OPO<sub>3</sub><sup>2-</sup> and refers to the identity of the group transferred, without implication about the mechanism of transfer.

<sup>2</sup> Abbreviations: Glc-1-P, α-D-glucose 1-phosphate; Glc-6-P, α-D-glucose 6-phosphate; Glc-P<sub>2</sub>, α-D-glucose 1,6-bisphosphate; E<sub>P</sub> and E<sub>D</sub>, the phospho and dephospho forms of rabbit muscle phosphoglucomutase; P<sub>i</sub>, inorganic phosphate; M, a bivalent metal ion activator (of phosphoglucomutase).

<sup>†</sup> This work was supported in part by a research grant from the U.S. Public Health Services (GM08963, to W.J.R.).

**Instrumentation.** The Varian XL200A NMR spectrometer described in the accompanying paper (Post et al., 1989) was used in the manner indicated.

**Procedures.** General procedures for handling dilute solutions of metal-free phosphoglucomutase have been described (Magneson et al., 1987). As in that reference, *all* components of solutions that contained the enzyme were specially treated; *all* containers and *all* equipment used for transferring the enzyme were plastic and also were specially treated. All pH values refer to 25 °C. Radioactive phosphate was quantified by scintillation counting of 0.5-mL aliquots in 10 mL of ACS cocktail.

Dilute solutions of [ $^{32}\text{P}$ ]Glc-6-P that also contained  $^{32}\text{P}$ -labeled inorganic phosphate were quantified after the latter was precipitated in the presence of 0.15 M  $\text{HClO}_4$ , 1.3 mM ammonium molybdate, and 10 mM  $\text{Et}_3\text{N}\cdot\text{HCl}$  [cf. Sugino and Mioshi (1964)] by making the solution 0.2 mM in  $\text{P}_i$ . After standing 5 min and centrifuging, a second addition of  $\text{P}_i$  was made (initial concentration 0.1 mM); the same centrifuge tube was used, and the soluble contents were rapidly mixed. After repeating the standing and centrifuging steps, most of the supernatant was transferred to a clean tube for a final centrifugation before aliquots were removed for counting. The total radioactivity initially present was assessed in samples that were treated identically but in the absence of  $\text{Et}_3\text{N}$ .

$^{31}\text{P}$  saturation-transfer studies were conducted in the manner described in the accompanying paper with solutions that initially contained 1.62 mM phosphoglucomutase,  $\text{Cd}^{2+}$  form, 1.62 mM Glc-1-P, 1.4–6.0 mM Glc- $\text{P}_2$ , and 20 mM Tris-HCl, pH 7.4. Resonances of the free bisphosphate (at 0.7 and –1.5 ppm) were irradiated, and loss of magnetization from both phosphates of the enzyme-bound Glc- $\text{P}_2$  (at –2.7 or +5.2 ppm) was monitored.

The specific activity of the  $\text{Cd}^{2+}$  enzyme at 24 °C under initial velocity conditions was determined by (a) treating the metal-free enzyme with 1 mM NTA/0.9 mM  $\text{Cd}^{2+}$  in the presence of gelatin, 1 mg/mL, and 20 mM Tris-HCl, pH 7.4, and (b) assaying aliquots of the treated enzyme in the presence of 4 mM Glc-1-P, 25  $\mu\text{M}$  Glc- $\text{P}_2$ , 10 mM EDTA, and 20 mM Tris-HCl, pH 7.4. Time aliquots were removed, quenched in 0.1 M acid, allowed to stand 5 min, and neutralized with excess Tris base containing sufficient EDTA to produce a final concentration of 10 mM. After centrifugation, aliquots were assayed for Glc-6-P by means of the dehydrogenase reaction (Michal, 1984). Rates observed in the time interval from 4 to 8 min were extrapolated to time zero by constructing plots of  $\ln \{([P]_2 - [P]_1)/(t_2 - t_1)\}$  versus  $t$ , where  $[P]_1$  and  $[P]_2$  are the concentrations of product at the reference time,  $t_1$  (4 min), and at a subsequent time  $t_2$ . [Such plots are linear with a slope of  $-k_d$ , where  $k_d$  is the rate constant for dissociation of  $\text{Cd}^{2+}$  from the enzyme (cf. Ray and Roscelli (1966)).]

Glucose [ $^{32}\text{P}$ ]-1-phosphate with a specific activity of about 200 mCi/mmol and essentially free of Glc-6-P and  $\text{P}_i$  was prepared via the sucrose phosphorylase reaction by using phosphorylase that previously had been exposed to 10 mM EDTA, pH 7.4, for 1 h at room temperature (to eliminate residual phosphoglucomutase activity). Initial concentrations in the reaction mixture were 0.08 M sucrose, 0.1 mM [ $^{32}\text{P}$ ]-phosphate, 0.2 M Bistris-HCl, pH 7, 10 mM EDTA, 12 mM cysteine-HCl adjusted to pH 7 just before use, and about 0.5 unit of sucrose phosphorylase: final volume, 0.1 mL. The reaction was terminated after about 0.85 of the initial [ $^{32}\text{P}$ ]-phosphate had been converted to organic phosphate (see assay procedure, below) by making the reaction mixture 0.15 M in  $\text{HClO}_4$ , 10 mM in  $\text{Et}_3\text{N}\cdot\text{HCl}$ , and 1.3 mM in ammonium

molybdate: total volume, 1.0 mL. [ $^{32}\text{P}$ ]Phosphate was precipitated as in the assay for [ $^{32}\text{P}$ ]Glc-6-P, above. The final supernatant was diluted to 20 mL with water, made basic with 1.2 mL of 1 M  $\text{NH}_4\text{OH}$ , and subjected to chromatography on a  $0.6 \times 1.5$  cm column of Spectra/Gel DEAE-resin,  $\text{HCO}_3^-$  form, using a gradient of  $\text{Et}_3\text{N}/\text{HCO}_3^-$  0–0.5 M in a total of 50 mL: flow rate, 1 mL/min. The  $\text{Et}_3\text{N}/\text{HCO}_3^-$  was removed by lyophilization; the [ $^{32}\text{P}$ ]Glc-1-P was dissolved in 1 mL of 1 mM Tris-HCl buffer, pH 7.4, lyophilized, and stored in the dry state at –20 °C until used.

Uniformly labeled [ $^{32}\text{P}$ ]Glc- $\text{P}_2$  was prepared by equilibrating [ $^{32}\text{P}$ ]Glc-1-P with a large excess of Glc- $\text{P}_2$  in the presence of phosphoglucomutase. The bisphosphate was isolated chromatographically, as above, except that the elution gradient was twice as steep.

The transfer of [ $^{32}\text{P}$ ]( $\text{PO}_3^-$ ) from labeled Glc-1-P to Glc-6-P at chemical equilibrium was initiated with enzyme that had been treated with the  $\text{Cd}^{2+}$ /NTA buffer solution described for initial velocity studies (see above). Assay concentrations were 3.8 mM Glc-6-P, 0.2 mM Glc-1-P, 0.025 mM Glc- $\text{P}_2$ , 10 mM EDTA, and 20 mM Tris-HCl, pH 7.4, at 24 °C. After 3.75 min, a trace of [ $^{32}\text{P}$ ]Glc-1-P was added. Time aliquots were removed at the 4-min point and every 45 s thereafter and were quenched by addition to a solution that contained reagents for precipitating  $\text{P}_i$  (see above). The production of acid-stable organic phosphate was determined via a modification of the procedure described for quantifying [ $^{32}\text{P}$ ]Glc-6-P (see above) in which the precipitating solution (0.15 M in  $\text{HClO}_4$ ) was heated at 100 °C for 10 min to hydrolyze all remaining [ $^{32}\text{P}$ ]Glc-1-P. After the solution was cooled to room temperature, the sequential precipitation procedure was conducted as described. Rates measured in the time interval of 4–8 min were extrapolated to time zero in the manner described above for initial velocity studies.

Metal-free,  $^{32}\text{P}$ -labeled phosphoglucomutase was prepared from an enzyme solution, 2 mL, at a concentration of about 5 mg/mL of enzyme, that had been extensively dialyzed against 10 mM Tris/MES buffer, pH 6.2, that contained  $10^{-4}$  M  $\text{Zn}^{2+}$ . After the addition of 0.1 equiv or less of [ $^{32}\text{P}$ ]Glc-1-P, the enzyme was adsorbed on a  $0.6 \times 1.5$  cm column of CM-Sephadex previously equilibrated with the same buffer (but in the absence of added  $\text{Zn}^{2+}$ ). The column was washed with 10 mL of buffer and the enzyme eluted (0.1 mL/min) with 8 mM Tris/6 mM (Tris) $_2\text{SO}_4$  adjusted to pH 6.7 with MES. The peak tubes were made 10 mM in EDTA and dialyzed for three 12-h intervals versus 10 mM EDTA/20 mM Tris-HCl, pH 7.4. Except for the dialysis steps (at 4 °C), all steps were conducted at room temperature.

The reaction of (freshly prepared) metal-free [ $^{32}\text{P}$ ]E $_p$  with a saturating concentration of Glc-1-P (20 mM) was conducted in the presence of 22 mM LiCl, 10 mM EDTA, and 1 mM Glc- $\text{P}_2$  at 24 °C and at an enzymic concentration of about 4 mg/mL of enzyme. Time aliquots (0.15 mL) were quenched by addition to 0.15 mL of a freshly prepared solution of 10% TCA containing 1  $\mu\text{mol}$  of Glc- $\text{P}_2$ , followed by brief sonication to disperse the precipitate. Water (2 mL) was added and the mixture again subjected to brief sonication. After centrifugation, the supernatant was removed, diluted with 8 mL of 3 mM Tris base (to produce a pH of about 4), and stored at 4 °C, if necessary, prior to analysis. The glucose monophosphate and bisphosphate fractions were separated by chromatography on a  $0.6 \times 4.5$  cm column of Spectra/Gel DEAE, formate form. After a washing with 2 mL of water, the monophosphates were eluted with 10 mL of 0.3 M formic acid adjusted to pH 3 with aqueous ammonia (1 mL/min) and

the bisphosphates with the same buffer at a concentration 5-fold higher. Fractions of 1 mL were collected. The phosphates were located by acid hydrolysis of 0.01-mL aliquots of these fractions and subsequent use of a modified Fiske SubbaRao procedure for inorganic phosphate [cf. Boltz (1972)]. The contents of phosphate-containing tubes were combined, lyophilized in scintillation vials, and subsequently counted.

The rate constant for the dissociation of Glc-P<sub>2</sub> from its complex with E<sub>P</sub>-Cd was measured in an equilibrium mixture that contained 4.8 mM Glc-6-P, 0.25 mM Glc-1-P, 0.09 mM Glc-P<sub>2</sub>, and 20 mM Tris-HCl, pH 7.4. Prior to the assay, the enzyme was treated as in initial velocity studies (see above) and was used at a final concentration of 3.2 μM; the assay also contained 0.5 mM NTA/0.45 mM Cd<sup>2+</sup>. In the case of Mg<sup>2+</sup>, the enzyme previously was treated with 1 mM EDTA/2 mM Mg<sup>2+</sup> and was used at a final concentration of 3.2 nM; the assay also contained 1 mM EDTA/2 mM Mg<sup>2+</sup>, and the concentrations of monophosphates and bisphosphate were twice and half the above values, respectively. After 20 min at 24 °C to achieve complete equilibrium, a trace of uniformly labeled [<sup>32</sup>P]Glc-P<sub>2</sub> was added, time aliquots were removed, and the acid-stable radioactivity was assessed as above.

The equilibrium distribution of enzyme/Li<sup>+</sup>/glucose phosphate complexes in a solution at 24 °C that initially contained 40 mg/mL of phosphoenzyme (0.66 mM) plus 20 mg/mL of dephosphoenzyme (0.33 mM), 0.25 mM Glc-1-P, 10 mM Tris-HCl, pH 7.4, 50 mM LiCl, 0.5 mM MgCl<sub>2</sub>, and 0.25 mM EDTA was estimated by quenching 0.2-mL aliquots in 2.5 mL of rapidly stirred 1 N HClO<sub>4</sub> according to the method of Ray and Long (1976a), but using the micro stirring apparatus described by Ray and Puvathingal (1986). After addition of EDTA and ammonium formate to concentrations of 10 and 20 mM, respectively, followed by pH adjustment to 4 with 10 N KOH, most of the KClO<sub>4</sub> was separated by precipitation and filtration via centrifugation. After lyophilizing the filtrate, the residue was suspended in 0.3 mL of cold 0.2 M Tris-HCl, pH 8.3, and additional KClO<sub>4</sub> separated as above. Aliquots (0.05 mL) of the filtrate were assayed sequentially for Glc-6-P, Glc-1-P, and Glc-P<sub>2</sub> by using a modification of the procedure described previously (Ray & Long, 1976b) in which the change in optical density at 340 nm instead of fluorescence change was used to quantify the NADP produced in the coupled assay.

## RESULTS

Most rate constants obtained for the processes described below are summarized in Table I.

**Initial Velocity and Isotope-Transfer Rate at Equilibrium for the Cd(II) Enzyme.** Because the Mg<sup>2+</sup> enzyme is approximately 100-fold more efficient than the Cd<sup>2+</sup> enzyme, a 0.1% contamination of the Cd<sup>2+</sup> enzyme by the Mg<sup>2+</sup> form produces a significant error in initial velocity studies. Even the use of a Cd<sup>2+</sup>/NTA buffer system to convert the demetallated enzyme to the Cd<sup>2+</sup> form did not entirely eliminate this problem, because NTA is a poor chelator for Mg<sup>2+</sup>. The Cd<sup>2+</sup>/NTA buffer also produced a reduced enzymic activity in the assay. Hence, both initial velocity and isotope exchange studies were conducted in the presence of excess EDTA (at saturating substrate concentrations). Under these conditions, the half-time for dissociation of Mg<sup>2+</sup> is about 25 s (Ray & Roscelli, 1966), while that for Cd<sup>2+</sup> is about 30 min (Ray, 1969). By ignoring the first 4 min of the reaction, during which time essentially all of the Mg<sup>2+</sup> dissociates from the enzyme, the activity of the Cd<sup>2+</sup> enzyme can be accurately assessed.

Table I: Measured and Calculated Rate Constants for Reactions of Phosphoglucomutase from Studies at High and Low Concentrations of Enzyme<sup>a</sup>

rate constants	enzyme concentration/form		
	low/Mg <sup>2+</sup> (s <sup>-1</sup> )	low/Cd <sup>2+</sup> (s <sup>-1</sup> )	high/Cd <sup>+</sup> (s <sup>-1</sup> )
$k_{cat}^f$	630 <sup>b</sup>	6.5 <sup>b</sup> {6.5–6.9} <sup>c</sup>	[10.0–10.6] <sup>d</sup>
$k_{cat}^r$	200 <sup>e</sup>	{7.0–7.3} <sup>c</sup>	[10.8–11.2] <sup>d</sup>
$k_{tr}^f$	84 <sup>f</sup>	1.6 <sup>b</sup> {1.6–1.7} <sup>c</sup>	[2.4], <sup>g</sup> [2.4–2.5] <sup>d</sup>
$k_{-1}^{app}$	[340–400] <sup>h</sup>	(7.8) <sup>i</sup>	12 <sup>j</sup>
$k_{-6}^{app}$	[960–1660] <sup>h</sup>	(7.1) <sup>i</sup>	11 <sup>j</sup>
$k_{t1}^+$	[≥ 1400(1 + p <sub>1</sub> )] <sup>k</sup>	{230(1 + p <sub>1</sub> )] <sup>k</sup>	
$k_{t6}^+$	[≥ 1500(1 + p <sub>6</sub> )] <sup>k</sup>	{100(1 + p <sub>6</sub> )] <sup>k</sup>	

<sup>a</sup> Listed parameters refer to steady-state processes:  $k_{cat}^f$  and  $k_{cat}^r$  describe initial velocities in the forward and reverse directions, respectively;  $k_{tr}^f$  describes transfer of [<sup>32</sup>P]phosphate from Glc-1-P to Glc-6-P at equilibrium;  $k_{-1}^{app}$  and  $k_{-6}^{app}$  describe processes defined in eq 1, 4, and 5. All constants were determined at pH 7.4 and a temperature close to 24 °C. Some constants were measured both at high enzyme concentration (100 mg/mL; 1.62 mM) and at low enzyme concentrations (< 10<sup>-4</sup> mg/mL). Values in parentheses were obtained by applying a scaling factor of 0.65 to values measured at high enzyme concentrations (see footnote 3 and Results), those in brackets were obtained indirectly from values measured at low enzyme concentrations, and those in braces were obtained indirectly from values measured at high enzyme concentration and scaled with a factor of 0.65. <sup>b</sup> Measured in this study. <sup>c</sup> Calculated as limits from values of  $k_{-1}^{app}$  and  $k_{-6}^{app}$  obtained via NMR saturation transfer at high enzyme (Post et al., 1989) plus the equilibrium distribution of central complexes (Ray & Long, 1976c) in the manner described in Appendix 1 and scaled by a factor of 0.65 (see Results). <sup>d</sup> Calculated as in b, but without scaling. <sup>e</sup> Calculated from the measured value of  $k_{cat}^f$  and the ratio of  $k_{cat}^f/k_{cat}^r$  reported by Ray and Roscelli (1966). <sup>f</sup> From Ray and Long (1976c) after correcting for an altered extinction coefficient (Ray et al., 1982). <sup>g</sup> Calculated from rates measured via NMR saturation transfer (Post et al., 1989) according to eq 2. <sup>h</sup> Calculated from values of  $k_{cat}^f$ ,  $k_{cat}^r$ , and the equilibrium distribution of central complexes (Ray & Long, 1976c) in the manner described in Appendix 2. <sup>i</sup> From the corresponding measured value at high enzyme (Post et al., 1988), after scaling by a factor of 0.65. <sup>j</sup> Values obtained via NMR saturation transfer (Post et al., 1989). <sup>k</sup> From eq 9–12, using either values or minimal estimates of  $k_{-1}^{app}$  or  $k_{-6}^{app}$  in rows 4 and 5 together with C<sub>1</sub> = 4.2 and C<sub>6</sub> = 1.6 or C<sub>1</sub> = 28 and C<sub>6</sub> = 14 for the Mg<sup>2+</sup> and Cd<sup>2+</sup> enzymes, respectively (Ray & Long, 1976); p<sub>1</sub> and p<sub>6</sub> are undetermined partition coefficients greater than zero for E<sub>P</sub>-M-Glc-1-P and E<sub>P</sub>-M-Glc-6-P, respectively, and differ for M = Mg<sup>2+</sup> or Cd<sup>2+</sup>.

Sufficiently low concentrations of enzyme were used so that during the time interval from 0 to 8 min an insufficient amount of Glc-1-P was converted to Glc-6-P to cause a decrease in catalytic rate. Hence, the downward curvature for product/time plots during the 4–8-min time interval was caused by the dissociation of Cd<sup>2+</sup>, which is “irreversible” because of the excess of EDTA present. Under these conditions the activity of the enzyme decreases in a first-order manner (Ray & Roscelli, 1966), which facilitates extrapolation of the activity of the Cd<sup>2+</sup> enzyme observed in the 4–8-min range to the activity that would have been observed at *t* = 0 in the absence of contamination by the Mg<sup>2+</sup> enzyme: extrapolation factor, 1.15 (see Experimental Procedures);  $k_{cat} = 6.5$  s<sup>-1</sup> (at 24 °C). Under similar conditions [EDTA affects the activity of the Mg<sup>2+</sup> enzyme only in its role as a metal ion scavenger (Ray & Roscelli, 1966)]  $k_{cat} = 630$  s<sup>-1</sup> for Mg<sup>2+</sup> enzyme. Hence, the ratio ( $k_{cat}$ )<sub>Cd</sub>/( $k_{cat}$ )<sub>Mg</sub> is 0.0103.

Isotope-exchange studies at equilibrium using [<sup>32</sup>P]Glc-1-P were conducted in the presence of excess EDTA in a manner similar to that described above for the initial velocity (see Experimental Procedures). Since the decrease with time in the label appearing in the Glc-6-P fraction (which was held to less than 40%) is produced not only by a decrease in the fraction of the enzyme in the Cd<sup>2+</sup> form but also by the approach to isotopic equilibrium, the slope of the plot used to extrapolate the exchange rate to zero time was somewhat





FIGURE 1: Energy level diagram showing the relative values of the Gibbs function for the enzymic complexes present in the phosphoglucomutase reaction; left and right, respectively, values for the  $\text{Cd}^{2+}$  and  $\text{Mg}^{2+}$  enzymes. The dashed lines show changes that would produce a monotonic decrease in Gibbs energy for the process  $\text{E}_P\text{M}\cdot\text{Glc-1-P} \rightarrow \text{E}_D\text{M}\cdot\text{Glc-P}_2 \rightarrow \text{E}_P\text{M}\cdot\text{Glc-6-P}$ . The vertical double-headed arrows show the difference between the Gibbs function for the  $\text{E}_D\text{M}\cdot\text{Glc-P}_2$  complex and that which would produce a monotonic decrease, viz., the  $\text{Cd}^{2+}$  hole and the  $\text{Mg}^{2+}$  hole, respectively (see Discussion). The heights of the effective energy barrier for  $\text{E}_D\text{M}\cdot\text{Glc-P}_2 \rightarrow \text{E}_P\text{M} + \text{Glc-1-P}$ ,  $(\text{EEB})_1$ , and for  $\text{E}_D\text{M}\cdot\text{Glc-P}_2 \rightarrow \text{E}_P\text{M} + \text{Glc-6-P}$ ,  $(\text{EEB})_6$ , are shown, based on the definition of EEB under Discussion, and values of  $k_{-1}^{\text{app}}$  and  $k_{-6}^{\text{app}}$  in columns 2 and 3 of Table I (see also Results).

larger than that stated previously: extrapolation factor, 1.21;  $k_{\text{tr}}^f = 1.6 \text{ s}^{-1}$ .

**Comparison of  $^{32}\text{P}$  Isotope-Transfer and  $^{31}\text{P}$  Saturation-Transfer Rates for the  $\text{Cd}(\text{II})$  Enzyme.** The accompanying paper (Post et al., 1989) provides values for the rate constants identified in eq 1 under conditions where the concentration



of enzyme approaches that of the substrate/product pair, viz., under conditions where the approach of Glc-1-P to magnetic equilibrium when Glc-6-P is irradiated is not a steady-state process. In fact, a steady-state approach of Glc-1-P to magnetic equilibrium (very small ratio of enzyme to substrate/product pair) could not be obtained under conditions where the time dependence of Glc-1-P magnetization could be measured, because of the overall inefficiency of the  $\text{Cd}^{2+}$  enzyme coupled with the  $T_1$  values for the various phosphates in the system (see accompanying paper). But the value of  $k_{\text{tr}}^f$ , the constant that specifies the steady-state flux of phosphorus nuclei in the forward direction, can be calculated from the results of saturation-transfer studies:

$$k_{\text{tr}}^f = \frac{k_{-6}^{\text{app}}}{2} \frac{1}{1 + k_6^{\text{app}}K_e/k_1^{\text{app}}} \frac{[\text{E}_D\cdot\text{Cd}\cdot\text{Glc-P}_2]}{[\text{E}_{\text{total}}]} = 2.4 \text{ s}^{-1} \quad (2)$$

[The value of  $k_{-6}^{\text{app}}$  in eq 2 is  $11 \text{ s}^{-1}$ , and the factor  $1/2$  is used because transfer requires two catalytic cycles [see accompanying paper, Post et al. (1989), eq 7 and 8]; the second factor, whose value is 0.52, describes the partition of  $(\text{E}_P\cdot\text{Cd})$  produced by the first catalytic cycle between reaction with Glc-1-P and with Glc-6-P at chemical equilibrium (where  $K_e$  is  $[\alpha\text{-Glc-6-P}]/[\text{Glc-1-P}]$ ), in both cases to give  $\text{E}_D\cdot\text{Cd}\cdot\text{Glc-P}_2$ . The final factor is the fraction of enzyme present as  $\text{E}_D\cdot\text{Cd}\cdot\text{Glc-P}_2$ , 0.85 (see accompanying paper).]

The value measured for  $k_{\text{tr}}^f$  via  $^{32}\text{P}(\text{PO}_3^-)$  transfer at chemical equilibrium and at a low enzyme/substrate ratio is  $1.6 \text{ s}^{-1}$  under conditions otherwise similar to those used for NMR saturation-transfer studies, except for enzyme concentration (0.01 instead of 100 mg/mL) and substrate concentration (4 instead of 32 mM). The  $^{32}\text{P}(\text{PO}_3^-)$ -transfer rate



thus is about 0.67 of that expected from the NMR studies, i.e.,  $1.6 \text{ s}^{-1}/2.4 \text{ s}^{-1}$ .

**Comparison of  $^{31}\text{P}$  Saturation-Transfer Rates with the Steady-State Catalytic Constant:  $\text{Cd}(\text{II})$  Enzyme.** The nature of the  $^{31}\text{P}$  saturation-transfer process, combined with the observed rate constants for the half-reactions involved in that transfer [accompanying paper, Post et al. (1989)], and the equilibrium concentration of the monophosphate and bisphosphate complexes of  $\text{Cd}^{2+}$  phosphoglucomutase (Ray & Long, 1976) yield a simple but approximate relationship between the rate constants for saturation exchange and the steady-state catalytic constant,  $k_{\text{cat}}$  (see Appendix 1). In the forward reaction

$$10 \text{ s}^{-1} \leq k_{\text{cat}}^f \leq 10.6 \text{ s}^{-1} \quad (3)$$

The measured value of  $k_{\text{cat}}$  for the  $\text{Cd}^{2+}$  enzyme, at  $5 \times 10^{-4}$  instead of 100 mg/mL and at 4 instead of 32 mM glucose phosphates but under conditions otherwise analogous to those used for saturation transfer (see above), is  $6.5 \text{ s}^{-1}$ . Thus, in accord with the results in the previous section, the rate constant obtained from studies at low enzyme concentration is about 0.63 as large as that obtained from measurements at much higher enzyme concentrations, i.e.,  $6.5 \text{ s}^{-1}/10.6 \text{ s}^{-1}$ .

In view of the comparisons at high and low enzyme concentration in this and the previous section,<sup>3</sup> the earlier estimates of 350 and  $150 \text{ s}^{-1}$  for  $k_{11}^+$  and  $k_{16}^+$  for the following processes (see accompanying paper) are reduced to 230 and  $100 \text{ s}^{-1}$ , respectively, for comparison with steady-state rate constants obtained at low enzyme concentration (see below).



<sup>3</sup> The somewhat larger values of the two rate constants calculated from data obtained at high enzyme concentration, relative to studies at low enzyme concentration, might arise from a difference in activity coefficients [cf. Nichol et al. (1983)] from a failure to adequately account for the relative concentrations of the various enzyme/glucose phosphate complexes (Ray & Long, 1976c), or from unrecognized differences in conditions.

*Comparison of Net Rate Constants for (PO<sub>3</sub><sup>-</sup>) Transfer by the Mg(II) and Cd(II) Enzymes.* Figure 1 shows the relative Gibbs energies for the intermediate complexes of the Cd<sup>2+</sup> and Mg<sup>2+</sup> enzymes, left and right, respectively. The Gibbs function for the E<sub>P</sub>·M·Glc-1-P complex is used as a reference since the equilibrium binding of Glc-1-P to E<sub>P</sub>·M is essentially independent of the identity of M<sup>2+</sup>, not only for Mg<sup>2+</sup> and Cd<sup>2+</sup> but also for Zn<sup>2+</sup>, Co<sup>2+</sup>, and Mn<sup>2+</sup> (Ray & Long, 1976c). Also shown in Figure 1 are the effective energy barriers in the two half-reactions of the Cd<sup>2+</sup> enzyme, EEB<sub>1</sub> and EEB<sub>6</sub>, obtained from the values of the net rate constants<sup>4</sup>  $k_{-1}^{app}$  and  $k_{-6}^{app}$  (7.8 and 7.2 s<sup>-1</sup>, respectively) that were measured via <sup>31</sup>P saturation transfer in the accompanying paper and scaled as in the preceding section (Post et al., 1989; see eq 1). Here, EEB is equal to  $-0.6 \ln (k^{app}/10^{13})$  (see Discussion).

Analogous saturation-transfer studies cannot be conducted with the Mg<sup>2+</sup> enzyme because of chemical exchange broadening. But values of  $k_{-1}^{app}$  and  $k_{-6}^{app}$  (see eq 1) for this form of the enzyme can be approximated from the equilibrium ratios of the various intermediates (Ray & Long, 1976c) plus values of the steady-state catalytic constant in the forward and reverse reactions [630 and 200 s<sup>-1</sup>; data of Ray and Roscelli (1964), after making a correction for both a difference in extinction coefficient of the enzyme (Ray et al., 1983) and a temperature difference (25 instead of 30 °C)]. The required operation (see Appendix 2) is more or less the inverse of that used for approximating the range of values for  $k_{cat}^f$  from the half-reactions of the Cd<sup>2+</sup> enzyme (Appendix 1). This analysis shows that for the Mg<sup>2+</sup> enzyme

$$340 \leq k_{-1}^{app} \leq 400 \quad (6)$$

and

$$960 \leq k_{-6}^{app} \leq 1660 \quad (7)$$

Thus, the data restrict the range of allowed values for  $k_{-1}^{app}$  more than for  $k_{-6}^{app}$  (see eq 3). Both EEB<sub>1</sub> and EEB<sub>6</sub> for the Mg<sup>2+</sup> enzyme, calculated as above from  $k_{-1}^{app}$  and  $k_{-6}^{app}$ , also are shown in Figure 1 (right), where stippling is used to indicate the range of values consistent with the above approximations. As opposed to the Cd<sup>2+</sup> enzyme, where  $k_{-1}^{app} \approx k_{-6}^{app}$  (see above), the Mg<sup>2+</sup> enzyme facilitates (PO<sub>3</sub><sup>-</sup>) transfer from the 1-position of the bound bisphosphate to the dephosphoenzyme more efficiently than from the 6-position, as is indicated by the relative heights of the effective energy barriers in Figure 1. This difference will be further considered under Discussion.

*Comparison of Rate Constants for Phosphate Transfer by the Mg(II) and Cd(II) Enzymes.* The rate constant for the (PO<sub>3</sub><sup>-</sup>)-transfer step involving the primary hydroxyl group (the 6-hydroxyl) of bound glucose phosphate when the transfer is in the thermodynamically favorable direction, E<sub>P</sub>·M·Glc-1-P  $\xrightarrow{k_{-1}^{app}}$  E<sub>P</sub>·M·Glc-P<sub>2</sub>, can be related to the net rate constant,  $k_{-1}^{app}$  (see eq 1), in the manner described in Appendix 3. A quantitative comparison of  $k_{-1}^{app}$  for the Mg<sup>2+</sup> and Cd<sup>2+</sup> enzymes would require a knowledge of how E<sub>P</sub>·M·Glc-6-P is partitioned between dissociation to E<sub>P</sub>·M + Glc-6-P and return to E<sub>P</sub>·M·Glc-P<sub>2</sub>. Nevertheless, the analysis in Appendix 3 suggests


that the (PO<sub>3</sub><sup>-</sup>) transfer described by  $k_{-1}^{app}$  may well differ for the Mg<sup>2+</sup> and Cd<sup>2+</sup> enzymes by no more than an order of magnitude and perhaps less.

*Rate Constant for Dissociation of Glucose 1,6-Bisphosphate from Its Complex with the Cd(II) and Mg(II) Enzymes.* The strategy used to determine the rate constant for dissociation of Glc-P<sub>2</sub> from E<sub>P</sub>·Cd·Glc-P<sub>2</sub> is as follows. An equilibrium mixture of enzyme, glucose monophosphates, and Glc-P<sub>2</sub> was established with the monophosphate in large excess over Glc-P<sub>2</sub>, but with sufficient Glc-P<sub>2</sub> present to saturate the enzyme and sufficient enzyme to rapidly equilibrate the monophosphates. After the addition of a trace of uniformly labeled [<sup>32</sup>P]Glc-P<sub>2</sub> (50% acid-stable phosphate), the rate of approach to equilibrium (95% acid-stable phosphate) was measured. Under these conditions, the rate at which Glc-P<sub>2</sub> dissociates from the enzyme limits the introduction of additional label into the acid-stable (Glc-6-P) fraction. Plots of  $\ln (P_t - P_0)$  versus  $t$ , where  $P$  refers to acid-stable phosphate and the subscripts refer to reaction times, were linear (except, in the case of the Cd<sup>2+</sup> enzyme, for a small initial lag). Values for  $k_d$  were obtained from the slopes of such plots after multiplying by the reciprocal fraction (1/ $f$ ) of the enzyme present as E<sub>P</sub>·M·Glc-P<sub>2</sub>: for the Cd<sup>2+</sup> enzyme,  $k_d = 0.11 \text{ s}^{-1}$  [ $1/f = 1.18$ ; see the accompanying paper, Post et al. (1988)]; for the Mg<sup>2+</sup> enzyme,  $k_d = 10.7 \text{ s}^{-1}$  ( $1/f = 1.85$ ; Ray & Long, 1976c).

In the case of the Cd<sup>2+</sup> enzyme the value of  $k_d$  also was obtained via NMR magnetization transfer as described in the accompanying paper, but here conducted in the presence of an excess of Glc-P<sub>2</sub> and in the essential absence of free glucose monophosphates (see Experimental Procedures). Phosphorus resonances of free Glc-P<sub>2</sub> were independently irradiated for periods of 10 or 15 s, and the loss of magnetization from the 1- and 6-phosphates of bound Glc-P<sub>2</sub> was assessed by comparison with spectra obtained with an off-resonance irradiation time [i.e., a recycle time equal to 2.3–3.3  $T_1$ ; see the accompanying paper, Post et al. (1989)]. Because dissociation of the bound Glc-P<sub>2</sub> is slow relative to the catalytic rate (and thus slow relative to saturation transfer between the phosphates of bound Glc-P<sub>2</sub>; see accompanying paper), the decrease in magnetization of both phosphates of bound bisphosphate was approximately equal, regardless of which phosphate of free Glc-P<sub>2</sub> was irradiated. The magnetization observed after long irradiation times, relative to that measured in the absence of irradiation, decreased to a plateau value of about 0.75. The rate constant for dissociation of the bisphosphate complex was approximated from the fractional steady-state magnetization and the known  $T_1$  value (see Appendix 2 of the accompanying paper), although the small measured differences produced a substantial uncertainty in the results and calculated rate constants: average rate constant for dissociation of E<sub>P</sub>·Cd·Glc-P<sub>2</sub> ( $\pm$ SE) from eight observations in four different experiments,  $0.12 \pm 0.05 \text{ s}^{-1}$ .<sup>5</sup>

*Rate Constant for Transfer of the Enzymic (PO<sub>3</sub><sup>-</sup>) Group of the Li(I) Enzyme to Bound Glucose 1-Phosphate.* The strategy for evaluating the rate constant  $k_{-1}^{app}$  for transfer of the enzymic (PO<sub>3</sub><sup>-</sup>) group of the Li<sup>+</sup> enzyme to bound Glc-1-P (cf. eq 4) to produce Glc-P<sub>2</sub> is as follows. <sup>32</sup>P-Labeled phosphoenzyme was treated with a saturating concentration

<sup>4</sup> In a simple system, such as



a net rate constant describes the rate at which A is converted to C when the process is in the steady state. Here, the net rate constant for  $A \rightarrow C$  is equal to  $k_1 k_2 / (k_{-1} + k_2)$ ; see Cleland (1975).

<sup>5</sup> Because saturation transfer between the phosphate groups of bound Glc-P<sub>2</sub> is much more rapid than dissociation, the apparent rate constant for dissociation of Glc-P<sub>2</sub> assessed by irradiating only one of the phosphates of free Glc-P<sub>2</sub> is half that expected if both phosphates of free Glc-P<sub>2</sub> had been irradiated. It is the latter rate constant, viz., the constant that correlates with the molecular dissociation of Glc-P<sub>2</sub>, as assessed above in the [<sup>32</sup>P]phosphate exchange studies, that is reported here.

of Glc-1-P in the presence of saturating  $\text{Li}^+$  and excess EDTA. A sufficient amount of Glc-P<sub>2</sub> was included in the reaction so that any [ $^{32}\text{P}$ ]Glc-P<sub>2</sub> formed by the above ( $\text{PO}_3^-$ ) transfer would have a very low probability of reacting further, viz., reacting with the dephosphoenzyme in processes analogous to the reverse of eq 4 and 5. Since the efficiency of ( $\text{PO}_3^-$ ) transfer by the  $\text{Li}^+$  enzyme is many orders of magnitude less than for transfer by the  $\text{Mg}^{2+}$  enzyme (see below), even minute traces of  $\text{Mg}^{2+}$  could render an assessment of ( $\text{PO}_3^-$ ) transfer by the  $\text{Li}^+$  enzyme impossible. But once  $\text{Mg}^{2+}$  is bound at the active site of phosphoglucumutase, it remains associated with the enzyme, on the average, for about 20000 catalytic cycles at saturating Glc-1-P (since at 24 °C  $k_{\text{cat}} = 630 \text{ s}^{-1}$  and  $k_d = 0.032 \text{ s}^{-1}$ ; Ray & Roscelli, 1966). Only the first of these cycles can involve [ $^{32}\text{P}$ ]E<sub>P</sub>, and the probability that the label transferred in this cycle will enter the bisphosphate pool approaches 0.02 [Viz., is equal to  $k_d/k_{-6}^{\text{app}}$  or  $10.6 \text{ s}^{-1}$  (see preceding section) divided by 630 (see eq 7)]. Hence, by measuring the rate at which the labeled enzymic phosphate group is transferred to the bisphosphate pool, one can measure the exceedingly slow ( $\text{PO}_3^-$ ) transfer that characterizes the  $\text{Li}^+$  enzyme. To accomplish this, the glucose monophosphates in quenched time aliquots were chromatographically separated from the bisphosphate. (The labeled Glc-6-P also was quantified after hydrolysis of the acid-labile phosphate in the glucose monophosphate sample and precipitation of the released P<sub>i</sub>; as expected, the radioactivity in the Glc-6-P fraction was less than 1% that in the Glc-P<sub>2</sub> fraction.) A plot (not shown) of  $\ln f$  versus time from aliquots removed during the 72-h reaction was used to estimate the initial transfer rate, where  $f$  is the fraction of the labeled phosphate from [ $^{32}\text{P}$ ]E<sub>P</sub> that had entered the bisphosphate pool. (The recoverable catalytic activity after conversion of the enzyme from its  $\text{Li}^+$  to its  $\text{Mg}^{2+}$  form decreased only by about 6% during the 72-h reaction.) From the rate of label transfer plus the concentration of enzyme used, the observed rate constant is about  $4 \times 10^{-6} \text{ s}^{-1}$  at 24 °C. Since such a slow rate is not likely limited by the rate of Glc-P<sub>2</sub> dissociation from the  $\text{Li}^+$  enzyme (the corresponding rate constant for dissociation of Glc-P<sub>2</sub> from the  $\text{Mg}^{2+}$  enzyme is about  $10 \text{ s}^{-1}$ ; see preceding section), the above value almost certainly defines  $k_{\text{cl}}^+$  (eq 4) for the  $\text{Li}^+$  enzyme. This value is compared with estimates of  $k_{\text{cl}}^+$  for the  $\text{Mg}^{2+}$  and  $\text{Cd}^{2+}$  enzymes under Discussion. (In a parallel experiment under otherwise identical conditions, except for the radioactive label, the rate of Glc-6-P production, presumably by contaminating  $\text{Mg}^{2+}$  enzyme, was estimated as equivalent to that which would be produced if  $7.5 \times 10^{-5}$  of the enzyme were in the  $\text{Mg}^{2+}$  form throughout the experiment; this fractional abundance of the  $\text{Mg}^{2+}$  enzyme in turn would be produced by about  $1.7 \times 10^{-6} \text{ M}$  free  $\text{Mg}^{2+}$  in the presence of 22 mM  $\text{Li}^+$ .)

**Rate Constant for Transfer of the 6-Phosphate of Bound Glucose Bisphosphate to the Li(I) Form of the Dephosphoenzyme.** Technical problems prevented the direct measurement of  $k_{\text{cl}}^-$  (eq 4) for the  $\text{Li}^+$  enzyme. The value of this constant, about  $5 \times 10^{-7} \text{ s}^{-1}$ , was obtained from those for  $k_{\text{cl}}^+$  and the equilibrium constant ( $\text{E}_D\text{-Li-Glc-P}_2$ )/( $\text{E}_P\text{-Li-Glc-1-P}$ ) (9). Although the  $\text{Li}^+$  enzyme is not sufficiently active to establish such an equilibrium in a reasonable time interval, the required equilibrium can be established by adding a trace of  $\text{Mg}^{2+}$  to the system, as was done previously (Ray & Long, 1976c) to establish the analogous equilibrium in the absence of (all but a trace of) a metal ion. (Although no more than 0.5% of the enzyme was present as its  $\text{Mg}^{2+}$  complex, this enzyme facilitated the formation of a mixture of complexes

of the  $\text{Li}^+$  enzyme involving Glc-1-P, Glc-P<sub>2</sub>, and Glc-6-P; the composition of the mixture did not vary with time from 10 min to 2.3 h after addition of Glc-1-P.) The observed equilibrium constant, 9, differs only marginally from the value observed in the presence of  $\text{Mg}^{2+}$ , 4.2, or that for the metal-free system, 11, relative to the enormous rate differences observed here (see Discussion).

## DISCUSSION

The primary objective of this study and that in the accompanying paper (Post et al., 1989) is to define the reaction sequence for  $\text{Cd}^{2+}$ -activated phosphoglucumutase with sufficient accuracy to determine the origin of the 100-fold difference in  $k_{\text{cat}}$  between this form of the enzyme and the  $\text{Mg}^{2+}$ -activated form and, if possible, to furnish support for using the  $\text{Cd}^{2+}$  enzyme to model some aspects of the enzymic reaction involving  $\text{Mg}^{2+}$ . A more interesting though less readily attainable objective is to specify the role of the bivalent metal ion activator in the ( $\text{PO}_3^-$ )-transfer process.

With regard to the first objective, the accompanying paper provides minimal estimates of rate constants for ( $\text{PO}_3^-$ ) transfer by the  $\text{Cd}^{2+}$ -activated enzyme by direct measurement, using  $^{31}\text{P}$  saturation-transfer techniques. The comparison of ( $\text{PO}_3^-$ ) transfer in the  $\text{Cd}^{2+}$ - and  $\text{Mg}^{2+}$ -activated systems, below, is based partly on these constants, partly on the relative thermodynamic stability of the various intermediate complexes of the enzyme (Ray & Long, 1976c) plus the value for  $k_{\text{cat}}^f/k_{\text{cat}}^i$  for the  $\text{Mg}^{2+}$  enzyme (Ray & Roscelli, 1964), and partly on data under Results. (A comparison of rate constants obtained in NMR studies at 100 mg/mL with constants obtained at enzyme concentrations approximately  $10^{-6}$  as large also is made under Results; cf. Table I.)

At chemical equilibrium in the  $\text{Mg}^{2+}$ -activated system, the relative abundance of the three intermediate complexes,  $\text{E}_P\text{-Mg-Glc-1-P}$ ,  $\text{E}_D\text{-Mg-Glc-P}_2$ , and  $\text{E}_P\text{-Mg-Glc-6-P}$ , is 13:54:33 (Ray & Long, 1976c); i.e., the central  $\text{E}_D\text{-Mg-Glc-P}_2$  complex predominates, but not by much, as one might expect for an enzyme that has evolved close to "perfection" (Albery & Knowles, 1976). However, in the  $\text{Cd}^{2+}$  system the relative abundance of the corresponding complexes is about 3:91:6. As is illustrated by Gibbs energy profiles of the reaction intermediates for both enzyme forms (Figure 1), what is a relatively small depression, amounting to about  $-0.6 \text{ kcal/mol}$  in a descending monotonic energy profile for the  $\text{Mg}^{2+}$  enzyme, becomes a more substantial "hole", amounting to about  $-1.8 \text{ kcal/mol}$  in the  $\text{Cd}^{2+}$  enzyme system; see the vertical double-headed arrows in the figure. One might well view this hole as the  $\text{Cd}^{2+}$ -induced thermodynamic hole. Part of the reason why the  $\text{Cd}^{2+}$  enzyme is less active than the  $\text{Mg}^{2+}$  enzyme under steady-state conditions is that during a catalytic cycle the enzymic system will fall into and must extricate itself from this hole, regardless of the direction in which the reaction is conducted.<sup>6</sup>

A second problem with the  $\text{Cd}^{2+}$  enzyme is its reduced efficiency in the half-reaction involving Glc-6-P. This reduced efficiency can be described in terms of the height of the effective energy barrier (EEB) for  $\text{E}_D\text{-Cd-Glc-P}_2 \xrightarrow{k_{-6}^{\text{app}}} \text{E}_P\text{-Cd} + \text{Glc-6-P}$ , where  $k_{-6}^{\text{app}}$  is the net rate constant<sup>4</sup> (Cleland, 1975) and  $\text{EEB}_6$  is equal to  $-0.6 \ln (k_{-6}^{\text{app}}/10^{13}) \text{ kcal/mol}$ , i.e.,

<sup>6</sup> Earlier observations indicate that the group transfer potential for the enzymic phosphate is too high for maximal catalytic efficiency when  $\text{Cd}^{2+}$  is the activator. Thus,  $\Delta G^\circ$  for the process  $\text{E}_P\text{-M} + \text{Glc-1-P} \rightleftharpoons \text{E}_D\text{-M} + \text{Glc-P}_2$  becomes more negative by the substitution of  $\text{Cd}^{2+}$  for  $\text{Mg}^{2+}$  to about the same extent as  $\Delta G^\circ$  for  $\text{E}_P\text{-M-Glc-1-P} \rightleftharpoons \text{E}_D\text{-M-Glc-P}_2$  (Ray & Long, 1976b).




is equal to  $\Delta G^\ddagger$  for a one-step unimolecular process [cf. Glasstone et al. (1941)] with a numerically identical rate constant. Since  $k_{-6}^{\text{app}}$  depends on the rate constants for both (PO<sub>3</sub><sup>-</sup>) transfer and product dissociation, the transition states for both processes contribute to  $\text{EEB}_6$  (Ray, 1983). But regardless of the relative contributions of these steps (see below), a comparison of the effective energy barriers for the half-reactions of both forms of the enzyme shows that when Cd<sup>2+</sup> replaces Mg<sup>2+</sup>, the increase in  $\text{EEB}_6$  is significantly greater than that in  $\text{EEB}_1$ . In fact, if  $k_{-6}^{\text{app}}$  for the half-reaction involving Glc-6-P were the same for the Cd<sup>2+</sup> enzyme as it is for the Mg<sup>2+</sup> enzyme, the value of  $(k_{\text{cat}}^\ddagger)_{\text{Cd}}$  (Table I), which of necessity would include the energetics of the (PO<sub>3</sub><sup>-</sup>)-transfer step for the half-reaction involving Glc-1-P, would be at least  $0.3(k_{\text{cat}}^\ddagger)_{\text{Mg}}$  instead of about  $0.01(k_{\text{cat}}^\ddagger)_{\text{Mg}}$ . [The minimum value of  $k_{\text{t1}}^+$  for  $\text{E}_\text{P}\cdot\text{Cd}\cdot\text{Glc-1-P} \rightarrow \text{E}_\text{D}\cdot\text{Cd}\cdot\text{Glc-P}_2$  at 25 °C is  $230 \text{ s}^{-1}$ , while  $k_{-6}^{\text{app}} \geq (k_{\text{cat}}^\ddagger)_{\text{Mg}} = 630 \text{ s}^{-1}$ .] It seems likely that the more severe problem in the half-reaction involving Glc-6-P is related to a more rigid steric requirement for (PO<sub>3</sub><sup>-</sup>) transfer to and from a secondary as opposed to a primary hydroxyl group, coupled with the larger size of Cd<sup>2+</sup> relative to Mg<sup>2+</sup>.<sup>7</sup>

A comparison of rate constants for the actual bond-making/bond-breaking process facilitated by the Cd<sup>2+</sup> and Mg<sup>2+</sup> forms of the enzyme is more tenuous. In the first place, metal ion substitution alters the thermodynamic stability of "ground-state" species, although not to a major extent (see Figure 1).<sup>8</sup> Hence, we have chosen to make efficiency comparisons based on (PO<sub>3</sub><sup>-</sup>) transfer in the thermodynamically favorable direction, i.e., on  $\text{E}_\text{P}\cdot\text{M}\cdot\text{Glc-1-P} \xrightarrow{k_{\text{t1}}^+} \text{E}_\text{D}\cdot\text{M}\cdot\text{Glc-P}_2$ . Thus, the minimum values of  $k_{\text{t1}}^+$  are 230 and 1600 s<sup>-1</sup>, respectively, for the Cd<sup>2+</sup> and Mg<sup>2+</sup> enzymes. Although estimating the true ratio of  $k_{\text{t1}}^+$  values for the Mg<sup>2+</sup> and Cd<sup>2+</sup> enzymes from net rate constants requires a knowledge about the partitioning of intermediates between (PO<sub>3</sub><sup>-</sup>) transfer and product dissociation (cf. eq 23), it seems unlikely that  $(k_{\text{t1}}^+)_{\text{Mg}}$  exceeds  $(k_{\text{t1}}^+)_{\text{Cd}}$  by much more than an order of magnitude (see Appendix 3) and possibly not by this much.

In view of the above comparison, it seems reasonable to conclude that in its complex with the phosphoenzyme Mg<sup>2+</sup> coordinates directly to the enzymic phosphate group, since direct coordination has been observed with <sup>113</sup>Cd<sup>2+</sup> (Rhyu et al., 1983). Although the identity of the coordinating phosphate oxygen or oxygens has not been established, subsequent substrate binding probably produces at most minor alterations in the metal ion-phosphate interaction.<sup>9</sup> We thus view metal ion coordination in the  $\text{E}_\text{P}\cdot\text{Mg}\cdot\text{Glc-1-P}$  complex, just prior to

Scheme I: Relative Binding of Li<sup>+</sup> and Mg<sup>2+</sup> in both Ground- and Transition-State Complexes for (PO<sub>3</sub><sup>-</sup>) Transfer Facilitated by the Li<sup>+</sup> and Mg<sup>2+</sup> Complexes of Phosphoglucumutase with Glc-1-P<sup>a</sup>



<sup>a</sup> An assessment of the relative rate constants for the Li<sup>+</sup> and Mg<sup>2+</sup> forms of the enzyme is described under Results; that of the relative ground-state binding of Li<sup>+</sup> and Mg<sup>2+</sup> is described in footnote 14.

(PO<sub>3</sub><sup>-</sup>) transfer, as involving one or more oxy ligands of the enzymic phosphate group, either the bridging or nonbridging oxygens, or both.<sup>10</sup>

Of the nonphysiological metal ion activators of phosphoglucumutase, Mn<sup>2+</sup>, Co<sup>2+</sup>, Ni<sup>2+</sup>, Zn<sup>2+</sup>, and Cd<sup>2+</sup> (Ray, 1969), Cd<sup>2+</sup> differs from the physiological activator, Mg<sup>2+</sup>, to the greatest extent in terms of size (diameter = 2.28 versus 1.32 Å, respectively; Weast, 1987), ligand preference (Mg<sup>2+</sup> is a "hard" and Cd<sup>2+</sup> a "soft" metal ion; Pearson, 1963, 1965), and water substitution rates (about 10<sup>5</sup> s<sup>-1</sup> for Mg<sup>2+</sup> and about 3 × 10<sup>8</sup> s<sup>-1</sup> for Cd<sup>2+</sup>; Eigen & Hammes, 1962).<sup>11</sup> It is not obvious which of these properties or which combination of properties is responsible for the difference in the thermodynamics of (PO<sub>3</sub><sup>-</sup>) transfer by the Mg<sup>2+</sup> and Cd<sup>2+</sup> complexes of the enzyme (Ray & Long, 1976c). But, as is suggested above, these differences apparently affect the efficiency of (PO<sub>3</sub><sup>-</sup>) transfer from the enzyme to bound glucose 1-phosphate to only a relatively minor extent.

The above conclusion presents an interesting contrast with comparisons between the Mg<sup>2+</sup> and Li<sup>+</sup> forms of phosphoglucumutase. Before making this comparison/contrast, it should be pointed out that a variety of published and unpublished evidence suggests that Mg<sup>2+</sup> and Li<sup>+</sup> bind at the same site in phosphoglucumutase. For example, Li<sup>+</sup> binds competitively with Mg<sup>2+</sup> and on binding elicits sizable changes in the UV absorbance of the enzyme similar to those produced by Mg<sup>2+</sup> (Ray, 1978; Ray et al., 1978). In addition, the binding of glucose monophosphates (Ray et al., 1978) as well as glucose bisphosphate (W. J. Ray, Jr., unpublished results) is relatively unaffected by the substitution of Li<sup>+</sup> for Mg<sup>2+</sup>. Both spectral changes (Ma & Ray, 1980) and changes in the <sup>31</sup>P NMR chemical shift of the enzymic phosphate group in the Mg<sup>2+</sup> and Li<sup>+</sup> enzymes produced by the subsequent binding of Glc-6-P and a Glc-6-P analogue (W. J. Ray, Jr., unpublished results) also are quite similar.<sup>12</sup> In fact, Li<sup>+</sup> is much closer in size to Mg<sup>2+</sup> (diameter = 1.36 versus 1.32 Å, respectively; Weast, 1978) than is Cd<sup>2+</sup> (see above), and its

<sup>7</sup> A persistent problem in interpreting relative rates observed in these studies arises from an ambiguity in the structure of the  $\text{E}_\text{P}\cdot\text{Cd}\cdot\text{Glc-P}_2$  complex observed in <sup>31</sup>P NMR studies [cf. the accompanying paper, Post et al. (1989)]: two such complexes should exist, and both should act as intermediates in the overall reaction (Ray et al., 1973; Ma & Ray, 1980). Thus, the observed complex might be a "single" species because of equilibrium effects, as is suggested by other studies (W. J. Ray, Jr., unpublished results), or might represent two complexes in rapid equilibrium. At present, we are unable to distinguish between these possibilities, although we have shown that interconversion of two such complexes cannot limit the catalytic process if these are present in similar amounts (C. B. Post, unpublished results).

<sup>8</sup> The equilibrium constant that relates the two expected  $\text{E}_\text{P}\cdot\text{M}\cdot\text{Glc-P}_2$  complexes (see footnote 7) also may be altered by substituting Cd<sup>2+</sup> for Mg<sup>2+</sup>.

<sup>9</sup> The lack of a significant change in the <sup>31</sup>P-<sup>113</sup>Cd NMR coupling constant produced on binding of 6-deoxyglucose 1-phosphate, a substrate analogue that induces large substrate-like changes in the near-UV spectrum of the enzyme (Ma & Ray, 1980) but which cannot act as (PO<sub>3</sub><sup>-</sup>) acceptor, suggests that substrate binding does not appreciably alter the metal ion-phosphate interaction (W. J. Ray, Jr., unpublished results).

<sup>10</sup> Whether or not metal ion-phosphate coordination in  $\text{E}_\text{P}\cdot\text{Mg}\cdot\text{Glc-1-P}$  is ruptured in the (PO<sub>3</sub><sup>-</sup>) transfer that produces  $\text{E}_\text{D}\cdot\text{Mg}\cdot\text{Glc-P}_2$  still is a moot question [cf. Rhyu et al. (1983)]. However, a comparison of the equilibrium constant for  $\text{E}_\text{P}\cdot\text{Mg}\cdot\text{Glc-1-P} \rightleftharpoons \text{E}_\text{D}\cdot\text{Mg}\cdot\text{Glc-P}_2$  (Ray & Long, 1976) with that for transfer involving free reactants and products (Peck et al., 1968) suggests that thermodynamically favorable interactions increase, not decrease, when (PO<sub>3</sub><sup>-</sup>) transfer involves bound reactants and products.

<sup>11</sup> Differences in water substitution rate between ions of the same charge presumably reflect differences in ability to alter coordination geometry, i.e., to accommodate an elimination-addition or addition-elimination process [cf. Cotton and Wilkinson (1967)].


<sup>12</sup> In this study, 1-deoxyglucose 6-phosphate was bound to the Mg<sup>2+</sup> enzyme [to prevent (PO<sub>3</sub><sup>-</sup>) transfer], while Glc-6-P was bound to the Li<sup>+</sup> enzyme.

ligand preference ( $\text{Li}^+$  is a hard metal ion; Pearson, 1963, 1965) also is closer to  $\text{Mg}^{2+}$  than to  $\text{Cd}^{2+}$ . In spite of these similarities we observe a greater than  $10^8$ -fold difference between  $(k_{11}^+)_{\text{Mg}}$  and  $(k_{11}^+)_{\text{Li}}$  (Results), viz., a  $10^8$ -fold difference in the ability of these forms of the phosphoenzyme to facilitate  $(\text{PO}_3^-)$  transfer to bound Glc-1-P. By default, this difference, or at least the major part of it, most likely arises from the unit difference in electrostatic charge. In addition, the small effect on the equilibrium  $\text{E}_\text{P}\cdot\text{M}\cdot\text{Glc-1-P} \rightleftharpoons \text{E}_\text{D}\cdot\text{M}\cdot\text{Glc-P}_2$ , produced by replacing  $\text{Mg}^{2+}$  by  $\text{Li}^+$  (see Results) shows that if electrostatic effects are important for  $(\text{PO}_3^-)$  transfer from the phosphoenzyme to bound substrate, they are approximately equally important for the reverse process. Possible restrictions on the role of the metal ion in the bond-breaking/bond-making process facilitated by phosphoglucomutase that might be proscribed on the basis of these observations are considered below.

**Significance.** Although similar in several of their properties (see above),  $\text{Mg}^{2+}$  binds to most chelons much more strongly than does  $\text{Li}^+$ , especially those with a net negative charge, e.g., about  $10^6$ -fold more strongly to  $\text{EDTA}^{4-}$ . A reduced charge on the chelon not only reduces  $\text{Mg}^{2+}$  binding (by about  $10^5$ -fold for the  $\text{EDTA}^{3-}/\text{EDTA}^{4-}$  pair) but also reduces the ratio of binding constants for  $\text{Mg}^{2+}$  and  $\text{Li}^+$ , e.g., in the case of  $\text{NTA}^{3-}$  to about  $10^3$ -fold (Martel & Smith, 1974). Such equilibrium binding effects could be related to an increased "transition-state binding" (Jencks, 1969, 1975; Wolfenden, 1976, 1987) of  $\text{Mg}^{2+}$  relative to  $\text{Li}^+$  of at least  $(4 \times 10^8)$ -fold<sup>12</sup> observed in the first  $(\text{PO}_3^-)$ -transfer step of the phosphoglucomutase reaction (parentheses, Scheme I). Thus,  $\text{Mg}^{2+}$  and  $\text{Li}^+$  bind to  $\text{E}_\text{P}\cdot\text{Glc-1-P}$  and  $\text{E}_\text{D}\cdot\text{Glc-P}_2$  equally, while the Gibbs energy difference between the transition states for  $(\text{PO}_3^-)$  transfer in the  $\text{E}_\text{P}\cdot\text{Mg}\cdot\text{Glc-1-P}$  and  $\text{E}_\text{P}\cdot\text{Li}\cdot\text{Glc-1-P}$  complexes is equivalent to a binding difference for these metal ions in their respective transition states of at least 11.9 kcal/mol.

Of course, part of the apparent transition-state-binding effect, above, could be produced by masked ground-state phenomena. For example, the binding of  $\text{Mg}^{2+}$  (but not  $\text{Li}^+$ ) to the  $\text{E}_\text{P}\cdot\text{Glc-1-P}$  complex might produce an energetically unfavorable conformational change that greatly facilitates  $(\text{PO}_3^-)$  transfer and also makes the ground-state binding of  $\text{Mg}^{2+}$  much weaker than otherwise [cf. Jencks (1975)]. Although we cannot state that masked ground-state changes do not contribute to the  $\text{Li}^+/\text{Mg}^{2+}$  rate effect, several lines of evidence make  $\text{Mg}^{2+}$ -induced ground-state effects seem less likely than a transition-state effect. Thus,  $\text{Mg}^{2+}$  not only binds to  $\text{E}_\text{P}$ ,  $\text{E}_\text{P}\cdot\text{Glc-1-P}$ , and  $\text{E}_\text{D}\cdot\text{Glc-P}_2$  with almost equal affinity,<sup>13</sup> but bound substrate produces only a minor effect on  $\text{Mg}^{2+}$  binding and dissociation rates. Moreover, bound  $\text{Mg}^{2+}$  produces only a minor effect on the standard chemical potential of the enzymic phosphate group in both  $\text{E}_\text{P}$  and  $\text{E}_\text{P}\cdot\text{Glc-1-P}$  (as assessed in terms of equilibria such as  $\text{E}_\text{P}\cdot\text{M} \rightleftharpoons \text{E}_\text{D}\cdot\text{M} + \text{P}_i$ ; Ray & Long, 1976b), as well as on the chemical potential of the phosphates of the (bound) bisphosphate.<sup>14</sup> As a con-

Scheme II: Possible Representations of Bond Breaking and Bond Making in the Transition State for  $(\text{PO}_3^-)$  Transfer between Hydroxyl Groups Whose Anions Are Highly Basic and Possible Metal Involvements in Such Processes<sup>a</sup>



<sup>a</sup> In the process illustrated in (a), bond breaking in the transition state is essentially complete with little compensating bond formation (see Discussion). In (b), a metal ion is associated with the donor (leaving) group in the forward process (left to right) and with the acceptor (entering) group in the reverse process. In (c), a transition state with a pentacoordinate structure and a metal ion interacting with a peripheral oxygen or oxygens is represented.

sequence,  $\text{Mg}^{2+}$  (as well as  $\text{Li}^+$ ) also has only a relatively small effect on the equilibrium constant for  $(\text{PO}_3^-)$  transfer, viz., on  $\text{E}_\text{P}\cdot\text{Glc-1-P} \rightleftharpoons \text{E}_\text{D}\cdot\text{Glc-P}_2$  (Ray & Long, 1976).

But if ground-state differences are minimized, it is even more difficult to rationalize how a simple electrostatic effect such as positioning a metal ion close to a developing negative charge in the transition state could produce a rate effect of the observed magnitude—especially in view of what is known about bond breaking/bond making from solution studies of  $(\text{PO}_3^-)$  transfer and what has been suggested about the mechanism of the phosphoglucomutase reaction [on the basis of the efficiency of a sulfur analogue of the normal substrate as a  $(\text{PO}_3^-)$  acceptor (Knight et al., 1984)]: extensive bond breaking in the transition state with little compensating bond making. How a direct electrostatic effect might operate in the phosphoglucomutase reaction is further complicated, relative to most model reactions, by the fact that both donor and acceptor oxygens derive from very weakly acidic alcohols and that two proton-transfer steps, whose timing is obscure, must be considered in the overall process. In fact, because of the timing of proton transfer, the reaction may well be borderline [between  $\text{S}_\text{N}1$  and  $\text{S}_\text{N}2$ ; cf. Williams (1987)] and thus particularly difficult to deal with. Nevertheless, how the activating metal ion does *not* act in the phosphoglucomutase reaction can be considered (see below) in terms of two extreme cases so as to restrict the ways in which it might act. In both cases a dissociative or  $\text{S}_\text{N}1$ -like mechanism (Jencks, 1981; Williams, 1987; Cullis, 1987), with extensive P–O bond breaking and relatively little bond making in the transition state, as suggested above, is considered. Arguments are based on the conclusion (Rhyu et al., 1985) that the enzymic phosphate in  $\text{E}_\text{P}\cdot\text{Li}$  and  $\text{E}_\text{P}\cdot\text{Cd}$ , as well as the sugar phosphates

<sup>13</sup> The binding of  $\text{Mg}^{2+}$  to a specific complex in the equilibrium mixture of complexes  $\text{E}_\text{P}\cdot\text{Glc-1-P}$ ,  $\text{E}_\text{D}\cdot\text{Glc-P}_2$ , and  $\text{E}_\text{P}\cdot\text{Glc-6-P}$  cannot be measured directly. However, an approximation which shows that binding is similar for all three complexes can be obtained from the weighted average of the  $\text{Mg}^{2+}$  dissociation constants for each species at equilibrium [the "isotope exchange constant" for  $\text{Mg}^{2+}$  (45  $\mu\text{M}$ ; Ray et al., 1965)] and the equilibrium distribution of complexes in both the presence and absence of  $\text{Mg}^{2+}$  (Ray & Long, 1976c):  $K_d$  for  $\text{E}_\text{P}\cdot\text{Mg}\cdot\text{Glc-1-P}$ , 24  $\mu\text{M}$ ; for  $\text{E}_\text{P}\cdot\text{Mg}\cdot\text{Glc-P}_2$ , 66  $\mu\text{M}$ ; and for  $\text{E}_\text{P}\cdot\text{Mg}\cdot\text{Glc-6-P}$ , 20  $\mu\text{M}$ . The relative binding of  $\text{Mg}^{2+}$  and  $\text{Li}^+$  to  $\text{E}_\text{P}\cdot\text{Glc-1-P}$  is taken as equal to the relative binding of  $\text{Mg}^{2+}$  and  $\text{Li}^+$  to the equilibrium mixture of monophosphate and bisphosphate complexes of the enzyme, about 1:1 (Ray et al., 1978).

<sup>14</sup> For example, since  $\text{Mg}^{2+}$  binds to  $\text{E}_\text{D}\cdot\text{Glc-1-P}$  almost as well as to  $\text{E}_\text{D}\cdot\text{Glc-P}_2$  [dissociation constants are 300  $\mu\text{M}$  (calculated from constants in Ray et al. (1966)) and 66  $\mu\text{M}$  (footnote 8), respectively], to a first approximation the equilibrium  $\text{E}_\text{D}\cdot\text{Glc-P}_2 \rightleftharpoons \text{E}_\text{D}\cdot\text{Glc-1-P} + \text{P}_i$ , which defines the chemical potential of the 6-phosphate of bound Glc-P<sub>2</sub>, also is relatively independent of  $\text{Mg}^{2+}$  binding.



in E<sub>C</sub>-Cd-Glc-P<sub>2</sub>, are dianionic because of the pH dependence of their <sup>31</sup>P NMR chemical shifts.

Model studies of simple organic phosphates show that  $\partial \log k / \partial \text{p}K_{\text{lg}}$ , or  $-\beta_{\text{lg}}$ , is in the neighborhood of 1 for (PO<sub>3</sub><sup>-</sup>) transfer when the reactant is a dianionic phosphate with a good leaving group but that  $-\beta_{\text{lg}}$  approaches zero when the leaving group is strongly basic. The decreased value of  $-\beta_{\text{lg}}$  is reasonable since in such cases partial protonation of the incipient oxyanion is necessary to assist bond breaking, as is illustrated in Scheme IIa [cf. Hall and Williams (1986)]. Thus, in an enzymic reaction where the substrate binding step juxtaposes the incipient leaving group with a bound metal ion, *and the metal ion serves primarily to lower the effective pK<sub>a</sub> of the leaving group*, as in Scheme IIb, a large metal ion induced rate effect is *not* expected when the leaving group has a high pK<sub>a</sub> and bond breaking is assisted by partial protonation of that group, other factors being equal [cf. Herschlag and Jencks (1987)]. In such a case, a failure of the monovalent Li<sup>+</sup> to lower the pK<sub>a</sub> of the leaving group as efficiently as would Mg<sup>2+</sup> should not produce nearly as large a difference in (PO<sub>3</sub><sup>-</sup>)-transfer rate as was observed.

An alternative possibility is that metal ion coordination lowers the effective pK<sub>a</sub> of the leaving group sufficiently that bond breaking in an S<sub>N</sub>1-like process does not require assistance from proton transfer and that the active-site serine leaves as a metal-coordinated anion. This possibility also seems unlikely, even though in such a case a large decrease in rate would be expected from a large increase in effective pK<sub>a</sub> of the leaving group produced by employing a metal ion that is a less effective Lewis acid, e.g., substituting Li<sup>+</sup> for Mg<sup>2+</sup>. But, when the leaving group departs as an anion, as in ROPO<sub>3</sub><sup>2-</sup> + H<sub>2</sub>O → RO<sup>-</sup> + HPO<sub>4</sub><sup>2-</sup> + H<sup>+</sup>, a change in the pK<sub>a</sub> of the leaving group produces an even larger change in the equilibrium constant for (PO<sub>3</sub><sup>-</sup>) transfer. For example,  $\partial \log K_{\text{e}} / \partial \text{p}K_{\text{lg}}$  for the above process is -1.35 (Bourne & Williams, 1984) while  $\partial \log k / \partial \text{p}K_{\text{lg}}$  is in the neighborhood of -1 (cf. Cullis, 1987).<sup>15</sup> Hence, *on the sole basis of differences in induced pK<sub>a</sub>*, other factors being equal, replacing bound Mg<sup>2+</sup> by Li<sup>+</sup> should effect the equilibrium constant for the transfer process to an even larger extent than its effect on the forward rate constant ( $k_{\text{t1}}^+$ ). But while the observed rate effect for Li<sup>+</sup>/Mg<sup>2+</sup> replacement is large [(4 × 10<sup>8</sup>)-fold], the equilibrium effect is negligible (about 2-fold).

On the other hand, if the P-O bond-breaking/bond-making process in the phosphoglucomutase reaction were much more nearly synchronized or S<sub>N</sub>2-like than in most model reactions, substantial formation of the new P-O bond and much less extensive breaking of the old P-O bond would characterize the transition state. In such a case, charge density in general and charge density particularly at the peripheral oxygens should increase in the transition state. An increase in charge on the peripheral oxygens also might be coupled with increased electrostatic interactions in a manner analogous to the way in which the serine proteases use an oxyanion binding pocket (Kraut, 1977), although no evidence for a mechanism of this type has been obtained from model studies [cf. Herschlag and Jencks (1987)]. Whether such a mechanism can provide a reasonable rationale for the efficiency of (PO<sub>3</sub><sup>-</sup>) transfer exhibited by the Mg<sup>2+</sup> form of phosphoglucomutase, as well as the enormous differential rate effect observed when Li<sup>+</sup> is

substituted for bound Mg<sup>2+</sup>, must wait further studies in this system.

## APPENDIX 1

*Comparison of Saturation-Transfer Rates with the Steady-State Catalytic Constant: Cd(II) Enzyme.* Minimal estimates for the rate constants  $k_{\text{t1}}^+$  and  $k_{\text{t6}}^+$  (eq 4 and 5), the measured values of  $k_{-1}^{\text{app}}$  and  $k_{-6}^{\text{app}}$  (eq 1) for saturation transfer, plus the equilibrium concentration of the various intermediates (Ray & Long, 1976b) can be used to restrict allowed values for the steady-state catalytic constants  $k_{\text{cat}}^{\text{f}}$  and  $k_{\text{cat}}^{\text{r}}$  in the forward and reverse reactions, respectively. The process is illustrated for  $1/k_{\text{cat}}^{\text{f}}$ , which can be expressed as a series of terms, each related to the contribution made by a transition state between E<sub>P</sub>-Cd-Glc-1-P and E<sub>P</sub>-Cd + Glc-6-P to the effective energy barrier for the overall process [cf. eq 39 and 40 of Ray (1983)]:

$$1/k_{\text{cat}}^{\text{f}} = 1/k_{\text{t1}}^+ + (1/k_{\text{t6}}^-)(1 + k_{\text{t1}}^-/k_{\text{t1}}^+) + (1/k_{\text{d6}})[1 + (k_{\text{t6}}^+/k_{\text{t6}}^-)(1 + k_{\text{t1}}^-/k_{\text{t1}}^+)] \quad (8)$$

(The rate constants in eq 8 are defined in eq 4 and 5.) To transform eq 8 into eq 16, various factors and terms are replaced by three types of parameters: C<sub>1</sub> and C<sub>6</sub> are equilibrium constants evaluated previously (Ray & Long, 1976c);  $k_{-1}^{\text{app}}$  and  $k_{-6}^{\text{app}}$  are rate constants (see eq 1) evaluated in the accompanying paper; (1 + p<sub>1</sub>) and (1 + p<sub>6</sub>) are steady-state partition coefficients whose values have not been established.

$$C_1 = k_{\text{t1}}^+/k_{\text{t1}}^- = (\text{E}_\text{P}\cdot\text{Cd}\cdot\text{Glc-P}_2)/(\text{E}_\text{P}\cdot\text{Cd}\cdot\text{Glc-1-P}) \quad (9)$$

$$C_6 = k_{\text{t6}}^+/k_{\text{t6}}^- = (\text{E}_\text{P}\cdot\text{Cd}\cdot\text{Glc-P}_2)/(\text{E}_\text{P}\cdot\text{Cd}\cdot\text{Glc-6-P}) \quad (10)$$

$$k_{\text{t1}}^+ = k_{-1}^{\text{app}}(1 + p_1)C_1 \quad (11)$$

$$k_{\text{t6}}^- = k_{-6}^{\text{app}}(1 + p_6) \quad (12)$$

$$k_{\text{d6}} = k_{-6}^{\text{app}}(1 + 1/p_6)C_6 \quad (13)$$

$$p_1 = k_{\text{t1}}^+/k_{\text{d1}} \quad (14)$$

$$p_6 = k_{\text{t6}}^+/k_{\text{d6}} \quad (15)$$

Substituting eq 9–15 into eq 8 gives

$$1/k_{\text{cat}}^{\text{f}} = C_1^{-1}[k_{-1}^{\text{app}}(1 + p_1)]^{-1} + (1 + C_1^{-1}) \times [k_{-6}^{\text{app}}(1 + p_6)]^{-1} + (1 + C_1^{-1} + C_6^{-1})[k_{-6}^{\text{app}}(1 + p_6^{-1})]^{-1} \quad (16)$$

By using the values below, eq 16 becomes

$$\frac{1}{k_{\text{cat}}^{\text{f}}} = \frac{1}{k_{-6}^{\text{app}}} \left[ \frac{0.03}{1 + p_1} + \frac{1.04}{1 + p_6} + \frac{1.11}{1 + 1/p_6} \right] \quad (17)$$

where C<sub>1</sub> = 28.3, C<sub>6</sub> = 14.2,  $k_{-1}^{\text{app}}$  = 12 s<sup>-1</sup>, and  $k_{-6}^{\text{app}}$  = 11 s<sup>-1</sup> [see the accompanying paper, Post et al. (1989)]. Thus, because of the relatively large equilibrium constant for E<sub>P</sub>-Cd-Glc-1-P ⇌ E<sub>P</sub>-Cd-Glc-P<sub>2</sub>, the transition state for the process in eq 9, above (first term on right-hand side of eq 17), makes a negligible contribution to the effective energy barrier for  $k_{\text{cat}}^{\text{f}}$ . Hence

$$k_{\text{cat}}^{\text{f}} \approx \frac{k_{-6}^{\text{app}}(1 + p_6)}{1.04 + 1.11p_6} \quad (18)$$

which gives rise to the restriction on  $k_{\text{cat}}^{\text{f}}$  shown in eq 3 under Results.

An analogous process for the reverse reaction provides the following relationship:

<sup>15</sup> The absolute value of β is expected to be greater for the overall equilibrium than for the forward rate constant since  $\partial \log K_{\text{e}} / \partial \text{p}K_{\text{lg}} = -(\beta_{\text{lg}} + \beta_{\text{r}})$ , where β<sub>r</sub> refers to the effect of the altered pK<sub>a</sub> on the ability of the leaving group to act as the nucleophile in the reverse transfer process.

$$\frac{1}{k_{\text{cat}}^r} = \frac{1}{k_{-1}^{\text{app}}} \left[ \frac{0.08}{1 + p_6} + \frac{1.07}{1 + p_1} + \frac{1.11}{1 + 1/p_1} \right] \quad (19)$$

Considerations analogous to those above produce the relationship

$$10.8 \text{ s}^{-1} \leq k_{\text{cat}}^r \leq 11.2 \text{ s}^{-1}$$


The range of values for the steady-state parameters in initial velocity studies,  $k_{\text{cat}}^f$  and  $k_{\text{cat}}^r$ , calculated in this manner are in good agreement with the more directly calculated value for the steady-state rate constant describing saturation transfer at chemical equilibrium, since

$$k_{\text{tr}}^f = 1/2(1/k_{\text{cat}}^f + 1/k_{\text{cat}}^r)^{-1} \quad (20)$$

Thus, the range of values for  $k_{\text{tr}}^f$  calculated with the above values of  $k_{\text{cat}}^f$  and  $k_{\text{cat}}^r$  is 2.5–2.6 s<sup>-1</sup>, while the value calculated more directly from saturation-transfer measurements (which eliminate any dependency of  $k_{\text{tr}}^f$  on assumed values of  $p_1$  and  $p_6$ ; see eq 2) is 2.4 s<sup>-1</sup>.

## APPENDIX 2

*Comparison of Net Rate Constants for Phosphate Transfer by the Mg<sup>2+</sup> Enzyme.* Values of the net rate constants for the processes in eq 21 were approximated in the following way.



Equation 16 for  $1/k_{\text{cat}}^f$  and the analogous equation for  $1/k_{\text{cat}}^r$  comprise two simultaneous equations with four unknowns:  $k_{-1}^{\text{app}}$ ,  $k_{-6}^{\text{app}}$ ,  $p_1$ , and  $p_6$ . Hence, an expression for  $k_{-6}^{\text{app}}$  can be obtained in which the only unknowns are  $p_1$  and  $p_6$  and similarly for  $k_{-1}^{\text{app}}$ . In the latter case the expression takes the form

$$k_{-1}^{\text{app}} = \frac{a(p_1 + b)}{(p_1 + 1)(p_6 + c)} + \frac{d(p_1 + e)}{(p_1 + 1)} \quad (22)$$

Although  $k_{\text{tl}}^+$  and  $k_{\text{d1}}$ , which determine  $p_1$ , can assume a wide range of values and thus produce a wide variation in  $p$  (see eq 14), the values of  $k_{-1}^{\text{app}}$ ,  $k_{\text{tl}}^+/(1 + p_1)$ , produced by such a variation are much more severely restricted, as are the values of the corresponding equation for  $k_{-6}^{\text{app}}$ .

In theory, values of both  $p_1$  and  $p_6$  can range from very small to very large. But in the phosphoglucomutase reaction the two half-reactions are more or less mirror images of each other, and it seems reasonable that the partition of  $\text{E}_P\text{Mg}\cdot\text{Glc-1-P}$  between  $(\text{PO}_3^-)$  transfer and dissociation,  $p_1$ , should not be vastly different from the analogous partition of  $\text{E}_P\text{Mg}\cdot\text{Glc-6-P}$ ,  $p_6$ , especially since the equilibrium binding of Glc-1-P and  $\alpha$ -Glc-6-P, as well as  $k_{\text{cat}}^f$  and  $k_{\text{cat}}^r$ , differ only by a fewfold. Hence, in a survey of possible values of  $k_{-1}^{\text{app}}$  and  $k_{-6}^{\text{app}}$ , we assume that  $0.1 \leq p_1/p_6 \leq 10$ , but consider individual values of  $p_1$  and  $p_6$  from very small to very large. With this restriction a comprehensive survey of possible values shows that  $340 \text{ s}^{-1} \leq k_{-1}^{\text{app}} \leq 400 \text{ s}^{-1}$  ( $p_1 = 0.17$ ;  $p_6 = 1.7$ ) and  $960 \text{ s}^{-1} \leq k_{-6}^{\text{app}} \leq 1660 \text{ s}^{-1}$  ( $p_1 = 3$ ;  $p_6 = 0.3$ ).

## APPENDIX 3

*Comparison of Rate Constants for Phosphate Transfer by the Mg(II) and Cd(II) Enzymes.* An expression for the rate constant  $k_{\text{tl}}^+$  for transfer of the  $(\text{PO}_3^-)$  group of the enzyme to the 6-position of bound Glc-1-P (eq 5) for both metal ion forms (subscript M) of the enzyme can be obtained from eq 9 and 11 plus known values of  $C_1$  for  $\text{Mg}^{2+}$  and  $\text{Cd}^{2+}$  forms

of the enzyme (see Appendix 1). The ratio of these expressions is

$$\frac{(k_{\text{tl}}^+)_{\text{Mg}}}{(k_{\text{tl}}^+)_{\text{Cd}}} \approx 7 \frac{1 + (p_1)_{\text{Mg}}}{1 + (p_1)_{\text{Cd}}} \quad (23)$$

The  $(\text{PO}_3^-)$ -transfer rate for  $\text{Mg}^{2+}$  will be no more than an order of magnitude larger than for  $\text{Cd}^{2+}$  if any of the following relationships hold.

(a) The partition of  $\text{E}_P\text{M}\cdot\text{Glc-1-P}$  between  $(\text{PO}_3^-)$  transfer ( $k_{\text{tl}}^+$ ) and dissociation of Glc-1-P ( $k_{\text{d1}}$ ) is small for both  $\text{Mg}^{2+}$  and  $\text{Cd}^{2+}$ ; i.e., in the process,  $\text{E}_P\text{Mg}\cdot\text{Glc-P}_2 \rightarrow \text{E}_P\text{Mg} + \text{Glc-1-P}$ ,  $(\text{PO}_3^-)$  transfer is solely rate limiting. This is possible, although it would require a value of about  $1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  for  $(k_{\text{a1}})_{\text{Mg}}$ , the bimolecular rate constant for binding of Glc-1-P.

(b) The values of both  $(p_1)_{\text{Mg}}$  and  $(p_1)_{\text{Cd}}$  are large and essentially the same; i.e., product dissociation in the above process is solely rate limiting and  $(k_{\text{d1}})_{\text{Mg}} \approx (k_{\text{d1}})_{\text{Cd}}$ . The latter possibility is in accord with, but certainly not required by, the observation that the equilibrium binding of Glc-1-P to  $\text{E}_P\text{M}$  is essentially independent of the identity of  $\text{M}^{2+}$ , not only for  $\text{Mg}^{2+}$  and  $\text{Cd}^{2+}$  but also for  $\text{Zn}^{2+}$ ,  $\text{Co}^{2+}$ , and  $\text{Mn}^{2+}$  (Ray & Long, 1976c).

(c) The value of  $(p_1)_{\text{Mg}}$  is larger than  $(p_1)_{\text{Cd}}$ ; i.e., product dissociation is more nearly rate limiting for the less efficient metal ion form of the enzyme. Although one normally expects product dissociation to be more nearly rate limiting for a good substrate than for a poor one, such a pattern does not necessarily hold when reduced efficiency is produced by a change in the identity of the activating metal ion. For example, in the present system the rate constant for dissociation of Glc-P<sub>2</sub> from  $\text{E}_D\text{M}\cdot\text{Glc-P}_2$  is decreased by about 0.01 by the substitution of  $\text{Cd}^{2+}$  for  $\text{Mg}^{2+}$  (see Results) without a significant decrease in the equilibrium binding constant of Glc-P<sub>2</sub> (Ray & Long, 1976b); i.e., the association rate is similarly affected. Similarly,  $\text{E}_P\text{M}\cdot\text{Glc-1-P}$  may well dissociate more slowly when  $\text{M} = \text{Cd}$  than when  $\text{M} = \text{Mg}$ , so that  $k_{\text{d}}$  dominates the expression for  $(k_{\text{tl}}^+)_{\text{Cd}}$  but not that for  $(k_{\text{tl}}^+)_{\text{Mg}}$ .

Only if product dissociation from the  $\text{Mg}^{2+}$  enzyme is more nearly rate limiting than from the  $\text{Cd}^{2+}$  enzyme, viz., if  $(p_1)_{\text{Mg}} > (p_1)_{\text{Cd}}$ , would  $(\text{PO}_3^-)$ -transfer efficiency involving the 6-hydroxyl group of bound Glc-1-P be more than an order of magnitude greater for the  $\text{Mg}^{2+}$  than the  $\text{Cd}^{2+}$  enzyme.

**Registry No.** Glc-1-P, 59-56-3; Glc-6-P, 15209-11-7;  $\text{Cd}^{2+}$ , 22537-48-0;  $\text{Mg}^{2+}$ , 22537-22-0;  $\text{Li}^+$ , 17341-24-1; phosphoglucomutase, 9001-81-4.

## REFERENCES

- Albery, W. J., & Knowles, J. R. (1976) *Biochemistry* 15, 5631–5640.
- Boltz, D. F. (1972) in *Analytical Chemistry of Phosphorus Compounds* (Halmann, H., Ed.) p 59, Wiley-Interscience, New York.
- Bourne, N., & Williams, A. (1984) *J. Org. Chem.* 49, 1200–1204.
- Cleland, W. W. (1975) *Biochemistry* 14, 3220–3224.
- Cotton, A. F., & Wilkinson, G. (1966) *Advanced Inorganic Chemistry*, 2nd ed., p 158, Wiley, New York.
- Cullis, P. M. (1987) in *Enzyme Mechanisms* (Page, M. J., & Williams, A., Eds.) pp 178–220, Royal Society of Chemistry, London.
- Douderoff, M. (1955) *Method Enzymol.* 1, 225–229.
- Eigen, M., & Hammes, G. G. (1962) *Adv. Enzymol. Relat. Areas Mol. Biol.* 25, 1–38.
- Glasstone, S., Laidler, K. J., & Eyring, H. (1941) *The Theory of Rate Processes*, McGraw-Hill, New York.

- Hall, A., & Williams, A. (1986) *Biochemistry* 25, 4784-4790.
- Herslag, D., & Jencks, W. P. (1987) *J. Am. Chem. Soc.* 109, 4665-4674.
- Jencks, W. P. (1975) *Adv. Enzymol. Relat. Areas Mol. Biol.* 43, 219-410.
- Jencks, W. P. (1981) *Chem. Soc. Rev.* 10, 345-375.
- Jencks, W. P. (1985) *Chem. Rev.* 85, 511-527.
- Kirby, A. J., & Varvoglis, A. G. (1967) *J. Am. Chem. Soc.* 89, 415-423.
- Knight, W. B., Rendina, A. R., & Cleland, W. W. (1984) *Fed. Proc., Fed. Am. Soc. Exp. Biol.* 43, 2011.
- Kraut, J. (1977) *Annu. Rev. Biochem.* 46, 331-335.
- Lang, G., & Michal, G. (1974) in *Methods of Enzymatic Analysis* (Bergmeyer, H. U., Ed.) p 1128, Verlag, Weinheim, FRG.
- Ma, C., & Ray, W. J., Jr. (1980) *Biochemistry* 19, 751-795.
- Magneson, G. R., Puvathingal, J. M., & Ray, W. J., Jr. (1987) *J. Biochem. (Tokyo)* 262, 11140-11148.
- Martel, A. E., & Smith, R. M. (1974) *Critical Stability Constants*, pp 139, 204, Plenum, New York.
- Michal, G. (1984) in *Methods of Enzymatic Analysis* (Bergmeyer, H. U., Ed.) 3rd ed., pp 191-198, Verlag, Basel.
- Nichol, L. W., Sculley, M. J., Ward, L. D., & Winzor, D. J. (1983) *Arch. Biochem. Biophys.* 222, 574-581.
- Pearson, R. G. (1963) *J. Am. Chem. Soc.* 85, 3533-3539.
- Pearson, R. G. (1965) *Chem. Eng. News* 43(May 31), 90-91.
- Pearson, R. G. (1967) *J. Am. Chem. Soc.* 89, 1827-1836.
- Peck, E. J., Jr., Kirkpatrick, D. S., & Ray, W. J., Jr. (1968) *Biochemistry* 7, 152-162.
- Post, C. B., Ray, W. J., Jr., & Gorenstein, D. G. (1989) *Biochemistry* (preceding paper in this issue).
- Ray, W. J., Jr. (1969) *J. Biol. Chem.* 244, 3740-3747.
- Ray, W. J., Jr. (1983) *Biochemistry* 22, 4625-4637.
- Ray, W. J., Jr. (1986) *J. Biol. Chem.* 261, 275-278.
- Ray, W. J., Jr., & Roscelli, G. A. (1964) *J. Biol. Chem.* 239, 3935-3941.
- Ray, W. J., Jr., & Rocelli, G. A. (1966) *J. Biol. Chem.* 241, 3499-3506.
- Ray, W. J., Jr., & Peck, E. J. (1972) *Enzymes (3rd Ed.)* 6, 407-477.
- Ray, W. J., Jr., & Long, J. W. (1976a) *Biochemistry* 15, 3990-3993.
- Ray, W. J., Jr., & Long, J. W. (1976b) *Biochemistry* 15, 3393-4006.
- Ray, W. J., Jr., & Long, J. W. (1976c) *Biochemistry* 15, 4018-4025.
- Ray, W. J., Jr., & Puvathingal, J. M. (1986) *J. Biol. Chem.* 261, 11544-11549.
- Ray, W. J., Jr., Mildvan, A. S., & Long, J. W. (1973) *Biochemistry* 12, 3724-3732.
- Ray, W. J., Jr., Hermodson, M. A., Puvathingal, J. M., & Mahoney, W. C. (1982) *J. Biol. Chem.* 258, 9166-9174.
- Rhyu, G. I., Ray, W. J., Jr., & Markley, J. L. (1984) *Biochemistry* 23, 252-260.
- Sugino, Y., & Miyoshi, Y. (1964) *J. Biol. Chem.* 239, 2360-2364.
- Weast, R. C., Ed. (1980) *Handbook of Chemistry and Physics*, 61st ed., Chemical Rubber Co., Cleveland, OH.
- Williams, A. (1987) in *Enzyme Mechanisms* (Page, M. J., & Williams, A., Eds.) pp 123-139, Royal Society of Chemistry, London.
- Wolfenden, R., & Frick, L. (1987) in *Enzyme Mechanisms* (Page, M. J., & Williams, A., Eds.) pp 97-122, Royal Society of Chemistry, London.

## Rat Brain Has the $\alpha 3$ Form of the $(\text{Na}^+, \text{K}^+)\text{ATPase}^\dagger$

Yen-Ming Hsu and Guido Guidotti\*

Department of Biochemistry and Molecular Biology, Harvard University, Cambridge, Massachusetts 02138

Received June 7, 1988; Revised Manuscript Received August 8, 1988

**ABSTRACT:** Multiple forms of the catalytic subunit of the  $(\text{Na}^+, \text{K}^+)\text{ATPase}$  have been identified in rat brain. While two of them ( $\alpha 1$  and  $\alpha 2$ ) have been well characterized, the third form ( $\alpha 3$ ) of these catalytic subunits only recently has been described by cDNA cloning; the corresponding polypeptide has not been isolated. In this paper it is shown that rat brain contains the  $\alpha 3$  chain. The catalytic subunits of the  $(\text{Na}^+, \text{K}^+)\text{ATPase}$  from rat brain axolemma were purified by SDS-PAGE and subjected to formic acid cleavage. Amino acid sequence analysis of the resulting fragments revealed that axolemma has the  $\alpha 3$  form of the catalytic subunit. In addition,  $\alpha 3$ -specific antiserum was raised in rabbits immunized with a synthetic peptide. Immunoblotting with this antiserum revealed that the  $\alpha 3$  form of the  $(\text{Na}^+, \text{K}^+)\text{ATPase}$  is present also in whole brain microsomes. In SDS-PAGE, the mobilities of the three catalytic subunits of brain  $(\text{Na}^+, \text{K}^+)\text{ATPase}$  follow the order  $\alpha 1 > \alpha 2 > \alpha 3$ . Determination of the ouabain-inhibitable ATPase activity indicates that if the  $\alpha 3$  form of the  $(\text{Na}^+, \text{K}^+)\text{ATPase}$  is able to hydrolyze ATP, it is present in a form of the enzyme with a high affinity for this cardiac glycoside and is similar to the  $\alpha 2$  form in this respect.

$(\text{Na}^+, \text{K}^+)\text{ATPase}$  [sodium and potassium ion activated ATP phosphohydrolase (EC 3.6.1.3)] is the enzyme in the plasma membrane of animal cells that establishes the sodium and potassium ion gradients across the plasma membrane at the expense of ATP. The enzyme is composed of two subunits,

a large catalytic subunit of  $M_r$  112000 ( $M_r$  100000 by SDS-PAGE) and a smaller glycoprotein subunit of  $M_r$  50000 of unknown function in a 1:1 stoichiometry (Cantley, 1986; Jorgensen, 1982). Biochemical studies revealed that there are two forms of the large subunit, which are called  $\alpha 1$  and  $\alpha 2$  chains (Sweadner, 1979; Schellenberg et al., 1981; Matsuda et al., 1984; Lytton et al., 1985). [In this paper, the  $\alpha$ ,  $\alpha+$ ,

<sup>†</sup>This work was supported by NIH Grant HL 08893 to G.G.