

Myosin Catalytic Domain Flexibility in MgADP[†]

Dražen Raucher,[‡] Cecilia P. Sár,[§] Kálman Hideg,[§] and Piotr G. Fajer^{*,‡}

Institute of Molecular Biophysics and Department of Biological Science, Florida State University, Tallahassee, Florida 32306-3015, and Central Research Laboratory, Chemistry, University of Pécs, Pécs, Hungary, H-7643

Received August 3, 1994; Revised Manuscript Received September 12, 1994[®]

ABSTRACT: Conventional EPR studies of muscle fibers labeled with a novel α -iodoketo spin label at Cys-707 of the myosin head revealed substantial internal domain reorganization on the addition of ADP to rigor fibers. The spin probes that are well-ordered in the rigor state become disordered and form two distinct populations. These orientational changes do not correspond to rotation of the myosin catalytic domain as a whole because other probes (maleimide and iodoacetamide nitroxides attached to the same Cys-707 of myosin head) report only a small ($5\text{--}10^\circ$) torsional rotation and little or no change in the tilt angle [Ajtai *et al.* (1992) *Biochemistry* 31, 207–17; Fajer (1994) *Biophys. J.* 66, 2039–50]. In the presence of ADP, the labeled domain becomes more flexible and executes large-amplitude microsecond motions, as measured by saturation-transfer EPR with rates ($\tau_r = 150\ \mu\text{s}$) intermediate between the rotations of detached ($\tau_r = 7\ \mu\text{s}$) and rigor heads ($\tau_r = 2500\ \mu\text{s}$). This finding contrasts with an absence of global motion of the myosin head in ADP ($\tau_r = 2200\ \mu\text{s}$) as reported by the maleimide spin label. Our results imply that the myosin head in a single chemical state (AM-ADP) is capable of attaining many internal configurations, some of which are dynamic. The presence of these slow structural fluctuations might be related to the slow release of the hydrolysis products of actomyosin ATPase.

The cyclic interaction of myosin and actin filaments forms the basis for muscle contraction (Huxley, 1969, 1974). The sequence of structural events beginning with the attachment of myosin heads to actin, followed by a structural change in the actomyosin complex, and subsequent detachment of the heads allows for force generation and length change of the cell. This mechanical cycle is driven by the chemical cycle of actomyosin ATPase. Actin and myosin form a strong complex in the absence of nucleotides, which can be dissociated by the binding of MgATP¹ to a site on the myosin head. The MgATP is subsequently cleaved by myosin, and the products of this hydrolysis are released after the myosin has rebound to actin. It is natural to assume that each of the chemical states might induce a different structure of the actomyosin complex, and much effort has been devoted to the structural characterization of muscle proteins in different chemical states. A recent determination of the atomic structure of actin (Kabsch *et al.*, 1990) and myosin (Rayment *et al.*, 1993a) and subsequent “docking” experiments in

which actin and the myosin head were brought together to fit a 3D reconstruction of the complex from electron microscopy have suggested large shape changes of the head that might occur on nucleotide and actin binding (Rayment *et al.*, 1993b). Such changes have been observed by a comparison of the length of the myosin head in rigor and ADP using low-angle X-ray scattering (Wakabayashi *et al.*, 1992) and electrical birefringence (Highsmith & Eden, 1992). ADP is known to induce changes in optical probes (IATR dichroism) (Borejdo *et al.*, 1982; Burghardt *et al.*, 1983) which most probably correspond to local changes in the head internal structure, because EPR has suggested that the head does not substantially reorient as a rigid body (Fajer *et al.*, 1990; Fajer, 1994a).

Internal fluctuations of myosin heads induced by nucleotide binding have been inferred previously from changes of sulfhydryl reactivity (Burke & Reisler, 1977; Schaub *et al.*, 1975), cross-linking studies (Wells & Yount, 1979; Lu & Wong, 1989), NMR (Shriver & Sykes, 1982), and EPR (Barnett & Thomas, 1987). Most of these studies were confined to isolated myosin in solution and were not suited to resolving or observing the heterogeneity of the structures within a complex of myosin and nucleotide.

In this paper, we report on local domain changes induced by ADP in muscle fibers labeled with a novel iodoketo spin label [IKSL: 3-(2-iodopropionyl)-1-oxy-2,5-dihydro-2,2,5,5-tetramethyl-1H-pyrrole]. The observation of two distinct populations while attached to the actin filament strongly implies that a multitude of local structures correspond to a single chemical state. Furthermore, these conformations are not static as in rigor; there is considerable domain mobility in the 100–200- μs range. Because MSL, which is more sensitive to global reorientation of the head and attached at the same Cys-707, does not sense these motions, we believe that these motions correspond to nucleotide-induced slow structural fluctuations that might be important in the release

[†] Research was sponsored by National Science Foundation IBN-9206658, an Initial Investigator Award of the American Heart Association (to P.F.), and Hungarian Research Foundation OTK-3/42/1990 (to K.H.).

* Address correspondence to this author; e-mail address: Fajer@sb.fsu.edu.

[‡] Florida State University.

[§] University of Pécs.

[®] Abstract published in *Advance ACS Abstracts*, October 15, 1994.

¹ Abbreviations: ADP, adenosine 5'-diphosphate; Ap₅A, P₁P₅-diadenosine 5'-pentaphosphate; ATP, adenosine 5'-triphosphate; DTNB, 5,5'-dithiobis(2-nitrobenzoic acid); EDTA, ethylenediaminetetraacetic acid; EGTA, ethylene glycol bis(β -aminoethyl ether)-N,N,N',N'-tetraacetic acid; EPR, electron paramagnetic resonance; IKSL, 3-(2-iodopropionyl)-1-oxy-2,5-dihydro-2,2,5,5-tetramethyl-1H-pyrrole; MOPS, 3-(N-morpholino)propanesulfonic acid; MSL, N-(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl)maleimide; NMR, nuclear magnetic resonance; S1, myosin subfragment 1; ST-EPR, saturation transfer electron paramagnetic resonance; FDNB, 2,4-dinitrofluorobenzene; IAA, iodoacetamide.

of products during the actomyosin ATPase cycle. The presence of different conformational substates in myoglobin has been linked to different reaction rates (Hong *et al.*, 1990; Frauenfelder *et al.*, 1991).

EXPERIMENTAL SECTION

Muscle fibers from rabbit psoas were harvested and glycerinated as described by Fajer *et al.* (1988). The glycerinated fibers were dissected into 0.5-mm bundles and washed with labeling buffer, LB (60 mM KAc, 20 mM MOPS, 5 mM MgCl₂, 1 mM EGTA, 1 mM NaN₃, pH 6.5). In order to block cysteines other than Cys-707 of myosin, we pretreated the fibers in rigor with 120 μ M DTNB (reversible sulfhydryl reagent) in LB for 120 min at pH 8.0 (Naber and Cooke, private communication). The labeling with 0.25 mM IKSL (α -iodopropionyl)pyrroline spin label or MSL [*N*-(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl)maleimide] was accomplished in LB at pH 6.5 in the presence of 5 mM MgATP, which detaches the myosin heads increasing the reactivity of Cys-707. Incubation time was 180 min for IKSL and 20 min for MSL. After labeling, the TNB blocking groups were removed by repeated thiolysis with 30 mM DTT in the presence of 5 mM MgATP at 20 °C for 30 min. The extent of labeling as estimated from the integrated intensity of the EPR signal and protein determination was 0.55 spin label per myosin head. The efficiency of TNB thiolysis was 80% as determined colorimetrically.

The rigor buffer (RB) consisted of 130 mM KPr, 2 mM MgCl₂, 1 mM EGTA, 20 mM MOPS, and 1 mM NaN₃, pH 7.0. Relaxation was achieved by the addition of 5 mM MgATP to the rigor solution. ADP solution was made from rigor solution with the addition of 4 mM MgADP plus 100 μ M Ap₅A to inhibit myofibrillar myokinase, 10 mM glucose, and 1 mg/mL hexokinase to remove contaminant ATP. All chemicals were obtained from Sigma Chemical Co. (St. Louis, MO).

Myosin subfragment 1 (S1) was prepared from myosin according to the method of Weeds and Taylor (1975) and was labeled as described by Fajer *et al.* (1988). Labeled S1 was covalently attached to isothiocyanate glass beads (Machleidt & Wachter, 1977). The beads were incubated overnight with labeled S1 at an equimolar ratio of protein and isothiocyanate in S1 buffer, 120 mM KCl, 10 mM MOPS, and 1 mM EDTA, pH 7.0. The unbound S1 was removed by repeated washes. Fiber decoration experiments were performed by a 20-min infusion of 10 mg/mL IKSL-S1 in RB, followed by the washout of unbound molecules with rigor solution.

EPR and ST-EPR spectra were obtained as previously described (Raucher & Fajer, 1994) with a Bruker ECS 106 spectrometer (Bruker Instruments, Billerica, MA). Conventional spectra of fibers parallel to the magnetic field were obtained in a modified TM₁₁₀ cavity; ST-EPR spectra of samples perpendicular to the field were collected in a TE₁₀₂ cavity. The experiments were performed at 20 °C. Effective rotational correlation times (τ_r) were obtained by comparison to spectra of hemoglobin tumbling in media of known viscosity (Thomas *et al.*, 1976). This comparison was accomplished on normalized integrated intensity of ST-EPR spectra which circumvents orientational effects in the spectra (Squire & Thomas, 1988).

Synthesis of α -Iodoketone Spin Label. A new spin label has been designed to achieve the specificity of the commonly




FIGURE 1: Structures of iodoacetamide spin label (left), α -iodoketo spin label (middle), and maleimide spin label (right). Arrows denote partial neutralization of the keto group electron withdrawing effect by the NH of iodoacetamide spin label.

used iodoacetamide spin label (IASL) but with steric restrictions of the IASL librational motion. The spacer between the cysteine reacting group (iodine) and the nitroxide has been reduced by one bond length; the methyl group was introduced next to the protein attachment point hindering *cis-trans* isomerization of the S-C bond, and finally the six-membered piperidine ring of nitroxide was substituted with the five-membered pyrrole ring, which is more rigid (Hideg & Hankovszky, 1989). This iodoketo spin label is more reactive than the corresponding iodoacetamide since the NH group neutralizes the electron-withdrawing effect of the keto group (see Figure 1).

The synthesis was accomplished as follows: to a solution of 3-(2-bromopropionyl)-2,5-dihydro-2,2,5,5-tetramethyl-1H-pyrrol-1-yloxy radical (1.0 mmol, 275.2 mg) in dry acetone (50 mL), NaI (2.0 mmol, 300.0 mg) was added and refluxed for 30 min (Hankovszky *et al.*, 1990). The mixture was then diluted with water, extracted with chloroform, dried (Na₂SO₄), and evaporated to dryness. The solid yellow residue was purified by preparative TLC with hexane-EtOAc (4:1) on Merck Silica Gel 60 F-254 plates to give 273.8 mg (85%) of 3-(2-iodopropionyl)-2,5-dihydro-2,2,5,5-tetramethyl-1H-pyrrol-1-yloxy radical as yellow crystals, mp 110–112 °C. Anal. Calcd for [C₁₁H₁₇INO₂ (322.17)]: C, 41.01; H, 5.32; N, 4.35; I, 39.39%. Found: C, 40.89; H, 5.22; N, 4.28; I, 39.30%.

RESULTS

Orientation. The spectra of IKSL-labeled fibers in rigor and relaxation (Figure 2) display the same characteristic features as the spectra of fibers labeled with MSL, implying a high degree of order in rigor and dynamic relaxation in ATP. In rigor, the spectra show three sharp resonances due to interaction with the ¹⁴N nucleus of the nitroxide. The splitting of the resonances, $2T_{eff} = 17.9 \pm 0.2$ G, is similar to that of MSL-labeled fibers (18.0 ± 0.2 G), suggesting that the orientation of the *z*-axis of the nitroxide is similar for the two labels [MSL spectra have been extensively published; e.g., Thomas and Cooke (1980), Fajer *et al.* (1991), and Raucher *et al.* (1994)]. The width of the low-field resonance at half-height is 3.9 ± 0.2 versus 3.6 ± 0.2 G for MSL, because of the high order of the labeled domains in rigor. The addition of ATP detaches the actin-bound heads, and the ensuing spectrum has a characteristic, nearly isotropic line shape indicating large head disorder (Thomas & Cooke, 1980; Thomas *et al.*, 1980). The similarity of the reported changes is not coincidental since both MSL and IKSL are covalently bound to the same site (Cys-707) on the myosin head. The identification of the IKSL-modified




FIGURE 2: Conventional EPR spectra of IKSL-labeled muscle fibers oriented parallel to the magnetic field: (a) rigor; (b) 4 mM MgADP; (c) fibers relaxed with 5 mM MgATP. Upper inset: Comparison of the line shapes of the ordered peak P_2 in rigor (dotted) and ADP (solid). Lower inset: Comparison of the disordered peaks P_1 in relaxation (dashed) and ADP (solid).

site was accomplished by competition with SH1-specific probes: IASL, IAA, and FDNB (Raucher *et al.*, 1994).

The spectral line shape in the presence of 4 mM MgADP is significantly different from the rigor or relaxed spectra. There is substantial spectral intensity at both P_1 (disordered peak) and P_2 (ordered peak), implying that a fraction of the probes is ordered at an average angle similar to that observed in rigor, while the remainder of the probes are disordered. These populations are, however, not identical to rigor and relaxation. The ADP spectrum cannot be simulated by a linear composition of relaxed and rigor spectra. A close inspection of the P_2 peak (Figure 2, upper inset) reveals that the ordered population is distinct from the rigor orientation. Likewise, a comparison of the P_1 regions of the disordered component and the relaxed spectrum in the lower inset of Figure 2 shows the differences between the disordered fraction in ADP and the relaxed heads.

Because disordering of the heads is often associated with detachment, e.g., PP_i and AMPPNP (Pate & Cooke, 1988; Fajer *et al.*, 1988), we have measured the detachment of IKSL-labeled S1 decorating unlabeled fibers. As for any other label, the spectrum of decorated IKSL-S1 in the absence of any nucleotide is identical to that of intrinsically labeled fibers in rigor (data not shown). The addition of ADP to decorated fibers resulted in changes identical to those in intrinsically labeled fibers (Figure 3, inset). Importantly, subsequent perfusion of the fiber with ADP solution resulted in little change of total intensity, indicating that the labeled heads remained bound in ADP. Over a period of 35 min, only 10% of the spins were lost to detachment and washed out from the muscle fiber (Figure 3). The myosin heads were not trapped with the fiber as the addition of 5 mM MgATP washed out all but 15% of the labeled S1 from the fiber within a 10-min period.

Therefore, the difference in the line shape at P_1 along with the strong binding of all of the heads in the presence of ADP implies that the labeled domain of the attached myosin heads is undergoing significant disordering. This behavior can be




FIGURE 3: Washout of IKSL-S1 from decorated muscle fibers. The amount of bound S1 was determined by integration of EPR signals collected at 3.5-min intervals. The first arrow denotes the addition of 4 mM MgADP to the elution buffer; the second arrow denotes the change to relaxing solution. Inset: EPR spectrum of decorating IKSL-S1.

contrasted with that of the MSL- and IASL-labeled heads; in both cases a high degree ($\pm 8^\circ$) of order is maintained in ADP (Fajer *et al.*, 1990; Fajer, 1994a). The difference is not due to possible structural changes induced by the iodoketo label since both the tension, steady-state ATPase, and the kinetic rates of the cycle were not affected by labeling (Raucher *et al.*, 1994). Thus, we conclude that the IKSL label is sensitive to local disordering of the immediate probe environment, whereas the other labels report the global behavior of the head.

Mobility. The saturation-transfer EPR spectra in Figure 4 demonstrate that the disordering of the IKSL probes is accompanied by the onset of rotational motion. The spectra in ADP are intermediate between the immobilized rigor heads, correlation time $\tau_r = 2-3$ ms, and the rapidly rotating relaxed heads, $\tau_r = 3-10$ μ s (Table 1) (calculated from the integrated intensities which avoids orientational effects in ST-EPR spectra). Because neither of the two distinct orientational populations is rigor- or relaxed-like, no rigorous attempt was made to assign any particular mobility to the ordered and disordered fractions. However, if we equate the conventional line shapes of the ordered and disordered fractions to rigor and relaxation respectively, then we can estimate the disordered fraction to be 60–70 mol %; similar analysis of ST-EPR spectra indicates the mobile (relaxed-like) fraction to be 40 mol %. The difference between the orientational and motional values points to a difference between the mobility of the disordered fraction and that of the relaxed heads. Because the heads in the ordered fraction are unlikely to be more immobilized than the rigor heads, the mobility of the disordered heads must be lower than that of the heads in the presence of ATP.

In contrast to IKSL, MSL-labeled fibers in the presence of ADP show no change compared to rigor spectra (Figure 4), which restricts the global movement of the head to the millisecond time range. At least parts of the head or the myosin head as a whole is very immobilized.

The question arises whether the disordering and the flexibility arise from librational motion of the label with respect to the protein. A standard method of assessing librational motions is to immobilize the protein on glass beads and look at the mobility of the label, as was done for the myosin head (Bottomley & Trayer, 1980; Thomas *et al.*, 1980), myosin light chains (Hambly & Cooke, 1990), troponin C (Li & Fajer, 1994), and tropomyosin (Szczesna and Fajer, private communication). The conventional EPR




FIGURE 4: Saturation-transfer EPR spectra of IKSL fibers (left) and MSL-labeled fibers (right). Top: In the absence of nucleotides, rigor; middle: in 4 mM ADP; bottom: relaxed with 5 mM ATP. Second harmonic, out-of-phase spectra (dotted) and their integrals (solid). The height of the arrows denotes relative restriction on mobility in each state.

Table 1: Rotational Correlation Times (τ_r) of IKSL- and MSL-Labeled Fibers^a

sample	IKSL	MSL
rigor	2000–3000	2000–3000
ADP	100–200	1800–2800
ATP	5–10	3–8

^a τ_r in microseconds.

and ST-EPR spectra of IKSL-S1 cross-linked to glass beads (Figure 5) were identical in rigor, ADP, and ATP. The hyperfine splitting of conventional spectra was 69.9 G irrespective of the nucleotide (Table 2), suggesting no nanosecond motion of the label with respect to the protein. This immobilization is further extended to the microsecond regime of ST-EPR spectra; spectral intensity and the diagnostic line height ratios in Table 2 are indicative of the absence of motions faster than $\tau_r = 1$ ms. Interestingly, no onset of microsecond mobility in the fibers is observed on the addition of ADP. The cross-linked S1 is enzymatically active and capable of ATP hydrolysis, excluding the possibility of a gross change in the structure due to covalent cross-linking.

DISCUSSION

Conventional EPR spectroscopy of IKSL-labeled myosin in skinned muscle fibers revealed significant ADP-induced local changes in both orientation and mobility of the domain containing Cys-707. The addition of ADP to rigor fibers induced the appearance of a bimodal orientational population with components of varying degrees of order, intermediate between rigor and relaxation. The more-ordered component was similar to that observed in rigor but had a wider orientational distribution; the disordered component was more ordered than the distribution observed for relaxed fibers. The mobility of the labeled domain as measured by ST-EPR was also intermediate between the microsecond mobility of relaxation and the millisecond mobility of rigor. The observed changes are consistent with a number of findings implying structural changes on the binding of ADP to the myosin head in solution. Our findings extend the presence of these structural changes to heads interacting with actin in muscle fibers and characterize these changes in terms of the orientation and mobility of the labeled domain.

Other Studies. In solution, the binding of ADP to myosin heads causes a number of well-documented changes [re-




FIGURE 5: Conventional EPR spectra (left) and ST-EPR of IKSL-S1 covalently bound to glass beads. Top: in the absence of nucleotides, rigor; middle: 4 mM ADP; bottom: 5 mM ATP.

Table 2: Mobility of IKSL-S1 Bound to Glass Beads

sample	STEPR intensity ^a (% rigor)	L''/L^b	H''/H^c	$2T_{\text{eff}}$ (G) ^d
no nucleotide	100	0.74	0.86	69.9
ADP	101	0.75	0.86	69.9
ATP	101	0.75	0.87	69.9

^a Double-integrated intensity of ST-EPR spectra normalized to rigor, error is $\pm 5\%$. ^b Error = ± 0.05 . ^c Error = ± 0.06 . ^d Error = ± 0.1 G.

viewed in Cooke and Highsmith (1983)]. The reactivity of Cys-707 and Cys-697 increases in the presence of ADP (Reisler *et al.*, 1977), the distance between these two cysteines decreases by 4–7 Å as probed by chemical cross-linkers (Burke & Reisler, 1977; Wells & Yount, 1979) and by fluorescence energy transfer (Dalbey *et al.*, 1983). The mobility of some spin probes (IASL) is sensitive to ADP binding, albeit not as much as in the M•ATP and M•ADP•P_i states (Seidel *et al.*, 1970; Barnett & Thomas, 1987). Since Cys-707 is near the active-site pocket in the crystal structure of the myosin head (Rayment *et al.*, 1993a), it should not be surprising that the probe environment is changing in the presence of nucleotides. Indeed, a change of shape due to the closure and opening of the active site induced by different states of bound nucleotide was postulated to result in force generation by muscle (Highsmith & Eden, 1992; Wakabayashi *et al.*, 1992; Rayment *et al.*, 1993b). Using low angle X-ray diffraction (Wakabayashi *et al.*, 1992) and electrical birefringence (Highsmith & Eden, 1992) changes were observed on ADP binding in the chord length of the myosin head in solution.

Whether the changes observed in solution studies also occur in fibers has been less certain. No changes have been observed in X-ray diffraction patterns (Rodger & Tregear, 1984), EM (Katayama, 1989), birefringence (Oborah & Irving, 1989), or linear dichroism of a probe attached to SH2 (Ajtai & Burghardt, 1989). Our own and others EPR studies using the maleimide and iodoacetamide derivatives of nitroxides attached to SH1 indicated a small torsional reorientation of the Cys-707 domain, putting an upper bound of 5–10° on the rotation of the myosin head as a rigid body (Fajer *et al.*, 1990; Ajtai *et al.*, 1992). However optical probes, especially IATR, did show reorientation of the probe axis on ADP binding (Borejdo *et al.*, 1982; Burghardt *et al.*, 1983). The lack of time correlation of tension change and spectroscopic change (Tanner *et al.*, 1992) and our recent EPR modeling studies, which excluded the possibility that the spin probe orientation might make EPR insensitive to head rotation (Fajer 1994a,b), both indicate that the optical probe movement is associated with an internal structural change rather than head rotation as a rigid body. The present study confirms this assertion since (a) two orientationally resolved conformations were observed; (b) one of them displayed a large orientational distribution; (c) the onset of microsecond fluctuations was observed. Any one of these phenomena would account for a change in the time- and spatially-averaged orientation of the optical probe. Why different labels attached to the same site are behaving differently is unclear (Fajer, 1994a). The size of the linkage arm might play a role; in the case of MSL, the labeled cysteine is separated by a succinamide ring from the reporting

nitroxide, whereas in IKSL the nitroxide is separated by three C—C bonds. However, IASL is similar in linkage structure to IKSL (Figure 1), and it also does not report any significant changes in ADP (Fajer *et al.*, 1990; Fajer, 1994a). This question might be answered when the atomic structure of the myosin head is made available, and steric restraints on different labels can be established.

In addition to identifying the nature of the structural change induced by ADP, we have established the coupling of this change to the interaction with actin. The myosin bound to glass beads did not reveal increased domain mobility in ADP as was found in the fibers. Apart from a possible change in the structure induced by binding to the glass beads, the difference between fibers and beads is the strong binding to actin. The flow experiment of Figure 3 indicates that the heads remain strongly bound in fibers in the presence of ADP.

The presence of two resolved populations of the probe signifies that the myosin head does not have a singular structure even in a single chemical state, which cautions against simplistic interpretation of static structures as derived from EM and crystallography. Conformational heterogeneity of the myosin head was first noted by Shriver and Sykes (1981a,b, 1982), who observed two resolved NMR resonances of the β -phosphate of ADP and two limiting values of chemical shift of a fluorine label attached to SH1. It was suggested that the equilibrium between these two different populations of the myosin head was determined by the chemical state of the nucleotide, and the transition between the two forms was responsible for force generation (Shriver & Sykes, 1981a). It is not clear whether the two orientational populations observed here correspond to the conformations observed in NMR because no change in mobility was observed in isolated myosin heads and no NMR was done in fibers. We would argue that the structural changes observed here correspond to optical probe changes which are not related to force generation (Tanner *et al.*, 1992) but which assist the release of ATP hydrolysis products.

Internal Mobility. The observed structural changes indicate that the head has a considerable flexibility, manifesting itself in the large orientational spread and the presence of microsecond mobility. The labeled-domain "breathing" motions are slower than the mobility of the detached myosin heads ($\tau_r = 100$ –200 and 5–10 μ s, respectively). An alternative explanation to structural fluctuations is that the pocket around Cys-707 created by ADP allows the label to orient itself at many different angles with respect to the protein. This possibility is unlikely because the accompanying mobility is 4–5 orders of magnitude slower than the rotation about the C—S bond of cysteine. Such a rotation is predicted to be in the range of 0.05–0.1 ns (Oton *et al.*, 1981). Because there is little steric hindrance (wide spread of orientations in the disordered component), there should be equally little restriction on the rate of reorientation.

Proteins have been described previously as dynamic systems (Linderstrom-Lang & Shellman, 1959; Weber, 1975) with many structural substates (Hong *et al.*, 1990; Frauenfelder *et al.*, 1991). For example, CO dipoles in carbon-monoxymyoglobin can have three different angles with respect to the heme plane. The transitions between these substates depend on the energy landscape of the conformational surface. Substates separated by low activation energies will exhibit fast transitions, slowing down with increasing

activation energies. Thus, the internal protein dynamics might exhibit a full range of time scales. The side chain motions might be on the picosecond time scale (Karplus & Petsko, 1990), but correlated motions involving large segments of protein structure are considerably slower in the nanosecond to millisecond regime (Frauenfelder *et al.*, 1991). Most domain motions are in the nanosecond range, e.g., in hemoglobin (Oton *et al.*, 1981), troponin C (Wang *et al.*, 1993; Potter *et al.*, 1975; Li & Fajer, 1994), tropomyosin (Graceffa & Lehrer, 1984; Szczesna and Fajer, private communication), or gramicidin A (North & Cross, 1993), but slower motions are also observed, e.g., disulfide bond isomerization; *cis-trans* isomerization of proline peptide bonds and aromatic ring flips are on the millisecond time scale (Otting *et al.*, 1993; Wutrich, 1976). The motions responsible for hydrogen exchange are thought to be even slower (Englander & Mauel, 1972). Internal motions in myosin have been described previously, but all of these were in the nanosecond region (Seidel *et al.*, 1970; Barnett & Thomas, 1987), and then are inhibited by the interaction with actin (Highsmith *et al.*, 1979). The internal motions observed here are qualitatively different, they are considerably slower (correlation times in the microsecond region), and they are only observed in the presence of actin.

These motions and the presence of conformational substates most certainly have functional implications. The "induced fit" model proposed by D. Koshland (1958) explicitly treated protein conformations and transitions between them as being at the core of catalytic activity. More recently, Frauenfelder and colleagues have shown that the conformational substates of myoglobin display different binding kinetics (Hong *et al.*, 1990). The oxygen affinity of hemoglobin has been correlated with the conformations and internal flexibility of the hemoglobin subunits (el Antri *et al.*, 1990). We can only speculate that the microsecond structural fluctuations observed here are of importance to enzymatic activity. ADP release, which is thought to limit the actomyosin ATPase in the fibers, has been shown to be 10 times slower than the rate of structural fluctuations. It is likely that the structural changes observed here assist ADP release (Siemankowski & White, 1984).

In conclusion, we have observed slow ($\tau_r = 150 \mu$ s) internal motions of the catalytic domain of the myosin head bound to actin, which are present in ADP, but absent in the rigor state. These motions result in a wide range of local conformations, as probed by the attached spin label.

ACKNOWLEDGMENT

We thank Elizabeth Fajer for help with the manuscript.

REFERENCES

- Ajtai, K., & Burghardt, T. P. (1989) *Biochemistry* 28, 2204–10.
- Ajtai, K., Ringler, A., & Burghardt, T. P. (1992) *Biochemistry* 31, 207–17.
- Barnett, V. A., & Thomas, D. D. (1984) *J. Mol. Biol.* 179, 83–102.
- Barnett, V. A., & Thomas, D. D. (1987) *Biochemistry* 26, 314–23.
- Borejdo, J., Assulin, O., Ando, T., & Putnam, S. (1982) *J. Mol. Biol.* 158, 391–414.
- Burghardt, T. P., Ando, T., & Borejdo, J. (1983) *Proc. Natl. Acad. Sci. U.S.A.* 80, 7515–9.

- Dalbey, R. E., Weiel, J., & Yount, R. G. (1983) *Biochemistry* 22, 4696–706.
- el Antri, S., Sire, O., & Alpert, B. (1990) *Eur. J. Biochem.* 191, 163–8.
- Englander, W. S., & Mael, C. (1972) *J. Biol. Chem.* 247, 2387–94.
- Fajer, P. G. (1994a) *Biophys. J.* 66, 2039–50.
- Fajer, P. G. (1994b) *Proc. Natl. Acad. Sci. U.S.A.* 91, 937–41.
- Fajer, P. G., Fajer, E. A., Brunsvold, N. J., & Thomas, D. D. (1988) *Biophys. J.* 53, 513–24.
- Fajer, P. G., Fajer, E. A., Matta, J. J., & Thomas, D. D. (1990) *Biochemistry* 29, 5865–71.
- Fajer, P. G., Fajer, E. A., Schoenberg, M., & Thomas, D. D. (1991) *Biophys. J.* 60, 642–9.
- Frauenfelder, H., Sligar, S. G., & Wolynes, P. G. (1991) *Science* 254, 1598–603.
- Graceffa, P., & Lehrer, S. S. (1994) *Biochemistry* 23, 2606–12.
- Hambly, B., & Cooke, R. (1990) *Biophys. J.* 57, 339a.
- Hambly, B., Franks, K., & Cooke, R. (1991) *Biophys. J.* 59, 127–38.
- Hideg, K., & Hankovský, O. H. (1989) *Biological magnetic resonance* (Berliner, L. J., & Reuben, J., Eds.) Vol. 8, pp 427–488, Plenum, New York.
- Highsmith, S., & Eden, D. (1990) *Biochemistry* 29, 4087–93.
- Highsmith, S., Akasaka, K., Konrad, M., Goody, R., Holmes, K., Wade-Jardetzky, N., & Jardetzky, O. (1979) *Biochemistry* 18 (19), 4238–44.
- Holmes, K. C., & Kabsch, W. (1991) *Curr. Opin. Struct. Biol.* 1, 270–80.
- Holmes, K. C., Popp, D., Gebherd, W., & Kabsch, W. (1990) *Nature* 347, 44–9.
- Hong, M. K., Braunstein, D., Cowen, B. R., Frauenfelder, H., Iben, I. E., Mourant, J. R., Ormos, P., Scholl, R., Schulte, A., Steinbach, P. I. & et al. (1990) *Biophys. J.* 58, 429–36.
- Huxley, A. F. (1974) *J. Physiol.* 243, 1–43.
- Huxley, H. E. (1969) *Science* 164, 1356–65.
- Ishiwata, S. B., Munuck, B. A., Seidel, J. C., & Gergely, J. (1986) *Biophys. J.* 49, 821–8.
- Kabsch, W., Mannherz, H. G., Suck, D., Pai, E. F., & Holmes, K. C. (1990) *Nature* 347, 37–44.
- Karplus, M., & Petsko, G. A. (1990) *Nature* 347, 631–9.
- Katayama, E. (1989) *J. Biochem. (Tokyo)* 106, 751–70.
- Li, H. & Fajer, P. G. (1994) *Biochemistry* (in press).
- Linderstrom-Lang, K. U., & Schellman, J. A. (1959) *Enzyme* 1, 443.
- Lu, R. C., & Wong, A. (1989) *Biochemistry* 28, 4826–9.
- Machleidt, W., & Wachter, E. (1977) *Methods Enzymol.* 47, 263–77.
- North, C., & Cross, T. (1993) *J. Magn. Reson. Ser. B* 101, 35–43.
- Oborah, O., & Irving, M. (1989) *Biophys. J.* 55, 9a.
- Ohnishi, S., Maruyama, K., & Ebashi, S. (1975) *J. Biochem.* 78, 73–81.
- Oton, J., Bucci, E., Steiner, R. F., Fronticelli, C., Franchi, D., Montemarano, J., & Martinez, A. (1981) *J. Biol. Chem.* 256, 7248–56.
- Otting, G., Liepinsh, E., & Wüthrich, K. (1993) *Biochemistry* 32, 3571–82.
- Pate, E., & Cooke, R. (1988) *Biophys. J.* 53, 561–73.
- Potter, J. D., Seidel, J. C., Leavis, P., Lehrer, S. S., & Gergely, J. (1976) *J. Biol. Chem.* 251, 7551–6.
- Raucher, D., & Fajer, P. G. (1994) *Biochemistry* 33, 11993–11999.
- Raucher, D., Fajer, E. A., Hideg, K., Sar, C., Kawai, M., Zhao, Y., & Fajer, P. G. (1994) *Biophys. J.* (in press).
- Rayment, I., Holden, H. M., Whittaker, M., Yohn, C. B., Lorenz, M., Holmes, K. C., & Milligan, R. A. (1993a) *Science* 261, 58–65.
- Rayment, I., Rypniewski, W. R., Schmidt, B. K., Smith, R., Tomchick, D. R., Benning, M. M., Winkelmann, D. A., Wesenberg, G., & Holden, H. M. (1993b) *Science* 261, 50–8.
- Reisler, E., & Applegate, D. (1984) *Biochemistry* 23, 4 779.
- Reisler, E., Burke, M., Himmelfard, S., & Harrington, W. F. (1974) *Biochemistry* 13, 3837.
- Rodger, C. D., & Tregear, R. T. (1974) *J. Mol. Biol.* 86, 495–7.
- Schaub, M. C., Watterson, J. G., & Waser, P. G. (1975) *Hoppe-Seyler's Z. Physiol. Chem.* 356, 325–39.
- Seidel, J. C., Chopek, M., & Gergely, J. (1970) *Biochemistry* 9, 3265–72.
- Shriver, J. W., & Sykes, B. D. (1981a) *Biochemistry* 20, 2004–12.
- Shriver, J. W., & Sykes, B. D. (1981b) *Biochemistry* 20, 6357–62.
- Shriver, J. W., & Sykes, B. D. (1982) *Biochemistry* 21, 3022–8.
- Siemankowski, R. F., & White, H. D. (1984) *J. Biol. Chem.* 259, 5045–53.
- Siemankowski, R. F., Wiseman, M. O., & White, H. D. (1985) *Proc. Natl. Acad. Sci. U.S.A.* 82, 658–62.
- Squier, T. C., & Thomas, D. D. (1986) *Biophys. J.* 46, 921–35.
- Tanner, J. W., Thomas, D. D., & Goldman, Y. E. (1992) *J. Mol. Biol.* 223, 185–203.
- Thomas, D. D., & Cooke, R. (1980) *Biophys. J.* 32, 891–906.
- Thomas, D. D., Dalton, L. R., & Hyde, J. S. (1976) *J. Chem. Phys.* 65, 3006–24.
- Thomas, D. D., Ishiwata, S., Seidel, J. C., & Gergely, J. (1980) *Biophys. J.* 32, 873–89.
- Wakabayashi, K., Tokunaga, M., Kohno, I., Sugimoto, Y., Hamanaka, T., Takezawa, Y., Wakabayashi, T., & Amemiya, Y. (1992) *Science* 258, 443–7.
- Wang, C. L., & Gergely, J. (1986) *Eur. J. Biochem.* 154, 225–8.
- Wang, C. K., Liao, R., & Cheung, H. C. (1992) *Biochim. Biophys. Acta* 1121, 16–22.
- Weber, G. (1975) *Adv. Protein Chem.* 29, 1.
- Weeds, A. G., & Taylor, R. S. (1975) *Nature (London)* 257, 54–6.
- Wells, J. A., & Yount, R. G. (1979) *Proc. Natl. Acad. Sci. U.S.A.* 76, 4966–70.
- Wüthrich, K. (1986) *NMR of Proteins and Nucleic Acids*, Wiley, New York.