

Corrections

Role of the Linker Region Connecting the Reductase and Heme Domains in Cytochrome P450_{BM-3}, by Shanthi Govindaraj and Thomas L. Poulos*, Volume 34, Number 35, September 5, 1995, pages 11221–11226.

Page 11223. In Table 1, the heme reduction rates were incorrectly calculated. They should read as follows: for wild type, $1742 \pm 19.4 \text{ min}^{-1}$; for Ala₃, $1100 \pm 18.3 \text{ min}^{-1}$ (63%); for Δ_3 , $681 \pm 11.5 \text{ min}^{-1}$ (39%); for Δ_6 , $153 \pm 2.4 \text{ min}^{-1}$ (8.7%).

BI963447M

Roles of Surface Hydrophobic Residues in the Interfacial Catalysis of Bovine Pancreatic Phospholipase A₂, by B.-I. Lee, E. T. Yoon, and W. Cho*, Volume 35, Number 13, April 2, 1996, pages 4231–4240.

Page 4233. Equation 1 is misprinted. The equation should read as follows:

$$f = (1/2[E]_0) \{ [E]_0 + K_d + [PL]_0/n - [(E]_0 + K_d + [PL]_0/n)^2 - 4[E]_0[PL]_0/n]^{1/2} \} \quad (1)$$

BI965005G