

# Small Molecule Inhibitors of $\alpha$ -Synuclein Filament Assembly<sup>†</sup>

Masami Masuda,<sup>‡,§</sup> Nobuyuki Suzuki,<sup>||</sup> Sayuri Taniguchi,<sup>‡,⊥</sup> Takayuki Oikawa,<sup>‡,§</sup> Takashi Nonaka,<sup>‡</sup> Takeshi Iwatsubo,<sup>⊥</sup> Shin-ichi Hisanaga,<sup>§</sup> Michel Goedert,<sup>@</sup> and Masato Hasegawa<sup>\*,‡</sup>

Department of Molecular Neurobiology, Tokyo Institute of Psychiatry, 2-1-8 Kamikitazawa, Setagaya-ku, Tokyo 156-8585, Japan, Molecular Neuroscience Laboratory, Graduate School of Science, Tokyo Metropolitan University, 1-1 Minami-Osawa, Hachioji-shi, Tokyo 192-0397, Japan, New Product Research Laboratory II, Daiichi Pharmaceutical Company Ltd., 16-13 Kita-Kasai 1-Chome, Edogawa-ku, Tokyo 134-8630, Japan, Department of Neuropathology and Neuroscience, Graduate School of Pharmaceutical Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan, and Medical Research Council Laboratory of Molecular Biology, Hills Road, Cambridge CB2 2QH, U.K.

Received January 12, 2006; Revised Manuscript Received March 17, 2006

**ABSTRACT:**  $\alpha$ -Synuclein is the major component of the filamentous inclusions that constitute defining characteristics of Parkinson's disease and other  $\alpha$ -synucleinopathies. Here we have tested 79 compounds belonging to 12 different chemical classes for their ability to inhibit the assembly of  $\alpha$ -synuclein into filaments in vitro. Several polyphenols, phenothiazines, porphyrins, polyene macrolides, and Congo red and its derivatives, BSB and FSB, inhibited  $\alpha$ -synuclein filament assembly with IC<sub>50</sub> values in the low micromolar range. Many compounds that inhibited  $\alpha$ -synuclein assembly were also found to inhibit the formation of A $\beta$  and tau filaments. Biochemical analysis revealed the formation of soluble oligomeric  $\alpha$ -synuclein in the presence of inhibitory compounds, suggesting that this may be the mechanism by which filament formation is inhibited. Unlike  $\alpha$ -synuclein filaments and protofibrils, these soluble oligomeric species did not reduce the viability of SH-SY5Y cells. These findings suggest that the soluble oligomers formed in the presence of inhibitory compounds may not be toxic to nerve cells and that these compounds may therefore have therapeutic potential for  $\alpha$ -synucleinopathies and other brain amyloidoses.

Filamentous inclusions made of the protein  $\alpha$ -synuclein in nerve cells or glial cells are the defining neuropathological feature of a group of neurodegenerative diseases which include Parkinson's disease (PD),<sup>1</sup> dementia with Lewy bodies (DLB), and multiple-system atrophy (MSA) (1). In these so-called " $\alpha$ -synucleinopathies",  $\alpha$ -synuclein is deposited in a hyperphosphorylated form (2–8). Missense mutations (A30P, E46K, and A53T) in the  $\alpha$ -synuclein gene cause familial forms of PD and DLB (9–11). Furthermore, multiplications (duplication and triplication) of a region on

the long arm of chromosome 4 that encompasses the  $\alpha$ -synuclein gene cause an inherited form of PD dementia (12–14), indicating that the simple overproduction of wild-type  $\alpha$ -synuclein is sufficient to cause PD dementia.

$\alpha$ -Synuclein is a 140-amino acid protein of unknown function that is abundantly expressed in brain, where it is concentrated in presynaptic nerve terminals (15, 16). The amino-terminal region of  $\alpha$ -synuclein (residues 7–87) consists of seven imperfect repeats, each 11 amino acids in length, with the consensus sequence KTKEGV. The repeats are continuous, except for a four-amino acid stretch between repeats 4 and 5, and partially overlap with a hydrophobic region (amino acids 61–95). The carboxy-terminal region (amino acids 96–140) is negatively charged.  $\alpha$ -Synuclein is a natively unfolded protein with little ordered secondary structure that binds to lipid membranes through its amino-terminal repeats, indicating that it may be a physiological lipid-binding protein (17–21). Upon binding to lipid membranes,  $\alpha$ -synuclein adopts structures rich in  $\alpha$ -helical character (18, 20–25).

Recombinant  $\alpha$ -synuclein readily assembles into filaments that share many of the morphological and ultrastructural characteristics of the filaments present in human brain (26–29). Assembly is a nucleation-dependent process and occurs through sequences located in the 100 amino-terminal amino acids of  $\alpha$ -synuclein (30, 31). The carboxy-terminal region, in contrast, inhibits assembly (26, 29, 32). Although  $\alpha$ -synuclein is largely unstructured, its negatively charged C-terminus makes long-range contacts with the central

<sup>†</sup> This work was supported by Grants-in-Aid for Scientific Research on Priority Areas (Research on Pathomechanisms of Brain Disorders) from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to T.I., S.H., and M.H.).

\* To whom correspondence should be addressed: Department of Molecular Neurobiology, Tokyo Institute of Psychiatry, 2-1-8 Kamikitazawa, Setagaya-ku, Tokyo 156-8585, Japan. Telephone: +81 3 3304 5701. Fax: +81 3 3329 8035. E-mail: masato@prit.go.jp.

<sup>‡</sup> Tokyo Institute of Psychiatry.

<sup>§</sup> Tokyo Metropolitan University.

<sup>||</sup> Daiichi Pharmaceutical Co. Ltd.

<sup>⊥</sup> The University of Tokyo.

<sup>@</sup> Medical Research Council Laboratory of Molecular Biology.

<sup>1</sup> Abbreviations: PD, Parkinson's disease; DLB, dementia with Lewy bodies; MSA, multiple-system atrophy; ThS, thioflavin S; HPLC, high-performance liquid chromatography; DMSO, dimethyl sulfoxide; PBS, phosphate-buffered saline; ThT, thioflavin T; DAPH, 4,5-dianilino-phthalimide; BSB, 1-bromo-2,5-bis(3-carboxystyryl)benzene; FSB, 1-fluoro-2,5-bis(3-hydroxycarbonyl-4-hydroxystyryl)benzene; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; PET, positron emission tomography; MRI, magnetic resonance imaging; FTDP-17, frontotemporal dementia and parkinsonism linked to chromosome 17.

hydrophobic region (33, 34). Perturbation of these interactions is likely to expose hydrophobic surfaces, facilitating filament formation.

Mutations E46K and A53T in  $\alpha$ -synuclein have been found to accelerate the rate of filament assembly (28, 35, 36). The A30P mutation has been reported to increase the total rate of aggregation of  $\alpha$ -synuclein (37–39) but to slow the rate of formation of mature filaments (40). X-ray fiber diffraction and electron diffraction analyses have shown that the transition from natively unfolded to cross- $\beta$  structure underlies the assembly of  $\alpha$ -synuclein into filaments (29). It is therefore appropriate to consider the  $\alpha$ -synucleinopathies a form of brain amyloidosis.

The conversion of a small number of soluble peptides and proteins into insoluble filaments is believed to be the central event in the pathogenesis of the most common neurodegenerative diseases (41, 42). Consequently, many current therapeutic strategies are aimed at inhibiting filament formation and at promoting filament clearance. They include the use of antibodies, synthetic peptides, molecular chaperones, and chemical compounds. Among these, small organic molecules have been extensively tested for their ability to inhibit filament formation in vitro, particularly in relation to A $\beta$  deposition (43), the formation of protease-resistant forms of the prion protein (44), the aggregation of huntingtin (45), and, more recently, the heparin-induced formation of tau filaments (46, 47). Less is known about small organic molecules that inhibit the formation of  $\alpha$ -synuclein filaments. Several studies have described inhibition of filament formation by catecholamines (48–50), with other studies reporting inhibition by the polyphenol baicalein (51), the porphyrin phthalocyanine (52), and the anti-tuberculosis drug rifampicin (53).

Here we have investigated the effects of 79 compounds belonging to 12 different chemical classes on  $\alpha$ -synuclein filament formation. The ability of these compounds to inhibit the assembly of A $\beta$  and tau protein was investigated in parallel. Filament formation was assessed by electron microscopy, thioflavin S (ThS) fluorescence, and the formation of sarkosyl-insoluble  $\alpha$ -synuclein. Compounds belonging to seven chemical classes (polyphenols, porphyrins, phenothiazines, polyene macrolides, rifamycins, Congo red and derivatives, and terpenoids) inhibited  $\alpha$ -synuclein filament formation. By SDS–PAGE, inhibition of assembly was reflected in a large reduction in the level of sarkosyl-insoluble  $\alpha$ -synuclein and the accumulation of soluble, SDS-resistant dimers and oligomers. Unlike  $\alpha$ -synuclein filaments and protofibrils, these soluble species did not reduce the viability of human dopaminergic neuroblastoma SH-SY5Y cells.

## EXPERIMENTAL PROCEDURES

**Expression and Purification of  $\alpha$ -Synuclein Proteins.** Human  $\alpha$ -synuclein(1–140) and  $\alpha$ -synuclein(1–120) were expressed in *Escherichia coli* BL21(DE3) cells and purified using boiling, Q-Sepharose (for 1–140) or SP-Sepharose (for 1–120) ion exchange chromatography, and ammonium sulfate precipitation, as described previously (54). During the course of the analysis of dimeric  $\alpha$ -synuclein, we noticed that a cysteine residue was incorporated at position 136 instead of a tyrosine in ~20% of bacterially expressed human  $\alpha$ -synuclein and that mutagenesis of codon 136 (TAC to

TAT) resulted in the expression of  $\alpha$ -synuclein lacking cysteine (55). Therefore, the Y136-TAT construct was used in all experiments.  $\alpha$ -Synuclein proteins were dialyzed against 30 mM Tris-HCl (pH 7.5) and cleared by centrifugation at 113000g for 20 min. Following separation by reverse phase high-performance liquid chromatography (HPLC) (Aquapore RP300 column), the absorbance at 215 nm was measured and compared with that of  $\alpha$ -synuclein of known concentration, to give the concentration of the freshly purified proteins. For immunoblotting, aliquots of reaction mixtures were separated by SDS–PAGE, blotted onto a PVDF membrane, blocked with 3% gelatin/PBS, and incubated overnight at room temperature with antibody Syn102 (1:1000 dilution), which recognizes residues 131–140 of human  $\alpha$ -synuclein, in a 10% fetal calf serum/PBS solution. Following washing, the blots were incubated for 2 h at room temperature with biotinylated secondary antibody (1:500) (Vector Laboratories). Following further washing, the blots were incubated with peroxidase-labeled avidin–biotin (Vector Laboratories) for 30 min at room temperature and developed with NiCl-enhanced diaminobenzidine (Sigma).

**$\alpha$ -Synuclein Filament Assembly and Inhibitor Testing.** The compounds listed in Table 1 were dissolved in dimethyl sulfoxide (DMSO) and kept as 20 mM stock solutions. Purified recombinant human  $\alpha$ -synuclein(1–140) and  $\alpha$ -synuclein(1–120) (2 mg/mL) were incubated with shaking (200 rpm) at 37 °C for 72 h in 50  $\mu$ L of 30 mM Tris-HCl (pH 7.5) containing 0.02% sodium azide, in the presence or absence of 200  $\mu$ M compound. For a quantitative assessment of filament formation, the amount of sarkosyl-insoluble  $\alpha$ -synuclein and the level of ThS fluorescence were measured. Sarkosyl-soluble and sarkosyl-insoluble  $\alpha$ -synuclein were prepared as follows. Aliquots (10  $\mu$ L) of assembly mixtures were removed and added to 40  $\mu$ L of 30 mM Tris-HCl (pH 7.5) containing 1% sarkosyl. This was followed by a 20 min centrifugation at 453000g. The supernatants (sarkosyl-soluble) were removed and the pellets (sarkosyl-insoluble) resuspended in 50  $\mu$ L of sample buffer, followed by SDS–PAGE. Following staining with Coomassie Brilliant Blue, the intensities of the sarkosyl-insoluble  $\alpha$ -synuclein bands were quantified by scanning densitometry, as described previously (47). The supernatants were stained with the silver stain MS kit (Wako). For ThS fluorescence, aliquots (10  $\mu$ L) of the assembly mixtures were removed and brought to 300  $\mu$ L with 5  $\mu$ M ThS (Sigma-Aldrich) in 20 mM MOPS (pH 6.8). Fluorimetry was performed using a Hitachi F4000 fluorescence spectrophotometer (set at 440 nm for excitation and 521 nm for emission), as described previously (47). The results were expressed as the percentage of  $\alpha$ -synuclein assembly in the absence of compound (taken to be 100%). Statistical analysis was carried out with an unpaired *t* test using Kai plot software, and the results were expressed as the means  $\pm$  the standard deviation. IC<sub>50</sub> values were calculated for each compound by quantifying the amounts of sarkosyl-insoluble  $\alpha$ -synuclein. For a semiquantitative assessment of filament formation, electron microscopy was used. Aliquots of assembly mixtures were placed on collodion-coated 300-mesh copper grids, stained with 2% phosphotungstate, and micrographs were recorded at a nominal magnification of 10000 $\times$  on a JEOL 1200EX electron microscope.

Table 1: Inhibitors of A $\beta$ ,  $\alpha$ -Synuclein, and Tau Filament Formation

IC <sub>50</sub> (μM)					IC <sub>50</sub> (μM)				
no.	compound	Aβ	α-synuclein	tau	no.	compound	Aβ	α-synuclein	tau
<b>polyphenols</b>					<b>lignans</b>				
A1	apigenin	>40	>80	>200	D1	magnolol	>40	>80	>200
A2	baicalein	4.5	8.2	2.7	D2	sesamin	>40	>80	>200
A3	(+)-catechin	>40	>80	>200	<b>phenothiazines</b>				
A4	(-)-catechin gallate	5	21.4	>200	E1	acetopromazine maleate salt	>40	>80	>200
A5	chlorogenic acid	>40	>80	>200	E2	azure A	0.4	>80	2.6
A6	curcumin	1.7	>80	>200	E3	azure C	0.2	>80	1.9
A7	cyanidin	4	10.3	33.3	E4	chlorpromazine hydrochloride	>40	>80	>200
A8	daidzein	>40	>80	>200	E5	lacmoid	1.4	14.7	31.2
A9	delphinidin	3	6.5	6.9	E6	methylene blue	2.3	>80	1.9
A10	2,2'-dihydroxybenzophenone	>40	>80	>200	E7	perphenazine	>40	38.7	>200
A11	4,4'-dihydroxybenzophenone	>40	>80	>200	E8	promazine hydrochloride	>40	>80	>200
A12	dopamine chloride	28.6	7.1	>200	E9	propionylpromazine hydrochloride	>40	>80	>200
A13	(-)-epicatechin	>40	29.9	>200	E10	quinacrine	8.4	>80	79.6
A14	(-)-epicatechin 3-gallate	3	14.5	4.5	E11	quinacrine mustard	1.2	>80	3.1
A15	epigallocatechin	7	10.6	8.4	<b>polyene macrolides</b>				
A16	epigallocatechin gallate	2	9.8	9.6	F1	amphotericin B	2.2	27.1	>200
A17	exifone	0.7	2.5	3.3	F2	filipin III	14.6	78.2	>200
A18	(-)-gallo catechin	7	8.9	13.3	<b>porphyrins</b>				
A19	(-)-gallo catechin gallate	1.5	3.6	1	G1	ferric dehydroporphyrin IX	0.2	9.7	1.4
A20	gingerol	25	>80	>200	G2	hematin (from bovine blood)	0.2	15	10.4
A21	gossypetin	1.3	5.6	2	G3	phthalocyanine tetrasulfonate	3.2	27.5	67
A22	hinokiflavone	5	8.1	>200	<b>rifamycin</b>				
A23	hypericin	0.9	7.5	26.8	H1	rifampicin	4.9	46.2	>200
A24	kaempferol	8	>80	>200	<b>steroids</b>				
A25	luteolin	3	28	>200	I1	taurochenodeoxycholic acid	>40	>80	>200
A26	myricetin	0.9	13.3	1.2	I2	taurohydroxycholic acid	>40	>80	>200
A27	naringenin	25	>80	>200	I3	tauroolithocholic acid	>40	>80	>200
A28	2,3,4,2',4'-pentahydroxybenzophenone	2.8	28.3	2.4	I4	tauroolithocholic acid 3-sulfate	>40	>80	>200
A29	procyanidin B1	14	7.3	27.5	I5	tauroursodeoxycholic acid	>40	>80	>200
A30	procyanidin B2	>40	4.3	>200	<b>Congo red and derivatives</b>				
A31	purpurogallin	0.5	12.9	5.6	J1	Congo red	0.9	2.3	2.2
A32	quercetin	5	20	>200	J2	chlorazol black E	0.3	16.4	>200
A33	rosmarinic acid	12	4.8	16.6	J3	BSB	6.4	4	18.2
A34	rutin	32	>80	>200	J4	FSB	1.9	12.4	35.7
A35	(+)-taxifolin	>40	>80	>200	J5	Ponceau SS	1.2	>80	>200
A36	2,2',4,4'-tetrahydroxybenzophenone	>40	>80	>200	<b>terpenoids</b>				
A37	theaflavine	2	5.8	7.9	K1	asiatic acid	>40	45	>200
A38	(+)-α-tocopherol	>40	10.9	107.7	K2	ginkgolide A	>40	>80	>200
A39	2,3,4-trihydroxybenzophenone	3.1	18.6	12.2	K3	ginkgolide B	11	>80	>200
<b>anthracycline</b>					K4	ginkgolide C	>40	>80	>200
B1	daunorubicin hydrochloride	1.4	>80	>200	<b>others</b>				
<b>benzothiazoles</b>					L1	4,5-dianilinophthalimide (DAPH)	2.9	>80	>200
C1	2-(4-aminophenyl)-6-methyl-benzothiazole	2	>80	>200	L2	methyl yellow	1.5	>80	>200
C2	basic blue 41	1.4	>80	>200					
C3	2-[4-(dimethylamino)phenyl]-6-methylbenzothiazole	2	>80	>200					
C4	3,3'-dipropyl thiocarbocyanine iodide (DTCI)	0.3	>80	>200					

*A $\beta$  and Tau Filament Assemblies and Inhibitor Testing.* Monomeric A $\beta$ (1–40) (1 mM, Peptide Institute Inc., Osaka, Japan) in distilled water was diluted with phosphate-buffered saline (PBS, pH 7.5) to a final concentration of 15  $\mu$ M and incubated at 37 °C for 24 h in 40  $\mu$ L of PBS, containing 5  $\mu$ M thioflavin T (ThT), in the presence or absence of compounds (0.3–40  $\mu$ M). Fluorimetry was performed using a Biolumin960 fluorescence spectrophotometer (set at 450 nm for excitation and 485 nm for emission, Amersham Biosciences) (56), and IC<sub>50</sub> values were calculated. For the analysis of oligomeric A $\beta$ , A $\beta$ (1–40) was incubated at 37 °C for 4 days in 40  $\mu$ L of PBS at a concentration of 75  $\mu$ M in the presence or absence of 200  $\mu$ M compound. Aliquots (10  $\mu$ L) of mixtures were removed and added to 40  $\mu$ L of 30 mM Tris-HCl (pH 7.5) containing 1% sarkosyl and the mixtures centrifuged at 25 °C for 20 min at 453000g. The supernatants (sarkosyl-soluble) were removed and the pellets

(sarkosyl-insoluble) resuspended in 50  $\mu$ L of sample buffer, run on a SDS–PAGE gel, and visualized by silver staining. The 412-amino acid isoform of human brain tau (57) was expressed from cDNA clone htau46 and purified, as described previously (47, 58). The effect of inhibitory compounds on the heparin-induced filament formation of tau was measured by quantifying the levels of Sarkosyl-insoluble tau (47).

*Preparation of  $\alpha$ -Synuclein Filaments, Protofibrils, and Soluble Oligomers.*  $\alpha$ -Synuclein filaments were obtained by pelleting the assembly mixtures at 113000g for 20 min. Protofibrils were prepared as described previously (59). Briefly, purified  $\alpha$ -synuclein was dialyzed against 20 mM NH<sub>4</sub>HCO<sub>3</sub>, lyophilized, and resuspended in PBS at 20 mg/mL. Following a 5 min centrifugation at 20000g, the supernatants were loaded onto a Superdex 200 gel filtration column (1 cm  $\times$  30 cm), eluted in 10 mM Tris-HCl (pH






FIGURE 1: Semiquantitative assessment by electron microscopy of  $\alpha$ -synuclein filament formation in the absence (Control) or presence of compounds. Small numbers of filaments or no filaments were seen in the presence of exifone (A17), hypericin (A23), lacmoid (E5), amphotericin B (F1), hematin (G2), and chlorazol black E (J2). Filament numbers similar to controls were observed in the presence of 3,3'-dipropylthiocarbocyanine iodide (DTCI) (C4). A typical experiment is shown. Similar results were obtained in three separate experiments.

7.5) containing 150 mM NaCl, and monitored at 215 nm. The material eluting in the void volume was defined as protofibrils. To prepare soluble oligomers, purified  $\alpha$ -synuclein (9 mg/mL) was incubated with 2 mM inhibitory compound for 30 days at 37 °C in 30 mM Tris-HCl containing 0.1% sodium azide. The samples were then centrifuged for 20 min at 113000g and the supernatants run on a Sephadex G-25 column, to separate oligomers from unbound inhibitor (52). The eluates were then fractionated on a Superdex 200 gel filtration column, as described above. Protein concentrations were determined using HPLC and the BCA protein assay kit (Pierce).

**Cytotoxicity Assay.** A solution (100  $\mu$ L) containing 10 000 SH-SY5Y cells in DMEM-F12 with 10% fetal calf serum was added to each well of a 96-well microtiter plate and left for 24 h in a humidified incubator at 37 °C and 5% CO<sub>2</sub>. The medium was then replaced with 100  $\mu$ L of serum-free medium, followed by the addition of the monomeric, soluble oligomeric, protofibrillar, and filamentous  $\alpha$ -synuclein. Following a 6 h incubation, the cytotoxic effect of  $\alpha$ -synuclein was assessed by measuring cellular redox activity with 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), following the manufacturer's (Sigma) instructions.

## RESULTS

**Inhibition of  $\alpha$ -Synuclein Filament Assembly.** Seventy-nine compounds belonging to 12 different chemical classes (listed in Table 1; see the Supporting Information for chemical structures) were tested for their ability to inhibit the assembly of full-length human  $\alpha$ -synuclein into filaments, which was assessed by electron microscopy (Figure 1), thioflavin S fluorescence, and sarkosyl insolubility (Figure 2). By SDS-PAGE, a large reduction in the level of sarkosyl-insoluble  $\alpha$ -synuclein and a corresponding increase in the level of sarkosyl-soluble protein were observed in the presence of several polyphenols (A17, A21, A26, and A31), phenothiazine E5, polyene macrolide F2, porphyrin G2, and Congo red derivative J2 (Figure 3). In the presence of inhibitors, dimeric and oligomeric, SDS-stable  $\alpha$ -synuclein appeared

in the sarkosyl-soluble fraction (Figure 3). Anthracyclines, benzothiazoles, lignans, steroids, 4,5-dianilinophthalimide (DAPI), and methyl yellow failed to inhibit  $\alpha$ -synuclein filament formation, and SDS-stable oligomers were not observed. A representative example of each class of compound is shown in Figure 3.

Filament formation was quantified by measuring the levels of sarkosyl-insoluble  $\alpha$ -synuclein and ThS fluorescence in the presence of each compound. In general, a good correlation was observed between the amounts of ThS fluorescence and sarkosyl-insoluble  $\alpha$ -synuclein. However, for some benzothiazoles (C2 and C4), phenothiazines (E2, E3, and E6), porphyrin G3, and Congo red derivative J5, a large reduction in ThS fluorescence was observed, with only a slight change in the levels of sarkosyl-insoluble  $\alpha$ -synuclein (Figure 2). The results observed by electron microscopy were in complete agreement with those from the sarkosyl experiments (Figure 1).

The IC<sub>50</sub> values of all 79 compounds for inhibiting  $\alpha$ -synuclein filament assembly were determined by quantifying the levels of sarkosyl-insoluble  $\alpha$ -synuclein (Table 1). The effects of the same compounds on the assembly of A $\beta$  and tau were assessed in parallel. For all three proteins, the inhibition of filament formation was concentration-dependent (data not shown). Strong inhibition of  $\alpha$ -synuclein filament assembly (IC<sub>50</sub> values of <10  $\mu$ M) was observed with the polyphenol compounds baicalein (A2), delphinidin (A9), dopamine chloride (A12), epigallocatechin gallate (A16), exifone (A17), (-)-galocatechin (A18), (-)-galocatechin gallate (A19), gossypetin (A21), hinokiflavone (A22), hypericin (A23), procyanidin B1 (A29), procyanidin B2 (A30), rosmarinic acid (A33) and theaflavine (A37), the porphyrin ferric dehydroporphyrin IX (G1), Congo red (J1), and its derivative 1-bromo-2,5-bis(3-carboxystyryl)benzene (BSB) (J3). Vitamin E ( $\alpha$ -tocopherol, A38) also exhibited a strong inhibitory effect, with an IC<sub>50</sub> value of 10.9  $\mu$ M. Other polyphenols (A4, A7, A13–A15, A25, A26, A28, A31, A32, A38, and A39), phenothiazines (E5 and E7), a polyene macrolide (F1), porphyrins (G2 and G3), and Congo red derivatives (J2 and J4) inhibited the assembly of  $\alpha$ -synuclein




FIGURE 2: Quantitation of  $\alpha$ -synuclein filament formation in the absence (Control) or presence of compounds using ThS fluorescence and sarkosyl insolubility. A significant reduction in the amount of sarkosyl-insoluble  $\alpha$ -synuclein (black bars) and ThS fluorescence (gray bars) was observed in the presence of polyphenols (A2, A4, A5, A7, A9, A12, A14–A19, A21–A23, A26, A28–A33, A37, and A39), phenothiazines (E5 and E7), polyene macrolides (F1 and F2), porphyrins (G1 and G2), Congo red and its derivatives (J1–J4), and terpenoid K1. The results are expressed as means  $\pm$  the standard error ( $n = 3$ ; asterisks denote  $p < 0.001$ ). With some compounds (C2, C4, E2, E3, E6, G3, and J5), ThS fluorescence was reduced, but the amount of sarkosyl-insoluble  $\alpha$ -synuclein was not changed relative to controls.

less strongly ( $IC_{50}$  values of 10–40  $\mu$ M). Rifampicin (H1) was weakly inhibitory ( $IC_{50}$  value of 46  $\mu$ M).

The role of the carboxy-terminal region of  $\alpha$ -synuclein was investigated by incubating  $\alpha$ -synuclein(1–120) with the following at 200  $\mu$ M: polyphenols dopamine chloride (A12) and exifone (A17), the phenothiazine lacmoid (E5), the porphyrin hematin (G2), and the Congo red derivative chlorazol black E (J2). All five compounds strongly inhibited the assembly of full-length  $\alpha$ -synuclein; however, unlike exifone and lacmoid, dopamine chloride, hematin, and chlorazol black E did not inhibit filament formation of  $\alpha$ -synuclein(1–120) (see the Supporting Information).

**Inhibition of  $A\beta$  Fibril Formation.** We investigated whether inhibition of  $A\beta$  fibril formation is accompanied by the formation of SDS-stable dimers and oligomers. As shown in Figure 4, both in the absence of compounds and in the presence of noninhibitory compounds, soluble dimer

and oligomer levels were very low. In the presence of compounds that inhibit  $A\beta$  and  $\alpha$ -synuclein assembly, such as some polyphenols (A17, A21, A26, and A33), the phenothiazine lacmoid (E5), and the porphyrin hematin (G2), a strong dimer band and more variable oligomer bands were observed in the sarkosyl-soluble fraction (Figure 4).

**Cytotoxicity Studies.** Protofibrils and soluble oligomeric species of  $\alpha$ -synuclein were prepared by gel filtration chromatography (Figures 5 and 6). Protofibrils were defined as the material eluting in the void volume (fraction 1) of the column, as proposed previously (59). SH-SY5Y cells were exposed for 6 h to monomeric, soluble oligomeric, protofibrillar, or filamentous human  $\alpha$ -synuclein. Cell viability was evaluated using reduction of MTT to MTT formazan and compared with that of vehicle-treated cells. In cells exposed to monomeric  $\alpha$ -synuclein, there was no significant reduction in cell viability. This stood in marked contrast to




FIGURE 3: Formation of soluble  $\alpha$ -synuclein oligomers in the presence of inhibitory compounds. Sarkosyl-insoluble (pellet, top panel) and sarkosyl-soluble (supernatant, bottom panel)  $\alpha$ -synuclein was prepared following incubation of  $\alpha$ -synuclein in the absence (Control) or presence of compounds (A10, A11, A17, A21, A26, A31, B1, C1, D1, E1, E5, F2, G2, I1, J2, and L2). Via SDS-PAGE, SDS-stable, high-molecular weight  $\alpha$ -synuclein (arrow-heads) was detected in the supernatants of samples incubated with inhibitory compounds, but not in the supernatants of controls or samples incubated with noninhibitory compounds. The presence of high-molecular weight  $\alpha$ -synuclein in the supernatant was accompanied by a reduction in the levels of monomeric  $\alpha$ -synuclein in the pellet. A typical experiment is shown. Similar results were obtained in three separate experiments.




FIGURE 4: Formation of soluble  $A\beta$  oligomers in the presence of inhibitory compounds. Sarkosyl-insoluble (pellet, top panel) and sarkosyl-soluble (supernatant, bottom panel)  $A\beta$  was prepared following incubation of  $A\beta$  in the absence (Control) or presence of compounds (A10, A11, A17, A21, A26, A33, D1, E1, E5, E8, G2, and I1). Via SDS-PAGE, SDS-stable, soluble oligomeric  $A\beta$  was detected in the presence of inhibitory compounds, but not in the presence of noninhibitory compounds. A typical experiment is shown. Similar results were obtained in three separate experiments.

$\alpha$ -synuclein filaments, which were cytotoxic. When they were used at 500 nM, approximately 60% of cells were killed, with  $\sim 50\%$  cell death at 50 nM filaments (Figure 5). Protofibrils were also cytotoxic, with 30% cell death at 500 nM and 10% cell loss at 50 nM. The toxicities of filaments and protofibrils were concentration-dependent (Figure 5). Next, we tested the cytotoxicity of soluble oligomeric species obtained following incubation of  $\alpha$ -synuclein in the presence of inhibitory compounds dopamine chloride (A12), exifone (A17), lacmoid (E5), and hematin (G2). In presence of each inhibitory compound, five protein peaks were resolved by gel filtration (Figure 6) and tested individually in the cytotoxicity assay. In contrast to filaments and protofibrils, none of the soluble oligomeric fractions of  $\alpha$ -synuclein that were tested exhibited cytotoxicity in this assay (Figure 6).

## DISCUSSION

Thioflavin S fluorescence, levels of sarkosyl-insoluble  $\alpha$ -synuclein, and electron microscopic examination were used




FIGURE 5: Cytotoxicity of  $\alpha$ -synuclein monomer, protofibrils, and filaments.  $\alpha$ -Synuclein protofibrils and soluble monomer were fractionated by gel filtration chromatography and analyzed by Western blotting (WB) using antibody syn102. The cytotoxicity of  $\alpha$ -synuclein monomers, protofibrils, and filaments at 50 nM (black bars) or 500 nM (white bars) was assessed using the MTT assay. Protofibrils and filaments exhibited dose-dependent cytotoxicity. The results are presented as percent MTT reduction, with the values obtained in the absence of added  $\alpha$ -synuclein taken as 100%. They are expressed as means  $\pm$  the standard error ( $n = 6$ ; one asterisk denotes  $p < 0.01$ , and two asterisks denote  $p < 0.001$ ).

to investigate the effects of 79 compounds belonging to 12 chemical classes on the assembly of  $\alpha$ -synuclein into filaments. Compounds belonging to seven classes (polyphenols, phenothiazines, polyene macrolides, porphyrins, rifamycins, Congo red and its derivatives, and terpenoids) inhibited filament assembly. The three methods used for monitoring filament assembly gave similar results in the presence of inhibitors, except for thioflavin S (a benzothiazole), which could not be used for testing inhibition by some benzothiazoles and phenothiazines. It could also not be used in the presence of Congo red and its derivatives.

A number of polyphenols [baicalein, delphinidin, dopamine chloride, epigallocatechin gallate, exifone, (-)-gallic acid, gossypetin, hinokiflavone, hypericin, procyanidins, rosmarinic acid, and theaflavine], the porphyrin compound ferric dehydroporphyrin IX, and Congo red and its derivative BSB inhibited  $\alpha$ -synuclein filament assembly with  $IC_{50}$  values of  $<10 \mu M$ . Other polyphenols [cyanidin, (-)-epicatechin 3-gallate, epigallocatechin, myricetin, purpurogallin, (+)- $\alpha$ -tocopherol, and 2,3,4-trihydroxybenzophenone], as well as the phenothiazine lacmoid, the porphyrin hematin, and the Congo red derivatives chlorazol black E and 1-fluoro-2,5-bis(3-hydroxycarbonyl-4-hydroxystyryl)benzene (FSB), inhibited filament formation with  $IC_{50}$  values of 10–20  $\mu M$ . The phenothiazine perphenazine, the polyene macrolides amphotericin B and filipin III, the terpenoid asiatic acid, and rifampicin were less potent. Anthracyclines, benzothiazoles, lignans, steroids, DAPH, and methyl yellow were without






FIGURE 6: Soluble  $\alpha$ -synuclein oligomers formed in the presence of inhibitory compounds do not reduce the viability of SH-SY5Y cells.  $\alpha$ -Synuclein soluble oligomers were fractionated by gel filtration chromatography and analyzed by WB using antibody syn102. In the presence of inhibitory compounds A12, A17, E5, and G2, SDS-resistant dimers (fraction 4) and oligomers (fractions 1–3) formed. Fraction 5 corresponds to the  $\alpha$ -synuclein monomer. In the absence of inhibitory compounds, oligomers did not form (not shown). The cytotoxicity of soluble oligomers was assessed using the MTT assay. In contrast to protofibrils and filaments, monomeric  $\alpha$ -synuclein and fractions 1–5 of oligomeric  $\alpha$ -synuclein at 50 nM (black bars) or 500 nM (white bars) were not cytotoxic. The results are presented as the percent MTT reduction, with the values obtained in the absence of added  $\alpha$ -synuclein taken as 100%. They are expressed as means  $\pm$  the standard error ( $n = 6$ ).

effect ( $IC_{50}$  values of  $>80 \mu M$ ). These findings establish that polyphenols constitute a major class of compounds that can inhibit the assembly of  $\alpha$ -synuclein. Many polyphenols are natural substances present in beverages obtained from plants, fruits, and vegetables. Besides inhibiting the aggregation of  $\alpha$ -synuclein,  $A\beta$ , and tau, they are also known to have neuroprotective effects in a number of paradigms (60). Of the 39 compounds that were tested, 26 inhibited  $\alpha$ -synuclein assembly, 29 inhibited  $A\beta$  assembly, and 19 inhibited tau filament formation. In particular, some catechins and vitamin E ( $\alpha$ -tocopherol) were inhibitory. We confirmed and extended recent work showing inhibition of  $\alpha$ -synuclein filament formation by baicalein, dopamine, and a number of catecholamines. Previous work suggested that two adjacent phenolic OH groups may be required for the inhibition of  $\alpha$ -synuclein filament formation via covalent modification (48–51). Of the 27 inhibitory polyphenols identified here, 23 had two adjacent phenolic OH groups. Furthermore, all polyphenols with three OH groups in the ring (a total of 15 compounds) were inhibitory (Supporting Information), suggesting that adjacent phenolic OH groups may indeed play an important role. However, some compounds (hinokiflavone, hypericin, and  $\alpha$ -tocopherol) lacking adjacent phenolic OH groups were also inhibitory. Additional studies are

needed to better understand these structure–activity relationships.

Porphyrins have previously been shown to inhibit the assembly of  $A\beta$  and tau (47). Prior to the work presented here, phthalocyanine tetrasulfonate was the only porphyrin known to inhibit  $\alpha$ -synuclein filament assembly (52). We found a weak inhibitory effect of phthalocyanine tetrasulfonate, with stronger inhibition by the porphyrins ferric dehydroporphyrin IX and hematin. In contrast to a previous report (52), we failed to detect disassembly of  $\alpha$ -synuclein filaments in the presence of phthalocyanine tetrasulfonate. Consistent with previous work (53), rifampicin inhibited  $\alpha$ -synuclein assembly, albeit not very potently. Prior to the study presented here, inhibition of  $\alpha$ -synuclein filament formation by phenothiazines had not been reported. Of the 11 compounds that were tested, lacmoid and perphenazine were inhibitory. As shown previously (47), compounds lacking a side chain at position 9 of the phenothiazine ring were inhibitory toward the assembly of  $A\beta$  and tau. However, this was not true of  $\alpha$ -synuclein aggregation, suggesting a difference in conformation between oligomers and filaments of  $\alpha$ -synuclein and those of  $A\beta$  and tau. Congo red was strongly inhibitory toward the assembly of  $\alpha$ -synuclein and tau. The same was true of  $A\beta$  fibril formation, confirming previous results (61, 62). Moreover, Congo red has also been shown to prevent the aggregation and infectivity of the prion protein (63, 64) and the aggregation of huntingtin (65, 66). Congo red derivatives BSB and FSB also inhibited the assembly of  $A\beta$ ,  $\alpha$ -synuclein, and tau. Unlike Congo red, they cross the blood–brain barrier and are currently being developed as reagents for visualizing  $A\beta$  deposits in vivo by positron emission tomography (PET) and magnetic resonance imaging (MRI) (67–69). BSB has also been shown to label some  $\alpha$ -synuclein deposits in brain sections from cases of MSA and DLB and some tau deposits in cases of Alzheimer's disease, progressive supranuclear palsy, and frontotemporal dementia and parkinsonism linked to chromosome 17 (FTDP-17) (70). Overall, a number of compounds inhibited the assembly of  $A\beta$ ,  $\alpha$ -synuclein, and tau, in line with findings suggesting that soluble amyloid oligomers can have structural features in common (71). Almost all the compounds that inhibited  $\alpha$ -synuclein assembly also inhibited  $A\beta$  assembly. Exceptions were (–)-epicatechin, procyanidin B2,  $\alpha$ -tocopherol, perphenazine, and asiatic acid. Conversely, several compounds inhibited  $A\beta$  assembly without interfering with  $\alpha$ -synuclein and tau filament formation. They included all the benzothiazoles that were tested, several polyphenols (curcumin, gingerol, kaempferol, naringenin, and rutin), the anthracycline daunorubicin hydrochloride, the Congo red derivative Ponceau S, the terpenoid ginkgolide B, and DAPH and methyl yellow.

Previous work has shown that the C-terminal region of  $\alpha$ -synuclein is a negative regulator of filament assembly (26, 29, 32), probably through long-range interactions with part of the hydrophobic region (33, 34). We therefore investigated the ability of dopamine chloride, exifone, lacmoid, hematin, and chlorazol black E to inhibit the assembly of  $\alpha$ -synuclein (1–120). In contrast to that of full-length  $\alpha$ -synuclein, the assembly of  $\alpha$ -synuclein(1–120) was not inhibited by dopamine chloride, hematin, or chlorazol black E. Exifone and lacmoid inhibited but did not abolish filament formation of  $\alpha$ -synuclein(1–120). These findings establish an important




FIGURE 7: Model for the inhibition of amyloid filament formation by small compounds.

role for the C-terminal region of  $\alpha$ -synuclein in the inhibition of filament assembly and suggest that the compounds used may bind to this region. They are in line with a recent study reporting that dopaminochrome inhibited assembly by binding to residues 125–129 of  $\alpha$ -synuclein (49). Consistent with this report, we also found that covalent modification of  $\alpha$ -synuclein was not required to prevent filament formation (not shown). Furthermore, binding of single-chain antibodies to the C-terminal region of  $\alpha$ -synuclein has been shown to inhibit filament assembly (72). It remains to be determined whether the binding of inhibitory compounds and antibodies influences the long-range interactions between the C-terminus of  $\alpha$ -synuclein and the hydrophobic region.

Analysis of the sarkosyl-soluble fraction of  $\alpha$ -synuclein assemblies by SDS–PAGE revealed the presence of dimeric and oligomeric protein in the presence of inhibitory compounds. This was in marked contrast to the sarkosyl-insoluble, SDS-sensitive  $\alpha$ -synuclein which formed in the absence of inhibitors. Dimers and oligomers were also observed upon incubation of A $\beta$  with inhibitory compounds. These findings are reminiscent of what was observed when heparin-induced filament formation of tau protein was inhibited by some of the same compounds (47). It suggests that the compounds investigated here may inhibit filament formation of  $\alpha$ -synuclein by stabilizing soluble, prefibrillar intermediates (Figure 7). The latter, whose precise mechanism of formation remains to be determined, may be intermediates in the pathway leading from monomeric to filamentous  $\alpha$ -synuclein. A recent study (73) has shown that heat shock protein Hsp70 inhibits  $\alpha$ -synuclein filament formation by binding to soluble prefibrillar intermediates, suggesting that small organic molecules and chaperone proteins may inhibit assembly of  $\alpha$ -synuclein through similar mechanisms. Since soluble amyloid oligomers of A $\beta$ ,  $\alpha$ -synuclein, and tau appear to have structural features in common (71), a number of compounds that inhibited the assembly of A $\beta$ ,  $\alpha$ -synuclein, and tau may recognize and interact with this common structure. Other compounds only inhibited A $\beta$  aggregation, indicating that they may recognize a conformation specific to A $\beta$ .

The cytotoxicity of soluble oligomers was investigated in SH-SY5Y cells and compared with that of monomeric  $\alpha$ -synuclein, protofibrils, and filaments. Monomers and

soluble oligomers did not reduce cell viability. By contrast, protofibrils and filaments were markedly cytotoxic. These results indicate that the compounds investigated here inhibited not only filament formation but also the toxicity of  $\alpha$ -synuclein. They demonstrate a link between filament formation and toxicity in this system, in agreement with earlier findings (74). Future experiments will investigate whether the same inhibitors have similar effects in model systems where  $\alpha$ -synuclein filaments are present intracellularly rather than being added from the outside. Currently, it is not clear which  $\alpha$ -synuclein species are the most detrimental to nerve cells and glial cells in Lewy body diseases and MSA, and in animal models thereof (75–77). This notwithstanding, these findings indicate that the soluble oligomeric species formed in the presence of inhibitory compounds may not be toxic to nerve cells and may hence have therapeutic potential.

## ACKNOWLEDGMENT

We thank M. Mikami for assistance with the SDS–PAGE analysis and Drs. A. L. Fink (University of California) and F. Kametani (Tokyo Institute of Psychiatry) for helpful discussions.

## SUPPORTING INFORMATION AVAILABLE

List of compounds tested and quantitation of  $\alpha$ -synuclein (1–120) filament formation in the absence or presence of inhibitory compounds using sarkosyl insolubility. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## REFERENCES

- Goedert, M. (2001)  $\alpha$ -Synuclein and neurodegenerative diseases, *Nat. Rev. Neurosci.* 2, 492–501.
- Spillantini, M. G., Schmidt, M. L., Lee, V. M.-Y., Trojanowski, J. Q., Jakes, R., and Goedert, M. (1997)  $\alpha$ -Synuclein in Lewy bodies, *Nature* 388, 839–840.
- Baba, M., Nakajo, S., Tu, P.-H., Tomita, T., Nakaya, K., Lee, V. M.-Y., Trojanowski, J. Q., and Iwatsubo, T. (1998) Aggregation of  $\alpha$ -synuclein in Lewy bodies of sporadic Parkinson's disease and dementia with Lewy bodies, *Am. J. Pathol.* 152, 879–884.
- Spillantini, M. G., Crowther, R. A., Jakes, R., Hasegawa, M., and Goedert, M. (1998)  $\alpha$ -Synuclein in filamentous inclusions of Lewy



- bodies from Parkinson's disease and dementia with lewy bodies, *Proc. Natl. Acad. Sci. U.S.A.* 95, 6469–6473.
5. Wakabayashi, K., Yoshimoto, M., Tsuji, S., and Takahashi, H. (1998)  $\alpha$ -Synuclein immunoreactivity in glial cytoplasmic inclusions in multiple system atrophy, *Neurosci. Lett.* 249, 180–182.
  6. Spillantini, M. G., Crowther, R. A., Jakes, R., Cairns, N. J., Lantos, P. L., and Goedert, M. (1998) Filamentous  $\alpha$ -synuclein inclusions link multiple system atrophy with Parkinson's disease and dementia with Lewy bodies, *Neurosci. Lett.* 251, 205–208.
  7. Tu, P. H., Galvin, J. E., Baba, M., Giasson, B., Tomita, T., Leight, S., Nakajo, S., Iwatsubo, T., and Trojanowski, J. Q. (1988) Glial cytoplasmic inclusions in white matter oligodendrocytes of multiple system atrophy brains contain insoluble  $\alpha$ -synuclein, *Ann. Neurol.* 44, 415–422.
  8. Fujiwara, H., Hasegawa, M., Dohmae, N., Kawashima, A., Masliah, E., Goldberg, M. S., Shen, J., Takio, K., and Iwatsubo, T. (2002)  $\alpha$ -Synuclein is phosphorylated in synucleinopathy lesions, *Nat. Cell Biol.* 4, 160–164.
  9. Polymeropoulos, M. H., Lavedan, C., Leroy, E., Ide, S. E., Dehejia, A., Dutra, A., Pike, B., Root, H., Rubenstein, J., Boyer, R., Stenroos, E. S., Chandrasekharappa, S., Athanassiadou, A., Papapetropoulos, T., Johnson, W. G., Lazzarini, A. M., Duvoisin, R. C., Di Iorio, G., Golbe, L. I., and Nussbaum, R. L. (1997) Mutation in the  $\alpha$ -synuclein gene identified in families with Parkinson's disease, *Science* 276, 2045–2047.
  10. Krüger, R., Kuhn, W., Müller, T., Woitalla, D., Graeber, M., Kösel, S., Przuntek, H., Eppelen, J. T., Schöls, L., and Riess, O. (1998) Ala30Pro mutation in the gene encoding  $\alpha$ -synuclein in Parkinson's disease, *Nat. Genet.* 18, 106–108.
  11. Zarranz, J. J., Alegre, J., Gómez-Esteban, J. C., Lezcano, E., Ros, R., Ampuero, I., Vidal, L., Hoenicka, J., Rodríguez, O., Atarés, B., Llorens, V., Gomez Tortosa, E., del Ser, T., Munoz, D. G., and de Yébenes, J. G. (2004) The new mutation, E46K, of  $\alpha$ -synuclein causes Parkinson and Lewy body dementia, *Ann. Neurol.* 55, 164–173.
  12. Singleton, A. B., Farrer, M., Johnson, J., Singleton, A., Hague, S., Kachergus, J., Hulihan, M., Peuralinna, T., Dutra, A., Nussbaum, R., Lincoln, S., Crawley, A., Hanson, M., Maraganore, D., Adler, C., Cookson, M. R., Muentner, M., Baptista, M., Miller, D., Blancato, J., Hardy, J., and Gwinn-Hardy, K. (2003)  $\alpha$ -Synuclein locus triplication causes Parkinson's disease, *Science* 302, 841.
  13. Chartier-Harlin, M. C., Kachergus, J., Roumier, C., Mouroux, V., Douay, X., Lincoln, S., Levecque, C., Lavoie, L., Andrieux, J., Hulihan, M., Waucquier, N., Defebvre, L., Amouyel, P., Farrer, M., and Destée, A. (2004)  $\alpha$ -Synuclein locus duplication as a cause of familial Parkinson's disease, *Lancet* 364, 1167–1169.
  14. Ibanez, P., Bonnet, A.-M., Débarges, B., Lohmann, E., Tison, F., Pollak, P., Agid, Y., Dürr, A., and Brice, A. (2004) Causal relation between  $\alpha$ -synuclein gene duplication and familial Parkinson's disease, *Lancet* 364, 1169–1171.
  15. Ueda, K., Fukushima, H., Masliah, E., Xia, Y., Iwai, A., Yoshimoto, M., Otero, D. A., Kondo, J., Ihara, Y., and Saitoh, T. (1993) Molecular cloning of cDNA encoding an unrecognized component of amyloid in Alzheimer disease, *Proc. Natl. Acad. Sci. U.S.A.* 90, 11282–11286.
  16. Jakes, R., Spillantini, M. G., and Goedert, M. (1994) Identification of two distinct synucleins from human brain, *FEBS Lett.* 345, 27–32.
  17. Weinreb, P. H., Zhen, W., Poon, A. W., Conway, K. A., and Lansbury, P. T. (1996) NACP, a protein implicated in Alzheimer's disease and learning, is natively unfolded, *Biochemistry* 35, 13709–13715.
  18. Davidson, W. S., Jonas, A., Clayton, D. F., and George, J. M. (1998) Stabilization of  $\alpha$ -synuclein secondary structure upon binding to synthetic membranes, *J. Biol. Chem.* 273, 9443–9449.
  19. Jensen, P. H., Nielsen, M. H., Jakes, R., Dotti, C. G., and Goedert, M. (1998) Binding of  $\alpha$ -synuclein to brain vesicles is abolished by familial Parkinson's disease mutation, *J. Biol. Chem.* 273, 26292–26294.
  20. Jo, E., McLaurin, J. A., Yip, C. M., St George-Hyslop, P., and Fraser, P. E. (2000)  $\alpha$ -Synuclein membrane interactions and lipid specificity, *J. Biol. Chem.* 275, 34328–34334.
  21. Perrin, R. J., Woods, W. S., Clayton, D. F., and George, J. M. (2000) Interaction of human  $\alpha$ -synuclein and Parkinson's disease variants with phospholipids. Structural analysis using site-directed mutagenesis, *J. Biol. Chem.* 275, 34393–34398.
  22. Eliezer, D., Kutluay, E., Bussell, R., and Browne, G. (2001) Conformational properties of  $\alpha$ -synuclein in its free and lipid-associated states, *J. Mol. Biol.* 307, 1061–1073.
  23. Chandra, S., Chen, X., Rizo, J., Jahn, R., and Südhof, T. (2003) A broken  $\alpha$ -helix in folded  $\alpha$ -synuclein, *J. Biol. Chem.* 278, 15313–15318.
  24. Jao, C. C., Der-Sarkissian, A., Chen, J., and Langen, R. (2004) Structure of membrane-bound  $\alpha$ -synuclein studied by site-directed spin labeling, *Proc. Natl. Acad. Sci. U.S.A.* 101, 8331–8336.
  25. Ulmer, T. S., Bax, A., Cole, N. B., and Nussbaum, R. L. (2005) Structure and dynamics of micelle-bound human  $\alpha$ -synuclein, *J. Biol. Chem.* 280, 9595–9603.
  26. Crowther, R. A., Jakes, R., Spillantini, M. G., and Goedert, M. (1998) Synthetic filaments assembled from C-terminally truncated  $\alpha$ -synuclein, *FEBS Lett.* 436, 309–312.
  27. El-Agnaf, O. M. A., Jakes, R., Curran, M. D., and Wallace, A. (1998) Effects of the mutations Ala30 to Pro and Ala53 to Thr on the physical and morphological properties of  $\alpha$ -synuclein protein implicated in Parkinson's disease, *FEBS Lett.* 440, 67–70.
  28. Conway, K. A., Harper, J. D., and Lansbury, P. T. (1998) Accelerated *in vitro* fibril formation by a mutant  $\alpha$ -synuclein linked to early-onset Parkinson disease, *Nat. Med.* 4, 1318–1320.
  29. Serpell, L. C., Berriman, J., Jakes, R., Goedert, M., and Crowther, R. A. (2000) Fiber diffraction of synthetic  $\alpha$ -synuclein filaments shows amyloid-like cross- $\beta$  conformation, *Proc. Natl. Acad. Sci. U.S.A.* 97, 4897–4902.
  30. Miake, H., Mizusawa, H., Iwatsubo, T., and Hasegawa, M. (2002) Biochemical characterization of the core structure of  $\alpha$ -synuclein filaments, *J. Biol. Chem.* 277, 19213–19219.
  31. Der-Sarkissian, A., Jao, C. J., Chen, J., and Langen, R. (2003) Structural organization of  $\alpha$ -synuclein fibrils studied by site-directed spin labeling, *J. Biol. Chem.* 278, 37530–37535.
  32. Murray, I. V., Giasson, B. I., Quinn, S. M., Koppaka, V., Axelsen, P. H., Ischiropoulos, H., Trojanowski, J. Q., and Lee, V. M.-Y. (2003) Role of  $\alpha$ -synuclein carboxy-terminus on fibril formation *in vitro*, *Biochemistry* 42, 8530–8540.
  33. Dedmon, M. K., Lindorff-Larsen, K., Christodoulou, J., Vendruscolo, M., and Dobson, C. M. (2005) Mapping long-range interactions in  $\alpha$ -synuclein using spin-label NMR and ensemble molecular dynamics simulations, *J. Am. Chem. Soc.* 127, 476–477.
  34. Bertoncini, C. W., Jung, Y.-S., Fernandez, C. O., Hoyer, W., Griesinger, C., Jovin, T. M., and Zweckstetter, M. (2005) Structural characterization of copper(II) binding to  $\alpha$ -synuclein: Insights into the bioinorganic chemistry of Parkinson's disease, *Proc. Natl. Acad. Sci. U.S.A.* 102, 1430–1435.
  35. Choi, W., Zibae, S., Jakes, R., Serpell, L. C., Davletov, B., Crowther, R. A., and Goedert, M. (2004) Mutation E46K increases phospholipid binding and assembly into filaments of human  $\alpha$ -synuclein, *FEBS Lett.* 576, 363–368.
  36. Greenbaum, E. A., Graves, C. L., Mishizen-Eberz, A. J., Lupoli, M. A., Lynch, D. R., Englander, S. W., Axelsen, P. H., and Giasson, B. I. (2005) The E46K mutation in  $\alpha$ -synuclein increases amyloid fibril formation, *J. Biol. Chem.* 280, 7800–7807.
  37. Giasson, B. I., Uryu, K., Trojanowski, J. Q., and Lee, V. M.-Y. (1999) Mutant and wild-type human  $\alpha$ -synucleins assemble into elongated filaments with distinct morphologies *in vitro*, *J. Biol. Chem.* 274, 7619–7622.
  38. Narhi, L., Wood, S. J., Steavenson, S., Jiang, Y., Wu, G. M., Anafi, D., Kaufman, S. A., Martin, F., Sitney, K., Denis, P., Louis, J. C., Wypych, J., Biere, A. L., and Citron, M. (1999) Both familial Parkinson's disease mutations accelerate  $\alpha$ -synuclein aggregation, *J. Biol. Chem.* 274, 9843–9846.
  39. Li, J., Uversky, V. N., and Fink, A. L. (2001) Effect of familial Parkinson's disease point mutations A30P and A53T on the structural properties, aggregation, and fibrillation of human  $\alpha$ -synuclein, *Biochemistry* 40, 11604–11613.
  40. Conway, K. A., Lee, S.-J., Rochet, J. C., Ding, T. T., Williamson, R. E., and Lansbury, P. T. (2000) Acceleration of oligomerization, not fibrillization, is a shared property of both  $\alpha$ -synuclein mutations linked to early-onset Parkinson's disease: Implications for pathogenesis and therapy, *Proc. Natl. Acad. Sci. U.S.A.* 97, 571–576.
  41. Goedert, M., Spillantini, M. G., and Davies, S. W. (1998) Filamentous nerve cell inclusions in neurodegenerative diseases, *Curr. Opin. Neurobiol.* 8, 619–632.
  42. Prusiner, S. B. (2001) Shattuck lecture: Neurodegenerative diseases and prions, *N. Engl. J. Med.* 344, 1516–1526.

43. Findeis, M. A. (2000) Approaches to discovery and characterization of inhibitors of amyloid  $\beta$ -peptide polymerization, *Biochim. Biophys. Acta* 1502, 76–84.
44. Cashman, N. R., and Caughey, B. (2004) Prion diseases: Close to effective therapy? *Nat. Rev. Drug Discovery* 3, 874–884.
45. Heiser, V., Engemann, S., Bröcker, W., Dunkel, I., Boeddrich, A., Waelter, S., Nordhoff, E., Lurz, R., Schugardt, N., Rautenberg, S., Herhaus, C., Barnickel, G., Böttcher, H., Lehrach, H., and Wanker, E. E. (2002) Identification of benzothiazoles as potential polyglutamine aggregation inhibitors of Huntington's disease by using an automated filter retardation assay, *Proc. Natl. Acad. Sci. U.S.A.* 99, 16400–16406.
46. Pickhardt, M., Gazova, Z., von Bergen, M., Khlistunova, I., Wang, Y., Hascher, A., Mandelkow, E. M., Biernat, J., and Mandelkow, E. (2005) Anthraquinones inhibit tau aggregation and dissolve Alzheimer's paired helical filaments *in vitro* and in cells, *J. Biol. Chem.* 280, 3628–3635.
47. Taniguchi, S., Suzuki, N., Masuda, M., Hisanaga, S.-I., Iwatsubo, T., Goedert, M., and Hasegawa, M. (2005) Inhibition of heparin-induced tau filament formation by phenothiazines, polyphenols, and porphyrins, *J. Biol. Chem.* 280, 7614–7623.
48. Conway, K. A., Rochet, J. C., Bieganski, R. M., and Lansbury, P. T. (2001) Kinetic stabilization of the  $\alpha$ -synuclein protofibril by a dopamine- $\alpha$ -synuclein adduct, *Science* 294, 1346–1349.
49. Norris, E. H., Giasson, B. I., Hodara, R., Xu, S., Trojanowski, J. Q., Ischiropoulos, H., and Lee, V. M.-Y. (2005) Reversible inhibition of  $\alpha$ -synuclein fibrillization by dopaminochrome-mediated conformational alterations, *J. Biol. Chem.* 280, 21212–21219.
50. Cappai, R., Leck, S.-L., Tew, D. J., Williamson, N. A., Smith, D. P., Galatis, D., Sharples, R. A., Curtain, C. C., Ali, F. E., Cherny, R. A., Culvenor, J. G., Bottomley, S. P., Masters, C. L., Barnham, K. J., and Hill, A. F. (2005) Dopamine promotes  $\alpha$ -synuclein aggregation into SDS-resistant soluble oligomers via a distinct folding pathway, *FASEB J.* 19, 1377–1379.
51. Zhu, M., Rajamani, S., Kaylor, J., Han, S., Zhou, F., and Fink, A. L. (2004) The flavonoid baicalein inhibits fibrillation of  $\alpha$ -synuclein and disaggregates existing fibrils, *J. Biol. Chem.* 279, 26846–26857.
52. Lee, E.-N., Cho, H.-J., Lee, C.-H., Lee, D., Chung, K. C., and Paik, S. R. (2004) Phthalocyanine tetrasulfonates affect the amyloid formation and cytotoxicity of  $\alpha$ -synuclein, *Biochemistry* 43, 3704–3715.
53. Li, J., Zhu, M., Rajamani, S., Uversky, V. N., and Fink, A. L. (2004) Rifampicin inhibits  $\alpha$ -synuclein fibrillation and disaggregates fibrils, *Chem. Biol.* 11, 1513–1521.
54. Nonaka, T., Iwatsubo, T., and Hasegawa, M. (2005) Ubiquitination of  $\alpha$ -synuclein, *Biochemistry* 44, 361–368.
55. Masuda, M., Dohmae, N., Nonaka, T., Oikawa, T., Hisanaga, S. I., Goedert, M., and Hasegawa, M. (2006) Cysteine misincorporation in bacterially expressed human  $\alpha$ -synuclein, *FEBS Lett.* 580, 1775–1779.
56. Naiki, H., and Nakakuki, K. (1996) First-order kinetic model of Alzheimer's  $\beta$ -amyloid fibril extension *in vitro*, *Lab. Invest.* 74, 374–383.
57. Goedert, M., Spillantini, M. G., Jakes, R., Rutherford, D., and Crowther, R. A. (1989) Multiple isoforms of human microtubule-associated protein tau: sequences and localization in neurofibrillary tangles of Alzheimer's disease, *Neuron* 3, 519–526.
58. Hasegawa, M., Smith, M. J., and Goedert, M. (1998) Tau proteins with FTDP-17 mutations have a reduced ability to promote microtubule assembly, *FEBS Lett.* 437, 207–210.
59. Ding, T. T., Lee, S.-J., Rochet, J. C., and Lansbury, P. T. (2002) Annular  $\alpha$ -synuclein protofibrils are produced when spherical protofibrils are incubated in solution or bound to brain-derived membranes, *Biochemistry* 41, 10209–10217.
60. Mandel, S., and Youdim, B. H. (2004) Catechin polyphenols: Neurodegeneration and neuroprotection in neurodegenerative diseases, *Free Radical Biol. Med.* 37, 304–317.
61. Lorenzo, A., and Yankner, B. A. (1994)  $\beta$ -Amyloid neurotoxicity requires fibril formation and is inhibited by congo red, *Proc. Natl. Acad. Sci. U.S.A.* 91, 12243–12247.
62. Esler, W. P., Stimson, E. R., Ghilardi, J. R., Felix, A. M., Lu, Y.-A., Vinters, H. V., Mantyh, P. W., and Maggio, J. E. (1997) A $\beta$  deposition inhibitor screen using synthetic amyloid, *Nat. Biotechnol.* 15, 258–263.
63. Caughey, B., and Race, R. E. (1992) Potent inhibition of scrapie-associated PrP accumulation by congo red, *J. Neurochem.* 59, 768–771.
64. Caughey, B., Ernest, D., and Race, R. E. (1993) Congo red inhibition of scrapie agent replication, *J. Virol.* 67, 6270–6272.
65. Heiser, V., Scherzinger, E., Boeddrich, A., Nordhoff, E., Lurz, R., Schugardt, N., Lehrach, H., and Wanker, E. E. (2000) Inhibition of huntingtin fibrillogenesis by specific antibodies and small molecules: Implications for Huntington's disease therapy, *Proc. Natl. Acad. Sci. U.S.A.* 97, 6739–6744.
66. Sánchez, I., Mahlke, C., and Yuan, J. (2003) Pivotal role of oligomerization in expanded polyglutamine neurodegenerative disorders, *Nature* 421, 373–379.
67. Skovronsky, D. M., Zhang, B., Kung, M.-P., Kung, H. F., Trojanowski, J. Q., and Lee, V. M.-Y. (2000) *In vivo* detection of amyloid plaques in a mouse model of Alzheimer's disease, *Proc. Natl. Acad. Sci. U.S.A.* 97, 7609–7614.
68. Sato, K., Higuchi, M., Iwata, N., Saito, T. C., and Sasamoto, K. (2004) Fluoro-substituted and  $^{13}\text{C}$ -labeled styrylbenzene derivatives for detecting brain amyloid plaques, *Eur. J. Med. Chem.* 39, 573–578.
69. Higuchi, M., Iwata, N., Matsuba, Y., Sato, K., Sasamoto, K., and Saido, T. C. (2005)  $^{19}\text{F}$  and  $^1\text{H}$  MRI detection of amyloid  $\beta$  plaques *in vivo*, *Nat. Neurosci.* 8, 527–533.
70. Schmidt, M. L., Schuck, T., Sheridan, S., Kung, M.-P., Kung, H., Zhuang, Z.-P., Bergeron, C., Lamarche, J. S., Skovronsky, D., Giasson, B. I., Lee, V. M.-Y., and Trojanowski, J. Q. (2001) The fluorescent Congo red derivative, (trans,trans)-1-bromo-2,5-bis-(3-hydroxycarbonyl-4-hydroxy)styrylbenzene (BSB), labels diverse  $\beta$ -pleated sheet structures in postmortem human neurodegenerative disease brains, *Am. J. Pathol.* 159, 937–943.
71. Kaye, R., Head, E., Thompson, J. L., McIntire, T. M., Milton, S. C., Cotman, C. W., and Glabe, C. G. (2003) Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis, *Science* 300, 486–489.
72. Emadi, S., Liu, R., Yuan, B., Schulz, P., McAllister, C., Lyubchenko, Y., Messer, A., and Sierks, M. R. (2004) Single chain variable fragments against  $\beta$ -amyloid (A $\beta$ ) can inhibit A $\beta$  aggregation and prevent A $\beta$ -induced neurotoxicity, *Biochemistry* 43, 2871–2878.
73. Dedmon, M. W., Christodoulou, J., Wilson, M. R., and Dobson, C. M. (2005) Heat shock protein 70 inhibits  $\alpha$ -synuclein fibril formation via preferential binding to prefibrillar species, *J. Biol. Chem.* 280, 14733–14740.
74. El-Agnaf, O. M. A., Jakes, R., Curran, M. D., Middleton, D., Ingenito, R., Bianchi, E., Pessi, A., Neill, D., and Wallace, D. (1998) Aggregates from mutant and wild-type  $\alpha$ -synuclein proteins and NAC peptide induce apoptotic cell death in human neuroblastoma cells by formation of  $\beta$ -sheet and amyloid-like filaments, *FEBS Lett.* 440, 71–75.
75. Volles, M. J., and Lansbury, P. T. (2003) Zeroing in on the pathogenic form of  $\alpha$ -synuclein and its mechanism of neurotoxicity in Parkinson's disease, *Biochemistry* 42, 7871–7878.
76. Giasson, B. I., and Lee, V. M.-Y. (2003) Are ubiquitination pathways central to Parkinson's disease? *Cell* 114, 1–8.
77. Chen, L., and Feany, M. B. (2005)  $\alpha$ -Synuclein phosphorylation controls neurotoxicity and inclusion formation in a *Drosophila* model of Parkinson disease, *Nat. Neurosci.* 8, 657–663.

BI0600749