

# Thermodynamics of Binding of a Low-Molecular-Weight CD4 Mimetic to HIV-1 gp120<sup>†</sup>

Arne Schön,<sup>‡</sup> Navid Madani,<sup>§</sup> Jeffrey C. Klein,<sup>‡</sup> Amy Hubicki,<sup>§</sup> Danny Ng,<sup>||</sup> Xinzhen Yang,<sup>§</sup> Amos B. Smith, III,<sup>||</sup> Joseph Sodroski,<sup>§</sup> and Ernesto Freire<sup>\*,‡</sup>

*Department of Biology, The Johns Hopkins University, Baltimore, Maryland 21218, Dana-Farber Cancer Institute, Harvard Medical School, Boston, Massachusetts 02115, and Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104*

*Received June 15, 2006; Revised Manuscript Received July 16, 2006*

**ABSTRACT:** NBD-556 and the chemically and structurally similar NBD-557 are two low-molecular weight compounds that reportedly block the interaction between the HIV-1 envelope glycoprotein gp120 and its receptor, CD4. NBD-556 binds to gp120 with a binding affinity of  $2.7 \times 10^5 \text{ M}^{-1}$  ( $K_d = 3.7 \text{ }\mu\text{M}$ ) in a process characterized by a large favorable change in enthalpy partially compensated by a large unfavorable entropy change, a thermodynamic signature similar to that observed for binding of sCD4 to gp120. NBD-556 binding is associated with a large structuring of the gp120 molecule, as also demonstrated by CD spectroscopy. NBD-556, like CD4, activates the binding of gp120 to the HIV-1 coreceptor, CCR5, and to the 17b monoclonal antibody, which recognizes the coreceptor binding site of gp120. NBD-556 stimulates HIV-1 infection of CD4-negative, CCR5-expressing cells. The thermodynamic signature of the binding of NBD-556 to gp120 is very different from that of another viral entry inhibitor, BMS-378806. Whereas NBD-556 binds gp120 with a large favorable enthalpy and compensating unfavorable entropy changes, BMS-378806 does so with a small binding enthalpy change in a mostly entropy-driven process. NBD-556 is a competitive inhibitor of sCD4 and elicits a similar structuring of the coreceptor binding site, whereas BMS-378806 does not compete with sCD4 and does not induce coreceptor binding. These studies demonstrate that low-molecular-weight compounds can induce conformational changes in the HIV-1 gp120 glycoprotein similar to those observed upon CD4 binding, revealing distinct strategies for inhibiting the function of the HIV-1 gp120 envelope glycoprotein. Furthermore, competitive and noncompetitive compounds have characteristic thermodynamic signatures that can be used to guide the design of more potent and effective viral entry inhibitors.

The entry of human immunodeficiency virus (HIV-1)<sup>1</sup> into target cells is mediated by the gp120 exterior envelope glycoprotein and the gp41 transmembrane envelope glycoprotein, which assemble into trimers on the virion surface (1, 2). Upon engaging the receptor, CD4, the gp120 glycoprotein undergoes extensive structural ordering; the resulting

conformation of gp120 can bind the second HIV-1 receptor, CCR5 or CXCR4 (3–7). Receptor binding induces further conformational changes in the HIV-1 envelope glycoproteins that allow the gp41 glycoproteins to mediate the fusion of the viral and cell membranes.

Thermodynamically, the binding of CD4 is characterized by large favorable enthalpy and large unfavorable entropy changes that reflect the structuring of gp120, a protein with large unstructured regions in its unligated form (8). CD4 mimetic miniproteins based on scyllatoxin containing the gp120 binding epitope (9) also elicit conformational changes in gp120 resembling those triggered by CD4 binding and, consequently, are characterized by similar thermodynamic signatures (10, 11).

Recently, two low-molecular-weight compounds that presumably interfere with viral entry of HIV-1 into cells were reported (12) (Figure 1). The studies presented here show that these compounds are competitive inhibitors of CD4 and that they induce conformational changes in gp120 similar to those induced by CD4. These compounds activate coreceptor binding and, in our studies, enhance the entry of HIV-1 into CD4-negative cells expressing CCR5. This behavior is in contrast to that of the potent viral entry inhibitor BMS-378806, which binds gp120 with a small

<sup>†</sup> Supported by National Institutes of Health Grants GM56550 (E.F., J.S., and A.B.S.) and AI41851 (J.S.) and National Science Foundation Grant MCB0131241 (E.F.). N.M. was supported by a NRSA postdoctoral fellowship (F32 NS43260 M) from the National Institutes of Health, and D.N. was supported by a NRSA postdoctoral fellowship (F32 GM072111) from the National Institutes of Health.

\* To whom correspondence should be addressed: Department of Biology, The Johns Hopkins University, Baltimore, MD 21218. Phone: (410) 516-7743. Fax: (410) 516-6469. E-mail: ef@jhu.edu.

<sup>‡</sup> The Johns Hopkins University.

<sup>§</sup> Harvard Medical School.

<sup>||</sup> University of Pennsylvania.

<sup>1</sup> Abbreviations: ITC, isothermal titration calorimetry; CD, circular dichroism; HIV-1, human immunodeficiency virus, type 1; sCD4, soluble form of the human CD4 receptor; Env, envelope glycoprotein; GS, glutamine synthetase; HT, hypoxanthine/thymidine; MSX, methionine sulfoximine; SDS–PAGE, sodium dodecyl sulfate–polyacrylamide electrophoresis; DMSO, dimethyl sulfoxide; EDC, *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride; HOBT, 1-hydroxybenzotriazole; THF, tetrahydrofuran; EtOAc, ethyl acetate; MeOH, methanol; ES<sup>+</sup>, electron ionization; A-MLV, amphotropic murine leukemia virus.




FIGURE 1: General structure of the two compounds, NBD-556 (X = Cl) and NBD-557 (X = Br).

favorable enthalpy change in an entropically driven process (13). BMS-378806 does not compete with CD4 and does not induce any major structural ordering of the gp120 molecule, exerting its antiviral action in an allosteric fashion. In this paper, we demonstrate experimentally that the binding mode and antiviral properties of these compounds are reflected in their binding thermodynamics and that the thermodynamic signature of a compound can be used in the design of more potent and effective viral entry inhibitors.

## MATERIALS AND METHODS

### Synthesis of Compounds

BMS-378806 was prepared according to a published procedure (14). The synthesis of NBD-557 and NBD-556 is outlined in Figure 2. Acylation of either 4-bromo- or 4-chloroaniline provided the corresponding ethyl oxalamate **1**. Hydrolysis of **1** with NaOH to the corresponding acid followed by amide formation with 4-amino-2,2,6,6-tetramethylpiperidine in the presence of EDC and HOBt furnished the desired oxalamides NBD-557 and NBD-556. Tetrahydrofuran (THF) was freshly distilled from sodium/benzophenone under argon. All reagents were purchased from Aldrich and used without further purification. Reaction mixtures were magnetically stirred and reactions carried out under an argon atmosphere. Flash chromatography was performed with silica gel 60 (particle size of 0.040–0.062 mm) supplied by Silicycle. Infrared spectra were recorded on a Jasco model FT/IR-480 Plus spectrometer. Proton and  $^{13}\text{C}$  spectra were recorded on a Bruker AMX-500 spectrometer. Chemical shifts are reported relative to chloroform ( $\delta$  7.26 for  $^1\text{H}$  NMR and  $\delta$  77.0 for  $^{13}\text{C}$  NMR). High-resolution mass spectra were recorded at the University of Pennsylvania Mass Spectrometry Service Center.

*N*-(4-Bromophenyl)-*N'*-(2,2,6,6-tetramethylpiperidin-4-yl)-oxalamide (NBD-557). Ethyl oxalyl chloride (0.20 mL, 1.74 mmol) was added to a round-bottom flask containing anhydrous THF (10 mL), and the resulting solution was cooled in an ice bath. Triethylamine (0.24 mL, 1.74 mmol) and 4-bromoaniline (300 mg, 1.74 mmol) were then added, and the resulting mixture was removed from the ice bath and allowed to come to room temperature. The reaction mixture was stirred for 6 h at room temperature, and the solid was removed by vacuum filtration. The filtrate was concentrated in vacuo, and the residue was redissolved in EtOAc (20 mL). The organic phase was washed with 1 M HCl (2  $\times$  10 mL) and saturated  $\text{NaHCO}_3$  (10 mL) and dried over  $\text{MgSO}_4$ . The solvent was removed under reduced pressure to give the desired ester (408 mg, 88%), which was used in the subsequent reaction without further purification: mp 153–154  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  8.90 (br s, 1H), 7.54 (d,  $J$  = 8.9 Hz, 2H), 7.48 (d,  $J$  = 8.9 Hz, 2H),

4.40 (q,  $J$  = 7.1 Hz, 2H), 1.41 (t,  $J$  = 7.1 Hz, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  160.7, 153.9, 135.4, 132.2, 121.3, 118.3, 63.8, 13.9; IR (thin film) 3334, 3113, 2980, 1729, 1710, 1542, 1491, 1477, 1302, 1176  $\text{cm}^{-1}$ ; high-resolution mass spectrum ( $\text{ES}^+$ )  $m/z$  270.9851 ( $\text{M}^+$ ; calcd for  $\text{C}_{10}\text{H}_{10}\text{BrNO}_3$ , 270.9844).

To a flask containing the above ester (400 mg, 1.47 mmol) were added a 1 M NaOH solution (7.4 mL, 7.40 mmol) and THF (8 mL), and the resulting mixture was stirred at room temperature for 20 h. EtOAc (20 mL) was added, and the organic layer was washed with 1 M NaOH (2  $\times$  10 mL). The combined aqueous layer was acidified to pH 2 with 2 M HCl and extracted with EtOAc (3  $\times$  20 mL). The combined organic layer was dried over  $\text{MgSO}_4$  and concentrated to give the crude acid (272 mg, 76%). The acid was redissolved in anhydrous THF (3 mL) and added to a flask containing 4-amino-2,2,6,6-tetramethylpiperidine (0.21 mL, 1.23 mmol). HOBt (196 mg, 1.45 mmol) and EDC (279 mg, 1.45 mmol) were added, and the resulting mixture was stirred at room temperature for 20 h. The reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (30 mL), and the organic layer was washed with water (4  $\times$  20 mL) and dried over  $\text{MgSO}_4$ . After the solvent was removed in vacuo, the crude product was chromatographed (1:1 EtOAc/MeOH) to give NBD-557 (176 mg, 41%):  $R_f$  = 0.3; mp 204–205  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  9.25 (br s, 1H), 7.53 (d,  $J$  = 9.0 Hz, 2H), 7.49 (d,  $J$  = 9.0 Hz, 2H), 7.29 (br d,  $J$  = 7.9 Hz, 1H), 4.20–4.33 (m, 1H), 1.92 (dd,  $J$  = 12.6, 3.8 Hz, 2H), 1.29 (s, 6H), 1.17 (s, 6H), 1.03–1.13 (m, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  158.8, 157.6, 135.5, 132.2, 121.3, 118.1, 51.0, 44.6, 43.8, 34.9, 28.5; IR (thin film) 3311, 3249, 2959, 2917, 1658, 1511  $\text{cm}^{-1}$ ; high-resolution mass spectrum ( $\text{ES}^+$ )  $m/z$  382.1137 [ $(\text{M} + \text{H})^+$ ; calcd for  $\text{C}_{17}\text{H}_{25}\text{BrN}_3\text{O}_2$ , 382.1130].

*N*-(4-Chlorophenyl)-*N'*-(2,2,6,6-tetramethylpiperidin-4-yl)-oxalamide (NBD-556). Ethyl oxalyl chloride (0.26 mL, 2.35 mmol) was added to a round-bottom flask containing anhydrous THF (10 mL), and the resulting solution was cooled in an ice bath. Triethylamine (0.33 mL, 2.35 mmol) and 4-chloroaniline (300 mg, 2.35 mmol) were then added, and the resulting mixture was removed from the ice bath and allowed to come to room temperature. The reaction mixture was stirred for 6 h at room temperature, and the solid was removed by vacuum filtration. The filtrate was concentrated in vacuo, and the residue was redissolved in EtOAc (20 mL). The organic phase was washed with 1 M HCl (2  $\times$  10 mL) and saturated  $\text{NaHCO}_3$  (10 mL) and dried over  $\text{MgSO}_4$ . The solvent was removed under reduced pressure to give the desired ester (505 mg, 94%), which was used in the subsequent reaction without further purification: mp 149–150  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  8.92 (br s, 1H), 7.59 (d,  $J$  = 8.9 Hz, 2H), 7.33 (d,  $J$  = 8.9 Hz, 2H), 4.40 (q,  $J$  = 7.2 Hz, 2H), 1.41 (t,  $J$  = 7.1 Hz, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  160.8, 153.9, 134.9, 130.6, 129.2, 121.0, 63.8, 13.9; IR (thin film) 3333, 3120, 2988, 1728, 1700, 1597, 1546, 1496, 1476, 1302, 1177  $\text{cm}^{-1}$ ; high-resolution mass spectrum ( $\text{ES}^+$ )  $m/z$  227.0338 ( $\text{M}^+$ ; calcd for  $\text{C}_{10}\text{H}_{10}\text{ClNO}_3$ , 227.0349).

To a flask containing the above ester (580 mg, 2.55 mmol) were added a 1 M NaOH solution (12.8 mL, 12.8 mmol) and THF (15 mL), and the resulting mixture was stirred at room temperature for 20 h. EtOAc (20 mL) was added, and the organic layer was washed with 1 M NaOH (2  $\times$  10 mL).




FIGURE 2: Scheme for the synthesis of NBD-556 and NBD-557.

The combined aqueous layer was acidified to pH 2 with 2 M HCl and extracted with EtOAc ( $3 \times 20$  mL). The combined organic layer was dried over  $\text{MgSO}_4$  and concentrated to give the crude acid (430 mg, 85%). The acid was redissolved in anhydrous THF (6 mL) and added to a flask containing 4-amino-2,2,6,6-tetramethylpiperidine (0.41 mL, 2.37 mmol). HOBt (379 mg, 2.80 mmol) and EDC (540 mg, 2.80 mmol) were added, and the resulting mixture was stirred at room temperature for 20 h. The reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (30 mL), and the organic layer was washed with water ( $4 \times 20$  mL) and dried over  $\text{MgSO}_4$ . After the solvent was removed in vacuo, the crude product was chromatographed (1:1 EtOAc/MeOH) to give NBD-556 (247 mg, 34%):  $R_f = 0.3$ ; mp 197–198 °C;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  9.35 (br s, 1H), 7.59 (d,  $J = 8.8$  Hz, 2H), 7.30–7.40 (m, 3H), 4.20–4.33 (m, 1H), 1.92 (dd,  $J = 12.6$ , 3.8 Hz, 2H), 1.27 (s, 6H), 1.17 (s, 6H), 1.03–1.13 (m, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  158.8, 157.6, 134.9, 130.4, 129.2, 121.0, 50.9, 44.5, 43.7, 34.8, 28.4; IR (thin film) 3297, 2959, 1660, 1592, 1512, 1403  $\text{cm}^{-1}$ ; high-resolution mass spectrum ( $\text{ES}^+$ )  $m/z$  322.1312 [ $(\text{M} - \text{CH}_3)^+$ ; calcd for  $\text{C}_{16}\text{H}_{21}\text{ClN}_3\text{O}_2$ , 322.1322].

#### Cell Lines

The codon-optimized Env sequences encoding the gp120 glycoprotein from the YU2 HIV-1 strain with a CD5 signal peptide and a carboxy-terminal six-His tag were cloned into the pCIG plasmid. This plasmid was modified from the pIRES-GS glutamine synthetase amplification vector (15). The pCIG-YU2 gp120 plasmid was transfected into CHO-K1 cells, which were subjected to selection for glutamine synthetase, as described previously (15). Strongly expressing clones were selected for production of the gp120 glycoprotein. CHO-K1 cells expressed YU2 gp120 in 100 mL suspension cultures of ProCHO5 (Cambrex) with 2% GS supplement (JRH), 1% HT Supplement (Gibco), 100 units/L penicillin G and 100 mg/L streptomycin (Gibco), 275  $\mu\text{M}$  MSX (Sigma), and 1 mL/L Protease Inhibitor Cocktail Set III (Calbiochem). 293T human embryonic kidney and Cf2Th canine thymocytes (ATCC) were grown at 37 °C and 5%  $\text{CO}_2$  in Dulbecco's modified Eagle's medium (Invitrogen) containing 10% fetal bovine serum (Sigma) and 100  $\mu\text{g}/\text{mL}$  penicillin and streptomycin (Mediatech, Inc.). Cf2Th cells stably expressing human CCR5 were grown in medium supplemented with 0.4 mg/mL G418 (Invitrogen).

#### Preparation and Purification of Protein

Full-length gp120 from the YU2 strain of HIV-1 with a C-terminal His tag was produced using CHO-K1 cells described above. Cell supernatants were dialyzed against 20 mM sodium phosphate and 500 mM NaCl (pH 7.4). Approximately 50 mL of dialyzed supernatant was passed through

a 5 mL HisTrap HP column (Amersham Bioscience), and the protein was eluted using a linear imidazole gradient (ÄKTA FPLC system, Amersham Bioscience). Eluted fractions were immediately neutralized by dilution in 200 mM Tris (pH 7.4) and then dialyzed into PBS (Roche Diagnostics). Approximately 30 mL of this solution was passed through a 5 mL HiTrap NHS-activated HP column (Amersham Bioscience) coupled to mAb F105 (Strategic Biosolutions), and the protein was eluted using a stepwise gradient of 100 mM glycine and 150 mM NaCl (pH 2.4) (ÄKTA FPLC system, Amersham Bioscience). Eluted fractions were immediately neutralized by dilution in 0.2 M Tris (pH 7.4) and then dialyzed into PBS. Purified gp120 was analyzed by SDS-PAGE (Invitrogen) and shown to be a single species of approximately 90 kDa, of which 37 kDa is due to glycosylation. The purified YU2 gp120 was stored at  $-20$  °C in 1 mL aliquots of 0.5 mg.

Soluble D1–D2 CD4 (sCD4) was kindly provided by R. Wyatt (National Institutes of Health Vaccine Research Center, Bethesda, MD). Monoclonal antibody 17b was produced by Strategic Biosolutions (Newark, DE). sCD4 and MAb 17b were both dialyzed into PBS and stored in approximately 6 and 5 mg aliquots, respectively.

#### Isothermal Titration Calorimetry

Isothermal titration calorimetric experiments were performed using a high-precision VP-ITC titration calorimetric system from MicroCal Inc. (Northampton, MA). For the direct determination of the binding of the inhibitors to gp120, the calorimetric cell ( $\sim 1.4$  mL), which contained gp120 dissolved in PBS (Roche Diagnostics GmbH) (pH 7.4) with 2% DMSO, was titrated with the inhibitors NBD-557, NBD-556, and BMS-378806 dissolved in the same buffer. The concentration of gp120 was  $\sim 3$   $\mu\text{M}$ , and the concentration of inhibitor in the injection syringe was between 100 and 175  $\mu\text{M}$  for NBD-557 and NBD-556 and 45  $\mu\text{M}$  for BMS-378806. The binding of each inhibitor was studied at different temperatures in the range of 10–25 °C for NBD-557 and NBD-556 and 15–37 °C for BMS-378806. Binding of CD4 to gp120 was studied in experiments where the calorimetric cell, which contained 3  $\mu\text{M}$  gp120, was titrated with a solution of 30  $\mu\text{M}$  sCD4. The effect of NBD-556 and BMS-378806 on CD4 binding was studied by equilibrating 3  $\mu\text{M}$  gp120 with 300  $\mu\text{M}$  NBD-556 or BMS-378806 prior to titration with 30  $\mu\text{M}$  CD4. Binding of MAb 17b to gp120 was studied by stepwise 15  $\mu\text{M}$  additions (30  $\mu\text{M}$  per Fab site) to the calorimetric cell containing 3  $\mu\text{M}$  gp120 by itself or equilibrated with 300  $\mu\text{M}$  NBD-556. The effect of 17b on the binding of NBD-556 to gp120 was studied by titration of gp120 pre-bound to 17b with NBD-556. To that end, 2.5  $\mu\text{M}$  gp120 equilibrated with 5  $\mu\text{M}$  17b was titrated with NBD-556 at a concentration of 100  $\mu\text{M}$ . All titrations were



performed by adding the titrant in steps of 10  $\mu\text{L}$ . All solutions were properly degassed to avoid any formation of bubbles in the calorimeter during stirring. The heat evolved upon each injection of inhibitor was obtained from the integral of the calorimetric signal. The heat associated with binding to gp120 in the cell was obtained by subtracting the heat of dilution from the heat of reaction. The individual heats were plotted against the molar ratio, and the enthalpy change ( $\Delta H$ ) and association constant ( $K_a = 1/K_d$ ) were obtained by nonlinear regression of the data. Because of the low affinity, the stoichiometry was fixed to a value of 1 for the binding of NDB-556 and NBD-557 to gp120.

### Circular Dichroism Spectroscopy

CD experiments were conducted using a Jasco J-710 spectropolarimeter. Wavelength scans were performed with 3.7  $\mu\text{M}$  gp120 in the presence of 0, 5, 10, 20, and 300  $\mu\text{M}$  NBD-556 in a 0.1 cm cuvette in a water-jacketed cell. NBD-556 was dissolved in DMSO and dispensed in test tubes. The test tubes were placed in a lyophilizer to remove any DMSO prior to the addition of gp120 prepared in PBS. The solution was mixed thoroughly and allowed to equilibrate for 1 h. Each complete spectrum was the average of five consecutive scans collected from 195 to 240 nm. The individual scans were recorded using a scan rate of 20 nm/min, a bandwidth of 1 nm, and a response time of 2 s/point. Buffer scans were accumulated and subtracted from the sample scans, and the mean residue ellipticity was computed. The temperature was kept constant at 25  $^{\circ}\text{C}$ .

### Recombinant Luciferase Viruses

293T human embryonic kidney cells were cotransfected with plasmids expressing the pCMV $\Delta\text{P1}\Delta\text{env}$  HIV Gag-Pol packaging construct (16), the R5 YU2 envelope glycoproteins, or the envelope of control amphotropic murine leukemia virus (A-MLV), and the firefly luciferase-expressing vector at a DNA weight ratio (micrograms) of 1:1:3 using the Effectene transfection reagent (Qiagen). Cotransfection produced single-round, replication-defective viruses. The virus-containing supernatants were harvested 24–30 h after transfection, filtered (0.45  $\mu\text{m}$ ), aliquoted, and frozen at  $-80^{\circ}\text{C}$  until further use. The reverse transcriptase (RT) activities of all viruses were measured as described previously (17).

### Infection by Single-Round Luciferase Viruses

Cf2Th-CCR5 target cells were seeded at a density of  $6 \times 10^3$  cells/well in 96-well luminometer-compatible tissue culture plates (Dynex) 24 h before infection. On the day of infection, NBD-556 (0–100  $\mu\text{M}$ ) was incubated with recombinant viruses (10 000 RT units) at 37  $^{\circ}\text{C}$  for 30 min. The mixtures were then added to the target cells and incubated for 48 h at 37  $^{\circ}\text{C}$ ; after this time, the medium was removed from each well, and the cells were lysed by the addition of 30  $\mu\text{L}$  of passive lysis buffer (Promega) and three freeze–thaw cycles. An EG&G Berthold Microplate Luminometer (LB 96V) was used to measure the luciferase activity of each well after the addition of 100  $\mu\text{L}$  of luciferin buffer [15 mM  $\text{MgSO}_4$ , 15 mM  $\text{KPO}_4$  (pH 7.8), 1 mM ATP, and 1 mM dithiothreitol] and 50  $\mu\text{L}$  of 1 mM D-luciferin potassium salt (BD Pharmingen).




FIGURE 3: Calorimetric titration of gp120 with NBD-556 at 25  $^{\circ}\text{C}$  in PBS (pH 7.4) with 2% DMSO. The concentration of gp120 was 3.3  $\mu\text{M}$ , and the syringe contained NBD-556 at a concentration of 175  $\mu\text{M}$ . The change in heat capacity was calculated from the temperature dependence of the binding enthalpy (inset).

### Radiolabeling of Env Glycoprotein

293T cells were seeded at a density of  $3.5 \times 10^6$  cells in a T75 tissue culture flask 1 day before transfection. Cells were cotransfected with 9  $\mu\text{g}$  of pSVIIIEnv (YU2) and 1  $\mu\text{g}$  of pLTR-Tat using the Polyfect transfection reagent (Qiagen). One day after transfection, the cells were labeled for 48 h with [ $^{35}\text{S}$ ]Express protein labeling mix (100  $\mu\text{Ci}/\text{mL}$ ) (Perkin-Elmer). The supernatants were harvested 48 h later, cleared by centrifugation at 2000 rpm for 5 min, and stored at 4  $^{\circ}\text{C}$ . The amount of labeled gp120 was quantitated by immunoprecipitation with AIDS patient sera and protein A–Sepharose beads (Amersham Bio-Sciences), followed by SDS–PAGE gels and autoradiography.

### Env–CCR5 Binding Assay

Cf2Th cells expressing high levels of CCR5 were lifted using 5 mM EDTA (pH 7.5). The cells were washed with serum-free DMEM, added to microcentrifuge tubes ( $2\text{--}3 \times 10^6$  cells/tube), and incubated with 500  $\mu\text{L}$  of labeled YU2 gp120 in the presence and absence of varying concentrations of CD4-Ig or NBD-556 at 37  $^{\circ}\text{C}$  for 1.5 h with gentle agitation. The supernatants were removed following incubation, and the cells were washed twice with cold DMEM before lysis in 0.5 mL of IP buffer containing 0.5 M NaCl, 10 mM Tris (pH 7.5), 0.5% (v/v) NP40, and a cocktail of protease inhibitors. The cells were incubated in IP buffer for 30 min at 4  $^{\circ}\text{C}$  with gentle agitation. The lysates were cleared by centrifugation at 14 000 rpm for 30 min at 4  $^{\circ}\text{C}$ , immunoprecipitated with AIDS patient sera and protein A–Sepharose beads, and visualized by autoradiography of a 3 to 8% SDS–PAGE gel.

## RESULTS

*Isothermal Titration Calorimetric Studies of Binding of Inhibitor to HIV-1 gp120.* Figure 3 shows a microcalorimetric




FIGURE 4: Dissection of the thermodynamics of binding of sCD4, NBD-556, and BMS-378806 to gp120 at 25 °C.

ric titration of gp120 from the YU2 strain of HIV-1 with NBD-556. The binding of NBD-556 to gp120 at 25 °C is characterized by a binding affinity of 3.7  $\mu$ M, which corresponds to a change in Gibbs energy of  $-7.4$  kcal/mol. The large favorable binding enthalpy change of  $-24.5$  kcal/mol is partially compensated by a large unfavorable entropy change of  $-57.4$  cal  $K^{-1}$   $mol^{-1}$ . Under similar conditions, sCD4 binds to gp120 with a binding enthalpy of  $-34.5$  kcal/mol and an entropy change of  $-79$  cal  $K^{-1}$   $mol^{-1}$ . The binding of NBD-557 to gp120 was also characterized by an enthalpy change and an affinity similar to those measured for NBD-556 (data not shown). The change in heat capacity upon binding of NBD-556 to gp120, calculated from the temperature dependence of the binding enthalpy, is  $-1000 \pm 50$  cal  $K^{-1}$   $mol^{-1}$  (Figure 3, inset). Under similar conditions, the heat capacity change associated with the binding of sCD4 to gp120 is  $-1800$  cal  $K^{-1}$   $mol^{-1}$ . As described in an earlier report (13), the binding of BMS-378806 to gp120 is characterized by an affinity of 43 nM and an enthalpy change that is close to zero at 25 °C. The change in heat capacity for the binding of BMS-378806 to gp120 is only  $-292$  cal  $K^{-1}$   $mol^{-1}$ . Thus, NBD-556 exhibits a thermodynamic signature similar to that of sCD4 and much different from that of BMS-378806. Figure 4 shows the thermodynamic profiles for sCD4, NBD-556, and BMS-378806. The large favorable enthalpy change coupled to a large unfavorable entropy and large negative heat capacity is characteristic of a binding process that induces a significant structuring in the target protein (8).

**CD of gp120 Bound to NBD-556.** The gp120 structuring is also reflected in the CD spectrum of gp120 bound to NBD-556 (Figure 6). The specific increase in the level of secondary structure induced by NBD-556 is most pronounced between the isodichroic points at 221 and 238 nm, suggesting a larger helical content. The concomitant reduction in the level of random coil is reflected in the positive increase in ellipticity around 200 nm (18). The effect of NBD-556 is similar to the change in the CD spectrum observed for gp120 upon binding to CD4 (19).

**Effect of NBD-556 and BMS-378806 on Binding of CD4 to gp120.** The effect of the binding of NBD-556 or BMS-378806 on binding of CD4 to gp120 was assessed by performing sCD4 binding experiments on gp120 prebound to either compound. The expected apparent affinity in the presence of a competitive inhibitor can easily be calculated using the following equation:

$$K_{a,app} = \frac{K_a}{1 + K_{a,I}[I]} \quad (1)$$

where  $K_a$  is the association constant for sCD4 in the absence of inhibitor,  $K_{a,app}$  is the expected apparent association constant for sCD4 in the presence of inhibitor,  $K_{a,I}$  is the association constant of the inhibitor, and  $[I]$  is the concentration of the inhibitor. Similarly, the expected enthalpy change in the presence of a competitive inhibitor ( $\Delta H_{app}$ ) is calculated according to the equation

$$\Delta H_{app} = \Delta H - F_b \Delta H_I \quad (2)$$

where  $\Delta H$  and  $\Delta H_I$  are the enthalpy changes for the binding of sCD4 and the inhibitor, respectively, and  $F_b$  is the degree of saturation of gp120 with the prebound inhibitor.

According to eqs 1 and 2, in the presence of 300  $\mu$ M NBD-556, sCD4 is expected to bind gp120 with an association constant of  $\sim 1 \times 10^6$   $M^{-1}$  ( $K_d = 1$   $\mu$ M) and a change in enthalpy of  $-10$  kcal/mol if NBD-556 is a competitive inhibitor. Figure 5 shows the microcalorimetric titrations of gp120 (3  $\mu$ M) with sCD4 in the absence of inhibitor and in the presence of 300  $\mu$ M NBD-556, corresponding to more than 99% saturation of gp120. The experimentally measured binding enthalpy for binding of sCD4 to gp120 in the presence of a saturating concentration of NBD-556 is reduced to  $-11 \pm 1$  kcal/mol, which agrees well with the expected enthalpy change calculated with eq 2. The experimentally determined affinity under these conditions is  $0.3 \pm 0.1$   $\mu$ M, which is close to the expected value for a purely competitive inhibitor; in terms of the Gibbs energy of binding, the difference is 0.65 kcal/mol, which is well within experimental error. Taken together, the binding enthalpy change and affinity of sCD4 for gp120 in the presence of NBD-556 indicate that NBD-556 and sCD4 compete for the same binding site of gp120. The results are completely different for BMS-378806; in this case, the titrations in which 3  $\mu$ M gp120 is incubated with 300  $\mu$ M BMS-378806 show an only 10-fold reduction in the binding affinity of sCD4. Under those conditions, if BMS-378806 were a competitive inhibitor, the apparent affinity of CD4 for gp120 would have been reduced 7000-fold instead of 10-fold. The results clearly indicate that BMS-378806 is not a competitive inhibitor of sCD4. Albeit 700 times weaker than the affinity reduction expected from a competitive inhibitor, the 10-fold reduction in the affinity of CD4 for gp120 in the presence of excess BMS-378806 is real and most likely originates from some nonspecific conformational effect.

**NBD-556 Enhances Binding of gp120 to CCR5 and to the 17b Antibody.** CD4 binding stabilizes a gp120 conformation that efficiently binds CCR5 and CD4-induced antibodies, such as 17b, that recognize the gp120 coreceptor-binding site (20, 21). To determine whether NBD-556 would induce similar changes in gp120, the binding of radiolabeled gp120 glycoprotein to CCR5-expressing cells was examined in the absence or presence of NBD-556. As expected, the binding of gp120 to the Cf2Th-CCR5 cells was strengthened by incubation with CD4-Ig (Figure 7). Likewise, both NBD-556 and NBD-557 enhanced binding of gp120 to the Cf2Th-CCR5 cells. The observed binding was inhibited by several different CCR5 ligands, including TAK-779, the 2D7 anti-CCR5 antibody, and Compound A (data not shown). In




FIGURE 5: Microcalorimetric titrations of gp120 (3  $\mu$ M) with sCD4 (a) in the absence of inhibitor and (b) in the presence of 300  $\mu$ M NBD-556. The concentration of sCD4 in the syringe was 30  $\mu$ M in both experiments.




FIGURE 6: CD spectra of gp120 alone and in the presence of NBD-556. The concentrations of NBD-556 were 0, 5, 10, 20, and 300  $\mu$ M for curves 1–5, respectively, as indicated in the figure. The concentration of gp120 was kept constant at 3.7  $\mu$ M in all experiments.




FIGURE 7: Effect of NBD-556 on binding of gp120 to CCR5. Radiolabeled gp120 from the YU2 strain of HIV-1 was incubated with Cf2Th-CCR5 cells in the presence of the indicated ligands. The gp120 bound to the cells is shown.

contrast to NBD-556, BMS-378806 exerted no effect on the interaction of gp120 with the CCR5-expressing cells. Neither NBD-557/NBD-556 nor BMS-378806 inhibited binding of gp120 to CCR5 in the presence of sCD4 (data not shown). We conclude that NBD-556 and NBD-557, but not BMS-378806, activate the CCR5-binding ability of HIV-1 gp120.

Isothermal titration calorimetry was used to test whether the presence of saturating concentrations of NBD-556 altered the affinity of 17b for gp120. Direct binding of 17b to gp120 (22) is characterized by an affinity of 5 nM and an enthalpy change of  $-28.9$  kcal/mol at 25  $^{\circ}$ C. The presence of NBD-556 increases the affinity of 17b for gp120 more than 7-fold and reduces the enthalpy change to  $-10.9$  kcal/mol. As expected, the presence of 17b also enhances the affinity of NBD-556 for gp120. The binding of NBD-556 to gp120 prebound to 17b is characterized by an affinity of 0.5  $\mu$ M and an enthalpy change of  $-6.5$  kcal/mol. The binding of NBD-556 and 17b to gp120 is presented as a thermodynamic cycle in Figure 8.

*NBD-556 and NBD-557 Enhance Viral Entry in the Absence of CD4.* The results described above suggest that NBD-556 might promote conformational changes in gp120 related to virus entry. To test this hypothesis, we examined the effect of NBD-556 on the infection of CD4-negative, CCR5-expressing cells by recombinant HIV-1 bearing the envelope glycoproteins from the YU2 strain. Both NBD-556 and NBD-557 had a dramatic enhancing effect on CD4-independent infection (Figure 9 and data not shown). The infection of Cf2Th-CCR5 cells by the HIV-1 (YU2) virus was increased by a factor of more than 3 orders of magnitude in the concentration range of 0–40  $\mu$ M of NBD-556. The effect of NBD-556 on HIV-1 entry is specific, as no enhancement was observed for an HIV-1 bearing the envelope glycoproteins of the unrelated amphotropic murine leukemia virus (A-MLV), under the same conditions. Thus, NBD-556 and NBD-557 both induce entry-related conformational changes in the HIV-1 envelope glycoproteins similar to those induced by CD4.

## DISCUSSION

Previous studies (8) have shown that unliganded gp120 is characterized by the presence of a significant number (up to 50%) of unstructured residues. Furthermore, the relevant binding events associated with its biological function are coupled to large structuring events. For example, the binding of gp120 to CD4, the first event in entry of HIV-1 into cells, is coupled to the structuring of approximately 130 gp120 residues, many of which define the coreceptor binding epitope. The so-called activation of the coreceptor binding site induced by CD4 binding is, at the thermodynamic level, a reduction in the conformational energy penalty associated with structuring the binding site. In the absence of CD4 or a surrogate molecule, the coreceptor-interactive region on gp120 is not in a binding-competent conformation, and consequently, the binding to coreceptor or a surrogate (e.g., 17b) is much weaker because part of the binding energy is used in structuring the site.

The thermodynamic signature characteristic of binding events that are coupled to large structuring processes includes large favorable enthalpies, large unfavorable entropies, and large negative heat capacity changes. For example, at 25  $^{\circ}$ C the binding of sCD4 to gp120 is characterized by an enthalpy change of  $-34.5$  kcal/mol, an entropy change of  $-79$  cal  $K^{-1}$   $mol^{-1}$ , and a heat capacity change of  $-1.8$  kcal  $K^{-1}$   $mol^{-1}$ . A negative change in heat capacity is associated with the burial of hydrophobic surface, a process also characterized by a positive entropy change. The experimental obser-






FIGURE 8: Thermodynamics of binding of NBD-556 and 17b to gp120. All values were determined by isothermal titration calorimetry at 25 °C, except values for  $\Delta G$  and  $-T\Delta S$  for the binding of 17b to gp120 with prebound NBD-556 that were calculated by completing the cycle. The structures of gp120, sCD4, and 17b correspond to those in PDB entry 1GC1 (27). NBD-556 is represented as a green sphere.




FIGURE 9: Effect of NBD-556 on viral entry. The effect of NBD-556 on the infection of Cf2Th-CCR5 cells by recombinant luciferase-expressing HIV-1 bearing the envelope glycoproteins of the YU2 strain of HIV-1 or the amphotropic murine leukemia virus (A-MLV) is shown. Virus infection is expressed as the level of infection (measured by luciferase activity in the target cells) observed in the presence of NBD-556 relative to the level of infection observed in the absence of the drug.

vation of an overall negative entropy indicates that the positive entropy associated with the burial of hydrophobic surface is more than compensated by the negative conformational entropy associated with structuring.

The two compounds, NBD-557 and NBD-556, bind gp120 with large favorable enthalpy and unfavorable entropy changes and with a large negative heat capacity change, resembling CD4 itself. This signature contrasts with that of

the low-molecular-weight inhibitor of viral entry, BMS-378806 (14, 23, 24), that binds gp120 with a small enthalpy change in a process that is mostly entropy-driven (13). The effect of NBD-556 on binding of gp120 to CCR5 and to the CD4-induced antibody, 17b, is qualitatively similar to the effect observed for CD4. Binding of NBD-556 to gp120 induces the structuring of the coreceptor-binding site, thus enhancing the affinity of CCR5 and 17b due to a much reduced conformational energy penalty. An estimate of the change in conformational entropy can be made from the measured entropy and heat capacity changes, as described in detail elsewhere (25). Analysis of the data presented here shows that the binding of NBD-556 to gp120 induces the structuring of approximately 67 residues. This number is not as large as the number observed for sCD4 but sufficient to enhance coreceptor binding and remarkable for a compound with a molecular weight of less than 400 g/mol (338 g/mol for NBD-556 and 382 g/mol for NBD-557). Conformational changes to a more structured state are further confirmed by CD spectroscopy, which shows an increase in the level of secondary structure in gp120 after binding to NBD-556.

The results from the cell-based assays of viral entry show that the sCD4-induced structuring of the coreceptor-binding site in gp120 measured microcalorimetrically is similar to that induced by CD4 on the cell surface, resulting in a gp120 conformation with a higher affinity for the CCR5 or CXCR4 coreceptor. NBD-556 acts like a CD4 surrogate, enhancing viral entry in the absence of CD4. This also supports the relevance of the calorimetry measurements, made on monomeric HIV-1 gp120, to the gp120 glycoprotein in the context of the functional envelope glycoprotein trimer on the virion. The implied conformational flexibility of the HIV-1

gp120 glycoprotein in the virion trimer is consistent with the relationship between the binding thermodynamics and neutralizing activity of anti-gp120 antibodies (22) and with recent cross-linking studies of HIV-1 envelope glycoproteins on the surface of infected cells (26). An understanding of this conformational flexibility will assist the rational design of drugs and vaccines targeting the HIV-1 envelope glycoproteins.

## ACKNOWLEDGMENT

We thank Ms. Yvette McLaughlin for manuscript preparation. J.S. also acknowledges support from the International AIDS Vaccine Initiative, the Bristol-Myers Squibb Foundation, the William A. Haseltine Foundation for the Arts and Sciences, and the late William F. McCarty-Cooper.

## REFERENCES

- Chan, D. C., Fass, D., Berger, J. M., and Kim, P. S. (1997) Core structure of gp41 from the HIV envelope glycoprotein, *Cell* 89, 263–73.
- Wyatt, R., and Sodroski, J. (1998) The HIV-1 envelope glycoproteins: Fusogens, antigens, and immunogens, *Science* 280, 1884–8.
- Choe, H., Farzan, M., Sun, Y., Sullivan, N., Rollins, B., Ponath, P. D., Wu, L., Mackay, C. R., LaRosa, G., Newman, W., Gerard, N., Gerard, C., and Sodroski, J. (1996) The  $\beta$ -chemokine receptors CCR3 and CCR5 facilitate infection by primary HIV-1 isolates, *Cell* 85, 1135–48.
- Deng, H., Liu, R., Ellmeier, W., Choe, S., Unutmaz, D., Burkhart, M., Di Marzio, P., Marmon, S., Sutton, R. E., Hill, C. M., Davis, C. B., Peiper, S. C., Schall, T. J., Littman, D. R., and Landau, N. R. (1996) Identification of a major co-receptor for primary isolates of HIV-1, *Nature* 381, 661–6.
- Doranz, B. J., Rucker, J., Yi, Y., Smyth, R. J., Samson, M., Peiper, S. C., Parmentier, M., Collman, R. G., and Doms, R. W. (1996) A dual-tropic primary HIV-1 isolate that uses fusin and the  $\beta$ -chemokine receptors CKR-5, CKR-3, and CKR-2b as fusion cofactors, *Cell* 85, 1149–58.
- Dragic, T., Litwin, V., Allaway, G. P., Martin, S. R., Huang, Y., Nagashima, K. A., Cayanan, C., Maddon, P. J., Koup, R. A., Moore, J. P., and Paxton, W. A. (1996) HIV-1 entry into CD4<sup>+</sup> cells is mediated by the chemokine receptor CC-CKR-5, *Nature* 381, 667–73.
- Feng, Y., Broder, C. C., Kennedy, P. E., and Berger, E. A. (1996) HIV-1 entry cofactor: Functional cDNA cloning of a seven-transmembrane, G protein-coupled receptor, *Science* 272, 872–7.
- Leavitt, S. A., Schon, A., Klein, J. C., Manjappara, U., Chaiken, I. M., and Freire, E. (2004) Interactions of HIV-1 proteins gp120 and Nef with cellular partners define a novel allosteric paradigm, *Curr. Protein Pept. Sci.* 5, 1–8.
- Vita, C., Drakopoulou, E., Vizzavona, J., Rochette, S., Martin, L., Menez, A., Roumestand, C., Yang, Y. S., Ylisastigui, L., Benjouad, A., and Gluckman, J. C. (1999) Rational engineering of a miniprotein that reproduces the core of the CD4 site interacting with HIV-1 envelope glycoprotein, *Proc. Natl. Acad. Sci. U.S.A.* 96, 13091–6.
- Zhang, W., Canziani, G., Plugariu, C., Wyatt, R., Sodroski, J., Sweet, R., Kwong, P., Hendrickson, W., and Chaiken, I. (1999) Conformational changes of gp120 in epitopes near the CCR5 binding site are induced by CD4 and a CD4 miniprotein mimetic, *Biochemistry* 38, 9405–16.
- Dowd, C. S., Leavitt, S., Babcock, G., Godillot, A. P., Van Ryk, D., Canziani, G. A., Sodroski, J., Freire, E., and Chaiken, I. M. (2002)  $\beta$ -Turn Phe in HIV-1 Env binding site of CD4 and CD4 mimetic miniprotein enhances Env binding affinity but is not required for activation of co-receptor/17b site, *Biochemistry* 41, 7038–46.
- Zhao, Q., Ma, L., Jiang, S., Lu, H., Liu, S., He, Y., Strick, N., Neamati, N., and Debnath, A. K. (2005) Identification of N-phenyl-N'-(2,2,6,6-tetramethyl-piperidin-4-yl)-oxalamides as a new class of HIV-1 entry inhibitors that prevent gp120 binding to CD4, *Virology* 339, 213–25.
- Si, Z., Madani, N., Cox, J. M., Chruma, J. J., Klein, J. C., Schon, A., Phan, N., Wang, L., Biorn, A. C., Cocklin, S., Chaiken, I., Freire, E., Smith, A. B., III, and Sodroski, J. G. (2004) Small-molecule inhibitors of HIV-1 entry block receptor-induced conformational changes in the viral envelope glycoproteins, *Proc. Natl. Acad. Sci. U.S.A.* 101, 5036–41.
- Wang, T., Zhang, Z., Wallace, O. B., Deshpande, M., Fang, H., Yang, Z., Zadajura, L. M., Tweedie, D. L., Huang, S., Zhao, F., Ranadive, S., Robinson, B. S., Gong, Y. F., Ricard, K., Spicer, T. P., Deminie, C., Rose, R., Wang, H. G., Blair, W. S., Shi, P. Y., Lin, P. F., Colonno, R. J., and Meanwell, N. A. (2003) Discovery of 4-benzoyl-1-[(4-methoxy-1H-pyrrolo[2,3-b]pyridin-3-yl)oxoacetyl]-2-(R)-methylpiperazine (BMS-378806): A novel HIV-1 attachment inhibitor that interferes with CD4-gp120 interactions, *J. Med. Chem.* 46, 4236–9.
- Pu, H., Cashion, L. M., Kretschmer, P. J., and Liu, Z. (1998) Rapid establishment of high-producing cell lines using dicistronic vectors with glutamine synthetase as the selection marker, *Mol. Biotechnol.* 10, 17–25.
- Parolin, C., Taddeo, B., Palu, G., and Sodroski, J. (1996) Use of cis- and trans-acting viral regulatory sequences to improve expression of human immunodeficiency virus vectors in human lymphocytes, *Virology* 222, 415–22.
- Rho, H. M., and Gallo, R. C. (1979) Characterization of reverse transcriptase from feline leukemia virus by radioimmunoassay, *Virology* 99, 192–6.
- Woody, R. W. (1995) Circular dichroism, *Methods Enzymol.* 246, 34–71.
- Myszka, D. G., Sweet, R. W., Hensley, P., Brigham-Burke, M., Kwong, P. D., Hendrickson, W. A., Wyatt, R., Sodroski, J., and Doyle, M. L. (2000) Energetics of the HIV gp120-CD4 binding reaction, *Proc. Natl. Acad. Sci. U.S.A.* 97, 9026–31.
- Sattentau, Q. J., Moore, J. P., Vignaux, F., Traincard, F., and Poignard, P. (1993) Conformational changes induced in the envelope glycoproteins of the human and simian immunodeficiency viruses by soluble receptor binding, *J. Virol.* 67, 7383–93.
- Thali, M., Moore, J. P., Furman, C., Charles, M., Ho, D. D., Robinson, J., and Sodroski, J. (1993) Characterization of conserved human immunodeficiency virus type 1 gp120 neutralization epitopes exposed upon gp120-CD4 binding, *J. Virol.* 67, 3978–88.
- Kwong, P. D., Doyle, M. L., Casper, D. J., Cicala, C., Leavitt, S. A., Majeed, S., Steenbeke, T. D., Venturi, M., Chaiken, I., Fung, M., Katinger, H., Parren, P. W., Robinson, J., Van Ryk, D., Wang, L., Burton, D. R., Freire, E., Wyatt, R., Sodroski, J., Hendrickson, W. A., and Arthos, J. (2002) HIV-1 evades antibody-mediated neutralization through conformational masking of receptor-binding sites, *Nature* 420, 678–82.
- Lin, P. F., Blair, W., Wang, T., Spicer, T., Guo, Q., Zhou, N., Gong, Y. F., Wang, H. G., Rose, R., Yamanaka, G., Robinson, B., Li, C. B., Fridell, R., Deminie, C., Demers, G., Yang, Z., Zadajura, L., Meanwell, N., and Colonno, R. (2003) A small molecule HIV-1 inhibitor that targets the HIV-1 envelope and inhibits CD4 receptor binding, *Proc. Natl. Acad. Sci. U.S.A.* 100, 11013–8.
- Guo, Q., Ho, H. T., Dicker, I., Fan, L., Zhou, N., Friberg, J., Wang, T., McAuliffe, B. V., Wang, H. G., Rose, R. E., Fang, H., Scarnati, H. T., Langley, D. R., Meanwell, N. A., Abraham, R., Colonno, R. J., and Lin, P. F. (2003) Biochemical and genetic characterizations of a novel human immunodeficiency virus type 1 inhibitor that blocks gp120-CD4 interactions, *J. Virol.* 77, 10528–36.
- Luque, I., and Freire, E. (1998) Structure-based prediction of binding affinities and molecular design of peptide ligands, *Methods Enzymol.* 295, 100–27.
- Yuan, W., Bazick, J., and Sodroski, J. (2006) Characterization of the multiple conformational states of free monomeric and trimeric human immunodeficiency virus (HIV-1) envelope glycoproteins after fixation by crosslinker, *J. Virol.* 80, 6725–37.
- Kwong, P. D., Wyatt, R., Robinson, J., Sweet, R. W., Sodroski, J., and Hendrickson, W. A. (1998) Structure of an HIV gp120 envelope glycoprotein in complex with the CD4 receptor and a neutralizing human antibody, *Nature* 393, 648–59.

BI061193R