

# DNA Binding, Annealing, and Strand Exchange Activities of Brh2 Protein from *Ustilago maydis*<sup>†</sup>

Nayef Mazloum, Qingwen Zhou, and William K. Holloman\*

Department of Microbiology and Immunology, Cornell University Weill Medical College, New York, New York 10021

Received February 27, 2007; Revised Manuscript Received April 17, 2007

**ABSTRACT:** Brh2 is the *Ustilago maydis* ortholog of the BRCA2 tumor suppressor. It functions in repair of DNA by homologous recombination by controlling the action of Rad51. A critical aspect in the control appears to be the recruitment of Rad51 to single-stranded DNA regions exposed as lesions after damage or following a disturbance in DNA synthesis. In previous experimentation, Brh2 was shown to nucleate formation of the Rad51 nucleoprotein filament that becomes the active element in promoting homologous pairing and DNA strand exchange. Nucleation was found to be initiated at junctions of double-stranded and single-stranded DNA. Here we investigated the DNA binding specificity of Brh2 in more detail using oligonucleotide substrates. We observed that Brh2 prefers partially duplex structures with single-stranded branches, flaps, or D-loops. We found also that Brh2 has an inherent ability to promote DNA annealing and strand exchange reactions on free as well as RPA-coated substrates. Unlike Rad51, Brh2 was able to promote DNA strand exchange when preincubated with double-stranded DNA. These findings raise the notion that Brh2 may have roles in homologous recombination beyond the previously established Rad51 mediator activity.

Individuals with a mutation in the gene encoding BRCA2 are predisposed to breast, ovarian, and other cancers. The function of BRCA2 is in DNA repair by homologous recombination where it mediates interactions of Rad51, a protein needed for proficiency in homologous pairing and DNA strand exchange (1, 2). BRCA2 associates with Rad51 through a BRC repeat domain located medially plus a second unrelated domain at the extreme C-terminus (3) and interacts with DNA through a domain comprised of a helix rich region and tandem OB<sup>1</sup> folds closely related to those present in RPA for association with DNA (4). In studies using the conserved but smaller homologue (Brh2) from *Ustilago maydis* as a model system for understanding interaction of BRCA2 with Rad51, it was found that Brh2 directs Rad51 to junctions of single-stranded (ss) and double-stranded (ds) DNA to initiate filament formation (5). Assembly of Rad51 on ss DNA to form an extended presynaptic filament is a necessary prerequisite for efficient homologous pairing and strand exchange (6, 7). The overall reaction is dependent on ATP, but joint molecule formation and strand exchange over significant lengths do not depend on ATP hydrolysis, only that it be present as a cofactor (8). As ds DNA is readily bound by Rad51 (7, 9, 10), pairing reactions can be inhibited

by excess ds DNA since this competes with unbound ss DNA for the available free Rad51 (11). Such inhibition can be circumvented by precoating ss DNA with Rad51 before addition of ds DNA (7).

Discovery of mediator activity associated with Brh2 followed in the tracks of the earlier pioneering studies demonstrating mediator function of Rad52 protein in promoting Rad51-catalyzed strand exchange (12–15). Yeast lacks a protein orthologous to BRCA2 but relies instead on Rad52 for all aspects of homology-directed repair and recombination events (16). In studies performed in vitro, Rad52 was found to stimulate Rad51-promoted strand exchange by enabling Rad51 to displace RPA and access DNA, thus facilitating assembly into a filament (17, 18). This mediator activity stems from Rad52's highly specific ss DNA binding activity and from its ability to interact physically with Rad51 and RPA. Besides its mediator function, Rad52 has the capacity to promote homologous pairing as measured by the level of D-loop formation and oligonucleotide DNA strand exchange. It greatly accelerates annealing of complementary single strands (19–21) and carries out the homologous pairing reactions in the absence of ATP in contrast to Rad51 (22, 23). Rad52's innate capacity to promote recombination reactions as well as to mediate loading of Rad51 on DNA has been interpreted as evidence for multifunctionality underlining its pivotal role in homologous recombination in yeast.

BRCA2 and Rad52 are unrelated at the sequence level, and their DNA binding domains bear no relationship to each other. The three OB folds within the DNA binding domain of BRCA2 are arranged in a tandem linear order (4), while with Rad52, a ss DNA-binding groove is thought to form around the outside of an 11-subunit ringlike complex (24,

<sup>†</sup> This work was supported in part by a grant from the National Institutes of Health.

\* To whom correspondence should be addressed. Telephone: (212) 746-6510. Fax: (212) 746-8587. E-mail: wkholl@med.cornell.edu.

<sup>1</sup> Abbreviations: BH2, Black Hole 2 dark quencher; ds DNA, double-stranded DNA; DTT, dithiothreitol; FAM, 6-carboxyfluorescein; FRET, fluorescence resonance energy transfer; HEPES, *N*-(hydroxyethyl)piperazine-*N'*-2-ethanesulfonic acid; His, hexahistidine affinity tag; IB, Iowa Black RQ dark quencher; IPTG, isopropyl  $\beta$ -D-thiogalactoside; MBP, maltose binding protein; *n*mer, oligomer with *n* residues; NTA, nitrilotriacetate; OB, oligonucleotide/oligosaccharide binding; RPA, replication protein A; ss DNA, single-stranded DNA.




FIGURE 1: Proteins in the analysis. SDS-polyacrylamide gels with approximately 15 pmol each of the indicated proteins run alongside a ladder of Precision Protein standards (Bio-Rad Laboratories) were stained with Coomassie blue: (A) MBP-Brh2/His-Dss1 and Rad51, (B) MBP-Rad52, and (C) RPA heterotrimer.

25). The Rad51-interacting region in Rad52 has been localized by deletion mapping (26). There is no sequence motif reminiscent of BRC, but the interaction region is not yet defined at the structural level. Nevertheless, there is an interesting functional parallel. In studies with Brh2 and with a truncated form of BRCA2, it was found that these proteins recruit Rad51 to RPA-coated single-stranded DNA and facilitate the nucleation of the filament with concomitant displacement of RPA (5, 27). Thus, there seems to be a measure of functional conformity between Brh2 and Rad52 in the Rad51 loading or mediator function. Here, using oligonucleotide substrates, we examined the DNA binding specificity of Brh2 and tested its ability to promote DNA transactions involving complementary strands in comparison with both Rad51 and Rad52.

## EXPERIMENTAL PROCEDURES

**Protein Expression and Purification.** All proteins used in this study were recombinant proteins that were purified after being overproduced in *Escherichia coli* (Figure 1). As previously described, Brh2 was purified as a complex with Dss1 (5) with the exception that here Brh2 was purified as a MBP fusion protein in a complex with His-tagged Dss1 protein. The Brh2-Dss1 complex (Figure 1A) was used in the experimentation described below and is termed Brh2 for the sake of simplicity. The Brh2 cDNA was cloned into pMAL-C2 (New England Biolabs) such that the fusion begins at Brh2 residue 106. The Dss1 cDNA was cloned into pACYC Duet vector (Novagen) for expression with an N-terminal hexahistidine affinity tag. The Brh2 and Dss1 fusion constructs are completely functional in complementing the radiation sensitivity of the *U. maydis* *brh2* and *dss1* mutants, respectively. The plasmids were sequentially transformed into *E. coli* BL21(DE3) cells and cultures grown in LB medium containing ampicillin and chloramphenicol. Cell cultures (3 L) were grown at 37 °C to an  $A_{600}$  of 0.6, induced by addition of IPTG to 0.1 mM, but maintained at 18 °C after induction for 10 h. After being harvested, cells were resuspended in 50 mL of BA buffer [25 mM Tris-HCl (pH 7.5), 200 mM KCl, 1 mM DTT, 10% glycerol, and 0.1% NP40] with 20 mM imidazole, crushed by two passes through a French pressure cell at 20 000 psi, and centrifuged at 16 000

rpm in a Sorvall SS34 rotor for 30 min. The supernatant was recovered and centrifuged at 27 000 rpm for 1 h in a Beckman SW28 rotor. Subsequent procedures were all performed at 0–4 °C. The supernatant was passed onto a column (6 mL bed volume) of NTA agarose (Novagen) charged with  $\text{Ni}^{2+}$  and equilibrated in the same buffer. The column was washed with 40 mL of buffer and the protein eluted with 25 mL of BA buffer and 200 mM imidazole. The eluate was loaded on a 5 mL column of cross-linked amylose which was washed with 40 mL of BA buffer followed by 25 mL BA buffer and 10 mM maltose. The fractions containing Brh2 protein were pooled, diluted to 100 mM KCl, and applied to a 1 mL MonoQ column equilibrated in buffer BB [25 mM Tris-HCl (pH 7.5), 100 mM KCl, 1 mM DTT, and 10% glycerol]. After being washed with 5 mL of buffer, the protein was eluted with a linear gradient (20 mL) from 0.1 to 0.6 M KCl. Peak fractions were pooled and dialyzed against buffer BB and stored frozen at –80 °C.

For overproduction of Rad51 in *E. coli* BL21(DE3), the *U. maydis* cDNA encoding Rad51 was cloned in a pET vector derivative (Novagen) under control of the T7 promoter so no affinity tags would be included. Cultures (6 L) were grown at 37 °C and induced as described above and cells harvested 5 h later after continued incubation at 37 °C. Cells were resuspended in 50 mL of ice-cold buffer A [25 mM Tris-HCl (pH 7.5), 1 mM DTT, 1 mM EDTA, 10% glycerol, and 0.1% NP40] and 0.5 M KCl, crushed, and centrifuged as described above. Subsequent procedures were all performed at 0–4 °C. The high-speed supernatant was diluted with buffer RA [25 mM Tris-HCl (pH 7.5), 1 mM DTT, 1 mM EDTA, and 10% glycerol], to a conductivity equivalent to 100 mM KCl, mixed with 200 mL of gravity-settled DEAE-cellulose slurry (Whatman DE52), stirred for 45 min, and then passed onto a Buchner funnel with a sintered glass filter. The resin was washed batchwise on the funnel under gentle vacuum with 2 L of buffer RA with 100 mM KCl, followed by a wash with 200 mL of buffer A with 2 M KCl. The 2 M eluate was diluted 4-fold and then loaded onto a column (20 mL bed volume) of Affi-Gel Blue (Bio-Rad Laboratories) equilibrated in buffer RA and 0.2 M KCl. Following the method described for yeast Rad51 (10), the column was eluted with a linear salt gradient (200 mL) from 0.2 to 1.5 M KCl, and peak fractions were pooled after identification of those containing Rad51 by SDS gel electrophoresis and passed over a column (20 mL bed volume) of hydroxyapatite equilibrated with buffer RB [25 mM potassium phosphate (pH 7.5), 500 mM KCl, 1 mM DTT, and 10% glycerol] which was then eluted with a linear gradient (200 mL) from 25 mM to 0.8 M potassium phosphate. Peak fractions were pooled and dialyzed against buffer RC [25 mM Tris-HCl (pH 7.5), 100 mM KCl, 1 mM DTT, and 10% glycerol], then applied to a 1 mL Mono Q column (Amersham Biosciences), washed with 10 mL of RC buffer, and eluted with a linear gradient (20 mL) from 0.1 to 1 M KCl. Peak fractions that appeared to be >95% homogeneous were pooled (Figure 1A) and dialyzed against buffer RD [25 mM Tris-HCl (pH 7.5), 0.5 M KCl, 1 mM DTT, 0.1 mM EDTA, and 20% glycerol], snap-frozen in liquid nitrogen, and stored at –80 °C.

For overproduction of Rad52 in *E. coli* BL21(DE3), the *U. maydis* cDNA encoding the Rad52 ortholog (GenBank

accession number XP761136) was amplified by PCR and cloned in pMAL-C2 (New England Biolabs) as a MBP fusion protein. Cultures (6 L) were induced at 23 °C and processed as described above for Rad51 protein. Extracts in buffer A and 0.5 M KCl were passed through a column (100 mL) of DEAE-cellulose. The flow-through fraction was collected and passed onto a 5 mL column of cross-linked amylose (New England Biolabs). The column was washed with 10 mL of buffer A with 0.5 M KCl and then eluted with the same buffer with 10 mM maltose. The eluate was passed onto successive columns of Affi-Gel Blue and hydroxyapatite and fractionated by salt and phosphate gradient chromatography, respectively, as described above with Rad51. Rad52 prepared in this way was very sensitive to contaminating protease activity, and the purified protein preparations quickly accumulated a significant amount of proteolytic fragments deduced by their amylose binding capacity and mobility on SDS gel electrophoresis (Figure 1B) that had been truncated by approximately 300 residues at the C-terminus. We estimate that the most purified fractions contained 70–80% of the full-length protein. However, the *U. maydis* Rad52 at 722 residues is much larger than the yeast (504 residues) and human (418 residues) orthologs with an unusually long extension at the C-terminal end, a feature not uncommon among *U. maydis* proteins (28). The conserved DNA binding domain is located within the first 300 residues of the N-terminus so it seems likely that even in the most severely truncated proteolytic products the DNA binding domain was still intact. To date, we have been unsuccessful in ridding the preparations of the trace protease activity that appears to be responsible for generating Rad52-truncated fragments.

For overproduction of RPA in *E. coli* BL21(DE3), the cDNAs for RPA70 were cloned into pET28a (Novagen) to fuse a hexahistidine tag to the N-terminus and the cDNAs for RPA32 and RPA14 were cloned into pACYC Duet (Novagen). The plasmids were transformed sequentially into BL21(DE3). Cultures grown with shaking in a water bath (New Brunswick G76) were induced as described above but with a decreasing temperature gradient from 32 to 23 °C over the course of 5 h effected by switching off the water bath's heating element. Extracts from 3 L of culture were prepared using BA buffer, and protein was loaded on a NTA agarose column and eluted as described above. The eluate was diluted to a conductivity equal to that of 0.1 M KCl loaded onto a 1 mL MonoQ column and eluted with a linear gradient (30 mL) from 0.1 to 0.6 M KCl. Fractions eluting at 0.3 M KCl containing RPA subunits were pooled, diluted to a conductivity equal to that of 0.1 M KCl, loaded onto a heparin–Sephacrose column (1 mL HiTrap Heparin HP, Amersham Biosciences), and eluted with a linear gradient (20 mL) from 0.1 to 0.4 M KCl. Fractions eluting at 0.2 M KCl appeared to contain equimolar amounts of all three RPA subunits (Figure 1C) and were pooled and stored frozen at –80 °C.

**DNA Substrates.** Unless indicated otherwise, all DNA concentrations are expressed as moles of DNA molecules. Oligonucleotide sequences are in general based on pBluescript II DNA SK+ (Stratagene) and were synthesized and gel purified by Integrated DNA Technologies. The 100mer (and complementary strand) corresponds to pBluescript II SK+ nucleotides 2–101. After 30 pmol of 100mer (+) or

(–) strand had been labeled at the 5'-end with <sup>32</sup>P using T4 polynucleotide kinase and 60 μCi of [ $\gamma$ -<sup>32</sup>P]ATP, it was annealed with appropriate complementary strands in the presence of 20 mM KCl by being heated to 80 °C and then slowly cooled to 22 °C. Substrates were prepared by annealing the following combination of oligomers: ds–A/B; ds nicked–C/E/B; gap ds–F/G/B; 3'-overhang–A/D; 5'-overhang–C/B; ss–B; Holliday junction–L/M/N/O; fork–H/B; ds fork–H/I/C/B; 3'-flap–H/I/B; 5'-flap–H/C/B; bubble–J/B; D-loop–J/K/B; 5'-flap D-loop–J/B/I. Gel purification was performed when required by electrophoresis of annealed products in 12% polyacrylamide gels. The following oligonucleotides designated A–O were utilized: (A) 100mer (–), 5'-TAA ATT GTA AGC GTT AAT ATT TTG TTA AAA TTC GCG TTA AAT TTT TGT TAA ATC AGC TCA TTT TTT AAC CAA TAG GCC GAA ATC GGC AAA ATC CCT TAT A-3'; (B) 100mer (+), 5'-TAT AAG GGA TTT TGC CGA TTT CGG CCT ATT GGT TAA AAA ATG AGC TGA TTT AAC AAA AAT TTA ACG CGA ATT TTA ACA AAA TAT TAA CGC TTA CAA TTT A-3'; (C) 49mer (–), 5'-AA ATT GTA AGC GTT AAT ATT TTG TTA AAA TTC GCG TTA AAT TTT TGT TA-3'; (D) 49mer (+), 5'-TAA CAA AAA TTT AAC GCG AAT TTT AAC AAA ATA TTA ACG CTT ACA ATT T-3'; (E) pBSII (52–101) (–), 5'-AA ATC AGC TCA TTT TTT AAC CAA TAG GCC GAA ATC GGC AAA ATC CCT TAT A-3'; (F) pBSII (2–42) (–), 5'-TAA ATT GTA AGC GTT AAT ATT TTG TTA AAA TTC GCG TTA AA-3'; (G) pBSII (62–101) (–), 5'-TTT TTT AAC CAA TAG GCC GAA ATC GGC AAA ATC CCT TAT A-3'; (H) A HJ-pBSII 51–101 (–), 5'-GAC GCT GCC GAA TTC TGG CTT GCT AGG ACA TCT TTG CCC ACG TTG ACC CGA ATC AGC TCA TTT TTT AAC CAA TAG GCC GAA ATC GGC AAA ATC CCT TAT A-3'; (I) 1A HJ rev comp, 5'-CG GGT CAA CGT GGG CAA AGA TGT CCT AGC AAG CCA GAA TTC GGC AGC GTC-3'; (J) pBSII (2–42) 1A HJ (2–21) (62–101), 5'-TAA ATT GTA AGC GTT AAT ATT TTG TTA AAA TTC GCG TTA AAC GCT GCC GAA TTC TGG CTT TTT TTT AAC CAA TAG GCC GAA ATC GGC AAA ATC CCT TAT A-3'; (K) PBS II (44–59) (–), 5'-T TTT TGT TAA ATC AGC TCA-3'; (L) 1A HJ, 5'-GAC GCT GCC GAA TTC TGG CTT GCT AGG ACA TCT TTG CCC ACG TTG ACC CG-3'; (M) 2A HJ, 5'-CGG GTC AAC GTG GGC AAA GAT GTC CTA GCA ATG TAA TCG TCT ATG ACG TC-3'; (N) 3A HJ, 5'-GAC GTC ATA GAC GAT TAC ATT GCT AGG ACA TGC TGT CTA GAG ACT ATC GC-3'; and (O) 4A HJ, 5'-GCG ATA GTC TCT AGA CAG CAT GTC CTA GCA AGC CAG AAT TCG GCA GCG TC-3'. The sequence of the heterologous 80mer derived from pUC19 was 5'-TTA ACT ATG CGG CAT CAG AGC AGA TTG TAC TGA GAG TGC ACC ATA TGC GGT GTG AAA TAC CGC ACA GAT GCG TAA GGA GA-3'.

**Gel Mobility Shift.** Oligonucleotide DNA substrate at 2 nM was incubated with protein for 10 min at 37 °C in reaction mixtures (15 μL) with buffer containing 25 mM HEPES (pH 7.5) and 1 mM DTT. Glutaraldehyde was then added to a final concentration of 0.2%, incubation continued for 10 min to cross-link protein–DNA complexes, and Tris-HCl (pH 8.0) added to a final concentration of 120 mM to quench the cross-linking reaction. Products were resolved by electrophoresis on 1% agarose gels run at 4 °C. Gels were



dried on DE81 paper and radiolabeled bands identified using a Molecular Dynamics phosphorimager. Relative amounts of substrate and shifted product bands were determined using Image Quant.

**Single-Strand Annealing.** Annealing of complementary 100mer oligonucleotides was monitored by a fluorimetric method based on the procedure described by the Kowalczykowski laboratory (29) but modified to utilize PicoGreen (Molecular Probes) instead of DAPI to signal formation of duplex DNA as the former has a higher fluorescence yield (30). Measurements were performed in microtiter plates in 100  $\mu$ L reaction mixtures containing 25 mM Tris-HCl (pH 7.5), 25 mM KCl, 1 mM DTT, and 12 nM each A (–) strand and B (+) strand 100mer oligonucleotides at 37 °C in the presence of 0.15  $\mu$ M PicoGreen unless otherwise indicated. Reactions were set up with A strand ss 100mer without or with RPA at a ratio of nucleotide to protein of 25:1. The RPA/DNA mixture was deposited on one side of a 96-well white microtiter dish chamber as a small microdroplet (25  $\mu$ L) and the Brh2 or Rad52 solution as a separate microdroplet (25  $\mu$ L) on the opposite side of the chamber with care taken to avoid mixing. Reactions were started by addition of a solution (50  $\mu$ L) with the complementary B strand 100mer plus PicoGreen using a multichannel pipettor such that mixing of the microdroplets was rapid. Fluorescence measurements were taken at 10–30 s intervals with excitation and emission wavelengths set at 485 and 538 nm, respectively, using a SpectraMax Gemini fluorimetric plate reader (Molecular Devices). Due to the manual handling, several seconds elapsed before the first readings could be recorded. Therefore, the initial time points were not zero. Percent annealing was normalized to the fluorescence intensity of standards consisting of 12 nM 100mer ds DNA made by thermal annealing (100%) and 12 nM 100 ss DNA (0%).

**DNA Strand Exchange.** For strand exchange monitored by FRET, reactions were performed in two different formats depending on the instrumentation that was available for detection. In one using oligonucleotides modified with a Cy5 fluorophore donor and Iowa Black RQ dark quencher (Integrated DNA Technologies), reaction mixtures contained 10 nM ss 100mer (oligonucleotide B) and 10 nM ds 49mer (oligonucleotides C and D) in which the strand complementary to the ss 100mer (oligonucleotide C) was labeled at the 5'-end with Cy5 and the strand with identity to the ss 100mer (oligonucleotide D) was labeled at the 3'-end with the fluorescence quencher Iowa Black RQ. Reactions (100  $\mu$ L) were performed in 96-well white flat-bottom microtiter plates (Costar) and were started by addition of the ds 49mer to solutions containing the ss 100mer and proteins. Fluorescence measurements were monitored using the SpectraMax Gemini fluorimetric plate reader as described above with excitation and emission wavelengths set at 643 and 667 nm, respectively. Percent strand exchange was normalized to the fluorescence intensity of standards consisting of 10 mM ds 5'-Cy5-49mer/3'-Iowa Black-49mer (0%) and 10 nM ss 5'-Cy5-49mer (100%). The second format utilized a FAM fluorophore donor and Black Hole Quencher 2 (Integrated DNA Technologies) in which oligonucleotide C was labeled at the 5'-end with FAM and oligonucleotide D was labeled at the 3'-end with Black Hole Quencher 2. Reaction mixtures (15  $\mu$ L) containing 25 mM Tris-HCl (pH 7.5), 25 mM KCl,

1 mM DTT, and 30 nM ds 5'-FAM/3'-BH2 49mer were run at 37 °C in MicroAmp optical 96-well PCR plates (Applied Biosystems) and monitored in a TaqMan 7500 Real Time PCR System (Applied Biosystems) at 38 s intervals with excitation set at 488 nm and emission monitored with Filter A in place. Percent release from quenching, a metric interpreted as being equivalent to strand exchange, was normalized to the fluorescence intensity of standards consisting of 30 nM ds 5'-FAM/3'-BH2 49mer with 200 nM Brh2 (0%) and 30 nM ss 5'-FAM-49mer with 200 nM Brh2 (100%). *E. coli* exonuclease III (Exo III, New England Biolabs) was used as a control with units of activity specified by the manufacturer.

For more conventional strand exchange reactions with radiolabeled substrates, 60 nM ss 100mer (oligonucleotide B) was paired with 6 nM ds 49mer (oligonucleotides C and D) in which the strand complementary to the ss 100mer (oligonucleotide C) was labeled at the 5'-end with  $^{32}$ P using T4 polynucleotide kinase and [ $\gamma$ - $^{32}$ P]ATP. Preincubations were performed with the first DNA (60 nM ss DNA or 6 nM ds DNA, plus the protein as appropriate for the experiment), and then reactions were started by addition of the second DNA substrate (60 nM ss DNA or 6 nM ds DNA, respectively). In the standard reaction with Rad51, 60 nM ss 100mer was preincubated with Rad51 protein at 37 °C in a 15  $\mu$ L reaction mixture in a buffer containing 25 mM Tris-HCl (pH 7.5), 20 mM KCl, 1 mM DTT, 4 mM CaCl<sub>2</sub>, and 2 mM ATP. After 15 min, reactions were started by addition of 6 nM  $^{32}$ P-labeled ds 49mer DNA, and incubation was continued for additional time. In the standard Brh2 reaction, conditions were the same except that CaCl<sub>2</sub> and ATP were omitted and preincubation with ss 100mer was not performed unless otherwise indicated. Reactions were stopped by addition of SDS and proteinase K to concentrations of 1.2% and 1.7 mg/mL, respectively, for 30 min. EDTA (20 mM) and loading dye were added, and DNA components were resolved by electrophoresis on 10% polyacrylamide gels in 90 mM Tris-borate (pH 7.9) and 1 mM EDTA buffer. Dried gels were exposed to PhosphorImager plates, and developed images were quantified using ImageQuant as described above. The percentage of product was calculated as the radiolabeled product divided by the total amount of radiolabeled DNA in each lane.

## RESULTS

**Brh2 Binds Preferentially to DNA Structures with D-Loops or Branches.** The Brh2–Dss1 complex (Figure 1A) was used in the experimentation described below and is termed Brh2 for the sake of simplicity. In previous studies, it was established that Brh2 can recruit Rad51 to DNA to facilitate filament nucleation at a junction between ds DNA and ss DNA (5). The specificity for 3'-protruding single-strand regions implied Brh2 has an inherent ability to recognize junctions with the polarity of a processed DNA double-strand break. Support for this notion was obtained from measurement of the affinity for binding 3'-tailed versus 5'-tailed oligonucleotides using a gel mobility shift assay in which a 2.5-fold preference for 3'-tailed molecules was determined. We were interested in exploring additional DNA structural features that might constitute determinants contributing to Brh2 binding and used a similar methodology to measure specificity for an expanded series of DNA substrates with a




FIGURE 2: Specificity of Brh2 in DNA binding. (A) Reaction mixtures containing oligonucleotide substrates shown schematically and increasing concentrations of Brh2 (samples from left to right in each frame, 0, 2, 4, 10, 20, 100, and 200 nM) were incubated for 10 min and then analyzed by gel electrophoresis as described in Experimental Procedures. Samples were run in triplicate, but a single representative is shown. The sample with the 3'-flap substrate ran crooked, and the image was straightened using Photoshop. (B) The level of substrate shifted relative to the total signal in the lane was determined and is shown graphically. For the sake of clarity, the substrates are listed in order according to their binding affinity and are divided arbitrarily into two groups arranged so that all curves are visible. The left-hand figure contains the data represented in the top panel of mobility shift assays (A), while the right-hand figure contains the data represented in the bottom panel.

variety of different conformations. The set was framed around a 100mer oligonucleotide derived from a natural sequence stretch within pBluescript II plasmid DNA. By annealing oligomers with various runs of noncomplementary and complementary sequences to the 100mer, we prepared the various substrates shown schematically in Figure 2. The substrates end-labeled with  $^{32}\text{P}$  were incubated with increasing concentrations of Brh2, and protein-bound complexes were then detected by shifts in the mobility of substrates after gel electrophoresis. Tailed molecules with 3'- or 5'-overhangs, forked molecules, and flap molecules were designed to be 100 residues long and consist of half base-paired duplex and half tails, flaps, etc. The 100mer molecules of gapped ds DNA, bubble, and D-loop structures contained 20 internal residues that were unpaired.

Upon being tested with Brh2, the substrates appeared to fall into three general classes according to binding preference. Linear ds DNA molecules that were completely base paired were bound the weakest, linear ss DNA and linear molecules with single-stranded regions bound substantially better, and ds DNA molecules with forked, branched, or bubble structures bound the best. Binding preference did not strictly correlate with branches as the four-armed Holliday junction appeared to be somewhat anomalous in its binding. By a slight margin, the D-loop substrate appeared to be the top candidate with half of the input DNA shifted at a Brh2

concentration of  $\sim 4$  nM under these conditions. It was notable that a ladder of up to perhaps four distinctly migrating complexes became evident with an increasing protein concentration using several of the branched DNA substrates. Formation of each more slowly migrating complex appeared to be dependent on accumulation of the previous complex, suggesting the higher-order complexes resulted from protein-protein interaction.

**Brh2 Anneals ss DNA Complexed with RPA.** In the course of examining the association of Brh2 with various combinations of oligonucleotides, we noticed that oligomers with complementary stretches annealed quickly when mixed in the presence Brh2. To investigate this annealing activity in more detail, we used a fluorimetric assay to assess reaction in real time rather than tracking with radiolabeled oligomers by gel electrophoresis. Since spontaneous annealing of complementary oligonucleotides can be fast, monitoring reaction in real time has the advantage of eliminating high background and artifacts that could arise in electrophoretic gel-based assays, in which end points are determined after manipulations that might promote annealing. The fluorimetric assay is based on a method developed to measure annealing by yeast Rad52 protein, which monitors the increase in DAPI fluorescence upon binding annealed ds DNA (18), except that we substituted PicoGreen for DAPI to enhance sensitivity. PicoGreen has been developed as a selective reagent for determining ds DNA concentrations (30).

A 100mer oligonucleotide and its complement were used as substrates in annealing reactions. Since Rad52 has been reported to promote annealing of complementary ss DNA oligonucleotides (19), we used the *U. maydis* Rad52 protein as a positive control for annealing. To reduce the rate of spontaneous annealing of the oligonucleotides so that the effects of added protein could be measured, we manipulated the DNA concentration and temperature of reaction. Under conditions where the spontaneous reaction was slow, there was a substantial rate enhancement with added Brh2, although by comparison Brh2 was less active on a molar basis than Rad52 (Figure 3A,B). Annealing was delayed by 2–3 min when 4 mM  $\text{MgCl}_2$  was added to reaction mixtures (Figure 3C) but then proceeded with almost identical kinetics.

Because RPA binds tightly to ss DNA, it creates a barrier to Rad51 in DNA strand exchange. Rad52 and Brh2 have been demonstrated to serve as mediators in strand exchange by enabling Rad51 to gain access to ss DNA coated with RPA, thus facilitating displacement of RPA (5, 12, 14, 15). We tested whether Brh2 could promote annealing of the oligonucleotides when precoated with RPA. Spontaneous annealing was strongly inhibited when the ss 100mers were precoated with RPA at a saturating level of 25 nucleotides per RPA (Figure 3D). However, Brh2, like Rad52 (29, 31), was still able to promote annealing (Figure 3E,F). In the case of Brh2, there was a marked lag before annealing was initiated followed by an increasing rate indicative of overcoming a kinetic barrier in reaction. These results reinforce the previous observations that Brh2 has the capacity for specific interaction with RPA-coated ss DNA (5) but also suggest it is capable of enabling displacement of the RPA. Evidence for annealing of RPA-coated oligonucleotides by CeBRC-2, the BRCA2-related protein from *Caenorhabditis elegans*, was also presented recently but with the use of a gel assay measuring end points (32).




FIGURE 3: Brh2 anneals complementary single strands. The 100mer oligonucleotide (+) and (−) strands at 12 nM each were incubated at 30 °C with the indicated levels of (A) Brh2 or (B) Rad52 in the presence of 0.15  $\mu$ M PicoGreen as described in Experimental Procedures. (C) Reaction mixtures as in panel A with or without 4 mM MgCl<sub>2</sub> were incubated at 37 °C. (D) Reaction mixtures containing 100mer (+) and (−) strands at 40 nM each were incubated at 37 °C with or without RPA at a ratio of nucleotide to protein of 25:1. Reaction mixtures containing 100mer (+) and (−) strands at 12 nM each with (E) Brh2 at 200 nM or (F) Rad52 at 100 nM with or without RPA at a nucleotide to protein of 25:1 were incubated at 37 °C.

**DNA Strand Exchange Monitored by FRET.** To test whether Brh2 could promote a pairing transaction more complex than complementary strand annealing, we investigated its activity in reaction involving three DNA strands, namely, oligonucleotide strand exchange. Here the substrates were based on the ss 100mer used above and a ds 49mer homologous to the distal end of the ss 100mer. In an approach similar to that taken above to enable monitoring reaction in real time, we developed a fluorimetric assay based on FRET to follow DNA strand exchange. The basis of the assay was the release of emission by a fluorescent reporter from FRET-mediated dark quenching upon strand exchange. The 49mer (−) strand complementary to the ss 100mer (+) strand was labeled at the 5′-end with the fluorophore Cy5, and the 49mer (+) strand was labeled at the 3′-end with the fluorescence quencher Iowa Black (IB). When the 49mer strands were paired, the fluorescent signal of the Cy5 donor was quenched by FRET with the Iowa Black energy dissipater due to the proximity. However, when the Cy5-49mer strand was exchanged with the complementary 100mer, the FRET quenching was abrogated and fluorescence emission of Cy5 revealed. Thus, release from the quencher under these conditions is a measure of DNA strand exchange.




FIGURE 4: FRET assay for DNA strand exchange. (A) Reaction mixtures containing 30 nM ds 5′-FAM-49mer/3′-BHQ2-49mer, 120 nM homologous ss100mer, or heterologous ss 80mer were started by addition of 200 nM Brh2, and the increase in fluorescence was monitored. (B) Reaction mixtures were the same as those in panel A but containing 30 nM ds 5′-FAM-49mer/3′-BHQ2-49mer only, 4 mM MgCl<sub>2</sub>, and the indicated amounts of Exo III. (C) Reaction mixtures containing 10 nM ss 100mer were preloaded with 80 nM RPA for 10 min where indicated followed by 200 nM Brh2 or 100 nM Rad52. Reactions were started by addition of ds 5′-Cy5-49mer/3′-IB-49mer.


Under the same conditions that were used in the annealing reactions with equimolar concentrations of substrates, there was a substantial increase in fluorescence emission upon addition of Brh2 indicative of Cy5 fluorescence emission released from quenching to a level approaching 20% of the ss Cy5-49mer (Figure 4A). The increased fluorescence was dependent on addition of homologous ss 100mer, although it made no difference whether the (+) strand 100mer or (−) strand 100mer was used. Furthermore, no increase was observed when a heterologous ss 80mer was used in place of the ss 100mer or when ss DNA was omitted altogether (Figure 4B). On the other hand, addition of *E. coli* exonuclease III resulted in an immediate increase in fluorescence as would be expected upon removal of the 3′-IB quencher given Exo III's 3′ to 5′ polarity on ds DNA. These controls imply that the increase in fluorescence in the three-DNA strand reaction was due to strand transfer of the 5′-Cy5-49mer from the 5′-Cy5/3′-IB ds 49mer to the ss 100mer and not to nucleolytic removal of the quencher from the 5′-Cy5/3′-IB ds 49mer or to ss DNA-dependent unwinding activity.



When the ss DNA was precoated with a saturating amount of RPA as described above (25 nucleotide residues per RPA heterotrimer), the reaction proceeded with almost the same rate and to nearly the same extent as in the absence of RPA (Figure 4C). However, when the analysis was performed with Rad52 under conditions optimal for its annealing, only a low level of strand exchange activity was detectable. This modest activity was slightly elevated when the ss DNA was precoated with RPA. Thus, Brh2 is active in pairing a ss DNA with a complementary strand that is free or duplexed when RPA is present. Moreover, while Brh2 is less potent in the two-strand annealing reaction than Rad52, it is more active in the three-strand reaction.


**Brh2 Strand Exchange Is Insensitive to the Ordering of Components.** Oligonucleotide DNA strand exchange could potentially result from several different mechanisms. In one mode exemplified by Rad51 and RecA, it is thought that the protein induces an alteration in DNA structure upon binding to enable base pair switching between ds DNA and ss DNA, facilitating homologous pairing and strand exchange (33). A hallmark of the Rad51 class of mechanism is the requirement for formation of the nucleoprotein filament on ss DNA. Thus, DNA strand exchange is stimulated when Rad51 is preincubated with ss DNA but inhibited when preincubated with ds DNA. An alternative mode suggested from studies on the paralog Rad51C is that duplex oligomers are melted by the action of the protein and the separated strands can change partners by spontaneously annealing with other strands of complementary sequence that are present (34). We performed experiments to investigate the mechanism of DNA strand exchange observed with Brh2. Here we used the ss 100mer and ds 49mer substrates above but with conventional methodology following the transfer of the  $^{32}\text{P}$ -labeled (–) strand to the ss 100mer (+) strand by gel electrophoresis. Strand exchange is evident in the shift in label from the faster running ds 49mer to the slower  $^{32}\text{P}$ -labeled 49mer–100mer heteroduplex. Strand exchange promoted by Brh2 (Figure 5A) was dose-dependent and evident regardless of whether Brh2 was preincubated with ss 100mer or ds 49mer (Figure 5B), unlike the reaction promoted by Rad51, which proceeded only when Rad51 was preincubated with ss 100mer. Thus, in contrast to Rad51 strand exchange, Brh2 strand exchange was insensitive to the order of addition of components. We considered the possibility that the observed strand exchange might be an artifact arising from trace exonuclease contamination. Here, exposure of a single-stranded tract in ds DNA in the presence of a complementary strand and an annealing activity could lead to an apparent strand exchange (35). However, when we examined Brh2 directly for exonuclease activity by incubating it with the  $^{32}\text{P}$ -labeled ds 49mer and  $\text{Mg}^{2+}$ , a cofactor required for many *E. coli* deoxyribonucleases, and then analyzing the products on a DNA sequencing gel in which release of a single nucleotide could be detected, there was no indication of exonuclease contamination (Figure 5C). By comparison, controls in which Exo III was added exhibited a ladder of bands indicative of exonucleolytic degradation.

To test whether the mechanism of Brh2 strand exchange involved melting of the 49mer duplex followed by spontaneous annealing with the ss 100mer, we followed the procedure used in analysis of Rad51C by Kowalczykowski's group (34). Reaction mixtures containing Brh2 and the ds 49mer



**FIGURE 5:** Oligonucleotide strand exchange. (A) Brh2 or Rad51 was preincubated with 6 nM  $^{32}\text{P}$ -labeled ds 49mer (label on the minus strand) for 15 min (4 mM  $\text{CaCl}_2$  and 2 mM ATP added for Rad51 reactions), and then 60 nM ss 100mer was added and incubation continued for 15 min. (B) Brh2 or Rad51 was preincubated with 60 nM ss 100mer. After 15 min, 6 nM  $^{32}\text{P}$ -labeled ds 49mer was added and incubation continued for an additional 15 min. Concentrations for the samples of Brh2 in panels A and B were 0.05, 0.1, 0.25, 0.5, 1, and 2  $\mu\text{M}$ , respectively. For Rad51, concentrations were 0.06, 0.15, 0.3, 0.6, 1.5, and 3  $\mu\text{M}$  in panel A and 0.06, 0.15, 0.3, 0.6, 1.5, and 3  $\mu\text{M}$  in panel B, respectively. (C) Brh2 (1  $\mu\text{M}$ ) was incubated with 6 nM  $^{32}\text{P}$ -labeled ds 49mer (label on the minus strand) for 30 min in 25 mM Tris-HCl (pH 7.5), 20 mM KCl, 4 mM  $\text{MgCl}_2$ , 1 mM DTT, and 2 mM ATP. Similarly, reactions were run with 0.004 or 1 unit of Exo III. Samples were mixed with tracking dye stop solution containing formamide and EDTA to yield final concentrations of 50% and 10 mM, respectively, heated at 50  $^\circ\text{C}$  for 5 min, and then loaded on an 8% polyacrylamide gel containing 7 M urea that had been prerun for 30 min to warm the gel.

were incubated for a set time, then mixed with a 10-fold excess of ss 100mer, and either deproteinized immediately or incubated further and then deproteinized. If separated strands of the 49mer were generated by incubation with Brh2, then the labeled ss 49mer should anneal when the reaction was quenched upon addition of an excess of complementary ss 100mer, thus forming the strand exchange product with different mobility. No further increase in the amount of strand exchange product would be expected upon continued incubation with ss 100mer. However, no strand exchange above the background level was apparent after the initial incubation period of ds 49mer with Brh2 followed by the quench that included addition of excess ss 100mer and immediate deproteinization (Figure 6A). Strand exchange became apparent only when there was continued incubation with the added ss 100mer. These results imply that the Brh2 strand exchange reaction occurs without the generation of free ss DNA from the ds DNA as an intermediate. No strand exchange took place when the same procedure was used with Rad51 (Figure 6C), as expected. In this case, strand exchange



**FIGURE 6:** Order of addition in strand exchange. Brh2 or Rad51 was preincubated with one DNA substrate. Strand exchange reactions were then started by addition of the second DNA substrate. In panels A–C, proteins were mixed with the first DNA substrate as indicated and preincubated for 15 min. Reactions were started by addition of the second DNA substrate and terminated at 0, 0.5, 1, 1.5, 2, 3, 5, 7.5, 10, 15, 20, and 30 min. These 12 time points are represented in order from left to right by the 12 lanes in each frame. (A) Brh2 (0.5  $\mu$ M) was preincubated with 6 nM  $^{32}$ P-labeled ds 49mer for 15 min, followed by 60 nM ss 100mer. Samples were then taken at the times indicated above and deproteinized. Control lanes at the far left include  $^{32}$ P-labeled ss 49mer or ds 49mer with no protein. These are followed by control mixtures of  $^{32}$ P-labeled ds 49mer and ss 100mer with no protein but incubated for the indicated times before loading. A low level of slower moving DNA, which is a heteroduplex formed by spontaneous annealing of some residual free  $^{32}$ P-labeled ss 49mer with ss 100mer, is apparent in all three lanes. (B) Rad51 (1.5  $\mu$ M) was preincubated with 60 nM ss 100mer for 15 min. Reactions were initiated by addition of 6 nM  $^{32}$ P-labeled ds49mer and samples taken at the indicated times. (C) Rad51 (1.5  $\mu$ M) was preincubated with 6 nM  $^{32}$ P-labeled ds 49mer for 15 min. Reactions were initiated by addition of 60 nM ss 100mer and samples taken at the times indicated above. In panels D and E, proteins were mixed with the first DNA substrate as indicated and preincubated for variable lengths of time (0, 0.5, 1, 1.5, 2, 3, 5, 7.5, 10, 15, 20, and 30 min). Reactions were started by addition of the second DNA substrate and then terminated after 15 min. (D) Brh2 (0.5  $\mu$ M) or (E) Rad51 (1.5  $\mu$ M) was preincubated with 6 nM  $^{32}$ P-labeled ds 49mer for variable times given above, followed by addition of 60 nM ss 100mer for 15 min. Control lanes at the far left include mixtures of  $^{32}$ P-labeled ds 49mer and ss 100mer with no protein but incubated for the indicated times before loading. (F) Data from panels A–E are summarized with percentage strand exchange plotted for Rad51 and Brh2 reactions preincubated with ss or ds DNA.

with Rad51 became efficient only with extended preincubation time with the ss 100mer (Figure 6C). Conversely, the extent of Brh2-promoted strand exchange was undiminished regardless of the time of incubation of Brh2 with ds49-mer as long as this was followed by a second incubation with ss 100mer (Figure 6D,E), in contrast to the Rad51 reaction (Figure 6E,F). Thus, oligonucleotide strand exchange by Brh2 proceeds by a mechanism that does not involve duplex melting and does not require prior formation of a complex with ss DNA.

Given that ds DNA is less stable in the absence of divalent cation, we considered the possibility that the apparent strand exchange might be explained as a protein-enhanced reaction initiated in response to increased spontaneous breathing of the duplex. If so, there should be no reaction when a physiological level of  $Mg^{2+}$  is present. Therefore, we performed reactions in the presence of 4 mM  $Mg^{2+}$ , the concentration optimal for the Rad51-promoted strand exchange activity.<sup>2</sup> Under these conditions, there was robust

strand exchange promoted by Brh2 and the activity was slightly stimulated by addition of RPA (Figure 7, lanes c and k). These findings argue against the notion that the observed Brh2-promoted strand exchange results from spontaneous breathing of the ds DNA but, on the contrary, support the idea that the strand exchange activity of Brh2, regardless of the molecular basis, could represent a physiologically significant aspect of its biological function.

In the yeast system, it has been established that Rad52 forms a complex with Rad51 and can stimulate Rad51-catalyzed strand exchange. However, the ratio of Rad52 to Rad51 is critical. When the level of Rad52 protomers exceeds that of Rad51 by 1:3, strand exchange activity is strongly inhibited (17, 36). In agreement, we found that when Rad51 was at suboptimal levels for strand exchange (1 protomer per 18 nucleotides of ss DNA) there was almost total inhibition of activity upon addition of Rad52 at a ratio of

<sup>2</sup> N. Mazloun and W. K. Holloman, unpublished results.






FIGURE 7: Multiple components in strand exchange. All reactions were carried out in the presence of 4 mM  $\text{MgCl}_2$  and 2 mM ATP. In lanes b–i, protein components (500 nM Brh2, 330 nM Rad51, and 330 nM Rad52) were mixed together as indicated with 60 nM ss 100mer. After 15 min, 6 nM  $^{32}\text{P}$ -labeled ds 49mer was added and incubation continued for an additional 15 min. In lanes j–q, 60 nM ss 100mer was preincubated with 160 nM RPA for 10 min. Other proteins were added to the mix as indicated for an additional 15 min. Then 6 nM  $^{32}\text{P}$ -labeled ds 49mer was added and incubation continued for a further 15 min.

1:1 regardless of the presence or absence of RPA (Figure 7, lanes f and h and lanes n and p). Rad52 also inhibited the Brh2 strand exchange reaction (Figure 7, lanes e and m). Inhibition of Rad51 by Rad52 was completely relieved when Brh2 was also present (Figure 7, lanes i and q). These results suggest that precise control of the interplay among Brh2, Rad51, and Rad52 is essential for well-ordered DNA strand exchange.

## DISCUSSION

Previously, Brh2 was shown to be able to nucleate Rad51 filament formation on RPA-coated DNA at a ds DNA–ss DNA junction (5), thus demonstrating its activity as a mediator. Here we followed up on an aspect of that work by focusing more closely on the interplay of Brh2 with DNA using an oligonucleotide-based model system. Three principal conclusions can be drawn from these studies. First, Brh2 prefers binding to partially ds DNA oligonucleotide structures that contain ss DNA regions as forks, branches, or D-loops. Second, Brh2 can promote annealing of complementary single-strand oligonucleotides coated with RPA. Third, Brh2 can promote a type of DNA strand exchange using oligonucleotide substrates without regard for the order of addition of components. These findings raise the possibility that Brh2 might contribute to the homologous recombination system by means not previously considered in the studies on its mediator activity.

A paradoxical feature of Rad51 revealed by in vitro analysis is its sensitivity to ds DNA during the presynaptic stage. Efficient Rad51-promoted strand exchange requires a preloading step on ss DNA for circumventing the powerful inhibitory effects of ds DNA on strand exchange. How then can the function of Rad51 in vivo be reconciled in view of the largely duplex conformation of DNA within the nuclear environment? In the preceding work establishing that Brh2 could assist Rad51 in forming nucleoprotein filaments on a resected DNA molecule (5), the evidence suggested that Brh2 recognized ds DNA–ss DNA junctions and nucleated Rad51 filament formation in the appropriate polarity for subsequent strand exchange. That Brh2 can promote an oligonucleotide strand exchange reaction, albeit over a limited stretch of residues, with ds DNA and RPA-coated ss DNA present

raises the idea that this pairing activity might contribute to the initiation of Rad51-catalyzed pairing reactions. Is it possible that Brh2 sets up recombination events by establishing an initial synaptic step involving a limited stretch of residues and then recruits Rad51 to potentiate and extend pairing, thus forming a stable heteroduplex joint? The binding preference of Brh2 for D-loops and branched structures might be a clue to the mechanism underlying the oligonucleotide strand exchange. Perhaps the Brh2 strand exchange reaction is driven by stabilization of a branched or D-loop intermediate. Alternatively or in addition, perhaps the preferential binding of Brh2 to Holliday junctions suggests a role in stabilizing the cruciform or “chickenfoot” junctions formed at a stalled replication fork possibly enabling lesion bypass through template switching.

More experimentation will be required to establish the mechanism underlying the Brh2-promoted strand exchange, but it seems likely that it will involve some degree of duplex deformation, opening, or unwinding and concerted pairing of the third strand. With its intrinsic ability to promote single-strand annealing, and DNA strand exchange, Brh2 joins a set of proteins found in eukaryotes as well as prokaryotes providing similar functions central to homologous recombination. In prokaryotes, these include RecA family members, RecO (37), and RecT (38), and in eukaryotes Rad51 family members and paralogs such as Dmc1 (39, 40) and Xrcc3-Rad51C (34, 41), Hop2 (42), and Rad52. An emerging paradigm linking these proteins with a common mechanism is the formation of a presynaptic complex with ss DNA that enables switching of base pairs (33, 43). Brh2 seems to stand apart in being able to initiate DNA strand exchange in the presence of ds DNA, but that feature by no means rules out the possibility that its mechanism for DNA strand invasion proceeds through formation of a nucleoprotein structure favorable for base pair switching. As the *U. maydis* system offers the powerful advantage of genetic manipulation, it should be possible to generate Brh2 alleles defective in DNA annealing and strand exchange. Analysis of the corresponding mutant phenotypes would be decisive in elucidating the biological significance of these activities.

It seems to be a remarkable example of evolutionary convergence in function that Brh2 in *U. maydis* and Rad52 in *Saccharomyces cerevisiae* perform similar mediator reactions in vitro and provide the pivotal genetic control governing homologous recombination in vivo, yet they bear no structural resemblance. That a conserved Rad52 is present in *U. maydis*, yet relegated to a relatively minor role in recombination,<sup>3</sup> while a Brh2/BRCA2 ortholog is absent in *S. cerevisiae* raises the interesting questions of how these proteins evolved to similarly prominent positions in the homologous recombination systems of their respective organisms and what role is played by the *U. maydis* Rad52.

## ACKNOWLEDGMENT

This work greatly benefited from critical comments offered by Steve Kowalczykowski (University of California, Davis, CA) and Lorraine Symington (Columbia University, New York, NY).

<sup>3</sup> M. Kojic, Q. Zhou, and W. K. Holloman, unpublished results.

## REFERENCES

- Pellegrini, L., and Venkitesan, A. (2004) Emerging functions of BRCA2 in DNA recombination, *Trends Biochem. Sci.* 29, 310–316.
- West, S. C. (2003) Molecular view of recombination proteins and their control, *Nat. Rev. Mol. Cell Biol.* 4, 1–11.
- Esashi, F., Christ, N., Gannon, J., Liu, Y., Hunt, T., Jasin, M., and West, S. C. (2005) CDK-dependent phosphorylation of BRCA2 as a regulatory mechanism for recombinational repair, *Nature* 434, 598–604.
- Yang, H., Jeffrey, P. D., Miller, J., Kinnucan, E., Sun, Y., Thoma, N. H., Zheng, N., Chen, P. L., Lee, W. H., and Pavletich, N. P. (2002) BRCA2 function in DNA binding and recombination from a BRCA2-DSS1-ssDNA structure, *Science* 297, 1837–1848.
- Yang, H., Li, Q., Fan, J., Holloman, W. K., and Pavletich, N. P. (2005) The BRCA2 homologue Brh2 nucleates RAD51 filament formation at a dsDNA-ssDNA junction, *Nature* 433, 653–657.
- Benson, F. E., Stasiak, A., and West, S. C. (1994) Purification and characterization of the human Rad51 protein, an analogue of *E. coli* RecA, *EMBO J.* 13, 5764–5771.
- Sung, P., and Roberson, D. L. (1995) DNA strand exchange mediated by a RAD51-ssDNA nucleoprotein filament with polarity opposite to that of RecA, *Cell* 82, 453–461.
- Chi, P., Van Komen, S., Sehorn, M. G., Sigurdsson, S., and Sung, P. (2006) Roles of ATP binding and ATP hydrolysis in human Rad51 recombinase function, *DNA Repair* 5, 381–391.
- Baumann, P., Benson, F. E., and West, S. C. (1996) Human Rad51 protein promotes ATP-dependent homologous pairing and strand transfer reactions in vitro, *Cell* 87, 757–766.
- Zaitseva, E. M., Zaitsev, E. N., and Kowalczykowski, S. C. (1999) The DNA binding properties of *Saccharomyces cerevisiae* Rad51 protein, *J. Biol. Chem.* 274, 2907–2915.
- Sigurdsson, S., Trujillo, K., Song, B., Stratton, S., and Sung, P. (2001) Basis for avid homologous DNA strand exchange by human Rad51 and RPA, *J. Biol. Chem.* 276, 8798–8806.
- Sung, P. (1997) Function of yeast Rad52 protein as a mediator between replication protein A and the Rad51 recombinase, *J. Biol. Chem.* 272, 28194–28197.
- Benson, F. E., Baumann, P., and West, S. C. (1998) Synergistic actions of Rad51 and Rad52 in recombination and DNA repair, *Nature* 391, 401–404.
- Shinohara, A., and Ogawa, T. (1998) Stimulation by Rad52 of yeast Rad51-mediated recombination, *Nature* 391, 404–407.
- New, J. H., Sugiyama, T., Zaitseva, E., and Kowalczykowski, S. C. (1998) Rad52 protein stimulates DNA strand exchange by Rad51 and replication protein A, *Nature* 391, 407–410.
- Symington, L. S. (2002) Role of RAD52 epistasis group genes in homologous recombination and double-strand break repair, *Microbiol. Mol. Biol. Rev.* 66, 630–670.
- Song, B., and Sung, P. (2000) Functional interactions among yeast Rad51 recombinase, Rad52 mediator, and replication protein A in DNA strand exchange, *J. Biol. Chem.* 275, 15895–15904.
- Sugiyama, T., and Kowalczykowski, S. C. (2002) Rad52 protein associates with replication protein A (RPA)-single-stranded DNA to accelerate Rad51-mediated displacement of RPA and presynaptic complex formation, *J. Biol. Chem.* 277, 31663–31672.
- Mortensen, U. H., Bendixen, C., Sunjevaric, I., and Rothstein, R. (1996) DNA strand annealing is promoted by the yeast Rad52 protein, *Proc. Natl. Acad. Sci. U.S.A.* 93, 10729–10734.
- van Dyck, E., Stasiak, A. Z., Stasiak, A., and West, S. C. (2001) Visualization of recombination intermediates produced by RAD52-mediated single-strand annealing, *EMBO Rep.* 2, 905–909.
- Wu, Y., Sugiyama, T., and Kowalczykowski, S. C. (2006) DNA annealing mediated by Rad52 and Rad59 proteins, *J. Biol. Chem.* 281, 15441–15449.
- Kagawa, W., Kurumizaka, H., Ikawa, S., Yokoyama, S., and Shibata, T. (2001) Homologous pairing promoted by the human Rad52 protein, *J. Biol. Chem.* 276, 35201–35208.
- Bi, B., Rybalchenko, N., Golub, E. I., and Radding, C. M. (2004) Human and yeast Rad52 proteins promote DNA strand exchange, *Proc. Natl. Acad. Sci. U.S.A.* 101, 9568–9572.
- Kagawa, W., Kurumizaka, H., Ishitani, R., Fukai, S., Nureki, O., Shibata, T., and Yokoyama, S. (2002) Crystal structure of the homologous-pairing domain from the human Rad52 recombinase in the undecameric form, *Mol. Cell* 10, 359–371.
- Singleton, M. R., Wentzell, L. M., Liu, Y., West, S. C., and Wigley, D. B. (2002) Structure of the single-strand annealing domain of human, *Proc. Natl. Acad. Sci. U.S.A.* 99, 13492–13497.
- Krejci, L., Song, B., Bussen, W., Rothstein, R., Mortensen, U. H., and Sung, P. (2002) Interaction with Rad51 is indispensable for recombination mediator function of Rad52, *J. Biol. Chem.* 277, 40132–40141.
- San Filippo, J., Chi, P., Sehorn, M. G., Etchin, J., Krejci, L., and Sung, P. (2006) Recombination mediator and Rad51 targeting activities of a human BRCA2 polypeptide, *J. Biol. Chem.* 281, 11649–11657.
- Kamper, J., Kahmann, R., Bolke, M., Ma, L. J., Belfort, T., Saville, B. J., Banuett, F., Kronstad, J. W., Gold, S. E., Muller, O., Perlin, M. H., Wosten, H. A., de Vries, R., Ruiz-Herrera, J., Reynaga-Pena, C. G., Snetselaar, K., McCann, M., Perez-Martin, J., Feldbrugge, M., Basse, C. W., Steinberg, G., Ibeas, J. I., Holloman, W. K., Guzman, P., Farman, M., Stajich, J. E., Sentandreu, R., Gonzalez-Prieto, J. M., Kennell, J. C., Molina, L., Schirawski, J., Mendoza-Mendoza, A., Greiling, D., Munch, K., Rossel, N., Scherer, M., Vranes, M., Ladendorf, O., Vincon, V., Fuchs, U., Sandrock, B., Meng, S., Ho, E. C., Cahill, M. J., Boyce, K. J., Klose, J., Klosterman, S. J., Deelstra, H. J., Ortiz-Castellanos, L., Li, W., Sanchez-Alonso, P., Schreier, P. H., Hauser-Hahn, I., Vaupel, M., Koopmann, E., Friedrich, G., Voss, H., Schluter, T., Margolis, J., Platt, D., Swimmer, C., Gnirke, A., Chen, F., Vysotskaia, V., Mannhaupt, G., Guldener, U., Munsterkotter, M., Haase, D., Oesterheld, M., Mewes, H. W., Mauceli, E. W., DeCaprio, D., Wade, C. M., Butler, J., Young, S., Jaffe, D. B., Calvo, S., Nusbaum, C., Galagan, J., and Birren, B. W. (2006) Insights from the genome of the biotrophic fungal plant pathogen *Ustilago maydis*, *Nature* 444, 97–101.
- Sugiyama, T., New, J. H., and Kowalczykowski, S. C. (1998) DNA annealing by RAD52 protein is stimulated by specific interaction with the complex of replication protein A and single-stranded DNA, *Proc. Natl. Acad. Sci. U.S.A.* 95, 6049–6054.
- Singer, V. L., Jones, L. J., Yue, S. T., and Haugland, R. P. (1997) Characterization of PicoGreen reagent and development of a fluorescence-based solution assay for double-stranded DNA quantitation, *Anal. Biochem.* 249, 228–238.
- Shinohara, A., Shinohara, M., Ohta, T., Matsuda, S., and Ogawa, T. (1998) Rad52 forms ring structures and cooperates with RPA in single-strand DNA annealing, *Genes Cells* 3, 145–156.
- Petalcorin, M. I., Sandall, J., Wigley, D. B., and Boulton, S. J. (2006) CeBRC-2 stimulates D-loop formation by RAD-51 and promotes DNA single-strand annealing, *J. Mol. Biol.* 361, 231–242.
- Nishinaka, T., Shinohara, A., Ito, Y., Yokoyama, S., and Shibata, T. (1998) Base pair switching by interconversion of sugar puckers in DNA extended by proteins of RecA-family: A model for homology search in homologous genetic recombination, *Proc. Natl. Acad. Sci. U.S.A.* 95, 11071–11076.
- Lio, Y. C., Mazin, A. V., Kowalczykowski, S. C., and Chen, D. J. (2003) Complex formation by the human Rad51B and Rad51C DNA repair proteins and their activities in vitro, *J. Biol. Chem.* 278, 2469–2478.
- Kmiec, E. B., and Holloman, W. K. (1994) DNA strand exchange in the absence of homologous pairing, *J. Biol. Chem.* 269, 10163–10168.
- Arai, N., Ito, D., Inoue, T., Shibata, T., and Takahashi, H. (2005) Heteroduplex joint formation by a stoichiometric complex of Rad51 and Rad52 of *Saccharomyces cerevisiae*, *J. Biol. Chem.* 280, 32218–32229.
- Luisi-DeLuca, C., and Kolodner, R. (1994) Purification and characterization of the *Escherichia coli* RecO protein. Renaturation of complementary single-stranded DNA molecules catalyzed by the RecO protein, *J. Mol. Biol.* 236, 124–138.
- Noirot, P., Gupta, R. C., Radding, C. M., and Kolodner, R. D. (2003) Hallmarks of homology recognition by RecA-like recombinases are exhibited by the unrelated *Escherichia coli* RecT protein, *EMBO J.* 22, 324–334.
- Hong, E. L., Shinohara, A., and Bishop, D. K. (2001) *Saccharomyces cerevisiae* Dmc1 protein promotes renaturation of single-strand DNA (ssDNA) and assimilation of ssDNA into homologous super-coiled duplex DNA, *J. Biol. Chem.* 276, 41906–41912.

40. Sehorn, M. G., Sigurdsson, S., Bussen, W., Unger, V. M., and Sung, P. (2004) Human meiotic recombinase Dmc1 promotes ATP-dependent homologous DNA strand exchange, *Nature* 429, 433–437.
41. Kurumizaka, H., Ikawa, S., Nakada, M., Eda, K., Kagawa, W., Takata, M., Takeda, S., Yokoyama, S., and Shibata, T. (2001) Homologous-pairing activity of the human DNA-repair proteins Xrcc3.Rad51C, *Proc. Natl. Acad. Sci. U.S.A.* 98, 5538–5543.
42. Petukhova, G. V., Pezza, R. J., Vanevski, F., Ploquin, M., Masson, J. Y., and Camerini-Otero, R. D. (2005) The Hop2 and Mnd1 proteins act in concert with Rad51 and Dmc1 in meiotic recombination, *Nat. Struct. Mol. Biol.* 12, 449–453.
43. Gupta, R. C., Folta-Stogniew, E., O'Malley, S., Takahashi, M., and Radding, C. M. (1999) Rapid exchange of A:T base pairs is essential for recognition of DNA homology by human Rad51 recombination protein, *Mol. Cell* 4, 705–714.

BI700399M