

FIGURE 1: Trapped persulfenate intermediate in the active site of CDO. (a) Stereoview of the active site of CDO with electron density for the ligand shown at $0.8\rho_{\text{rms}}$. Bonds to the iron (orange dashed lines) and potential hydrogen bonds (red dashed lines) are shown. See Figure S1 of the Supporting Information for a close-up of omit density for the ligand. (b) Schematic of the active site interactions and metalcenter geometry based on the same view shown in panel a. Close nonpolar approaches surrounding the ligand (double arcs) and potential hydrogen bonds have been inferred from the environment. The geometry is such that one of the two protons of the sp^3 -hybridized α -amino group points toward a bound water (Wat156) and the other points toward a place where it can make a bifurcated hydrogen bond to both the Tyr157 OH group and the α -carboxylate oxygen. This is the main interaction allowing us to propose that the Tyr157 hydroxyl donates a hydrogen bond to atom O1.


1). In support of this interpretation, we note that the Cys $\text{S}\gamma$ position has strong anomalous difference density consistent with it being a sulfur atom (Table S2 of the Supporting Information) and that the Cys position is very similar to that seen previously (Figure S2 of the Supporting Information). Also, the refined B -factors of all of the Cys atoms and O1 are similar to each other and to those of the surrounding protein atoms, indicating the ligand is at least 70% occupied. The lower electron density for the O2 atom is consistent with the assignment because the O2 atom is in a loosely packed environment with no strong hydrogen bonding partners and would be expected to have significantly higher mobility than the O1 atom which is well-fixed by interactions with both the iron and the sulfur. Given the presence in the crystals of just CDO, cysteine, dissolved air, and buffer components, there is no other constellation of atoms that provides a plausible interpretation of the density.

In this complex, the metalcenter has distorted octahedral coordination with ligation distances ranging from 2.05 Å (for O1) to 2.47 Å (for Cys $\text{S}\gamma$) (Table S2 of the Supporting Information). The most distorted aspect of the coordination involves the S and O1 atoms, involved in a three-membered ring. Precedent for such an Fe–S–O cycle is found in two SO_2 iron-containing cluster compounds (17, 18).

A broader look at the Cys persulfenate binding site explains the remarkably high substrate specificity of CDO. The active site undergoes little change in conformation upon ligand formation (Figures S2 and S3 of the Supporting Information), yet the bound Cys is fully buried with no room to spare for additional atoms and with the full hydrogen bonding potential of both the α -amino and α -carboxylate satisfied (Figure 1). The phenolic hydroxyl of Tyr157 (11, 14) is clearly very important as it is located within 3.2 Å of four potential hydrogen bond donors and acceptors (Figure 1b). This complex net of interactions makes the assignment of the Tyr157 hydrogen bonding interactions challenging, but a reasonable proposal can be made that supports its assignment as a hydrogen bond donor (and potential proton donor) to the proximal O1 atom (Figure 1b).

Is this a true catalytic intermediate or a dead-end complex, and by what pathway does it form? These questions will

Scheme 1: Proposed CDO Mechanism



take extensive study to answer definitively. However, the consideration of this complex as an intermediate is consistent with mechanistic studies to date, and its observation stimulates new thinking about the CDO mechanism and sulfur oxidation chemistry. Also, should it be a dead-end complex, it nevertheless gives important insight into the chemistry of the CDO active site.

One plausible proposal for the CDO-catalyzed reaction mechanism, based on assuming that the structure described here is a true catalytic intermediate, is shown in Scheme 1. Cysteine binding to the free enzyme alters the coordination of the iron and creates the oxygen binding site (13). Starting from this enzyme cysteine complex 1, the binding of molecular oxygen to the active site ferrous iron gives 3. This is converted to 4 by a nucleophilic addition mechanism. Addition of the negatively charged distal oxygen to the sulfur then gives thiadioxirane 5. This undergoes heterolytic cleavage of the O–O bond, facilitated by a hydrogen bond (or proton transfer) between Tyr157 and the proximal oxygen, to give 6. This reaction is possible because of the low-energy empty d orbitals on the sulfur. Product release completes the reaction. We favor this mechanism over an alternative involving heterolytic cleavage of the O–O bond in 4 followed by addition of hydroxide to the resulting electron-deficient sulfur because the relatively nonpolar active site, with the distal oxygen surrounded by Leu95, Cys93 $\text{S}\gamma$, and Ile133, is poorly suited to stabilize negative charge on the distal oxygen of 4. Assuming this mechanism, we note that the trapping of intermediate 4 suggests that the rate-limiting

step of the reaction, at least as the enzyme is cooled to cryo-temperatures, is the intramolecular cyclization to give the thiadioxirane **5**. Other mechanistic possibilities for the formation of intermediate **4** cannot yet be excluded. For example, **4** could be formed directly from **2** by the coupling of a thiyl radical with a radical on the proximal oxygen. Alternatively, **4** could be formed by addition of a thiyl radical to the distal oxygen of **2** followed by rearrangement or by addition of the thiolate to side-bound oxygen. Differentiating between these possibilities will require additional experimentation.

The conversion of **4** to product involves an intramolecular rearrangement, which implies a perfect incorporation of both atoms of molecular oxygen into the product. A previous labeling study using rat liver cytoplasm suggested that this is the case (4) but was not conclusive. We have repeated the oxygen labeling experiment using pure recombinant rat CDO, shorter reaction times, and a much more sensitive and rapid MS-based analysis. These experiments demonstrated unequivocally that both oxygen atoms of the sulfinic acid are derived from a single molecule of molecular oxygen and that no exchange with buffer is occurring (Figure S4 of the Supporting Information).

One model system that may be closely analogous to the CDO reaction is the Ni(II)-catalyzed oxidation of thiols to sulfinate. Those reactions also show the incorporation of both atoms of molecular oxygen into the sulfinate, suggesting that intramolecular rearrangement to sulfinate is an intrinsic property of metal-bound persulfenates (19, 20) and need not be a consequence of restricted exchange at the enzyme active site. Our mechanism also maintains a central role for Tyr157 as an active site acid facilitating O–O bond cleavage in agreement with other current models (11). As discussed previously, the cross-link between Tyr157 and Cys93 may enhance activity by modulating the precise positioning and possibly the pK_a of Tyr157.

While the chemistry of thiol and thioether oxidation has been extensively investigated, persulfenate intermediates have not previously been directly observed (3, 21). The stabilization of the cysteine persulfenate **4** in the CDO active site at cryo-temperatures is fortuitous. This was an intermediate in all four previously proposed mechanisms for CDO (11); however, none of the proposals predicted the observed coordination of this intermediate to the active site iron, nor did they suggest that the cysteine thiol would bind to the proximal (iron-bound) oxygen rather than to the distal oxygen. Thus, if this is a true intermediate, the CDO chemistry is different from what has been anticipated on the basis of existing paradigms for thiol oxidation. These potentially unique aspects of the CDO mechanism, compared with dioxygenases that oxidize C=C bonds (22–24), reflect the electron-rich character of the thiol functional group and the availability of low-energy empty d orbitals on the sulfur. The visualization of this putative intermediate and its unexpected coordination chemistry are likely to stimulate further studies on thiol oxidation chemistry.

ACKNOWLEDGMENT

Crystallographic data used in this study were measured at beamline X12b of the National Synchrotron Light Source whose financial support comes principally from the Offices of Biological and Environmental Research and of Basic Energy Sciences of the U.S. Department of Energy and from the National Center for Research Resources of the National Institutes of Health.

SUPPORTING INFORMATION AVAILABLE

Materials and methods, Figures S1–S3, and Tables S1–S3. This material is available free of charge via the Internet at <http://pubs.acs.org>.

REFERENCES

1. Giles, G. I. (2006) *Curr. Pharm. Des.* 12, 4427–4443.
2. Flohe, L., and Harris, J. R. (2007) *Subcell. Biochem.* 44, 1–25.
3. Winterbourn, C. C., and Metodieva, D. (1999) *Free Radical Biol. Med.* 27, 322–328.
4. Lombardini, J. B., Singer, T. P., and Boyer, P. D. (1969) *J. Biol. Chem.* 244, 1172–1175.
5. Yamaguchi, K., Hosokawa, Y., Kohashi, N., Kori, Y., Sakakibara, S., and Ueda, I. (1978) *J. Biochem.* 83, 479–491.
6. Stipanuk, M. H. (2004) *Annu. Rev. Nutr.* 24, 539–577.
7. Heafield, M. T., Fearn, S., Steventon, G. B., Waring, R. H., Williams, A. C., and Sturman, S. G. (1990) *Neurosci. Lett.* 110, 216–220.
8. Bradley, H., Gough, A., Sokhi, R. S., Hassell, A., Waring, R., and Emery, P. (1994) *J. Rheumatol.* 21, 1192–1196.
9. Aluri, S., and de Visser, S. P. (2007) *J. Am. Chem. Soc.* 129, 14846–14847.
10. Chai, S. C., Bruyere, J. R., and Maroney, M. J. (2006) *J. Biol. Chem.* 281, 15774–15779.
11. Joseph, C. A., and Maroney, M. J. (2007) *Chem. Commun.*, 3338–3349.
12. McCoy, J. G., Bailey, L. J., Bitto, E., Bingman, C. A., Aceti, D. J., Fox, B. G., and Phillips, G. N., Jr. (2006) *Proc. Natl. Acad. Sci. U.S.A.* 103, 3084–3089.
13. Pierce, B. S., Gardner, J. D., Bailey, L. J., Brunold, T. C., and Fox, B. G. (2007) *Biochemistry* 46, 8569–8578.
14. Simmons, C. R., Liu, Q., Huang, Q., Hao, Q., Begley, T. P., Karplus, P. A., and Stipanuk, M. H. (2006) *J. Biol. Chem.* 281, 18723–18733.
15. Ye, S., Wu, X., Wei, L., Tang, D., Sun, P., Bartlam, M., and Rao, Z. (2007) *J. Biol. Chem.* 282, 3391–3402.
16. Dominy, J. E., Jr., Hwang, J., Guo, S., Hirschberger, L. L., Zhang, S., and Stipanuk, M. H. (2008) *J. Biol. Chem.* 283, 12188–12201.
17. Eveland, R. W., Raymond, C. C., Albrecht-Schmitt, T. E., and Shriver, D. F. (1999) *Inorg. Chem.* 38, 1282–1287.
18. Karet, G. B., Stern, C. L., Norton, D. M., and Shriver, D. F. (1993) *J. Am. Chem. Soc.* 115, 9979–9985.
19. Farmer, P. J., Solouki, T., Mills, D. K., Soma, T., Russel, D. H., Reibenspies, J. H., and Dearensbourg, M. Y. (1992) *J. Am. Chem. Soc.* 114, 4601–4605.
20. Mirza, S. A., Pressler, M. A., Kumar, M., Day, R. O., and Maroney, M. J. (1993) *Inorg. Chem.* 32, 977–987.
21. Clennan, E. L. (2001) *Acc. Chem. Res.* 34, 875–884.
22. Bugg, T. D., and Ramaswamy, S. (2008) *Curr. Opin. Chem. Biol.* (in press).
23. Karlsson, A., Parales, J. V., Parales, R. E., Gibson, D. T., Eklund, H., and Ramaswamy, S. (2003) *Science* 299, 1039–1042.
24. Kovaleva, E. G., and Lipscomb, J. D. (2007) *Science* 316, 453–457.

BI801546N