

# $\alpha$ -Synuclein Conformation Affects Its Tyrosine-Dependent Oxidative Aggregation<sup>†</sup>

Rebecca A. S. Ruf,<sup>‡</sup> Evan A. Lutz,<sup>‡</sup> Imola G. Zigoneanu,<sup>‡</sup> and Gary J. Pielak<sup>\*,‡,§</sup>

Department of Chemistry and Department of Biochemistry and Biophysics, Program in Molecular Biology and Biotechnology, and Lineberger Comprehensive Cancer Research Center, University of North Carolina, Chapel Hill, North Carolina 27599

Received October 6, 2008; Revised Manuscript Received November 12, 2008

**ABSTRACT:** Oxidative stress and aggregation of the protein  $\alpha$ -synuclein are thought to be key factors in Parkinson's disease. Previous work shows that cytochrome *c* with H<sub>2</sub>O<sub>2</sub> causes tyrosine-dependent in vitro peroxidative aggregation of proteins, including  $\alpha$ -synuclein. Here, we examine the role of each of  $\alpha$ -synuclein's four tyrosine residues and how the protein's conformation affects covalent oxidative aggregation. When  $\alpha$ -synuclein adopts a collapsed conformation, tyrosine 39 is essential for wild-type-like covalent aggregation. This lone N-terminal tyrosine, however, is not required for wild-type-like covalent aggregation in the presence of a denaturant or when  $\alpha$ -synuclein is present in noncovalent fibrils. We also show that preformed oxidative aggregates are not incorporated into noncovalent fibrils. These data provide insight into how dityrosine may be formed in Lewy bodies seen in Parkinson's disease.

Parkinson's disease, the second most common neurodegenerative disorder (1), affects more than 1.5 million people in the United States, with more than 60000 new diagnoses each year (2). The cellular hallmark of Parkinson's disease is the presence of Lewy bodies, intracellular aggregates of  $\alpha$ -synuclein (3), cytochrome *c* (4), and other proteins (5–7).  $\alpha$ -Synuclein is a 140-amino acid, intrinsically disordered protein (8) with three distinct regions (Figure 1A). The N-terminal region is positively charged; the hydrophobic core (also known as the non-amyloid component) comprises residues 61–90, and the C-terminal region is negatively charged. Tyrosines are essential for  $\alpha$ -synuclein oxidative aggregation (9). As shown in Figure 1A, the protein has four unevenly distributed tyrosine residues, one (Y39) near the N-terminus and three (Y125, Y133, and Y136) near the C-terminus (the protein has no tryptophan residues). Although disordered, the protein adopts a compact state (10–13) in which the charged termini collapse around the hydrophobic core (Figure 1B). Little is known about the exact mechanism of Parkinson's disease at the molecular level, but both covalent and noncovalent aggregation of  $\alpha$ -synuclein are thought to play a key role (14).

The oxidative stress marker dityrosine is detected in brain hydrolysates of murine models of Parkinson's disease (15). Fenton chemistry-based oxidation systems (i.e., a transition metal with H<sub>2</sub>O<sub>2</sub>) have been used extensively in vitro in attempts to reproduce in vivo covalent aggregation of  $\alpha$ -synuclein, but these systems are probably not accurate mimics (16, 17). For instance, although dityrosine (tyrosines

with a covalent bond between the 3- and 5-carbon atoms) is observed in Parkinson's disease models, the covalent aggregation induced by several Fenton chemistry systems is tyrosine-independent (17). In fact, the HO<sup>•</sup> generated by Cu<sup>2+</sup> with H<sub>2</sub>O<sub>2</sub> impedes dityrosine formation in the Alzheimer's disease protein, A $\beta$  (16). The cytochrome *c*/H<sub>2</sub>O<sub>2</sub> oxidation system (4), which we call the peroxidative system, is a better model for  $\alpha$ -synuclein covalent aggregation for two reasons. First, the peroxidative system causes tyrosine-dependent covalent aggregation of  $\alpha$ -synuclein (9), similar to the dityrosine seen in murine models of Parkinson's disease. Second, the peroxidative system causes the direct transfer of a free radical from cytochrome *c* to the acceptor protein (18, 19) without an HO<sup>•</sup> intermediate.

In addition to covalent aggregation,  $\alpha$ -synuclein aggregates noncovalently into fibrils. The  $\alpha$ -synuclein fibrils observed in patients with Parkinson's disease are linear rods, 5–10 nm in diameter, much like those seen in other amyloid diseases (20). The fibrils comprise insoluble cross- $\beta$ -sheets, and their growth in vitro exhibits a sigmoidal time dependence (21). Prior to fibril growth there is a lag, the length of which depends on factors such as protein concentration and pH (22). It is likely that structured intermediates, called nuclei, are required for fibril growth (23, 24). Following the lag is a period of elongation in which fibril concentration increases exponentially and plateaus (20). Fibril growth can be monitored by using thioflavin-T, which experiences a shift in its excitation spectrum when bound to  $\beta$ -sheets in fibrils, allowing it to be selectively excited at 442 nm (25).

Here, we shed light on the process of both peroxidative and noncovalent aggregation of  $\alpha$ -synuclein and how these forms of aggregation may be related. Specifically, we address the role of each tyrosine residue and show how the protein's conformation affects aggregation.

## EXPERIMENTAL PROCEDURES

*Preparation of Wild-Type and Mutant or Variant Recombinant Human  $\alpha$ -Synucleins and Cytochromes *c*.* Twenty-

<sup>†</sup> This work was supported by a NIH Director's Pioneer Award (SDP10D783) to G.J.P. and a Foundation for Aging Research Glaxo-SmithKline Foundation Award to R.A.S.R.

\* To whom correspondence should be addressed: Kenan Laboratories, University of North Carolina, Chapel Hill, NC 27599-3290. Fax: (919) 962-2388. E-mail: gary\_pielak@unc.edu.

<sup>‡</sup> Department of Chemistry.

<sup>§</sup> Department of Biochemistry and Biophysics, Program in Molecular Biology and Biotechnology, and Lineberger Comprehensive Cancer Research Center.




FIGURE 1:  $\alpha$ -Synuclein structure and conformation. A schematic representation of  $\alpha$ -synuclein's primary structure showing relevant regions, net charges, and position numbers (A) and a cartoon of its collapsed conformation (B). The positions of the tyrosine residues are colored green.

four human  $\alpha$ -synuclein mutants in the pT7-7 vector were created by using a site-directed mutagenesis kit (Quick-Change, Stratagene, La Jolla, CA). The wild-type construct and the mutants with one, two, or three tyrosine codons converted to phenylalanine codons were created with and without the valine 3 codon converted to a cysteine codon for labeling purposes. Additionally, the wild-type construct was created with the valine 66 codon converted to a cysteine codon, also for labeling purposes. Wild-type and variant  $\alpha$ -synucleins were purified as previously described (26) except that the freeze–thaw step was eliminated. Protein concentrations were determined with the Lowry method (27) (Modified Lowry Protein Assay Kit, Pierce, Rockford, IL) using human recombinant cytochrome *c* (28) as a standard. After purification, proteins were aliquoted and lyophilized for storage at  $-80^{\circ}\text{C}$ .

Wild-type human cytochrome *c* in the pBTR vector (28) was produced with and without lysine 39 converted to cysteine for labeling purposes. The cytochrome *c* variants were purified as previously described (28). Protein concentrations were determined from the absorbance at 410 nm by using a molar extinction coefficient of  $106.1\text{ mM}^{-1}\text{ cm}^{-1}$  (29). After purification, proteins were aliquoted and lyophilized for storage at  $-80^{\circ}\text{C}$ .

**Alexa Fluor Labeling.** Sixty nanomoles of V3C  $\alpha$ -synuclein was reacted with 300 nmol of Alexa Fluor 633 C<sub>5</sub> maleimide (Invitrogen, Carlsbad, CA) in 20 mM Tris (pH 7.5) for 2 h at room temperature. The reaction was allowed to proceed for 18–66 h at  $4^{\circ}\text{C}$ . Unreacted cysteines were blocked with 3 mmol of iodoacetamide (Sigma, St. Louis, MO). Unreacted dye was removed with a two-step gel filtration chromatography process. First, the bulk of the free dye was removed by using a HiTrap desalting column (GE Healthcare, Uppsala, Sweden) in 20 mM phosphate buffer (pH 7.4) containing 150 mM NaCl and 30% (v/v) acetonitrile. A subsequent, more rigorous, purification was performed by using a Superdex200 10/300 column (GE Healthcare) in the same buffer. Fractions with no remaining unreacted dye were dialyzed (7000 molecular weight cutoff cassette, Pierce) into water, aliquoted, and lyophilized for storage at  $-80^{\circ}\text{C}$ . Alexa Fluor labeling efficiencies ranged from 31 to 85%, with most variants exhibiting efficiencies between 45 and 55%.

The K39C variant of cytochrome *c* was labeled and purified in the same way by using Alexa Fluor 488 C<sub>5</sub> maleimide (Invitrogen).

**Covalent Aggregation Assays.**  $\alpha$ -Synuclein (100  $\mu\text{M}$ , 9:1 unlabeled:Alexa Fluor-labeled for the wild type), cytochrome *c* (10  $\mu\text{M}$ , 1:3 unlabeled:Alexa Fluor-labeled), and  $\text{H}_2\text{O}_2$  (10

mM) were reacted in phosphate-buffered saline (PBS)<sup>1</sup> [20 mM  $\text{Na}_2\text{HPO}_4$  (pH 7.4) and 150 mM NaCl] at  $37^{\circ}\text{C}$  for 90 min. To compensate for the varying Alexa Fluor labeling efficiencies of  $\alpha$ -synuclein variants, the amount of labeled  $\alpha$ -synuclein was adjusted so that each variant had the same total fluorescence as the wild-type control sample. Control experiments show that this adjustment does not affect our conclusions. For the cytochrome *c* control reaction, 100  $\mu\text{M}$  cytochrome *c* (100% Alexa Fluor-labeled) and 10 mM  $\text{H}_2\text{O}_2$  were reacted under the same conditions described above. Samples were resolved by electrophoresis on 10 to 20% gradient SDS–PAGs (Criterion, Bio-Rad, Hercules, CA) for 75 min at 200 V. Gels were analyzed for fluorescence with a VersaDoc MP imager (Bio-Rad) and for total protein content with Coomassie Blue staining.

**Fibril Formation.**  $\alpha$ -Synuclein (200  $\mu\text{M}$ ) in PBS with 1 mM EDTA was shaken at  $37^{\circ}\text{C}$  and 225 rpm for 48 h. Fibril growth was quantified via thioflavin-T fluorescence in a 96-well plate by using the VersaDoc MP imager. Cytochrome *c* (final concentration of 20  $\mu\text{M}$ ) and  $\text{H}_2\text{O}_2$  (final concentration of 20 mM) were added to the reaction mixture either at the beginning or at the end of the shaking period. Fibrils were separated from  $\alpha$ -synuclein monomers and smaller aggregates by centrifugation at 17000g for 10 min. Fibrils (pellet) and small aggregates (supernatant) were treated with SDS and boiled for 10 min before being analyzed by SDS–PAG electrophoresis as described above.

## RESULTS

**Alexa Fluor Labeling and Tyrosine-Dependent  $\alpha$ -Synuclein Covalent Aggregation.** Labeling  $\alpha$ -synuclein and cytochrome *c* with different dyes, Alexa Fluor 633 (green) and 488 (red), respectively, allows us to follow each protein. The wild type and the variant containing no tyrosines were combined with cytochrome *c* and  $\text{H}_2\text{O}_2$ . Figure 2A shows the results of the reaction between cytochrome *c* and  $\text{H}_2\text{O}_2$  as a standard. As shown in Figure 2B, cytochrome *c* and  $\text{H}_2\text{O}_2$ , as well as the tyrosines in  $\alpha$ -synuclein, are required for oxidative aggregation. This result agrees with our previous work with Coomassie Blue and anti- $\alpha$ -synuclein antibody detection (9). In all peroxidative aggregation reactions, a population of  $\alpha$ -synuclein monomer is left unreacted, even after exposure to additional cytochrome *c* and  $\text{H}_2\text{O}_2$ . This result indicates that some portion of the  $\alpha$ -synuclein is rendered incapable of covalent aggregation after exposure to cytochrome *c* and  $\text{H}_2\text{O}_2$ . This effect has also been seen when human neuroglobin is exposed to peroxide (30). Changing the position

<sup>1</sup> Abbreviations: PAG, polyacrylamide gel; PBS, phosphate-buffered saline; SDS, sodium dodecyl sulfate.




FIGURE 2: Alexa Fluor labeling detects peroxidative aggregation. (A) Cytochrome *c* was incubated with and without H<sub>2</sub>O<sub>2</sub>. (B)  $\alpha$ -Synuclein, either the wild-type protein with the Alexa Fluor label at position 3 (WT, lanes 1–4) or a no tyrosine variant with the Alexa Fluor label at position 3 (noY, lanes 5–8), was combined with various combinations of cytochrome *c* and H<sub>2</sub>O<sub>2</sub> for 90 min, separated on a 10 to 20% gradient polyacrylamide gel, and visualized by fluorescence (green,  $\alpha$ -synuclein; red, cytochrome *c*).




FIGURE 3: Every tyrosine in  $\alpha$ -synuclein is reactive, but to varying extents. (A) Cytochrome *c* was incubated with H<sub>2</sub>O<sub>2</sub>. (B) The wild-type protein (WT) and single tyrosine-containing variants of  $\alpha$ -synuclein were reacted with cytochrome *c* and H<sub>2</sub>O<sub>2</sub> for 90 min and treated as described in the legend of Figure 2.

of the Alexa Fluor label in  $\alpha$ -synuclein from position 3 (Figure 2B, lanes 1–4) to position 66 (data not shown) does not change aggregation. We conclude that Alexa Fluor labels on  $\alpha$ -synuclein and cytochrome *c* allow facile detection of peroxidative covalent aggregation of  $\alpha$ -synuclein.

**Reactivity of the Tyrosines in  $\alpha$ -Synuclein to Covalent Aggregation.** To examine the chemical reactivity of each tyrosine, four variants, each containing only one tyrosine, were reacted with cytochrome *c* and H<sub>2</sub>O<sub>2</sub>. Our conditions do not promote noncovalent aggregation of  $\alpha$ -synuclein. Figure 3A shows the reaction between cytochrome *c* and H<sub>2</sub>O<sub>2</sub> as a standard. Figure 3B shows the differing ability of each tyrosine to form intermolecular dityrosine bonds. Tyrosines 133 and 136 are the most reactive, as shown by distinct dimer formation and little  $\alpha$ -synuclein degradation. Streaking of the  $\alpha$ -synuclein because of random backbone cleavage and large amounts of degraded  $\alpha$ -synuclein indicate that tyrosine 125 is less able to accept a radical from cytochrome *c*. A lack of  $\alpha$ -synuclein dimer and large amounts of degradation indicate that tyrosine 39 is the least reactive. Tyrosine 39 is so unreactive that it does not form

an intact heterodimer with cytochrome *c* as seen in the other variants. Instead, some of the free radicals generated cause degradation of the heterodimer. In summary, the chemical reactivities of the tyrosines in  $\alpha$ -synuclein to form dityrosines increase in the following order: 39, 125, 133/136.

**Tyrosine 39 Is Essential for Wild-Type-like Covalent Aggregation of Native  $\alpha$ -Synuclein.** To determine how each tyrosine contributes to covalent aggregation, seven variants, each containing two or three tyrosines, were reacted with cytochrome *c* and H<sub>2</sub>O<sub>2</sub>. The conditions used do not promote noncovalent aggregation of  $\alpha$ -synuclein. Figure 4A shows the reaction between cytochrome *c* and H<sub>2</sub>O<sub>2</sub> as a standard. Figure 4B shows the participation of each tyrosine in oxidative aggregation. When tyrosine 125, 133, or 136 is removed (Figure 4B, lane 3, 4, or 5, respectively), covalently aggregated species are populated in the same manner as for the wild-type protein (lane 1). When the least reactive tyrosine, tyrosine 39, is removed (Figure 4B, lane 2), even-numbered aggregates (dimer, tetramer, and perhaps hexamer and octamer) are favored over the trimer. In general, more uniform covalent aggregate populations are observed when  $\alpha$ -synuclein has at least one tyrosine at each end. This observation is supported by the data in Figure 4C, where variants with tyrosine 39 and one other tyrosine are examined.

**Changes in Covalent Aggregation Induced by a Denaturant.** If  $\alpha$ -synuclein were totally disordered, one would expect that wild-type-like covalent aggregation could be achieved if any two tyrosines are present. However, as shown in Figure 4, tyrosine 39 is essential for wild-type-like covalent aggregation. To determine if protein conformation was the basis of this observation, wild-type  $\alpha$ -synuclein or the variant with tyrosine 39 removed was reacted with cytochrome *c* and H<sub>2</sub>O<sub>2</sub> in the presence and absence of the denaturant, guanidine hydrochloride (300 mM). A control experiment (not shown) illustrates that 300 mM guanidine hydrochloride does not affect radical formation as assessed by the bands formed in cytochrome *c* covalent aggregation with and without guanidine hydrochloride. Coomassie Blue staining was used because guanidine hydrochloride interferes with Alexa Fluor fluorescence. In the presence of guanidine hydrochloride (Figure 5B,C, lane 2), large covalent aggregates are more heavily favored for both wild-type and Y39F  $\alpha$ -synuclein compared to the reaction in the absence of guanidine hydrochloride (Figure 5B,C, lane 1). In agreement with NMR-based experiments, these data show that  $\alpha$ -synuclein must have some structure that can be destroyed by a denaturant (11).

**Fibril Formation and Oxidative Aggregation.** To determine if covalent aggregation affects the conformation required for noncovalent fibril formation, fibrils were grown from the wild-type protein. Coomassie Blue staining was used because the fibril growth conditions interfere with Alexa Fluor fluorescence. As a control, either cytochrome *c* or H<sub>2</sub>O<sub>2</sub> was added either at the start or at the end of fibril growth. Under each of these conditions, no covalent aggregates were detected in the supernatant or in the fibrils (data not shown). Figure 6B shows the effects of exposing the fibrillization reaction mixture to cytochrome *c* and H<sub>2</sub>O<sub>2</sub> at the beginning or end of fibril growth. When peroxidation is introduced after fibril growth for 24 h (lanes 3 and 4), the covalent aggregates are observed in both the supernatant and the fibrils, and a






FIGURE 4: At least one tyrosine on each end is required for wild-type-like covalent aggregation of collapsed  $\alpha$ -synuclein. (A) Cytochrome *c* was incubated with  $H_2O_2$ . (B) The wild-type protein and variants containing three tyrosines or (C) two tyrosines were reacted with cytochrome *c* and  $H_2O_2$  for 90 min and treated as described in the legend of Figure 2. The legend at the top indicates which tyrosines are present in each variant.  $\alpha$ -Synuclein aggregate species are indicated on the right.




FIGURE 5: Tyrosine 39 is not required for full aggregation of denatured  $\alpha$ -synuclein. (A) Molecular weight markers. (B) The wild-type protein (WT) and (C) the variant with tyrosine 39 removed (Y39F) were reacted with cytochrome *c* and  $H_2O_2$  in the presence (lanes 2) or absence (lanes 1) of guanidine hydrochloride. Samples were separated on a 10 to 20% gradient PAG and visualized by Coomassie Blue staining.

large fibril pellet is observed. When peroxidation is introduced early, almost no fibrils are formed (as indicated by a miniscule pellet) and there is a nearly complete absence of covalent aggregates (lane 2). Instead, nearly all of the covalent aggregates are found in the supernatant (lane 1). Similar results were obtained for the variant containing only




FIGURE 6: Oxidative aggregation interferes with fibril formation. (A) Molecular weight markers. (B) The wild-type protein was reacted with cytochrome *c* and  $H_2O_2$  before (lanes 1 and 2) or after (lanes 3 and 4) fibril formation. Fibrils were isolated by centrifugation. Fibrils (lanes 2 and 4) and the supernatant (lanes 1 and 3) were boiled with SDS, separated on a 10 to 20% gradient PAG, and visualized with Coomassie Blue.

the tyrosine 39 to phenylalanine mutation (although the yield of fibrils is lower).

DISCUSSION

These studies examine the role of  $\alpha$ -synuclein conformation in its peroxidative aggregation. The conformations of




FIGURE 7:  $\alpha$ -Synuclein conformation and covalent aggregation. When  $\alpha$ -synuclein adopts a collapsed conformation, various covalently aggregated species are more equally populated when at least one tyrosine is present on each end of the protein (A). Even-numbered aggregates are favored when tyrosine 39 is removed (B and C). When  $\alpha$ -synuclein is completely disordered, the largest aggregate species are favored, even if tyrosine 39 is removed (D). Covalent  $\alpha$ -synuclein aggregates are unable to fold into the  $\beta$ -sheet aggregation nucleus, but once the folded monomers have assembled into protofibrils, the termini are stacked, allowing covalent aggregation of any of the tyrosines (E).

this intrinsically disordered protein have been characterized by NMR, and the shape shown in Figure 1B is just one of an ensemble of collapsed conformations adopted by  $\alpha$ -synuclein (11). While the protein does not have a rigid secondary structure, these conformations have similar characteristics, namely, that the charged termini collapse onto the hydrophobic core. In each conformation, the region surrounding tyrosine 39 abuts the core and C-terminus. This contact keeps tyrosine 39 partially protected from the solvent, which accounts for its low reactivity (Figure 3). NMR-detected amide proton exchange experiments also show that tyrosine 39 is protected from the solvent, as indicated by its low exchange rate (31). In each conformation, the region surrounding tyrosine 125 contacts three other regions (residues 11–13, 30–75, and 78–99) (12). Tyrosine 125 is in a region of increased rigidity compared to the rest of the protein (11, 12, 32), although the C-terminus as a whole is comparatively mobile. This synergy of rigidity and flexibility accounts for the decreased reactivity of tyrosine 125 compared to tyrosines 133 and 136, and its increased reactivity compared to tyrosine 39 (Figure 3).

The collapsed conformation of  $\alpha$ -synuclein explains why at least one tyrosine on each end of the protein is required to maintain a distribution of covalent aggregates similar to that observed for the wild-type protein (Figure 7A). When C-terminal tyrosines from two  $\alpha$ -synuclein monomers react, the collapsed conformation sterically hinders the two remaining C-terminal tyrosines from reacting (Figure 7B,C). Without a tyrosine near the N-terminus, i.e., tyrosine 39, dimers are favored (Figure 7B), but once a sufficient population of dimers accumulates, they tend to form tetramers, hexamers, and higher-order even-numbered aggregates (Figure 7C). The lower reactivities of tyrosines 39 and 125 (Figure 3) explain why variants with only these tyrosines (Figure 4C, lane 1) tend to form fewer higher-order covalent aggregates. When  $\alpha$ -synuclein is denatured with guanidine hydrochloride, an N-terminal tyrosine is no longer needed for wild-type-like covalent aggregation (Figure 7D) because the three C-terminal tyrosines are more accessible.

$\alpha$ -Synuclein must undergo a folding event to form an aggregation nucleus (23) before forming fibrils (Figure 7E). Covalently cross-linking the protein's tyrosines prevents it

from folding properly, so that, as shown in Figure 6, covalent aggregates are not incorporated into fibrils. In addition, when oxidation is introduced at the start of the fibrillization process, little to no fibril growth is observed because the concentration of monomer is too low to allow fibrillization in the time frame of the experiment. Once fibrils have assembled (Figure 7E), however, the termini are stacked (33) so that the tyrosines are solvent-exposed and accessible for oxidative cross-linking.<sup>2</sup> While tyrosine 39 is essential for fibril formation (34), an N-terminal tyrosine is not required for covalent aggregation within fibrils. This observation is similar to what is observed when the protein is denatured by guanidine hydrochloride. In summary, our data indicate that both covalent and noncovalent aggregation of  $\alpha$ -synuclein depend on its conformation and suggest that covalent aggregation occurs in Lewy bodies after the formation of fibrils.

## ACKNOWLEDGMENT

We thank the members of the Pielak group for insightful comments. We thank Sandy An and Melanie Wiley for technical assistance.

## REFERENCES

- Hague, S. M., Klaffke, S., and Bandmann, O. (2005) Neurodegenerative disorders: Parkinson's disease and Huntington's disease. *J. Neurol., Neurosurg. Psychiatry* 76, 1058–1063.
- National Parkinson's Foundation.
- Spillantini, M. G., Schmidt, M. L., Lee, V. M. Y., Trojanowski, J. Q., Jakes, R., and Goedert, M. (1997)  $\alpha$ -Synuclein in Lewy bodies. *Nature* 388, 839–840.
- Hashimoto, M., Takeda, A., Hsu, L. J., Takenouchi, T., and Masliah, E. (1999) Role of cytochrome *c* as a stimulator of  $\alpha$ -synuclein aggregation in Lewy body disease. *J. Biol. Chem.* 274, 28849–28852.
- D'Andrea, M. R., Ilyin, S., and Plata-Salaman, C. R. (2001) Abnormal patterns of microtubule-associated protein-2 (map-2) immunolabeling in neuronal nuclei and Lewy bodies in Parkinson's disease substantia nigra brain tissues. *Neurosci. Lett.* 306, 137–140.
- Gai, W. P., Power, J. H. T., Blumbergs, P. C., Culvenor, J. G., and Jensen, P. H. (1999)  $\alpha$ -Synuclein immunolocalization of glial inclusions from multiple system atrophy brain tissue reveals multiprotein components. *J. Neurochem.* 73, 2093–2100.
- Goldman, J. E., Yen, S. H., Chiu, F. C., and Peress, N. S. (1983) Lewy bodies of Parkinson's disease contain neurofilament antigens. *Science* 221, 1082–1084.
- Uversky, V. N. (2007) Neuropathology, biochemistry, and biophysics of  $\alpha$ -synuclein aggregation. *J. Neurochem.* 103, 17–37.
- Olteanu, A., and Pielak, G. J. (2004) Peroxidative aggregation of  $\alpha$ -synuclein requires tyrosines. *Protein Sci.* 13, 2852–2856.
- Morar, A. S., Olteanu, A., Young, G. B., and Pielak, G. J. (2001) Solvent-induced collapse of  $\alpha$ -synuclein and acid-denatured cytochrome *c*. *Protein Sci.* 10, 2195–2199.
- Bertoncini, C. W., Jung, Y. S., Fernandez, C. O., Hoyer, W., Greisinger, C., Jovin, T. M., and Zweckstetter, M. (2005) Release of long range tertiary interactions potentiates aggregation of natively unstructured  $\alpha$ -synuclein. *Proc. Natl. Acad. Sci. U.S.A.* 102, 1430–1435.
- Dedmon, M. M., Lindorff-Larsen, K., Christodoulou, J., Vendruscolo, M., and Dobson, C. M. (2005) Mapping long-range interactions in  $\alpha$ -synuclein using spin-label NMR and ensemble molecular dynamics simulations. *J. Am. Chem. Soc.* 127, 476–477.
- McNulty, B. C., Tripathy, A., Young, G. B., Orans, J., and Pielak, G. J. (2005) Temperature-induced reversible conformational change in the first 100 residues of  $\alpha$ -synuclein. *Protein Sci.* 15, 602–608.
- Schults, C. W. (2006) Lewy bodies. *Proc. Natl. Acad. Sci. U.S.A.* 103, 1661–1668.
- Pannathur, S., Jackson-Lewis, V., Przedborski, S., and Heinecke, J. W. (1999) Mass spectrometric quantification of 3-nitrotyrosine, ortho-tyrosine, and o,o'-dityrosine in brain tissue of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-treated mice, a model of oxidative stress in Parkinson's disease. *J. Biol. Chem.* 274, 34621–34628.
- Atwood, C. S., Perry, G., Zeng, H., Kato, Y., Jones, W. D., Ling, K., Huang, X., Moir, R. D., Wang, D., Sayre, L. M., Smith, M. A., Chen, S. G., and Bush, A. I. (2004) Copper mediates dityrosine cross-linking of Alzheimer's amyloid  $\beta$ . *Biochemistry* 43, 560–568.
- Norris, E. H., Giasson, B. I., Ischiropoulos, H., and Lee, V. M. (2003) Effects of oxidative and nitrate challenges on  $\alpha$ -synuclein fibrillogenesis involve distinct mechanisms of protein modifications. *J. Biol. Chem.* 278, 27230–27240.
- Barr, D. P., Gunther, M. R., Deterding, L. J., Tomer, K. B., and Mason, R. P. (1996) ESR spin-trapping of a protein-derived tyrosyl radical from the reaction of cytochrome *c* with hydrogen peroxide. *J. Biol. Chem.* 271, 15498–15503.
- Deterding, L. J., Barr, D. P., Mason, R. P., and Tomer, K. B. (1998) Characterization of cytochrome *c* free radical reaction with peptides by mass spectrometry. *J. Biol. Chem.* 273, 12963–12969.
- Fink, A. L. (2006) The aggregation and fibrillation of  $\alpha$ -synuclein. *Acc. Chem. Res.* 39, 628–634.
- Nielsen, L., Khurana, R., Coats, A., Frokjaer, S., Brange, J., Vyas, S., Uversky, V. N., and Fink, A. L. (2001) Effect of environmental factors on the kinetics of insulin fibril formation: Elucidation of the molecular mechanism. *Biochemistry* 40, 6036–6046.
- Lee, C. C., Nayak, A., Sethuraman, A., Belfort, G., and McRae, G. J. (2007) A three-stage kinetic model of amyloid fibrillation. *Biophys. J.* 92, 3448–3458.
- Shahi, P., Sharma, R., Sanger, S., Kumar, I., and Jolly, R. S. (2007) Formation of amyloid fibrils via longitudinal growth of oligomers. *Biochemistry* 46, 7365–7373.
- Lomakin, A., Chung, D. S., Benedek, G. B., Kirschner, D. A., and Teplow, D. B. (1996) On the nucleation and growth of amyloid  $\beta$  protein fibrils: Detection of nuclei and quantitation of rate constants. *Proc. Natl. Acad. Sci. U.S.A.* 93, 1125–1129.
- LeVine, H. (1999) Quantification of  $\beta$ -sheet amyloid fibril structures with thioflavin T. *Methods Enzymol.* 309, 274–284.
- Conway, K. A., Lee, S. J., Rochet, J. C., Ding, T. T., Williamson, R. E., and Lansbury, P. T., Jr. (2000) Acceleration of oligomerization, not fibrillization, is a shared property of both  $\alpha$ -synuclein mutations linked to early-onset Parkinson's disease: Implications for pathogenesis and therapy. *Proc. Natl. Acad. Sci. U.S.A.* 97, 571–576.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L., and Randall, R. J. (1951) Protein measurement with the folin phenol reagent. *J. Biol. Chem.* 193, 265–275.
- Olteanu, A., Patel, C. N., Dedmon, M. M., Kennedy, S., Linhoff, M. W., Minder, C. M., Potts, P. R., Deshmukh, M., and Pielak, G. J. (2003) Stability and apoptotic activity of recombinant human cytochrome *c*. *Biochem. Biophys. Res. Commun.* 312, 733–740.
- Margoliash, E., and Frohvirt, N. (1959) Spectrum of horse-heart cytochrome *c*. *Biochem. J.* 71, 570–572.
- Lardinois, O. M., Tomer, K. B., Mason, R. P., and Deterding, L. J. (2008) Identification of protein radicals formed in the human neuroglobin-H<sub>2</sub>O<sub>2</sub> reaction using immuno-spin trapping and mass spectrometry. *Biochemistry* 47, 10440–10448.
- Croke, R. L., Sallum, C. O., Watson, E., Watt, E. D., and Alexandrescu, A. T. (2008) Hydrogen exchange of monomeric  $\alpha$ -synuclein shows unfolded structure persists at physiological temperature and is independent of molecular crowding in *Escherichia coli*. *Protein Sci.* 17, 1434–1445.
- Bussell, R., Jr., and Eliezer, D. (2001) Residual structure and dynamics in Parkinson's disease associated mutants of  $\alpha$ -synuclein. *J. Biol. Chem.* 276, 45996–46003.
- Vilar, M., Chou, H.-T., Luhrs, T., Maji, S. K., Riek-Loher, D., Verel, R., Manning, G., Stahlberg, H., and Riek, R. (2008) The fold of  $\alpha$ -synuclein fibrils. *Proc. Natl. Acad. Sci. U.S.A.* 105, 8637–8642.
- Ulrich, N. P., Barry, C. H., and Fink, A. L. (2008) Impact of Tyr to Ala mutations on  $\alpha$ -synuclein fibrillation and structural properties. *Biochim. Biophys. Acta* 1782, 581–585.

<sup>2</sup> Studies are underway to determine if the covalent aggregation observed when H<sub>2</sub>O<sub>2</sub> and cytochrome *c* are reacted with fibrils is tyrosine-dependent.