

Kinetic Mechanism for Single-Stranded DNA Binding and Translocation by *Saccharomyces cerevisiae* Isw2[†]

Christopher J. Fischer,^{*,‡} Kazuhiro Yamada,[§] and Daniel J. Fitzgerald[§]

Department of Physics and Astronomy, University of Kansas, 1251 Wescoe Hall Drive, 1082 Malott Hall, Lawrence, Kansas 66045, and Institut für Molekularbiologie und Biophysik, ETH-Hönggerberg HPK, Zürich, Switzerland

Received November 17, 2008; Revised Manuscript Received February 6, 2009

ABSTRACT: The chromatin remodeling complex Isw2 from *Saccharomyces cerevisiae* (yIsw2) mobilizes nucleosomes through an ATP-dependent reaction that is coupled to the translocation of the helicase domain of the enzyme along intranucleosomal DNA. In this study, we demonstrate that yIsw2 is capable of translocating along single-stranded DNA in a reaction that is coupled to ATP hydrolysis. We propose that single-stranded DNA translocation by yIsw2 occurs through a series of repeating uniform steps with an overall macroscopic processivity (P) of 0.90 ± 0.02 , corresponding to an average translocation distance of 20 ± 2 nucleotides before dissociation. This processivity corresponds well to the processivity of nucleosome sliding by yIsw2, thus arguing that single-stranded DNA translocation or tracking may be fundamental to the double-stranded DNA translocation required for effective nucleosome mobilization. Furthermore, we find evidence that a slow initiation process, following DNA binding, may be required to make yIsw2 competent for DNA translocation. We also provide evidence that this slow initiation process may correspond to the second step of a two-step DNA binding mechanism by yIsw2 and a quantitative description of the kinetics of this DNA binding mechanism.

Members of the ISWI family of chromatin remodeling enzymes (termed remodelers) hydrolyze ATP to reposition nucleosomes along DNA (1–3). The ATP-hydrolyzing subunits possessed by members of this and other families of remodelers have significant homology to those of the helicase family of proteins (4–6). In fact, remodelers are classified as members of the SF2 superfamily of helicases on the basis of amino acid sequence comparison (7), although remodelers lack helicase activity (8). However, the ATP-dependent DNA translocation activity within the helicase derivative domains of remodelers is similar to that which is observed in helicases (9–18). This DNA translocation property plays a central role in certain models for ATP-dependent chromatin remodeling by members of both the ISWI and SWI/SNF subfamilies of chromatin remodeling enzymes (2, 4, 14, 15, 19–21). In one such model, the ATPase subunit of the remodeling enzyme translocates directionally along the DNA at an internal site on the nucleosome. This ATP-dependent DNA translocation results in the movement of the DNA around the nucleosome by drawing in DNA from one side of the nucleosome while simultaneously pumping it out the other side. More specif-

ically, the translocation results in the formation of a loop or bulge of DNA that then propagates around the surface of the octamer (4, 14, 19, 22, 23), thus resulting in a translation of the octamer relative to the DNA. The size of the propagated loop of DNA is believed to be related to the kinetic step size of DNA translocation by the remodeler, and estimates of it vary between enzymes: 1 bp for *Saccharomyces cerevisiae* RSC (19), 2 bp for *S. cerevisiae* Isw2 (2, 20), and ~50 bp for *S. cerevisiae* SWI/SNF (20).

yIsw2 comprises a heterodimer of the Itc1p (145 kDa) and Isw2p (130 kDa) proteins and is a representative member of the ISWI subfamily of the SWI2/SNF2 superfamily of chromatin remodeling proteins (21, 24); both Itc1 and Isw2p are required for the chromatin remodeling activity of the holocomplex (25). In addition, two small histone fold-containing proteins, Dpb4p and Dls1p, are present in at least a fraction of Isw2 complexes purified from yeast (26). yIsw2 has been demonstrated to affect in vivo repression of transcription of several genes (3, 27–29) and, together with Ino80, to promote replication fork progression (30). It has been suggested that yISW2 accomplishes these activities by modifying the spacing of sequential mononucleosomes along short, contiguous stretches of chromatin through nucleosome sliding (24, 31). Interestingly, yIsw2 is able to accomplish this sliding of nucleosomes without disrupting the integrity of the association of the DNA and the octamer (1, 32).

The ATPase activity of yIsw2 is stimulated in the presence of double-stranded DNA (21, 24); this is consistent with the hypothesis that yIsw2 is a double-stranded DNA translocase. As mentioned above, double-stranded DNA translocation by yIsw2 is also required for models of nucleosome mobilization by the enzyme (20, 21, 33). The ability of yIsw2 to slide

[†] This research was supported, in part, by startup funding from the University of Kansas (to C.J.F.) and by NIH Grant P20 RR17708 from the Institutional Development Award (IDeA) Program of the National Center for Research Resources (to C.J.F.).

* To whom correspondence should be addressed: Department of Physics and Astronomy, University of Kansas, 1251 Wescoe Hall Dr., 1082 Malott Hall, Lawrence, KS 66045. Telephone: (785) 864-4579. Fax: (785) 864-5262. E-mail: shark@ku.edu.

[‡] University of Kansas.

[§] ETH-Hönggerberg HPK.

¹ Abbreviations: NLLS, nonlinear least-squares; LLS, linear least-squares.

nucleosomes with a specific directional bias has led to the suggestion that DNA translocation by γ IsW2 is also directionally biased (2, 20, 32, 33). Furthermore, the ability of γ IsW2 to slide nucleosomes when either single- or double-stranded DNA is present in the extranucleosomal linker suggests that γ IsW2 is both a single- and double-stranded DNA translocase (33). Interestingly, the genetically related *Drosophila melanogaster* ISWI enzyme has been shown to be a 3' to 5' single-stranded DNA translocase. Taken together, this suggests that it is the single-stranded, directionally biased DNA translocation activity which is fundamental to the double-stranded DNA translocation activity of γ IsW2 and, by extension, to its nucleosome mobilization activity.

In spite of this, there has not yet been a quantitative description of the single- or double-stranded DNA translocation mechanism of γ ISW2. It is worth noting, however, that although kinetic studies of the single-stranded DNA translocation activity of the *D. melanogaster* ISWI enzyme have been published, an associated quantitative description of the mechanism of translocation was not presented (15). The results presented here thus represent the first quantitative characterization of the kinetic mechanism of single-stranded DNA translocation by a chromatin remodeling enzyme and, furthermore, offer insight not only into the relationship between single-stranded DNA binding and DNA translocation by γ IsW2 but also into the possible mechanistic similarities between subfamilies of chromatin remodeling enzymes.

MATERIALS AND METHODS

Buffers. Reaction buffers were prepared with reagent-grade chemicals using twice distilled water that was subsequently deionized with a Milli-Q purification system (Millipore Corp., Bedford, MA). All ATPase and DNA binding reactions were carried out in 10 mM HEPES (pH 7.0), 5% glycerol, 20 mM potassium acetate, 5 mM MgCl_2 , 0.5 mM dithiothreitol, and 0.1 mg/mL BSA at 37 °C.

ISW2 Protein and DNA. γ ISW2 was expressed and purified as described previously (20). Oligonucleotides were purchased from Integrated DNA Technologies (Coralville, IA) dialyzed extensively into Milli-Q water before use. Poly(dT) was purchased from Midland Reagent Co. (Midland, TX) and dialyzed extensively into Milli-Q water before use.

Radioactive ATPase Experiments. ATPase activity was determined by measuring the rate of conversion of ATP to ADP using [α - ^{32}P]ATP. All reactions were conducted at 37 °C; 140 nM ISW2 and 100 μM (nucleotide) single-stranded DNA were incubated together on ice for 5 min prior to the initiation of the translocation reaction. DNA translocation was initiated through the addition of ATP to the solution containing the preformed complexes of γ ISW2 and DNA. The final concentration of ATP was 1 mM, and the final reaction volume was 30 μL . Aliquots (4 μL) of the resulting solution were removed every 6 min and reactions quenched by adding an equal volume of 0.5 M EDTA. The extent of product formation at each time point was monitored by spotting 2 μL samples onto polyethyleneimine (PEI) cellulose TLC plates (Merck KGaA, Darmstadt, Germany) which were developed using 0.6 M KPO_4 (pH 3.4) and quantitatively

analyzed using a Bio-Rad (Hercules, CA) Molecular Imager FX. Spots corresponding to the radiolabeled ATP and ADP were quantified using the Quantity One software supplied by the manufacturer.

Stopped-Flow Experiments. Measurements of the production of P_i in the stopped-flow spectrometer were performed using the EnzChek phosphate assay kit (Invitrogen, Carlsbad, CA); this assay was first described by Webb (42). In the presence of P_i , the substrate 2-amino-6-mercapto-7-methylpurine riboside (MESG) is converted enzymatically by purine nucleoside phosphorylase (PNP) to ribose 1-phosphate and 2-amino-6-mercapto-7-methylpurine. The enzymatic conversion of MESG results in a spectrophotometric shift in the maximum absorbance from 330 nm for the substrate to 360 nm for the product which is monitored in the stopped-flow spectrometer. The k_{cat} for the reaction is 40 s^{-1} , which is sufficiently fast for monitoring the kinetics of ATPase by the translocating γ ISW2 complex. For the experiments shown in Figure 4, the concentration of γ ISW2 was 25 nM and the concentration of poly(dT) was 100 μM (nucleotide).

Measurements of the kinetics of γ ISW2 binding to dT₇₀ were performed in the stopped-flow spectrometer by monitoring the change in the intrinsic tryptophan fluorescence of γ ISW2 that occurs upon DNA binding. In these experiments, the tryptophan fluorescence was excited at 280 nm and emission monitored at wavelengths greater than 305 nm using a WG 305 Schott glass filter. BSA was omitted from these reaction mixtures since its tryptophan fluorescence created an overly large background signal. The final reaction concentration of γ ISW2 was 10 nM.

Analysis of Kinetic Data. The ATPase rate for each length of single-stranded DNA was determined from LLS¹ analysis of the corresponding ATPase time courses. Four independent measurements of V_{max} were taken for each DNA length.

All LLS and NLLS analyses were performed using Conlin, kindly provided by J. Williams. Fitting models and Monte Carlo simulation programs were written in the C++ computer language and compiled with the Microsoft C++ 6.0 compiler. The software library CNL50 (Visual Numerics Inc., Houston, TX) was used for the numerical calculation of the inverse Laplace transform. The uncertainties in all fitted parameters reported in this work represent 68% confidence limits (one standard deviation) and were determined by performing a 1000-cycle Monte Carlo simulation using the routine built into Conlin.

RESULTS

γ ISW2 Is an ATP-Dependent Single-Stranded DNA Translocase. We studied the kinetics of translocation of γ IsW2 along single-stranded DNA by analyzing the stimulation of γ IsW2's ATPase activity in the presence of single-stranded DNA of varying lengths. ATPase experiments were performed as described in Materials and Methods under solution conditions identical to those previously used to study the DNA and nucleosome stimulating ATPase activity of γ IsW2 (21). To minimize complications arising from secondary structure in the DNA, we performed our experiments using oligodeoxythmidylates. To ensure that our experiments were performed at DNA concentrations sufficient to elicit maximum steady-state ATPase stimulation, we first determined the K_M for the single-stranded DNA-stimulated ATPase




FIGURE 1: Time courses of ATP hydrolysis for yIsw2 in the presence of a saturating concentration of single-stranded DNA of varying length. In these experiments, yIsw2 and single-stranded DNA were incubated together before the addition of ATP. The lengths of the DNA are 16, 40, 60, and 100 nucleotides. A control experiment in which no DNA is included in the reaction is also shown.




FIGURE 2: Dependence of V_{\max} upon single-stranded DNA length. Each data point is the average of four separate measurements of V_{\max} , and the error bars represent the standard deviation of those measurements. The solid line is a NLLS fit of the data to eq 2 (Scheme 1). The NLLS fits using eqs 10 and 12 are identical. The parameter estimates associated with these fits are listed in Tables 1 and 2.

activity of yIsw2. We estimate K_M equals $5 \pm 2 \mu\text{M}$ (mononucleotide concentration), independent of the length of the DNA (data not shown).

We subsequently measured the stimulation of yIsw2's ATPase activity in the presence of DNA of varying lengths, but at a constant mononucleotide concentration (14, 18) of $100 \mu\text{M}$ (20-fold larger than K_M). As shown in Figures 1 and 2, we observe that the steady-state ATPase rate of yIsw2 increases with an increase in DNA length. These results are consistent with the previously published hypothesis that yIsw2 is a single-stranded DNA translocase (33). Furthermore, since these experiments were conducted under conditions of excess DNA concentration, the binding of single complexes of yIsw2 to the DNA is favored. This is further supported by equilibrium studies of DNA binding by yISW2 experiments which demonstrated that the occluded site size of yISW2 binding to DNA is larger than 100 bp (D. J. Fitzgerald et al., unpublished observations). Therefore, taken together, the results suggest that monomeric yIsw2 is functional as a single-stranded DNA translocase.

Kinetic Models for Single-Stranded DNA Translocation. The dependence of the V_{\max} of the single-stranded DNA-stimulated ATPase activity of yISW2 on DNA length shown

Scheme 1




Table 1: Estimates of Kinetic Parameters for Translocation of yIsw2 along Single-Stranded DNA According to Scheme 1 (eq 2)

P	0.90 ± 0.02
ck_t	$61 \pm 4 \text{ ADP yIsw2}^{-1} \text{ min}^{-1}$ (fixed)
d	$9.85 \pm 0.09 \text{ nucleotides}$
$m/(1 - P)$	$20 \pm 2 \text{ nucleotides}$
variance of fit	0.65


in Figure 2 is consistent with yIsw2 acting as a single-stranded DNA translocase. The simplest model for single-stranded DNA translocation by yIsw2 which is consistent with the data in Figures 1 and 2 is shown in Scheme 1. In this model, we assume that the yIsw2 complex binds randomly to one of its possible binding sites on the single-stranded DNA. Upon binding, the yIsw2 complex is i translocation steps away from the end of the DNA (the T_i state in Scheme 1). With subsequent repeating cycles of ATP binding, ATP hydrolysis, and release of ADP and inorganic phosphate, the yIsw2 complex translocates along the single-stranded DNA with directional bias (33) and finite processivity [$P = k_t/(k_d + k_t)$]. This translocation occurs in discrete steps of m nucleotides per step, with an associated translocation rate constant k_t and c molecules of ATP hydrolyzed per step; the constant c is thus the thermodynamic coupling efficiency for the process associated with the k_t rate constant, and m is the kinetic step size (34, 35). The rate constant for dissociation during translocation is k_d . The maximum number of translocation steps, n , for a given length of DNA, L , is related to the kinetic step size of translocation, m , through the approximate relation $n = (L - m)/d$, where d is the interaction site size of the protein (18, 34). Upon reaching the end of the DNA (the T_0 state in Scheme 1), the yIsw2 complex can no longer translocate and so dissociates with a rate constant k_d . We further assume that since our experiments are performed under conditions of excess DNA concentration there will be only one yIsw2 complex bound per DNA, and thus, we have ignored any potential protein-protein interactions.

All dissociated protein (and protein that was initially free in solution at the start of the reaction) will bind the DNA randomly at any available binding sites. The second-order rate constant for yIsw2 binding single-stranded DNA (and forming complex T_i) is k_1 . It is worth noting that we assume the binding affinity is the same for all binding sites, and thus, we ignore electrostatic effects associated with the ends of the DNA. Furthermore, in our experiments, the concentration of DNA is much larger than the concentration of protein, so we can treat the binding of yIsw2 to DNA as a pseudo-first-order process with associated rate constant $k_1[D]$, where $[D]$ is the nucleotide concentration of DNA in solution.

Table 2: Estimates of Kinetic Parameters for Translocation of yIsw2 along Single-Stranded DNA According to Scheme 2 (eq 10)

P	0.90 ± 0.02	0.90 ± 0.02
$ck_t[k_{np}/(k_d + k_{np})]$	$61 \pm 4 \text{ ADP yISW2}^{-1} \text{ min}^{-1} \text{ (fixed)}$	$61 \pm 4 \text{ ADP yISW2}^{-1} \text{ min}^{-1} \text{ (fixed)}$
$c_{np}k_d[k_{np}/(k_d + k_{np})]$	$0 \text{ ADP yISW2}^{-1} \text{ min}^{-1} \text{ (fixed)}$	$0.2 \pm 0.2 \text{ ADP yISW2}^{-1} \text{ min}^{-1}$
d	$9.85 \pm 0.09 \text{ nucleotides}$	$10 \text{ nucleotides (fixed)}$
$m/(1 - P)$	$20 \pm 2 \text{ nucleotides}$	$20 \pm 2 \text{ nucleotides}$
variance of fit	0.65	0.65

Scheme 2



It is also important to note that we have neglected any contributions from the potential futile hydrolysis of ATP by yIsw2 bound at the ends of the DNA. Such futile hydrolysis has been observed for helicase translocation along single-stranded DNA (35), but we cannot determine if yIsw2 is exhibiting similar behavior on the basis of our data. It is worth noting, however, that the contribution of this futile hydrolysis (if occurring) would decrease with an increase in DNA length (18), making its contribution to the dependence of the total observed steady-state ATPase rate on DNA length negligible.

The differential equations describing the concentration of protein bound at each position (T_i and free protein) in Scheme 1 can be solved to yield time-dependent equations for these protein concentrations that are functions of the maximum number of translocation steps, n , associated with a given DNA length; this is equivalent to finding expressions for these populations as a function of the length of the DNA and the kinetic step size of translocation (11, 18, 34). In this derivation, we assume an initial equilibrium between the bound and free complexes of yIsw2, as determined by k_1 , $[D]$, and k_d , exists before the addition of ATP so that it matches our experimental conditions (see Materials and Methods). These expressions can then be combined to form a time-dependent expression (eq 1) for the concentration of ATP hydrolyzed (or the concentration of ADP or P_i produced) during the translocation of yIsw2 along single-stranded DNA.

$$\text{ADP}(t) = L^{-1} \left[\frac{k_1[D]}{k_d + k_1[D]} \left(\frac{ck_t}{(n+1)s^2} \left\{ n + \frac{k_t}{k_d + s} \left[\left(\frac{k_t}{k_t + k_d + s} \right)^n - 1 \right] \right\} \right) \right] \quad (1)$$

Simulated time courses based upon this model are shown in Figure 3. As seen in Figure 3, these time courses consist of an initial rapid exponential increase in the amount of ADP produced (ATP hydrolyzed) by the translocating proteins, associated with their pre-steady-state translocation on the single-stranded DNA, followed by a linear phase corresponding to their steady-state translocation activity. The rate of the steady-state ATP hydrolysis by these translocating enzymes obeys Michaelis–Menten kinetics with V_{\max} and K_M shown in eqs 2 and 3, respectively.




FIGURE 3: Simulated ATPase time courses for protein translocating along single-stranded DNA of varying lengths (3, 10, and ∞ nucleotides) under the conditions of a saturating concentration of DNA. In these simulations, $k_t = 30$ steps per second, $k_d = 2 \text{ s}^{-1}$, $m = 1$ nucleotide per step, and $c = 1$ ATP per step. (A) Simulations according to Scheme 1. (B) Simulations according to Scheme 2 with a k_{np} of 0.01 s^{-1} .


$$V_{\max} = \frac{ck_t[n(P-1) - P(P^n-1)]}{(n+1)(P-1)} \quad (2)$$

$$K_M = \frac{k_d}{k_1} \quad (3)$$

The variable P in eq 2 is the previously defined processivity of translocation. It is interesting to note that this model (and corresponding eq 3) predicts that K_M is independent of the maximum number of steps (n), or equivalently independent of the length of the DNA. This is consistent with measurements of K_M of yIsw2 translocating along single-stranded DNA.

To obtain a better estimate of the kinetic parameters from our NLLS analysis of the data in Figure 2, we also measured the ATPase time course for the translocation of yIsw2 along poly(dT). Poly(dT) can effectively mimic infinitely long DNA, and thus, the value of V_{\max} determined for the translocation of yIsw2 along poly(dT) can serve as a constraint in the analysis of the data in Figure 2 using eq 2 (18). Specifically, in the limit of infinitely long DNA, eq 2 simplifies to eq 4.

Scheme 3



We determined $V_{\max, \infty} = 61 \pm 4$ ADP yISW2⁻¹ min⁻¹ from the analysis of the ATPase time courses obtained in the presence of poly(dT). We then determined estimates of P and d through NLLS analysis of the dependence of V_{\max} on DNA length (Figure 2) using eq 2 with our estimate of $V_{\max, \infty}$ as a fixed constraint. In this analysis, we further assumed that $m = 2$ nucleotides since an estimate of 2 nucleotides for the kinetic step size of double-stranded DNA translocation by yISW2 was recently reported (20). The parameter estimates obtained from this analysis are presented in Table 1, and the solid line overlaying the data in Figure 2 is a simulation using eq 2 and these parameter estimates.


As described above and shown in Figure 3, the ATPase time courses associated with Scheme 1 contain an initial exponential phase of activity associated with the pre-steady-state translocation of the proteins along the DNA. It is worth noting that within the resolution of our measurements we could not detect this phase; rather, we determine a zero intercept, within error, on the ordinate of the time courses shown in Figure 1. A simple explanation for this discrepancy would be that the rate of dissociation from the end of the DNA (from the T_0 state) is much faster than k_t and not equal to k_d . Such a kinetic mechanism would be consistent with the lack of any observed pre-steady-state translocation (and corresponding ATPase) activity of the yISW2 complex; however, this mechanism would also suggest that there would be no dependence of V_{\max} upon DNA length. Since we do see a dependence of V_{\max} on DNA length, we can reject this hypothesis.

To improve the time resolution of our measurements of the ATPase activity of translocating yISW2 and thus also specifically improve the resolution of the pre-steady-state kinetics of translocation, we next performed single-stranded DNA translocation experiments using a stopped-flow technique which monitors the production of P_i produced by yISW2 during translocation. P_i in solution was measured spectrophotometrically using the commercially available EnzChek phosphate assay (see Materials and Methods).

Since the rate of the single-stranded DNA-stimulated steady-state ATPase activity of yISW2 is low, we measured the kinetics of P_i release in the presence of a saturating concentration of poly(dT) to maximize the observed fluorescence signal; representative time courses from these experiments are shown in Figure 4. Linear analysis of the data yielded a hydrolysis rate of 63 ± 2 P_i yISW2⁻¹ min⁻¹, consistent with the estimate of 61 ± 4 ADP yISW2⁻¹ min⁻¹ determined from our radioactivity-based assay, and an intercept on the ordinate of -2 ± 1 P_i yISW2⁻¹. As shown in Figure 3, the time course of P_i release should be linear with a zero intercept on the ordinate if Scheme 1 is correct.

Alternative Model for DNA Translocation. The deviation from a zero intercept for the time course of release of P_i by yISW2 translocating along poly(dT) is inconsistent with the kinetic model outlined in Scheme 1. Although the deviation is small, we nevertheless consider it important to consider

Scheme 4



alternative models for single-stranded DNA translocation by yISW2. For example, it is worth noting that a previous study of the double-stranded DNA translocation by RSC (18) also detected no initial exponential phase of ATPase activity that could be associated with pre-steady-state kinetics. On the basis of this observation, a kinetic model for double-stranded DNA translocation was proposed that included a slow process, following DNA binding by the RSC, which was required to make RSC competent for double-stranded DNA translocation. In this model, shown in Scheme 2, the protein is i translocation steps away from the end of the DNA upon binding the DNA but is unable to initiate translocation (it is in the NP_i state in Scheme 2). The rate constant for dissociation from the NP_i state is k_2 and may be different from k_d . Upon addition of ATP, the protein undergoes a slow process, described by the rate constant k_{np} , before becoming competent for DNA translocation (the T_i state in Scheme 2). The thermodynamic coupling efficiency for the k_{np} rate constant is given by c_{np} .

The differential equations describing the concentration of protein bound at each position (NP_i , T_i , and free protein) in Scheme 2 can be solved to produce time-dependent equations for these protein concentrations that are functions of the maximum number of translocation steps (n) associated with a given DNA length. As before, these expressions can then be combined to form a time-dependent expression (eq 5) for the concentration of ATP hydrolyzed (or ADP or P_i produced) during translocation of yISW2 along single-stranded DNA.

$$\text{ADP}(t) = L^{-1} \left[\left(\frac{[(k_1[D] + k_2 + s)k_{np}k_1[D]]}{\{(k_1[D] + k_2)[(k_d + s)(k_2 + k_{np} + s) + k_1[D](k_{np} + k_d + s)]\}} \right) \left(\frac{ck_t}{(n+1)s^2} \left(n + \frac{k_t}{k_d + s} \left(\left(\frac{k_t}{k_t + k_d + s} \right)^n - 1 \right) \right) \right) \right] \quad (5)$$

Simulated time courses based upon this model are shown in Figure 3. As seen in Figure 3, these time courses consist of an initial slow exponential increase in the amount of ADP produced (ATP hydrolyzed) by the translocating proteins, associated with their pre-steady-state translocation along the single-stranded DNA, followed by a linear phase corresponding to their steady-state translocation activity. It is worth noting that the presence of this slow initial phase is consistent with the linear analysis of time course for P_i release shown in Figure 4. Specifically, the presence of this slow phase would result in an estimate of a negative intercept on the ordinate from a linear analysis of the time course.

The rate of steady-state ATP hydrolysis by these translocating enzymes obeys Michaelis–Menten kinetics with V_{\max} and K_M values shown in eqs 6 and 7, respectively.

$$V_{\max} = \left(\frac{k_{np}}{k_{np} + k_d} \right) \left\{ \frac{ck_t[n(P-1) - P(P^n-1)]}{(n+1)(P-1)} \right\} \quad (6)$$

$$K_M = \frac{(k_{np} + k_2)k_d}{(k_{np} + k_d)k_1} \quad (7)$$

As expected, eqs 6 and 7 are equal to eqs 2 and 3, respectively, in the limit of large k_{np} . If ATP hydrolysis is associated with the k_{np} step, then an additional term (eq 8) must be included in our expression for ADP produced by the translocating protein.

$$\text{ADP}(t) = L^{-1} \{ [c_{np}k_{np}(k_1[D] + s)(k_d + s)] / [(k_2 + k_{np} + s)(k_d + s) + k_1[D](k_{np} + k_d + s)] \} \quad (8)$$

This expression is dominated by a linear term which obeys Michaelis–Menten kinetics with a value of K_M identical to that shown in eq 7 and a value of V_{\max} given in eq 9

$$V_{\max} = \frac{c_{np}k_{np}k_d}{k_{np} + k_d} \quad (9)$$

Therefore, we can write the expression for the total amount of ADP produced during translocation in eq 10

$$V_{\max} = \left(ck_t \left[\frac{n(P-1) - P(P^n-1)}{(n+1)(P-1)} \right] + c_{np}k_d \right) \left(\frac{k_{np}}{k_{np} + k_d} \right) \quad (10)$$

The parameter estimates obtained from analysis of the data in Figure 2 using eq 10 are listed in Table 2. In this analysis, it is not possible to simultaneously determine estimates of c_{np} and d since these variables are strongly correlated in eq 10 (18). Thus, we first analyzed the data in Figure 2 assuming that $c_{np} = 0$ and then reanalyzed the data assuming $d = 10$ nucleotides since this corresponded to the shortest length of single-stranded DNA that stimulated the ATPase activity of yISW2. It is important to note that since eqs 2 and 10 predict an identical dependence of V_{\max} on n analysis of the dependence of V_{\max} on DNA length in Figure 2 using either model yields an identical estimate of P (0.90 ± 0.02) and describes the data equally well.

Kinetics of Single-Stranded DNA Binding by yIsW2. It is worth noting that the estimate of c_{np} obtained in the analysis of the data in Figure 2 using eq 10 is, within error, equal to zero (Table 2). This suggests that the k_{np} step in Scheme 2 may not be associated with significant hydrolysis of ATP. Since this step follows single-stranded DNA binding in Scheme 2, we decided to investigate whether it might correspond to the second step of a two-step single-stranded DNA binding mechanism by yIsW2.

In our single-stranded DNA binding experiments, we monitored the binding of yIsW2 to excess concentrations of dT₇₀ (Figure 5). In the absence of DNA, the observed fluorescence time course decayed monoexponentially. The rate of this signal change was independent of the fluorescence excitation wavelength or the bandwidth of the excitation monochromator, suggesting that it was not associated with photobleaching of the tryptophan residues but rather may result from the nonspecific binding of yIsW2 to the walls of the spectrometer flow cell. This slow signal change was also observed in the presence of the DNA, and the magnitude of the rate constant associated with this signal change was




FIGURE 4: Time courses of P_i production during single-stranded DNA translocation along poly(dT) monitored using the EnzChek phosphate assay as described in Materials and Methods. A linear analysis of the time course resulted in an estimate for V_{\max} of 63 ± 2 P_i yISW 2^{-1} min $^{-1}$, consistent with the estimate of 61 ± 4 P_i yISW 2^{-1} min $^{-1}$ determined from our radioactivity-based assay, and an intercept on the ordinate of -1 ± 0.5 P_i yISW 2^{-1} . This negative intercept on the ordinate is consistent with the kinetic model depicted in Scheme 4. A control experiment in which no DNA was included with the reaction mixture is also shown. A linear analysis of this time course resulted in an estimate for V_{\max} of 2 ± 1 P_i yISW 2^{-1} min $^{-1}$, consistent with the results from our radioactivity-based assay, and an intercept on the ordinate of 0.0 ± 0.1 .




FIGURE 5: Kinetics of yIsW2 binding to an excess concentration of dT₇₀, monitored by the quenching of the intrinsic tryptophan fluorescence of yIsW2. The time courses were observed in the stopped flow instrument upon mixing 10 nM yIsW2 (final concentration) with 0, 50, 250, and 500 nM dT₇₀. The solid lines overlaying each time course are NLLS squares fits obtained from global NLLS analysis of the time courses using eq 11.

independent of the concentration of the DNA (data not shown); this suggests that this process is not associated with the binding of yIsW2 to the single-stranded DNA. In the presence of DNA, however, two additional exponential phases are observed. The first observed rate constant exhibited a linear dependence on DNA concentration, whereas the second observed rate constant exhibited a hyperbolic dependence on DNA concentration (data not shown).

The minimal mechanism associated with these time courses and the associated dependence of these three observed rate constants on DNA concentration are shown in Scheme 3. The equation that describes the time depen-

Table 3: Estimates of the Kinetic Parameters for Single-Stranded DNA Binding by yIsW2 According to Scheme 3 (eq 11)

k_1 (M ⁻¹ s ⁻¹)	$(2.8 \pm 0.5) \times 10^6$	k_3 (s ⁻¹)	0.009 ± 0.002
k_{-1} (s ⁻¹)	1.40 ± 0.02	k_{-1}/k_1 (M)	$(4.9 \pm 0.2) \times 10^{-7}$
k_2 (s ⁻¹)	0.013 ± 0.002	k_{-2}/k_2	0.08 ± 0.01
k_{-2} (s ⁻¹)	0.157 ± 0.001	B/A	6.0 ± 0.8

dence of the intrinsic tryptophan fluorescence associated with the binding mechanism shown in Scheme 3 is given in eq 11.

$$f(t) = L^{-1} \left[\frac{1}{s + k_3} \left(C \frac{k_3}{s} + \{k_1[D][Bk_2 + A(s + k_{-2} + k_3)]\} / [k_{-1}(s + k_{-2} + k_3) + k_1[D](s + k_2 + k_{-2} + k_3) + (s + k_3)(s + k_2 + k_{-2} + k_3)] \right) \right] \quad (11)$$

In this equation, the parameter A is the fluorescence signal change associated with the first step of the binding, the parameter B is the fluorescence signal change associated with the second step of the binding, the parameter C is the fluorescence signal change associated with the nonspecific binding of the protein to the walls of the flow cell, and the rate constants k_1 , k_{-1} , k_2 , k_{-2} , and k_3 are as shown in Scheme 3. Since the slowest signal change occurs at a rate that is independent of the DNA concentration, we have assumed that an identical process affects all three protein species in Scheme 3. The parameter estimates obtained from the NLLS analysis of the time courses in Figure 5 using eq 11 are listed in Table 3.

Model of DNA Translocation Combined with DNA Binding. It is interesting to note that the equilibrium constant for the second step in the binding mechanism is highly unfavorable ($k_2/k_{-2} = 0.08 \pm 0.01$). This is consistent with the requirement that the slow conformation change in Scheme 2 is actually the second, and unfavorable, step of the two-step binding mechanism for yIsW2 to single-stranded DNA (see also ref 18). Naturally, the rate constants for single-stranded DNA binding (k_1 , k_{-1} , k_2 , and k_{-2}) will be affected by the presence of nucleotide, but the presence of this second step in the single-stranded DNA binding mechanism by yISW2 should nevertheless be incorporated into our kinetic model for single-stranded DNA translocation by yISW2.

A revised kinetic model for single-stranded DNA translocation by yIsW2 is shown in Scheme 4. In this scheme, the initial conformational change required to make yIsW2 competent for DNA translocation (the k_{np} step in Scheme 2) has been replaced with a second reversible process associated with single-stranded DNA binding by yIsW2. As before for Scheme 2, the differential equations describing the concentration of protein bound at each position (NP_i , T_i , and free protein) in Scheme 4 can be solved and subsequently combined to form a time-dependent expression for the concentration of ATP hydrolyzed (or ADP or P_i produced) during the translocation of yIsW2 along single-stranded DNA. This equation is too cumbersome to reproduce here; however, it does predict a linear term associated with the steady-state single-stranded DNA translocation activity of the enzyme. The magnitude of the rate constant associated with this steady-state ATPase activity follows Michaelis–Menten kinetics with V_{\max} and K_M as defined in eqs 12 and 13, respectively.

$$V_{\max} = \left[\frac{k_3(k_d + k_t)}{k_3(k_d + k_t) + k_d(k_4 + k_d + k_t)} \right] \left\{ \frac{ck_t[n(P - 1) - P(P^n - 1)]}{(n + 1)(P - 1)} \right\} \quad (12)$$

$$K_M = \frac{k_d[k_3(k_d + k_t) + k_2(k_4 + k_d + k_t)]}{k_1[k_3(k_d + k_t) + k_d(k_4 + k_d + k_t)]} \quad (13)$$

Not surprisingly, eqs 2, 6, and 12 exhibit an identical dependence of P on n . Therefore, analysis of data in Figure 2 using eq 12 also yields an estimate for P of 0.90 ± 0.02 and is equally as good at describing the data as the results of the analysis of the data using eq 2 or 6.

DISCUSSION

The chromatin remodeling activity of yIsW2 is manifest in the ability of the enzyme to directionally slide nucleosomes along DNA without disrupting the integrity of the nucleosome itself (1, 2, 31, 32). The current model for nucleosome mobilization by yISW2 involves the directionally biased translocation of the DNA on the surface of the nucleosome by yISW2 in an ATP-dependent reaction (15, 20, 31, 33). In this paper, we provide the first quantitative characterization of mechanisms of both single-stranded DNA translocation and single-stranded DNA binding by yISW2 and show how these two activities may be interrelated. Our direct observation of the single-stranded DNA translocation activity of yISW2 confirms previous indirect measurements (33) and offers a mechanism by which both double-stranded DNA translocation and nucleosome sliding can be directionally biased.

yISW2 Translocase Activity for Single-Stranded DNA. The dependence on single-stranded DNA length of the single-stranded DNA-stimulated ATPase activity of yIsW2 is consistent with yIsW2 being a single-stranded DNA translocase (Figure 2). Furthermore, this single-stranded DNA translocation activity may form the basis for double-stranded DNA translocation and nucleosome sliding by yIsW2 (2, 20, 33) and, perhaps, other remodelers as well (14, 18, 19).

Our estimate of the processivity of single-stranded DNA translocation by yIsW2 ($P = 0.90 \pm 0.02$) is consistent with enzyme moving approximately $m/(1 - P) = 20 \pm 2$ nucleotides on average before dissociation. This processivity is quite low compared to what is observed for helicase translocation along single-stranded DNA (11, 13, 35, 36) but similar to the processivity of double-stranded DNA translocation by the RSC chromatin remodeling complex from *S. cerevisiae* (18) and similar to the processivities of single- and double-stranded DNA translocation by *D. melanogaster* ISWI (15).

It is worth noting that *D. melanogaster* ISWI is proposed to translocate along single-stranded DNA with a 3' to 5' directional bias, whereas either a 5' to 3' directional bias, based upon modifications to the extranucleosomal linker DNA (33), or no directional bias, based upon modifications to the internucleosomal DNA (20), has been proposed for yIsW2. As mentioned previously, the ability of yISW2 to slide nucleosomes with directional bias implies that the enzyme also possesses directionally biased DNA translocation activity. Thus, it is possible that the interaction of one domain of yIsW2 with the extranucleosomal DNA dictates the directional bias to nucleosome sliding and the translocating motor domain moves either 3' to

5' or 5' to 3', accordingly, along a single strand of the internucleosomal DNA to accomplish the directionally biased sliding. In light of this, an important future experiment would be to directly test the directional bias of both single- and double-stranded DNA translocation by yIsw2 using a spectrophotometric assay (11, 34). Since either directional bias (3' to 5' or 5' to 3') of the single-stranded DNA translocation by yIsw2 would be consistent with the assumptions of our model, the resolution of this issue is not necessary to support the conclusions we draw here.

Translocation Initiation Is Likely Coupled to DNA Binding. The linear analysis of the ATPase time course shown in Figure 4 suggests that a slow initiation step might be present, following single-stranded DNA binding by yISW2, to make the enzyme competent for single-stranded DNA translocation. We propose that the slow initiation process may actually be the second step of a two-step mechanism for single-stranded DNA binding by yIsw2. Although yIsw2 is able to bind single-stranded DNA with reasonable affinity (Table 3), only a small fraction of the total population of bound protein has completed the second process in the binding reaction and thus is competent for DNA translocation. This hypothesis is consistent with previously published results which also demonstrated that although yIsw2 could bind DNA efficiently (25), its ATPase activity was not strongly stimulated by DNA binding (21).

Similarities to Other Motors. The kinetic mechanism for single-stranded DNA translocation proposed here for yIsw2 is similar to the mechanism recently proposed for the mechanism of double-stranded DNA translocation by the *S. cerevisiae* RSC chromatin remodeling complex (18). In both cases, a slow initiation step is required after DNA binding to make the enzyme competent for DNA translocation. In addition, slow initiation steps have also been proposed for mechanisms of single-stranded DNA translocation by the *Bacillus stearothermophilus* PcrA helicase (37, 38) and the T7 DNA helicase (36). The initiation steps for the PcrA and T7 DNA helicases were proposed to be ATP-dependent conformational changes (36–38), whereas no conclusion regarding the ATP dependence of the initiation step for RSC was proposed (18). However, it is worth noting that quantitative descriptions of the kinetic mechanisms for single-stranded DNA binding by these helicases have yet to be determined. Thus, it is not possible to further resolve the dependence of single-stranded DNA translocation by these enzymes on their single-stranded DNA binding characteristics.

In contrast, the kinetic mechanism of single-stranded DNA binding by monomers of the *Escherichia coli* Rep helicase has been determined (39). *E. coli* Rep is a member of the SF1 superfamily of helicases and is a known single-stranded DNA translocase (9). Interestingly, the second step in the two-step mechanism for single-stranded DNA binding by Rep is associated with a favorable equilibrium constant (39), and the kinetic mechanism for single-stranded DNA translocation by Rep does not include or require the presence of a slow initiation process to make the monomer competent for single-stranded DNA translocation (9). This correlation between the differences in both the mechanisms of single-stranded DNA binding and translocation for *E. coli* Rep and yIsw2 may further support our hypothesis that these two processes are linked, i.e., that the slow initiation process required to make yIsw2 competent for single-stranded DNA translocation is actually the second step of a two-step mechanism for single-stranded DNA binding by yIsw2. Similarly, the lower affinity of single-stranded DNA

binding by yISW2 when compared to that of Rep likely correlates with the lower processivity of single-stranded DNA translocation by yIsw2 when compared to Rep.

Implications for Nucleosome Mobilization. yIsw2 is able to slide nucleosomes without disrupting the integrity of the nucleosome (1, 2, 32). This suggests that the loops (or bulges) of DNA formed on the surface of the nucleosome as a result of the translocation of yIsw2 are likely small in size (2). The low processivity of single-stranded DNA translocation by yIsw2 reported here is consistent with the ability of yIsw2 to form small loops of DNA during its translocation (18). Although our studies here can provide no direct measurement of the average loop size or kinetic step size of translocation, we can safely assume that both are well below the processivity limit of the enzyme (~20 nucleotides); as mentioned above, a kinetic step size of 2 nucleotides has been proposed for single-stranded DNA translocation by yIsw2 based upon the enzyme's nucleosome sliding ability (20). As with the question of directional bias to single-stranded DNA translocation by yIsw2, the determination of the kinetic step size could likely be determined in future work using a spectrophotometric assay for single-stranded DNA translocation (11, 34).

Perhaps more interesting is the similarity between our estimate of the processivity of single-stranded DNA translocation and reported step sizes of nucleosome sliding by yIsw2 (20). The reported step size of 9–11 bp for nucleosome sliding (20) is smaller than our estimate of yISW2 translocating 20 ± 2 nucleotides on average before dissociation. One could therefore propose that the processivity of single-stranded DNA translocation may be ultimately limiting for the double-stranded DNA translocation and associated nucleosome sliding activity of the enzyme. Of course, the processivity of translocation for double-stranded DNA translocation may well be different from the processivity of single-stranded DNA translocation and could also be affected by the presence of and binding to the nucleosome. Nevertheless, this model is also consistent with recent *in vivo* studies of the chromatin remodeling activity of yIsw2 that suggest that the enzyme is bound to the chromatin only transiently during the specific period of nucleosome mobilization (1, 40). Furthermore, these studies also suggest that chromatin remodeling by yIsw2 involves the directional sliding of a few nucleosomes and, contrary to what is observed *in vitro* (32, 41), does not result in regularly spaced arrays of nucleosomes on the chromatin. Although *in vitro* studies of mononucleosome sliding by yIsw2 confirm that sliding is directionally biased from the ends of the DNA toward its center (31, 40), the differences between the results obtained *in vitro* and *in vivo* suggest that other cellular factors play a role in regulating the nucleosome mobilization activity of yIsw2 and also determining its directionality. Naturally, it would be interesting to determine what these factors are and how they also affect the single- and double-stranded DNA translocation activity of yIsw2.

ACKNOWLEDGMENT

We thank Dr. Timothy J. Richmond (ETH-Hönggerberg HPK) for providing samples of yISW2 for this work and for his critical review of the manuscript.

REFERENCES

1. Fazio, T. G., and Tsukiyama, T. (2003) Chromatin remodeling *in vivo*: Evidence for a nucleosome sliding mechanism. *Mol. Cell* 12, 1333–1340.

2. Gangaraju, V. K., and Bartholomew, B. (2007) Mechanisms of ATP dependent chromatin remodeling. *Mutat. Res.* 618, 3–17.
3. Kent, N. A., Karabetsov, N., Politis, P. K., and Mellor, J. (2001) In vivo chromatin remodeling by yeast ISWI homologs Isw1p and Isw2p. *Genes Dev.* 15, 619–626.
4. Flaus, A., and Owen-Hughes, T. (2001) Mechanisms for ATP-dependent chromatin remodelling. *Curr. Opin. Genet. Dev.* 11, 148–154.
5. Boyer, L. A., Logie, C., Bonte, E., Becker, P. B., Wade, P. A., Wolffe, A. P., Wu, C., Imbalzano, A. N., and Peterson, C. L. (2000) Functional delineation of three groups of the ATP-dependent family of chromatin remodeling enzymes. *J. Biol. Chem.* 275, 18864–18870.
6. Eisen, J. A., Sweder, K. S., and Hanawalt, P. C. (1995) Evolution of the SNF2 family of proteins: Subfamilies with distinct sequences and functions. *Nucleic Acids Res.* 23, 2715–2723.
7. Gorbalenya, A. E., and Koonin, E. V. (1993) Helicases: Amino acid sequence comparisons and structure-function relationships. *Curr. Opin. Struct. Biol.* 3, 419–429.
8. Cote, J., Peterson, C. L., and Workman, J. L. (1998) Perturbation of nucleosome core structure by the SWI/SNF complex persists after its detachment, enhancing subsequent transcription factor binding. *Proc. Natl. Acad. Sci. U.S.A.* 95, 4947–4952.
9. Brendza, K. M., Cheng, W., Fischer, C. J., Chesnik, M. A., Niedziela-Majka, A., and Lohman, T. M. (2005) Autoinhibition of *Escherichia coli* Rep monomer helicase activity by its 2B subdomain. *Proc. Natl. Acad. Sci. U.S.A.* 102, 10076–10081.
10. Dillingham, M. S., Soultanas, P., Wiley, P., Webb, M. R., and Wigley, D. B. (2001) Defining the roles of individual residues in the single-stranded DNA binding site of PcrA helicase. *Proc. Natl. Acad. Sci. U.S.A.* 98, 8381–8387.
11. Fischer, C. J., Maluf, N. K., and Lohman, T. M. (2004) Mechanism of ATP-dependent translocation of *E. coli* UvrD monomers along single-stranded DNA. *J. Mol. Biol.* 344, 1287–1309.
12. Soultanas, P., Dillingham, M. S., Wiley, P., Webb, M. R., and Wigley, D. B. (2000) Uncoupling DNA translocation and helicase activity in PcrA: Direct evidence for an active mechanism. *EMBO J.* 19, 3799–3810.
13. Niedziela-Majka, A., Chesnik, M. A., Tomko, E. J., and Lohman, T. M. (2007) *Bacillus stearothermophilus* PcrA monomer is a single-stranded DNA translocase but not a processive helicase in vitro. *J. Biol. Chem.* 282, 27076–27085.
14. Saha, A., Wittmeyer, J., and Cairns, B. R. (2002) Chromatin remodeling by RSC involves ATP-dependent DNA translocation. *Genes Dev.* 16, 2120–2134.
15. Whitehouse, I., Stockdale, C., Flaus, A., Szczelkun, M. D., and Owen-Hughes, T. (2003) Evidence for DNA translocation by the ISWI chromatin-remodeling enzyme. *Mol. Cell. Biol.* 23, 1935–1945.
16. Lia, G., Praly, E., Ferreira, H., Stockdale, C., Tse-Dinh, Y. C., Dunlap, D., Croquette, V., Bensimon, D., and Owen-Hughes, T. (2006) Direct observation of DNA distortion by the RSC complex. *Mol. Cell* 21, 417–425.
17. Zhang, Y., Smith, C. L., Saha, A., Grill, S. W., Mihardja, S., Smith, S. B., Cairns, B., Peterson, C. L., and Bustamante, C. (2006) DNA Translocation and Loop Formation Mechanism of Chromatin Remodeling by SWI/SNF and RSC. *Mol. Cell* 24, 559–568.
18. Fischer, C. J., Saha, A., and Cairns, B. R. (2007) Kinetic model for the ATP-dependent translocation of *Saccharomyces cerevisiae* RSC along double-stranded DNA. *Biochemistry* 46, 12416–12426.
19. Saha, A., Wittmeyer, J., and Cairns, B. R. (2005) Chromatin remodeling through directional DNA translocation from an internal nucleosomal site. *Nat. Struct. Mol. Biol.* 12, 747–755.
20. Zofall, M., Persinger, J., Kassabov, S. R., and Bartholomew, B. (2006) Chromatin remodeling by ISW2 and SWI/SNF requires DNA translocation inside the nucleosome. *Nat. Struct. Mol. Biol.* 13, 339–346.
21. Fitzgerald, D. J., DeLuca, C., Berger, I., Gaillard, H., Sigrist, R., Schimmele, K., and Richmond, T. J. (2004) Reaction cycle of the yeast Isw2 chromatin remodeling complex. *EMBO J.* 23, 3836–3843.
22. Lorch, Y., Cairns, B. R., Zhang, M., and Kornberg, R. D. (1998) Activated RSC-nucleosome complex and persistently altered form of the nucleosome. *Cell* 94, 29–34.
23. Leschziner, A. E., Saha, A., Wittmeyer, J., Zhang, Y., Bustamante, C., Cairns, B. R., and Nogales, E. (2007) Conformational flexibility in the chromatin remodeler RSC observed by electron microscopy and the orthogonal tilt reconstruction method. *Proc. Natl. Acad. Sci. U.S.A.* 104, 4913–4918.
24. Tsukiyama, T., Palmer, J., Landel, C. C., Shiloach, J., and Wu, C. (1999) Characterization of the imitation switch subfamily of ATP-dependent chromatin-remodeling factors in *Saccharomyces cerevisiae*. *Genes Dev.* 13, 686–697.
25. Gelbart, M. E., Rechsteiner, T., Richmond, T. J., and Tsukiyama, T. (2001) Interactions of Isw2 chromatin remodeling complex with nucleosomal arrays: Analyses using recombinant yeast histones and immobilized templates. *Mol. Cell. Biol.* 21, 2098–2106.
26. McConnell, A. D., Gelbart, M. E., and Tsukiyama, T. (2004) Histone fold protein Dls1p is required for Isw2-dependent chromatin remodeling in vivo. *Mol. Cell. Biol.* 24, 2605–2613.
27. Fazzio, T. G., Kooperberg, C., Goldmark, J. P., Neal, C., Basom, R., Delrow, J., and Tsukiyama, T. (2001) Widespread collaboration of Isw2 and Sin3-Rpd3 chromatin remodeling complexes in transcriptional repression. *Mol. Cell. Biol.* 21, 6450–6460.
28. Goldmark, J. P., Fazzio, T. G., Estep, P. W., Church, G. M., and Tsukiyama, T. (2000) The Isw2 chromatin remodeling complex represses early meiotic genes upon recruitment by Ume6p. *Cell* 103, 423–433.
29. Sherriff, J. A., Kent, N. A., and Mellor, J. (2007) The Isw2 chromatin-remodeling ATPase cooperates with the Fkh2 transcription factor to repress transcription of the B-type cyclin gene CLB2. *Mol. Cell. Biol.* 27, 2848–2860.
30. Vincent, J. A., Kwong, T. J., and Tsukiyama, T. (2008) ATP-dependent chromatin remodeling shapes the DNA replication landscape. *Nat. Struct. Mol. Biol.* 15, 477–484.
31. Kagalwala, M. N., Glaes, B. J., Dang, W., Zofall, M., and Bartholomew, B. (2004) Topography of the ISW2-nucleosome complex: Insights into nucleosome spacing and chromatin remodeling. *EMBO J.* 23, 2092–2104.
32. Kassabov, S. R., Henry, N. M., Zofall, M., Tsukiyama, T., and Bartholomew, B. (2002) High-resolution mapping of changes in histone-DNA contacts of nucleosomes remodeled by ISW2. *Mol. Cell. Biol.* 22, 7524–7534.
33. Zofall, M., Persinger, J., and Bartholomew, B. (2004) Functional role of extranucleosomal DNA and the entry site of the nucleosome in chromatin remodeling by ISW2. *Mol. Cell. Biol.* 24, 10047–10057.
34. Fischer, C. J., and Lohman, T. M. (2004) ATP-dependent translocation of proteins along single-stranded DNA: Models and methods of analysis of pre-steady state kinetics. *J. Mol. Biol.* 344, 1265–1286.
35. Tomko, E. J., Fischer, C. J., Niedziela-Majka, A., and Lohman, T. M. (2007) A Nonuniform Stepping Mechanism for *E. coli* UvrD Monomer Translocation along Single-Stranded DNA. *Mol. Cell* 26, 335–347.
36. Kim, D. E., Narayan, M., and Patel, S. S. (2002) T7 DNA helicase: A molecular motor that processively and unidirectionally translocates along single-stranded DNA. *J. Mol. Biol.* 321, 807–819.
37. Dillingham, M. S., Wigley, D. B., and Webb, M. R. (2000) Demonstration of unidirectional single-stranded DNA translocation by PcrA helicase: Measurement of step size and translocation speed. *Biochemistry* 39, 205–212.
38. Dillingham, M. S., Wigley, D. B., and Webb, M. R. (2002) Direct measurement of single-stranded DNA translocation by PcrA helicase using the fluorescent base analogue 2-aminopurine. *Biochemistry* 41, 643–651.
39. Bjornson, K. P., Moore, K. J., and Lohman, T. M. (1996) Kinetic mechanism of DNA binding and DNA-induced dimerization of the *Escherichia coli* Rep helicase. *Biochemistry* 35, 2268–2282.
40. Gelbart, M. E., Bachman, N., Delrow, J., Boeke, J. D., and Tsukiyama, T. (2005) Genome-wide identification of Isw2 chromatin-remodeling targets by localization of a catalytically inactive mutant. *Genes Dev.* 19, 942–954.
41. Langst, G., and Becker, P. B. (2001) Nucleosome mobilization and positioning by ISWI-containing chromatin-remodeling factors. *J. Cell Sci.* 114, 2561–2568.
42. Webb, M. R. (1992) A continuous spectrophotometric assay for inorganic phosphate and for measuring phosphate release kinetics in biological systems. *Proc. Natl. Acad. Sci. U.S.A.* 89, 4884–4887.