

Mechanism of Inactivation of Human Ribonucleotide Reductase with p53R2 by Gemcitabine 5'-Diphosphate[†]

Jun Wang,[‡] Gregory J. S. Lohman,[‡] and JoAnne Stubbe^{*,‡,§}

[‡]Departments of Chemistry and [§]Biology, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Received September 10, 2009; Revised Manuscript Received November 6, 2009

ABSTRACT: Ribonucleotide reductases (RNRs) catalyze the conversion of nucleoside 5'-diphosphates to the corresponding deoxynucleotides supplying the dNTPs required for DNA replication and DNA repair. Class I RNRs require two subunits, α and β , for activity. Humans possess two β subunits: one involved in S phase DNA replication (β) and a second in mitochondrial DNA replication (β' or p53R2) and potentially DNA repair. Gemcitabine (F₂C) is used clinically as an anticancer agent, and its phosphorylated metabolites target many enzymes involved in nucleotide metabolism, including RNR. The present investigation with α (specific activity of 400 nmol min⁻¹ mg⁻¹) and β' (0.6 Y \cdot / β' 2 and a specific activity of 420 nmol min⁻¹ mg⁻¹) establishes that F₂CDP is a substoichiometric inactivator of RNR. Incubation of this α/β' with [1'-³H]-F₂CDP or [5-³H]-F₂CDP and reisolation of the protein by Sephadex G-50 chromatography resulted in recovery 0.5 equiv of covalently bound sugar and 0.03 equiv of tightly associated cytosine to α 2. SDS-PAGE analysis (loaded without boiling) of the inactivated RNR showed that 60% of α migrates as a 90 kDa protein and 40% as a 120 kDa protein. Incubation of [1'-³H]-F₂CDP with active site mutants C444S/A, C218S/A, and E431Q/D- α and the C-terminal tail C787S/A and C790S/A mutants reveals that no sugar label is bound to the active site mutants of α and that, in the case of C218S- α , α migrates as a 90 kDa protein. Analysis of the inactivated wt- α/β' RNR by size exclusion chromatography indicates a quaternary structure of $\alpha\beta\beta'6$. A mechanism of inactivation common with h α/β is presented.

Gemcitabine (2',2'-dideoxydifluorocytidine, F₂C)¹ is a drug that is used clinically in the treatment of nonsmall cell lung carcinomas and advanced pancreatic cancer (2–6). The mechanism of its cytotoxicity is multifactorial where its metabolites, the mono-, di-, and triphosphates (F₂CMP, F₂CDP, and F₂CTP), inhibit a variety of steps in nucleic acid metabolism (4, 7). The essential step in apoptosis of the cells is F₂CTP inhibition of DNA polymerase by its incorporation into the growing polymer chain, resulting in chain termination (3). Potentiation of the effects of F₂CTP results from the inhibition of ribonucleotide reductase(s) (RNR), the enzymes that make deoxynucleoside 5'-diphosphates (dNDPs) from nucleoside diphosphates (8). RNRs are stoichiometrically inhibited by F₂CDP (9–12). This inhibition leads to a reduction in dNDP pools and consequently a

reduction in dNTP pools. Reduced concentrations of dNTPs reduce the competition for F₂CTP to become incorporated into DNA by DNA polymerase.

Human RNRs belong to class Ia RNRs and are composed of α and β subunits existing in complex quaternary structures $\alpha_n(\beta_2)_m$ ($n = 2, 4, 6$ and $m = 1, 3$) (13, 14). α houses the active site for nucleotide reduction and the binding sites for ATP/dNTP allosteric effectors that control the specificity and rate of nucleotide reduction. α contains five essential cysteines for catalysis: C429, the site of the thiyl radical that initiates catalysis, C218 and C444 that provide the reducing equivalents during dNDP formation, and C787 and C790 at the C-terminus that rereduces the active site disulfide subsequent to dCDP production. β houses the essential diferric-tyrosyl radical (Y \cdot) cofactor (15, 16). Recently, we have shown that hRNR, involved in DNA replication, is inactivated by 0.5 equiv of F₂CDP per α (11). Our studies unexpectedly revealed that the inhibition resulted from tight association of the two subunits and formation of an RNR quaternary structure determined by size exclusion chromatography (SEC) to be $\alpha\beta\beta'6$. Recently, a second RNR small subunit was discovered in humans (17, 18). It was designated p53R2 as its production is induced by p53 (17). Throughout the paper we will designate p53R2 as β' . In the present paper we report on the mechanism of inactivation of hRNR, $\alpha_n(\beta'2)_m$, by F₂CDP and compare the results with the replicative RNR, $\alpha_n(\beta2)_m$.

The biology and biochemistry of β' are actively being investigated. The biological function(s) of β' is (are) complex and appear(s) to be different in quiescent cells and cells which have been subjected to DNA damage. Recent studies of Pontarin et al. (19) suggest that β' is localized to the cytosol and is not shuttled to the nucleus as previously suggested (17, 20, 21). They

[†]This work was supported by an NIH grant (GM29595 to J.S.).

*To whom correspondence should be addressed. Tel: (617) 253-1814. Fax: (617) 258-7247. E-mail: stubbe@mit.edu.


¹Abbreviations: RNR(s), ribonucleotide reductase(s); α , ribonucleotide reductase large subunit; β , ribonucleotide reductase small subunit, also called R2; β' , the second ribonucleotide reductase small subunit, also called p53R2; h, human; hRNR, human RNR; h β , human β ; dNTP, deoxynucleoside 5'-triphosphate; dNDP, deoxynucleoside 5'-diphosphate; NDP, nucleoside 5'-diphosphate; equiv, equivalent; DTT, dithiothreitol; β -ME, β -mercaptoethanol; IPTG, isopropyl β -D-thiogalactopyranoside; wt, wild type; Y \cdot , tyrosyl radical; SEC, size exclusion chromatography; Amp, ampicillin; CM, chloramphenicol; PMSF, phenylmethanesulfonyl fluoride; F₂C, 2',2'-dideoxydifluorocytidine or gemcitabine; F₂CDP, 5'-diphosphate of F₂C; F₂CTP, 5'-triphosphate of F₂C; TR, thioredoxin; TRR, thioredoxin reductase; ATM, ataxia telangiectasia mutated; ERK, extracellular signal-regulated kinase; assay buffer, 50 mM Hepes (pH 7.6), 15 mM MgCl₂, 1 mM EDTA; buffer A, 25 mM Tris, pH 7.6, 10% glycerol, 30% (w/v) (NH₄)₂SO₄; buffer B, 25 mM Tris (pH 7.6), 10% glycerol, 5% (w/v) (NH₄)₂SO₄; buffer C, 50 mM Tris, pH 7.6, 5% glycerol, 1 mM EDTA, 1 mM PMSF.

proposed that β' plays a primary role in supplying the dNTPs required for mitochondrial DNA replication. This role is supported by identification of children with mutations in β' that experience severe mitochondrial DNA depletion (22).

While p53 mediates transcriptional induction of β' and as a consequence led to the proposal of its involvement in DNA repair, the induction process is not sufficiently rapid to supply dNTPs given the rate of damage repair (19, 23). Recent observations, however, have shown that β' interacts with and is phosphorylated by ATM (ataxia telangiectasia mutated), a DNA damage-inducible kinase, and is also found in the MRE11 (double strand break repair proteins) complex, involved in DNA double strand break repair. β' has also been shown to interact with ERK1/2 (extracellular signal-regulated kinase) and to inhibit MEK (a kinase that phosphorylates ERK)—ERK signaling and prevent cancer invasion, demonstrating an important role in genomic stability (23, 24).

β' is composed of 351 amino acids and is 80% homologous to human β ($h\beta$). Both proteins contain an essential diferric- $Y\cdot$ cofactor and are active in nucleotide reduction in the presence of α . $h\beta'$ has been reported to have activity 1.4-fold lower than β and a $Y\cdot$ content of $0.8/\beta^2$, 67% of that of β^2 (25, 26). However, the activity of $h\beta$ reported (25) is substantially lower (14-fold) than the activity recently reported from our laboratory (11) for reasons unknown. β' has also been proposed to have catalase activity in contrast to $h\beta$ and to have a 2.5-fold less susceptibility to inactivation by hydroxyurea (27).

As noted above, F_2CDP is a potent inhibitor of RNRs, and its demonstrated efficacy against a range of solid tumors warrants a detailed investigation of the mechanism of inhibition. Studies have been carried out on the class Ia RNRs from *Escherichia coli* and human $\alpha_n(\beta_2)_m$ and the class II RNR from *Lactobacillus leichmannii*. The *L. leichmannii* RNR is a monomeric enzyme that uses adenosylcobalamin as a cofactor. These studies have provided us with the following generalizations about the mechanism of inactivation. Using $[5-^3H]$ - and $[1'-^3H]$ - F_2CDP (F_2CTP) the stoichiometry of inhibition is 1 F_2CDP/α_2 for the class Ia RNRs (11) and 1 F_2CTP/α for the class II RNR (28–30). Despite the stoichiometry of inactivation, as with all mechanism-based inhibitors of RNR, at least two pathways are responsible for inhibition (1, 31). Covalent modification by a sugar moiety of F_2CDP is responsible for inactivation in one of these pathways. Destruction of the cofactor ($Y\cdot$ in the class Ia and adenosylcobalamin in the class II RNRs) (28) is responsible for the remaining inactivation. Subsequent to complete inactivation, stoichiometric amounts of cytosine are released, as are two fluoride ions. With the *E. coli* and *L. leichmannii* RNRs a new nucleotide radical (1) has been recently identified (10, 12, 28–30). This information has been used to formulate mechanisms of inactivation by the two pathways. Finally, with the oligomeric class Ia RNRs we have recently made the unexpected observation that the basis for their complete inactivation by 0.5 F_2CDP/α is that the α and β subunits form a tight complex proposed to be $\alpha_2\beta_2$ for *E. coli* RNR and $\alpha_6\beta_6$ for hRNR based on SEC analysis (11).



The prevalence of gemcitabine as a clinically useful drug and literature evidence suggesting that controlling the levels of β' may

increase potency of coadministered genotoxic therapeutics mandate a better understanding of the enzymology of β' to elucidate its function (32). The present communication reports that F_2CDP is an inactivator of β' , although the binding of F_2CDP to the α/β' complex appears to be weaker than that observed with the corresponding α/β complex. Studies with $[5-^3H]$ - and $[1'-^3H]$ - F_2CDP reveal that > 90% inactivation occurs with 0.5 equiv of inhibitor/ α , that 0.5 sugar is attached covalently to α , and that 0.65 cytosine is released per inactivation. SDS–PAGE analysis of the inactivated RNR reveals that α has been covalently modified. Forty percent of α migrates as a 120 kDa protein relative to unmodified α which migrates as a 90 kDa protein. Site-directed mutagenesis studies of all of the residues involved in catalysis in α and studies of the effect of each mutation on the inactivation of RNR composed of $h\beta$ or β' suggest that C218, one of the cysteines required to deliver reducing equivalents to make dNDP, and a cysteine (C790 or C787) from the C-terminus of α , involved in rereduction of the disulfide in the active site subsequent to dNDP formation, are likely responsible for covalent modification of α and the unusual migratory properties of α by SDS–PAGE analysis. SEC suggests that subsequent to inactivation RNR migrates as a $\alpha_6\beta'_6$ complex. β' thus can also be a target of gemcitabine and play a role in its cytotoxicity.

MATERIALS AND METHODS

Competent *E. coli* BL21(DE3)-RIL cells were purchased from Stratagene. Complete EDTA-free protease inhibitor tablets and calf alkaline phosphatase (20 units/ μ L) were purchased from Roche Biochemicals. $[^3H]$ -CDP (17 Ci/ μ mol) is from ViTrax Radiochemicals. p53R2 (26) containing the gene for β' was a gift from Lars Thelander (Department of Medical Biosciences, Medical Biochemistry, Umea University, SE-901 87 Umea, Sweden). phRRM1 containing the gene for α and phRRM2 containing the gene for β' was from Dr. Yun Yen (City of Hope National Medical Center, Duarte, CA) (25). Protein concentrations were determined using extinction coefficients (ϵ_{280nm}) per monomer (62000 $M^{-1} cm^{-1}$ for β' , 119160 $M^{-1} cm^{-1}$ for α). *E. coli* thioredoxin (TR, specific activity of 40 units/mg) and thioredoxin reductase (TRR, specific activity of 1320 units/mg) were isolated as previously described (33, 34).

Expression and Purification of Human β' . The expression vector pET3a- β' , encoding for untagged β' under the control of an IPTG inducible promoter, was transformed into BL21 Codon Plus (DE3)-RIL cells (Stratagene), plated on LB agar plates with 100 μ g/mL ampicillin (Amp) and 34 μ g/mL chloramphenicol (CM). A single colony was added to a culture (20 mL of LB in a 150 mL flask) and grown at 37 °C to saturation overnight. The culture was then diluted into 2 L of LB in a 6 L flask containing 100 μ g/mL Amp and 34 μ g/mL CM and grown at 37 °C to OD_{600nm} 0.7. 1,10-Phenanthroline (100 μ M) was then added to the media, and the cells were grown for an additional 15 min (35). Isopropyl β -D-thiogalactopyranoside (IPTG, 400 μ M) was then added, and the cells were grown at 37 °C for an additional 5 h. The cells were harvested and frozen in liquid N_2 , typically yielding 2 g/L.

For the isolation of the apo β' , 16 g of cells was resuspended (4 volumes/g) in 50 mM Tris-HCl, pH 7.6, 1 mM phenylmethanesulfonyl fluoride (PMSF), and 1 mM EDTA at 4 °C. The suspension was passed through the French press at 14000 psi. The cell lysate was centrifuged at 27000g for 30 min at 4 °C. A solution of 10% (w/v) of streptomycin sulfate was added to the

supernatant over 10 min while stirring at 4 °C, to a final concentration of 2.5%. After an additional 10 min of stirring, the pellet was removed by centrifugation (27000g for 30 min, 4 °C). Solid ammonium sulfate (0.243 g/mL, 40% saturation) was added to the supernatant over 10 min at 4 °C. After an additional 30 min of stirring, the precipitate was recovered by centrifugation (27000g for 30 min, 4 °C).

The pellet was dissolved in 4 mL of extraction buffer and loaded directly onto a phenyl-Sepharose 6 fast-flow column (20 mL, 2.5 × 10 cm) that was preequilibrated with 200 mL of 25 mM Tris, pH 7.6, 10% glycerol, and 30% (w/v) (NH₄)₂SO₄ (buffer A). The column was washed with buffer A (100 mL) and then 300 mL of 25 mM Tris (pH 7.6), 10% glycerol, and 5% (w/v) (NH₄)₂SO₄ (buffer B). Finally, β' was eluted with 50 mM Tris, pH 7.6, 5% glycerol, 1 mM EDTA, and 1 mM PMSF (buffer C). Fractions (2 mL) containing β' (based on Bradford assay) were collected.

The pooled protein fractions were then loaded onto a Q-Sepharose column (20 mL, 2.5 × 10 cm), which was preequilibrated with 100 mL of buffer C at 4 °C. The column was washed with 200 mL of buffer C and 150 mM KCl. The protein was eluted with a 70 mL × 70 mL linear gradient of 150–400 mM KCl in buffer C. Fractions (2 mL) containing β' (based on Bradford assay and 10% SDS–PAGE analysis) were collected. The apo β' eluted around 220 mM KCl. A typical yield of 0.4 mg of β' /g of cells was obtained.

Construction of the Mutant Y138F- β' Gene and Expression and Isolation of Y138F- β' . The gene for the mutant Y138F- β' was generated by site-directed mutagenesis using the QuickChange kit (Stratagene). The gene for β' was amplified by PCR using PfuUltra II polymerase (Stratagene) with primers for the Y138F mutant (Table S1 of the Supporting Information). The sequence of the mutant plasmid was confirmed by sequencing at the MIT Biopolymers Laboratory. Y138F- β' was expressed and purified as described for β' with a similar yield.

Conversion of Apo $\beta'2$ to Holo $\beta'2$. $\beta'2$ (60 μ M) in 500 μ L of 50 mM Hepes, 100 mM KCl, 10% glycerol, pH 7.6, and 2.5 mM ascorbate acid was deoxygenated by six cycles of evacuation (for 3 × 10 s) followed by argon flushing (2 min) on a Schlenk line. The deoxygenated $\beta'2$ solution was brought into the glovebox (M. Braun, Stratham, NH), and 6 equiv of Fe(II) (deoxygenated ferrous ammonium sulfate in 50 mM Tris and 100 mM KCl, pH 7.6) per $\beta'2$ was added. The resulting mixture was incubated at 4 °C for 30 min. The protein was then removed from the glovebox, and 100 μ L of ice-cold O₂ saturated 50 mM Tris and 100 mM KCl, pH 7.6, was added. O₂(g) was also blown over the surface of the protein solution for 1 min. Excess iron was removed by Sephadex G-25 chromatography (40 mL, 2.5 × 30 cm), and the protein fractions were pooled to give a concentration of 3.5 mg/mL. An activity assay was carried out immediately, and 250 μ L of the protein solution was placed in an EPR tube and frozen in liquid N₂ for measurement of the Y·. Typically 0.6 Y·/ $\beta'2$ was observed with a specific activity of 420 nmol min⁻¹ mg⁻¹.

Construction of Active Site and C-Terminal Tail Mutant Genes of α and Expression and Purification of the Mutant Proteins. Mutants of α [C218S(A), C429S(A), C444S(A), C787S(A), C790S(A), E431Q(D)] were generated by site-directed mutagenesis using the QuickChange kit (Stratagene). For each mutant, the gene was amplified by PCR using PfuUltra II polymerase (Stratagene) with primers (Table S1 of the Supporting Information). The sequence of each mutant plasmid was

confirmed by the MIT Biopolymers Laboratory. The mutants were expressed and purified as described below for wt- α with similar yields.

Purification of α and α Mutants by Ni-NTA and dATP Affinity Chromatography. Cells (15 g) were suspended (5 volumes/g) in 50 mM NaH₂PO₄, pH 7.0, 0.1% Triton X-100, and 10 mM β -mercaptoethanol (β -ME) with the complete protease inhibitor (Roche). The suspension was passed through the French press at 14000 psi. The cell lysate was centrifuged at 20000g for 30 min. Streptomycin sulfate (10%) was added to the supernatant over 10 min to a final concentration of 1% (w/v) and stirred for an additional 10 min. After the pellet was removed by centrifugation, the supernatant was incubated with Ni-NTA agarose resin (1 mL/g of cells; Qiagen) at 4 °C for 1 h and then loaded into a column (2.5 × 10 cm). The column was washed with 30 column volumes of 50 mM NaH₂PO₄, 800 mM NaCl, 50 mM imidazole, pH 7.0, 0.1% Triton X-100, and 10 mM β -ME. The protein was eluted with 50 mM NaH₂PO₄, 300 mM NaCl, and 125 mM imidazole, pH 7.0. The fractions containing protein were identified using the Bradford assay, pooled, and concentrated to < 10 mL, and then the imidazole was removed by Sephadex G-25 chromatography (200 mL, 2.5 × 50 cm) using 50 mM Tris, pH 7.6, 5% glycerol, and 1 mM DTT as eluent. The pooled protein fractions were added to dATP affinity resin (30 mL, equilibrated with 50 mM Tris, pH 7.6, 5% glycerol, 1 mM DTT) (36) and gently mixed for 2 h at 4 °C in a 50 mL falcon tube. The resin was then loaded into a column (2.5 × 10 cm) and washed with 300 mL of 50 mM Tris, pH 7.6, 5% glycerol, 300 mM KCl, and 1 mM DTT. The protein was eluted with 50 mM Tris, pH 7.6, 5% glycerol, 100 mM KCl, 5 mM DTT, and 10 mM ATP. The fractions containing protein were identified using the Bradford assay, pooled, and concentrated to < 0.5 mL. The ATP was removed by Sephadex G-25 chromatography (40 mL, 1.5 × 30 cm). α was stored in 50 mM Tris, 100 mM KCl, 15 mM MgCl₂, 5 mM DTT, pH 7.6, and 5% glycerol with a typical yield of 1 mg/g of cells after the Ni affinity column and 0.2 mg/g of cell pellet after dATP affinity chromatography. High quality α purified by dATP chromatography was used in SEC studies only. All other experiments were carried out using α (wt or mutants) purified by Ni affinity chromatography only.

Activity Assay. The reaction mixture contained the following in a final volume of 350 μ L: 50 mM Hepes (pH 7.6), 15 mM MgCl₂, 1 mM EDTA (assay buffer), 0.3 μ M (or 3 μ M) α , 3 μ M (or 0.3 μ M) β' , 3 mM ATP, 1 mM [³H]-CDP (specific activity 5115 cpm/nmol), 100 μ M *E. coli* TR, 1.0 μ M TRR, and 2 mM NADPH. The assay mixture was preincubated at 37 °C for 2 min, and the reaction was initiated by the addition of [³H]-CDP. Aliquots (30 μ L) were removed over a 15 min time period and quenched in a boiling water bath for 2 min. dC production was analyzed, subsequent to dephosphorylation with alkaline phosphatase as previously described (37), and analyzed by the method of Steeper and Stuart (38).

Time-Dependent Inactivation Assays. The inactivation mixture contained the following in a final volume of 100 μ L: 6 μ M α , 6 μ M β' , 3 mM ATP, 5 mM DTT, and assay buffer. The reaction was initiated by addition of 0.5 or 5 equiv of F₂CDP (3 μ M, 30 μ M) and incubated at 37 °C. Three types of assays were carried out. In assay 1, aliquots (10 μ L) from the reaction mixture were removed at several time points from 30 s to 23 min and diluted 5-fold into 50 μ L of assay buffer. dCDP production was measured as described above. A control experiment under identical conditions with the omission of F₂CDP was always

carried out to assess the stability of β' . In assays 2 and 3, the activity of α or β' was measured, respectively. Aliquots (2.5 μ L) of the inactivation mixture were diluted into 50 μ L of assay buffer containing 3 μ M (10-fold excess) second subunit. dCDP production was measured as described above.

Quantitation of Covalent Labeling of Human $\alpha_n(\beta'2)_m$ with [$1'$ - 3 H]- F_2 CDP and [5 - 3 H]- F_2 CDP. A typical reaction mixture contained the following in a final volume of 200 μ L: 8.5 μ M α and β' with 5 mM DTT and 3 mM ATP in assay buffer. The reaction was initiated with 10.6 μ M [$1'$ - 3 H]- F_2 CDP (5889 cpm/nmol) or 10.6 μ M [5 - 3 H]- F_2 CDP (6643 cpm/nmol). After 8 min at 37 °C, an aliquot of 195 μ L was either directly loaded onto a Sephadex G-50 column (1 cm \times 20 cm, 20 mL) preequilibrated in assay buffer or mixed with guanidine hydrochloride to a final concentration of 6 M, incubated for 5 min, and then loaded onto a Sephadex G-50 column in assay buffer containing 2 M guanidine hydrochloride. Fractions (1 mL) were collected and assayed for protein by Bradford assay, and 500 μ L of each fraction was analyzed by scintillation counting.

Quantitation of Cytosine Released during the Inactivation of Human $\alpha_n(\beta'2)_m$ by [5 - 3 H]- F_2 CDP. The reaction mixture contained in 500 μ L 1.2 μ M α , β' and 1.2 μ M [5 - 3 H]- F_2 CDP (6643 cpm/nmol, 2 equiv/ α 2). The reaction was incubated for 15 min at 37 °C. The mixture was then filtered through an YM-30 Centricon device (Millipore) at 4 °C. F_2 C (60 nmol) and cytosine (120 nmol) were added as carriers before filtration. The flow-through was treated with 30 units of alkaline phosphatase (Roche) for 3 h at 37 °C and filtered through a second YM-30 Centricon device. The flow-through was analyzed using a Waters 2480 HPLC with an Altech Adsorbosphere nucleotide nucleoside C-18 column (250 mm \times 4.6 mm) at a flow rate of 1 mL/min. The elution buffer contained the following: buffer I, 10 mM NH_4OAc , pH 6.8; buffer II, 100% methanol. A 10 min isocratic elution with buffer I was followed by a linear gradient to 40% buffer II over 30 min. A linear gradient was then run to 100% buffer II over 5 min. Fractions (1 mL) were collected, and 200 μ L of each was analyzed by scintillation counting. The recovery of [5 - 3 H]-cytosine and [5 - 3 H]- F_2 C was calculated based on the UV spectrum (cytosine, λ_{267nm} , ϵ = 6100 M $^{-1}$ cm $^{-1}$; F_2 C, λ_{268nm} , ϵ = 9360 M $^{-1}$ cm $^{-1}$) and normalized for carrier added. The radioactivity recovered with [5 - 3 H]-cytosine and [5 - 3 H]- F_2 C was analyzed by scintillation counting.

SDS-PAGE of the Inactivation Mixture without Boiling. The inactivation mixture contained the following in a final volume of 35 μ L: 6 μ M α or C218S- α , 6 μ M β' , 3 mM ATP, with or without 5 mM DTT, and assay buffer. The reaction was initiated by addition of 5 equiv of F_2 CDP (30 μ M) and incubated at 37 °C for 5 min. The inactivation mixture (8 μ L) was mixed with 8 μ L of 2 \times loading buffer \pm β -ME. [Note loading buffer contains SDS.] The samples were either heated at 90 °C for 2 min or not heated before loading on a 10% SDS-PAGE gel. The proteins were visualized with Coomassie blue staining. The band intensities were quantified using Bio-Rad Quantity One software.

Incubation of [$1'$ - 3 H]- F_2 CDP with C218S(A)-, C429S(A)-, C444S(A)-, C787S(A)-, C790S(A)-, and E431Q(D)- α and β' (β) and Analysis for Covalent Labeling by Sephadex G-50 Chromatography. A typical reaction mixture contained the following in a final volume of 200 μ L: 8.5 μ M α mutant and β' (β), 5 mM DTT, and 3 mM ATP in assay buffer. The reaction was initiated with 10.6 μ M [$1'$ - 3 H]- F_2 CDP (5889 cpm/nmol). After 8 min at 37 °C, an aliquot of 195 μ L was loaded onto a Sephadex G-50 column (1 cm \times 20 cm, 20 mL)

that was preequilibrated with the assay buffer. Fractions (1 mL) were collected and assayed for protein by the Bradford assay, and 500 μ L of each fraction was analyzed by scintillation counting.

SEC To Examine the Quaternary Structure of $\alpha_n(\beta'2)_m$ Subsequent to Inactivation by F_2 CDP. SEC was performed using a Superdex 200 column (10 \times 300 mm; GE Healthcare) attached to a Waters 2480 HPLC. Gel filtration molecular mass standards (GE healthcare) were as follows: ovalbumin, 43 kDa; conalbumin, 75 kDa; aldolase, 158 kDa; catalase 232 kDa; ferritin, 440 kDa; thyroglobulin, 669 kDa; blue dextran, 2000 kDa. The elution buffer was 50 mM Hepes (pH 7.6), 15 mM $MgCl_2$, 1 mM EDTA, and 0.5 mM ATP. Molecular mass standards were run at the beginning of each experiment. The reaction mixture (400 μ L) contained 15 μ M α and β' (or Y138F- β'), 5 mM DTT, 3 mM ATP, and 37.5 μ M F_2 CDP or [$1'$ - 3 H]- F_2 CDP (2.5 equiv) in assay buffer. α was purified using Ni affinity and dATP affinity chromatography and had a specific activity of 232 nmol min $^{-1}$ mg $^{-1}$ (measured with β 2). After 10 min incubation, 390 μ L was injected onto the column using a 500 μ L loop. The elution rate was 0.5 mL/min, 0.5 mL fractions were collected, and 300 μ L aliquots of each fraction were analyzed by scintillation counting.

Quantitative Analysis of the Subunits of $\alpha_n(\beta'2)_m$ by SDS-PAGE. The fractions collected from the SEC analysis above were analyzed directly (8 μ L from each fraction) by 10% SDS-PAGE and compared with concentrations of α and β from *E. coli* RNR (0.05–0.4 μ M) as standards. To analyze control experiments with Y138F- β' / α and with $\alpha/\beta'/ATP$, each fraction of the first SEC peak was concentrated before loading on the gel. The proteins were visualized with Coomassie blue staining. The band intensities were quantified using Bio-Rad Quantity One software. The concentrations of α and β' in the complex were determined from standard curves made with *E. coli* subunits as described above.

RESULTS

Purification and Reconstitution of the $\beta'2$. Growth of $\beta'2$ was carried out in the presence of 1,10-phenanthroline, resulting in apoprotein (35). This procedure, subsequent to reconstitution of cluster, gave higher recovery of $Y\cdot$ and activity than standard growth conditions with addition of ferrous ammonium sulfate and ascorbate to the crude lysate prior to purification (39). $\beta'2$ was purified by addition of phenyl-Sepharose 6 fast-flow and Q-Sepharose fast-flow column chromatographies to the procedure of Thelander (26). These steps were required to obtain homogeneous protein by SDS-PAGE. A variety of methods to assemble the essential diferric-tyrosyl radical ($Y\cdot$) cluster of $\beta'2$ were examined including use of tagged and nontagged versions of $\beta'2$ and varying the ratio of Fe^{2+}/β' . The detailed in vitro reconstitution procedure described in Materials and Methods gave 0.63 $Y\cdot/\beta'2$ and an activity of 420 nmol min $^{-1}$ mg $^{-1}$. The procedure reported by the Thelander group gave 1 $Y\cdot/\beta'2$ (nontagged), but the activity was 95 nmol min $^{-1}$ mg $^{-1}$ (26). A second purification procedure reported by Yen and co-workers using a tagged $\beta'2$ gave 0.8 $Y\cdot/\beta'2$ and a specific activity of 50 nmol min $^{-1}$ mg $^{-1}$ (25). The low activity and high $Y\cdot$ content reported by these groups are at odds with activity being directly proportional to $Y\cdot$ (40, 41). The basis for the discrepancy is not understood. In our hands we have been unable, after many attempts with identical constructs to those used by Thelander and Yen, to obtain higher levels of $Y\cdot/\beta'2$. However, with our




FIGURE 1: Time-dependent inactivation assay of human β'/α by F_2CDP at 30 μM (black \blacktriangle), 3 μM (red \blacksquare), and 0 μM (blue \blacklozenge). (A) Aliquots were removed from the inactivation mixture at various times and diluted 5-fold for determination of RNR activity. (B) Aliquots were removed and diluted 20-fold for determination of α (black \blacktriangle) or β' (red \blacksquare) activity in the presence of a 10-fold excess of β' or α , respectively. The activity of β' (red \blacksquare) was adjusted using results from the control experiment (blue \blacklozenge) in the absence of inhibitor.

previous purification of $h\beta$, our group did obtain the same amount of radical ($1 Y \cdot / \beta_2$) as these groups, but with 7–14-fold higher activity. Our ability to only obtain $0.6 Y \cdot / \beta_2$ makes the quantitation of products from the F_2CDP inactivation studies more complex as there is always 40% of β_2 and α that have not reacted. In the quantitations reported below we have placed numbers normalized to $1 Y \cdot / \beta_2$ in parentheses.

Time-Dependent Inactivation Studies of α , β_2 by F_2CDP/ATP . Our previous studies with F_2CDP on human $\alpha_n(\beta_2)_m$ and *E. coli* $\alpha_2\beta_2$ showed that F_2CDP is a substoichiometric mechanism-based inhibitor resulting in covalent binding of 1 equiv of the sugar moiety from $[1'-^3H]-F_2CDP$ per α_2 . To determine if β_2 behaves in a similar fashion, inactivation studies were carried out with α , β' (6 μM each) and 0.5 or 5 equiv of F_2CDP/α in the presence of reductant (either DTT or TR/TRR/NADPH). Three different assays were carried out. In the first assay (Figure 1A) the enzyme was diluted 5-fold and assayed for dCDP production. Both concentrations of F_2CDP resulted in $>90\%$ inactivation with the reaction containing a 5-fold excess of inhibitor reaching completion in 4 min (black \blacktriangle). Even under these conditions, it is unlikely that the enzyme is saturated with F_2CDP . As with $h\beta$, substoichiometric amounts of inhibitor result in $>90\%$ inactivation, although the inactivation is slow (red \blacksquare). The control with β_2 in the absence of inhibitor reveals that $Y \cdot$ is lost, precluding longer time points (blue \blacklozenge). The half-life of $Y \cdot$ loss is similar to our observations with $h\beta$ (11).

Because of the weak interactions between the two subunits in the class I RNRs, assays of the individual subunits, α (β'), are typically carried out with a 10-fold excess of the second subunit,

Table 1: Covalent Labeling of Human $\alpha_n(\beta_2)_m$ with $[1'-^3H]-F_2CDP$ and $[5-^3H]-F_2CDP$ Analyzed by SEC

protein	$[x-^3H]-F_2CDP$, $x =$	Sephadex G-50	$[^3H]/\alpha_2^b$
human $\beta' \alpha$	1'	native	0.5 (0.83)
	1'	denaturing	0.5 (0.83)
	5	native	0.03 (0.05)

^a0.6 $Y \cdot / \beta_2$. ^bNormalized 0.6 $Y \cdot$ to 1 $Y \cdot / \beta_2$.

β' (α), present in the assay mixture. The result of a typical experiment under these conditions (0.5 equiv of F_2CDP) is shown in Figure 1B. Under these conditions α is 95% inactivated (black \blacktriangle), while β' retained 40% of its activity (red \blacksquare) after correction for β' stability (blue \blacklozenge). These results are similar to those for the *E. coli* RNR and the hRNR with β . The results suggest that excess α can facilitate subunit dissociation and liberate the remaining active β' . The implications of these results will be addressed subsequently.

Quantitation of Labeled RNR and Small Molecules Generated by Inactivation with $[1'-^3H]-$ and $[5-^3H]-F_2CDP$. With $[1'-^3H]-F_2CDP$, our previous studies with $h\beta_2$ containing 1.2 $Y \cdot / \beta_2$ identified 0.9 equiv/ α_2 of covalently bound sugar. A similar experiment using α and β' at 8.5 μM and 2.5 equiv of $[1'-^3H]-F_2CDP$ was carried out for 8 min at 37 °C. The reaction mixture was analyzed by Sephadex G-50 column chromatography in the presence or absence of denaturant (guanidine hydrochloride). The results are summarized (Table 1) and reveal 0.5 equiv of sugar/ α_2 (normalization for $Y \cdot$ gives 0.83 equiv/ α_2). When the experiment was repeated with the $[5-^3H]-F_2CDP$ under identical conditions in the absence of denaturant, 0.03 equiv/ α_2 (0.05) was detected with protein (Table 1).

To test if 0.5 equiv of sugar/inactivated enzyme is accompanied by 0.5 equiv of cytosine release, 2 equiv of $[5-^3H]-F_2CDPs$ (1.2 μM) was incubated with $\alpha_2\beta_2$ (0.6 μM) for 15 min at 37 °C. Subsequent to inactivation the nucleotides were recovered by ultrafiltration and analyzed by HPLC. The analysis revealed that 0.65 equiv (1.08) of cytosine was released and 1.35 equiv of F_2C was recovered (Figure S1 of the Supporting Information). These results are in agreement with our previous studies on inactivation of human $\alpha_n(\beta_2)_m$ and *E. coli* RNR and reveal that the extent of reaction correlates with the amount of $Y \cdot$ in β or β' .

Analysis of the Inactivation Mixture of wt- and C218S- α by SDS-PAGE without Boiling. Our recent studies with *E. coli* RNR and *L. leichmannii* RNR revealed that when the enzyme inactivated with the inhibitor was analyzed by an SDS-PAGE gel without boiling, 30–50% of α migrated as a larger species (~110 kDa for *E. coli* and 110 kDa for *L. leichmannii*) (12, 29, 30). The α with the aberrant migratory properties is likely associated with the covalently bound sugar moiety from F_2CDP . To determine if the substoichiometric labeling of hRNR was accompanied by production of a modified α , the reaction mixture subsequent to inactivation was analyzed directly without boiling by SDS-PAGE (Figure 2). A similar experiment was carried out with β (Figure S2 of the Supporting Information). Twenty-four percent (40%) and 33% of the α migrated as a larger species (~120 kDa) with β' and $h\beta$, respectively. This number in each case corresponds roughly to the amount of covalently bound sugar. With all RNRs examined to date, even the monomeric adenosylcobalamin RNR, production of an α with altered migratory properties is associated with covalent modification by the sugar moiety of $F_2CD(T)P$.




FIGURE 2: SDS-PAGE analysis of the α/β' (6 μ M) inactivated by F_2 CDP (30 μ M) and ATP (3 mM) at 37 °C for 5 min. Each sample was mixed with 2 \times loading buffer \pm β -ME or boiled for 2 min before loading as indicated. A band at 120 kDa is observed in lanes 2 and 3 (inactivation in the presence of DTT, without boiling) and lanes 5 and 6 (inactivation in the absence of DTT, without boiling).

The same experiment was repeated with a mutant of one of the two active site cysteines involved directly in nucleotide reduction (C218S- α) based on observations reported in the next section. The results are also shown in Figure 2 (lanes 7–12) for β' and Figure S2 of the Supporting Information for β . With this mutant no altered conformation of α is observed, suggesting that this residue may play an important role in its production. A proposal for these observations is presented subsequently.

Identification of Potential Sites of Covalent Modification of Human RNR Inactivated by [$1'$ - 3 H]- F_2 CDP Using Site-Directed Mutants of Residues in α Involved in Catalysis. Our recent studies with *L. leichmannii* RTPR have established that 10% of the label associated with RNR subsequent to inactivation by [$1'$ - 3 H]- F_2 CTP is due to alkylation of the sugar of the nucleotide by both cysteines at the C-terminus of the protein (29, 30). All efforts to identify the major labeled species failed, due to instability and rearrangement, even when $NaBH_4$ was used to stabilize the alkylated protein. Furthermore, studies with the C225S mutant of *E. coli* RNR suggested that this residue plays a key role in covalent modification of the enzyme by F_2 CDP (12). To identify the site of covalent modification by the sugar derived from F_2 CDP, [$1'$ - 3 H]- F_2 CDP was incubated with active site mutants of human α : C429S/A, E431Q/D, C787S/A, C790S/A, C444S/A, C218S/A (the equivalent of C225 in *E. coli*), and β or β' in the presence of DTT. Each mutant was purified as described for wt- α with similar recovery. The extent of covalent labeling measured subsequent to Sephadex G-50 chromatography (in the absence of denaturant) was determined by scintillation counting and compared to the labeling of wt-RNR. The results are summarized in Table 2.

As expected, no label is associated with RNR when the radical initiator on α (C429), essential for removal of the 3'-H for all mechanism-based inhibitors, is changed to S or A. Two other mutants, E431Q/D- α and C218S/A- α , also showed no labeling. The E431 mutant is likely not labeled as this glutamate plays an important role (although not essential role) in the first two steps of the reduction process by removal of a proton from the 3'-OH of F_2 CDP (28, 42, 43). Thus the rate of chemistry on any nucleotide is likely to be greatly reduced when the E is changed to a Q/D. In addition, a recent structure of the *Saccharomyces cerevisiae* α soaked with F_2 CDP showed that E431 is 6.2 Å from

Table 2: Quantitation of Sugar Covalently Attached to hRNR Mutants Incubated with [$1'$ - 3 H]- F_2 CDP Analyzed by SEC

protein	[3 H]/ $\alpha 2\beta' 2$	[3 H]/ $\alpha 2\beta 2$
wt- α	0.5 (0.8) ^b	0.8–0.9
C429S- α (C439) ^a	0	0.008
C429A- α (C439)	0	0.01
E431Q- α (E441)	0	0.01
E431D- α (E441)	0	0.08
C787S- α (C754)	0.3 (0.5)	0.8
C787A- α (C754)	0.26 (0.4)	0.3
C790S- α (C759)	0.3 (0.5)	0.6
C790A- α (C759)	0	0.1
C444S- α (C462)	0.03 (0.05)	0.4
C444A- α (C462)	0.2 (0.3)	0.7
C218S- α (C225)	0	0.1
C218A- α (C225)	0	0.1
Y138F- β' (Y122)	0	

^aThe equivalent residues of *E. coli* RNR are given in parentheses. ^bThe values normalized to 1 Y- $\beta' 2$ are given in parentheses.

the 3'-HO of the sugar and that a water molecule intervenes (44). Thus, it is unlikely that this E is the site of labeling.

C225 on *E. coli* α , on the other hand, has been implicated as playing a key role in the inactivation pathway involving covalent modification of the *E. coli* RNR (12). Our previous studies with the C225S mutant of *E. coli* have demonstrated that the first few steps in catalysis are not impaired by this mutation (45, 46). Thus lack of labeling with C218S/A-h α makes this a likely candidate for alkylation. Consistent with this proposal, the *S. cerevisiae* structure shows that this is the only nucleophilic residue close to the sugar (44). The caveat is, however, that the second subunit, essential for catalysis, is missing in the structure. Thus our favored model is that C218 is covalently modified by a sugar moiety derived from F_2 CDP. In line with this proposal, the SDS-PAGE gel (Figure 2 and Figure S2 of the Supporting Information) revealed that with the C218S- α mutant and either h β or β' no α with modified migratory properties is observed.

The C-terminal tail mutants of α , C787S/A and C790S/A, retain variable amounts of label associated with hRNR (with either β or β') (Table 2). These cysteines are involved in rereduction of the active site disulfide generated during the synthesis of dNDP (47, 48). This C-terminus must be able to enter the active site to reduce the disulfide between C218 and C444 and then must swing out of the active site to react with the protein reductant TR. The variability in labeling suggests that one of these cysteines may be involved in covalent interaction with the sugar moiety as well. Labeling of the active site C218 and a C (787 or 790) in the tail might give rise to the observed conformational change found by SDS-PAGE (Figure 2 and Figure S2 of the Supporting Information).

SEC To Examine the Quaternary Structure of hRNR (with β') Inactivated by F_2 CDP. Our previous studies have shown that a tight $\alpha 6\beta 6$ complex is generated when human RNR is inactivated by F_2 CDP. To examine the quaternary structure of $\alpha_n(\beta' 2)_m$ subsequent to inactivation with [$1'$ - 3 H]- F_2 CDP, the inactivated complex was examined by SEC with 0.5 mM ATP in the elution buffer. The A_{280nm} revealed protein elution at 18.6 min and at 25.6 min (Figure 3). Comparison of the retention times with a standard curve generated with globular proteins of known molecular weight (Figure S3 of the Supporting Information) suggested that the protein eluting at 18.6 min has a mass (794 kDa) more consistent with $\alpha 6\beta' 6$ than $\alpha 6\beta' 2$ (798 vs




FIGURE 3: SEC on a S200 column to detect complex formation of α/β' inactivated by $[1\text{'-}^3\text{H}]\text{-F}_2\text{CDP}$ ($37.5\text{ }\mu\text{M}$). The elution buffer contained 0.5 mM ATP. (A) The elution profile monitored by $A_{280\text{nm}}$ and scintillation counting (red ■). (B) Analysis of the ratio of α : β (red —) using standard curves generated from known amounts of *E. coli* α and β . (C) SDS-PAGE of fractions 18–20 from (A). (D) The elution profile in the absence of F_2CDP .

633 kDa). Control experiments with α (88 kDa), $\alpha 2$ (189 kDa with TTP), and $\beta'2$ (116 kDa) (Table 3 and Figure S4 of the Supporting Information) suggest that the peak eluting at 25.6 min may be composed of $\beta'2$ and α . Each fraction from the SEC column was examined by SDS-PAGE with the amount of α and β' determined from a standard curve made with known concentrations of *E. coli* α and β (Figure S5 of the Supporting Information). Analysis of protein in fractions 18, 19, and 20 gave a ratio of α : β' as 1.2, 1.0, and 1.0, respectively, and an average of 1.06 ± 0.15 (Figure 3B,C). This analysis supports the formation of an $\alpha\beta'6$ complex in the presence of F_2CDP and ATP. Each fraction was also analyzed for radioactivity, and fractions 18 and 19 contained 0.57 and 0.54 labels/ $\alpha 2$, respectively.

As discussed above, $\beta'2$ contains only $0.6\text{ Y}\cdot$ and thus gives rise to protein that does not form a complex. The relative amounts of the two peaks (Figure 3) support the proposal that the percentage of active quaternary structure correlates with the amount of the $\text{Y}\cdot$. Together with the observations of 0.5 equiv of sugar/ $\alpha 2$ attached and 0.6 equiv of cytosine released after inactivation, these results support a model where 1 equiv of F_2CDP inactivates the two active sites of $\alpha 2$ by enhancing subunit interactions.

Given the poor resolution of the SEC method, a control experiment in which F_2CDP was omitted was carried out under identical conditions (Figure 3D). The mixture analyzed by SEC gave two protein peaks: one with a retention time of 20.9 min and an apparent molecular mass of 543 kDa and the second with a retention time of 26.2 min and an apparent molecular mass of 109 kDa. The SDS-PAGE analysis of the first peak showed only the presence of α , which corresponds to $\alpha 6$, and the second peak showed the presence of both α and β' corresponding to α and $\beta'2$ based on the apparent MW.

An Additional Control: SEC To Examine the Quaternary Structure of hRNRs with Y138F- β' in the Presence of F_2CDP and ATP. To gain further insight into the quaternary structure of hRNR without covalent modification in the presence of F_2CDP , Y138F- β' in which the essential $\text{Y}\cdot$ of $\beta'2$ was replaced with an F was studied. The reaction mixture was identical to that described above except that Y138F- β' replaced wt- β' . Analysis of the reaction mixture by SEC showed that a small portion of protein migrated between 20 and 22.5 min, with apparent MWs ranging from 389 to 712 kDa followed by a second protein peak with a retention time at 26.9 min and an apparent MW of 88 kDa (Figure 4, Table 3). No radioactivity eluted with either peak when $[1\text{'-}^3\text{H}]\text{-F}_2\text{CDP}$ was used. SDS-PAGE analysis revealed that both α and β' were present in the peak that eluted from 20 to 22.5 min, but the ratio of α : β' ranged from 5.8 to 9.1 by comparison with standards (Figure S5 of the Supporting Information). The results suggest a complex equilibrium between nucleotide analogue/ATP and subunits, consistent with the existence of $\alpha\beta'2$, $\alpha 6$, and other species prior to the chemistry of inactivation.

DISCUSSION

The mechanism of inactivation of RNRs by F_2CDP is obviously very complex. However, despite this complexity, the RNRs from *E. coli*, *L. leichmannii*, and the two RNRs from humans share common and mechanistically/structurally informative features. Stoichiometric amounts of inhibitor are sufficient for complete inactivation (10, 11). In the case of the class I RNRs, this inhibition prevents the second α/β pair from reaction because of the tight and unusual subunit interactions (11). In all cases, the protein becomes covalently modified in a chemically fragile state by a sugar moiety generated from F_2CDP , and the cofactor ($\text{Y}\cdot$ or adenosylcobalamin) is partially inactivated. In all cases, if one waits long enough, two fluoride ions and one cytosine are released, requiring ketone formation at the C3' position of the nucleotide during the inactivation process. Unexpectedly, the α subunit is observed to have two conformations in about an approximately 1:1 ratio on SDS-PAGE analysis in the absence of heat. Mutagenesis studies suggest that the second conformation is likely the result of a cross-link between a cysteine in the C-terminus of α and its active site C218, directly involved in water loss in the normal reduction

Table 3: Molecular Mass Determination of Inactivated hRNR by SEC

protein (effector)	retention time (min)	apparent mass (kDa)	calcd mass (kDa)	oligomeric state
β'	26	116	81	$\beta'2$
α	25.6 ^a	88	92	α
α (100 μ M TTP)	24 ^a	189	184	$\alpha2$
α , β' (ATP)	20.9, 26.2	543, 109	552, 184, 81	$\alpha6$, $\alpha2$, $\beta'2$
α , β' inactivated by F ₂ CDP	18.6, 25.6	794, 116	798	$\alpha6\beta'6$ complex ^b
α , Y138F- β' inactivated by F ₂ CDP	20–22.5, 26.9	389–712, 88	633, 552, 184, 81	$\alpha6\beta'2$, $\alpha6$, $\alpha2$, $\beta'2$

^aSuperose 12 column was used. ^bUnreacted $\alpha2$, $\beta'2$ were also present.




FIGURE 4: SEC on a S200 column to detect complex formation of α /Y138F- β' incubated with [$1'$ -³H]-F₂CDP. The elution buffer contained 0.5 mM ATP. (A) The elution profile monitored by $A_{280\text{nm}}$ and scintillation counting shows no radiolabel. (B) Fractions 20–23 in (A) were monitored by 10% SDS–PAGE. Lane 1–8 are *E. coli* α and β (0.05–0.4 μ M) standards at different concentrations analyzed by SDS–PAGE to determine the ratio of α , Y138F- β' .

process. The key to the function of RNR is understanding the gymnastics of the structurally floppy, and thus structurally inaccessible, C-terminus of β/β' and C-terminus of α .

From our present observations information about the subunit architecture/interactions results. The altered α is likely to be formed from an intramolecular cross-link, rather than a cross-link from the C-terminus of the adjacent α in the $\alpha2$ dimer. If the cross-link were intermolecular, α should migrate on the SDS–PAGE gel as a dimer and would likely be sensitive to β -ME treatment. Rereduction of the active site disulfide in one α by the C-terminal tail of the second α in the $\alpha2$ complex had previously been suggested by pre-steady-state *in vitro* studies on *E. coli* RNR by Ericksson (49) and *in vivo* studies of *S. cerevisiae* RNR by Huang and co-workers (50).

Our previous pre-steady-state experiments (51) with mutations of the C-terminal cysteines were not inconsistent with the Ericksson proposal. In the Huang studies, the kinetics of rereduction of the active site disulfide is not measured due to the *in vivo* nature of the experiments. We have previously shown

that the two cysteines in the C-terminal tail of *E. coli* RNR can function as a thioredoxin, and this observation is also likely to be applicable to the *S. cerevisiae* RNR and could account for their *in vivo* results (47).

A second observation of interest is that the C-terminus of α must have access to the active site of the same α , while β remains bound. If our interpretation of the altered migratory properties of α is correct, a recent structure of a complex of a class Ib RNR from *Salmonella typhimurium* at 4.5 Å resolution may be informative in this regard (52). In this structure, the two subunits are present with a 360 Å² interface. The interface, however, is very informative with respect to the function of the C-termini of each subunit. One can observe the end of β (residue 285), a gap of 20 amino acids, and the last 11 amino acids of β bound in a hydrophobic patch formed by 3 helices ($\alpha10$, $\alpha1$, and αD) within α . One can also locate the C-terminus (residue 699 missing the last 15 amino acids) of α within the structure and its position relative to β . One β is interacting with one α in a fashion that, with tightening, could align the radical transfer pathway between the two subunits in a chemically competent fashion. In this $\alpha\beta$ the active site cysteines are oxidized, and some electron density exists for the ADP substrate. The second α has no substrate bound and no contact with the second β . Examination of the locations of the C-termini of the subunits suggests that the tails of α and β could be intertwined and that complete dissociation of the subunits might require a push from an interaction with a second $\alpha2$ interacting with the dangling $\beta2$. This model would be consistent with the kinetics of inactivation by F₂CDP in all the class I RNRs where a 1:1 complex of subunits is inactive, while assaying for activity of one subunit in the presence of an excess of the other subunit leads to some recovery of $\beta2$ activity as all the Y• has not been destroyed.

Uhlin has suggested that the structure of the class Ib RNR complex is a snapshot of one of the steps during the RNR-mediated reduction: either an initial step in subunit interaction or a final step after the nucleotide reduction process. The unusual conformation of α in the F₂CDP-inactivated RNR complex suggests that it might be trapping the enzyme in an asymmetrical configuration as well. The tight interaction between the subunits subsequent to inactivation by F₂CDP suggests that crystallization of the complex may be possible. Recently, we have obtained crystals of the *E. coli* RNR that diffract to 6 Å resolution (Stock and Drennan, unpublished work). Obviously, higher resolution structures are essential to figure out the key role of the tails of the subunits in the mechanism of ribonucleotide reduction. Despite the absence of molecular understanding, our studies suggest that both human RNRs can be targeted by F₂CDP and that the consequences of inactivation may differ depending on growth conditions.

ACKNOWLEDGMENT

We thank Dr. Yun Yen (City of Hope National Medical Center) for providing phRRM1 and phRRM2, Dr. Lars Thelander (Umea University, Sweden) for providing the p53R2, and Eli Lilly for providing [5-³H]-F₂C and 2-deoxy-3,5-di-*O*-benzoyl-3,3-difluororibonolactone. We thank Ellen Minnihan in our laboratory for thoughtful comments.

SUPPORTING INFORMATION AVAILABLE

Table 1, the primers for mutants of β' and α ; Figure S1, analysis of products generated during inactivation of $\alpha_n(\beta')_m$ by [5-³H]-F₂CDP; Figure S2, SDS-PAGE analysis of the α/β inactivated by F₂CDP and ATP; Figure S3, gel filtration molecular mass standards; Figure S4, SEC controls with $\beta'2$, α , and $\alpha2$ with TTP; Figure S5, *E. coli* α and β standards at different concentrations analyzed by SDS-PAGE to determine the ratio of α , β' in Figure 3B. This material is available free of charge via the Internet at <http://pubs.acs.org>.

REFERENCES

- Stubbe, J., and van der Donk, W. A. (1998) Protein radicals in enzyme catalysis. *Chem. Rev.* 98, 705–762.
- Hertel, L. W., Boder, G. B., Kroin, J. S., Rinzel, S. M., Poore, G. A., Todd, G. C., and Grindey, G. B. (1990) Evaluation of the antitumor-activity of gemcitabine (2',2'-difluoro-2'-deoxycytidine). *Cancer Res.* 50, 4417–4422.
- Huang, P., Chubb, S., Hertel, L. W., Grindey, G. B., and Plunkett, W. (1991) Action of 2',2'-difluorodeoxycytidine on DNA-synthesis. *Cancer Res.* 51, 6110–6117.
- Plunkett, W., Huang, P., and Gandhi, V. (1997) Gemcitabine: actions and interactions. *Nucleosides Nucleotides* 16, 1261–1270.
- Rivera, F., López-Tarruella, S., Vega-Villegas, M. E., and Salcedo, M. (2009) Treatment of advanced pancreatic cancer: from gemcitabine single agent to combinations and targeted therapy. *Cancer Treat. Rev.* 35, 335–339.
- Danesi, R., Altavilla, G., Giovannetti, E., and Rosell, R. (2009) Pharmacogenomics of gemcitabine in non-small-cell lung cancer and other solid tumors. *Pharmacogenomics* 10, 69–80.
- Bergman, A. M., Pinedo, H. M., and Peters, G. J. (2002) Determinants of resistance to 2',2'-difluorodeoxycytidine (gemcitabine). *Drug Resist. Updates* 5, 19–33.
- Nordlund, N., and Reichard, P. (2006) Ribonucleotide reductases. *Annu. Rev. Biochem.* 75, 681–706.
- Baker, C. H., Banzon, J., Bollinger, J. M., Stubbe, J., Samano, V., Robins, M. J., Lippert, B., Jarvi, E., and Resvick, R. (1991) 2'-deoxy-2'-methylenecytidine and 2'-deoxy-2',2'-difluorocytidine 5'-diphosphates—potent mechanism-based inhibitors of ribonucleotide reductase. *J. Med. Chem.* 34, 1879–1884.
- van der Donk, W. A., Yu, G. X., Perez, L., Sanchez, R. J., Stubbe, J., Samano, V., and Robins, M. J. (1998) Detection of a new substrate-derived radical during inactivation of ribonucleotide reductase from *Escherichia coli* by gemcitabine 5'-diphosphate. *Biochemistry* 37, 6419–6426.
- Wang, J., Lohman, G. J., and Stubbe, J. (2007) Enhanced subunit interactions with gemcitabine-5'-diphosphate inhibit ribonucleotide reductases. *Proc. Natl. Acad. Sci. U.S.A.* 104, 14324–14329.
- Artin, E., Wang, J., Lohman, G. J. S., Yokoyama, K., Yu, G., Griffin, R. G., Barr, G., and Stubbe, J. (2009) Insight into the mechanism of inactivation of ribonucleotide reductase by gemcitabine 5'-diphosphate in the presence and absence of reductant, *Biochemistry* (DOI 10.1021/bi901590q).
- Kashlan, O. B., and Cooperman, B. S. (2003) Comprehensive model for allosteric regulation of mammalian ribonucleotide reductase: refinements and consequences. *Biochemistry* 42, 1696–1706.
- Rofougaran, R., Vodnala, M., and Hofer, A. (2006) Enzymatically active mammalian ribonucleotide reductase exists primarily as an $\alpha(6)\beta(2)$ octamer. *J. Biol. Chem.* 281, 27705–27711.
- Sjöberg, B. M., and Reichard, P. (1977) Nature of free-radical in ribonucleotide reductase from *Escherichia coli*. *J. Biol. Chem.* 252, 536–541.
- Sjöberg, B. M., Reichard, P., Gräslund, A., and Ehrenberg, A. (1978) The tyrosine free radical in ribonucleotide reductase from *Escherichia coli*. *J. Biol. Chem.* 253, 6863–6865.
- Tanaka, H., Arakawa, H., Yamaguchi, T., Shiraishi, K., Fukuda, S., Matsui, K., Takei, Y., and Nakamura, Y. (2000) A ribonucleotide reductase gene involved in a p53-dependent cell-cycle checkpoint for DNA damage. *Nature* 404, 42–49.
- Nakano, K., Balint, E., Ashcroft, M., and Vousden, K. H. (2000) A ribonucleotide reductase gene is a transcriptional target of p53 and p73. *Oncogene* 19, 4283–4289.
- Pontarin, G., Fijolek, A., Pizzo, P., Ferraro, P., Rampazzo, C., Pozzan, T., Thelander, L., Reichard, P. A., and Bianchi, V. (2008) Ribonucleotide reduction is a cytosolic process in mammalian cells independently of DNA damage. *Proc. Natl. Acad. Sci. U.S.A.* 105, 17801–17806.
- Yamaguchi, T., Matsuda, K., Saqiya, Y., Iwade, M., Fujino, M. A., Nakamura, Y., and Arakawa, H. (2001) p53R2-dependent pathway for DNA synthesis in a p53-regulated cell cycle checkpoint. *Cancer Res.* 61, 8256–8262.
- Xue, L., Zhou, B., Liu, X., Qiu, W., Jin, Z., and Yen, Y. (2003) Wild-type p53 regulates human ribonucleotide reductase by protein-protein interaction with p53R2 as well as hRRM2 subunits. *Cancer Res.* 63, 980–986.
- Bourdon, A., Minai, L., Serre, V., Jais, J., Sarzi, E., Aubert, S., Chretien, D., Lonlay, P., Paquis-fluckinger, V., Arakawa, H., Nakamura, Y., Munnich, A., and Rötig, A. (2007) Mutation of RRM2B, encoding p53-controlled ribonucleotide reductase (p53R2), causes severe mitochondrial DNA depletion. *Nat. Genet.* 39, 776–780.
- Chang, L., Zhou, B., Hu, S., Guo, R., Liu, X., Jones, S. N., and Yen, Y. (2008) ATM-mediated serine 72 phosphorylation stabilizes ribonucleotide reductase small subunit p53R2 protein against MDM2 to DNA damage. *Proc. Natl. Acad. Sci. U.S.A.* 105, 18519–18524.
- Piao, C., Jin, M., Kim, H. B., Lee, S. M., Amatya, P. N., Hyun, J. W., Chang, I. Y., and You, H. J. (2009) Ribonucleotide reductase small subunit p53R2 suppresses MEK-ERK activity by binding to ERK kinase 2. *Oncogene* 28, 2173–2184.
- Shao, J. M., Zhou, B. S., Zhu, L. J., Qiu, W. H., Yuan, Y. C., Xi, B. X., and Yen, Y. (2004) In vitro characterization of enzymatic properties and inhibition of the p53R2 subunit of human ribonucleotide reductase. *Cancer Res.* 64, 1–6.
- Guittet, O., Håkansson, P., Voevodskaya, N., Fridt, S., Gräslund, A., Arakawa, H., Nakamura, Y., and Thelander, L. (2001) Mammalian p53R2 protein forms an active ribonucleotide reductase in vitro with the R1 protein, which is expressed both in resting cells in response to DNA damage and in proliferating cells. *J. Biol. Chem.* 276, 40647–40651.
- Liu, X., Xue, L., and Yen, Y. (2008) Redox property of ribonucleotide reductase small subunit M2 and p53R2. *Methods Mol. Biol.* 477, 195–206.
- Silva, D. J., Stubbe, J., Samano, V., and Robins, M. J. (1998) Gemcitabine 5'-triphosphate is a stoichiometric mechanism-based inhibitor of *Lactobacillus leichmannii* ribonucleoside triphosphate reductase: Evidence for thyl radical-mediated nucleotide radical formation. *Biochemistry* 37, 5528–5535.
- Lohman, G. J. S., and Stubbe, J. (2009) Inactivation of *Lactobacillus leichmannii* ribonucleotide reductase by F₂CTP: covalent modification (part I). *Biochemistry* (manuscript in preparation).
- Lohman, G. J. S., Gerfen, G. J., and Stubbe, J. (2009) Inactivation of *L. leichmannii* ribonucleotide reductase by F₂CTP: adenosylcobalamin destruction and formation of a nucleotide based radical, *Biochemistry* (manuscript in preparation).
- Stubbe, J., and van der Donk, W. A. (1995) Ribonucleotide reductases: radical enzymes with suicidal tendencies. *Chem. Biol.* 12, 793–801.
- Wang, X., Zhenchuk, A., Wiman, K. G., and Albertioni, F. (2009) Regulation of p53R2 and its role as potential target for cancer therapy. *Cancer Lett.* 276, 1–7.
- Lunn, C. A., Kathju, S., Wallace, B. J., Kushner, S. R., and Pigiet, V. (1984) Amplification and purification of plasmid-encoded thioredoxin from *Escherichia coli* K12. *J. Biol. Chem.* 259, 469–474.
- Russell, M., and Model, P. (1985) Direct cloning of the trxB gene that encodes thioredoxin reductase. *J. Bacteriol.* 163, 238–242.
- Parkin, S. E., Chen, S., Ley, B. A., Mangravite, L., Edmondson, D. E., Huynh, B. H., and Bollinger, J. M. Jr. (1998) Electron injection through a specific pathway determines the outcome of oxygen activation at the diiron cluster in the F208Y mutant of *Escherichia coli* ribonucleotide reductase protein R2. *Biochemistry* 37, 1124–1130.
- Berglund, O., and Eckstein, F. (1974) ATP- and dATP-substituted agaroses and the purification of ribonucleotide reductases. *Methods Enzymol.* 34, 253–261.
- Perlstein, D. L., Ge, J., Ortigosa, A. D., Robblee, J. H., Zhang, Z., Huang, M., and Stubbe, J. (2005) The active form of the

- Saccharomyces cerevisiae* ribonucleotide reductase small subunit is a heterodimer in vitro and in vivo. *Biochemistry* 44, 15366–15377.
38. Steeper, J. R., and Steuart, C. C. (1970) A rapid assay for CDP reductase activity in mammalian cell extracts. *Anal. Biochem.* 34, 123–130.
39. Salowe, S. P., and Stubbe, J. (1986) Cloning, overproduction, and purification of the B2 subunit of ribonucleoside-diphosphate reductase. *J. Bacteriol.* 165, 363–366.
40. Ortigosa, A. D., Hristova, D., Perlstein, D. L., Zhang, Z., Huang, M. X., and Stubbe, J. (2006) Determination of the in vivo stoichiometry of tyrosyl radical per beta beta' in *Saccharomyces cerevisiae* ribonucleotide reductase. *Biochemistry* 45, 12282–12294.
41. Hristova, D., Wu, C. H., Jiang, W., Krebs, C., and Stubbe, J. (2008) Importance of the maintenance pathway in the regulation of the activity of *Escherichia coli* ribonucleotide reductase. *Biochemistry* 47, 3989–3999.
42. van der Donk, W. A., Yu, G., Silva, D. J., and Stubbe, J. (1996) Inactivation of ribonucleotide reductase by (*E*)-2'-fluoromethylene-2'-deoxycytidine 5'-diphosphate: a paradigm for nucleotide mechanism based inhibitors. *Biochemistry* 35, 8381–8391.
43. Zipse, H., Artin, E., Wnuk, S., Lohman, G. J., Martino, D., Griffin, R. G., Kacprzak, S., Kaupp, M., Hoffman, B., Bennati, M., Stubbe, J., and Lees, N. (2009) Structure of the nucleotide radical formed during reaction of CDP/TTP with the E441Q- α 2 β 2 of *E. coli* ribonucleotide reductase. *J. Am. Chem. Soc.* 131, 200–211.
44. Xu, H., Faber, C., Uchiki, T., Racca, J., and Dealwis, C. (2006) Structures of eukaryotic ribonucleotide reductase I define gemcitabine diphosphate binding and subunit assembly. *Proc. Natl. Acad. Sci. U.S.A.* 103, 4028–4033.
45. Mao, S. S., Johnston, M. I., Bollinger, J. M., and Stubbe, J. (1989) Mechanism-based inhibition of a mutant *Escherichia coli* ribonucleotide reductase (cysteine-225-serine) by its substrate CDP. *Proc. Natl. Acad. Sci. U.S.A.* 86, 1485–1489.
46. Mao, S. S., Holler, T. P., Bollinger, J. M. Jr., Yu, G. X., Johnston, M. I., and Stubbe, J. (1992) Interaction of C225SR1 mutant subunit of ribonucleotide reductase with R2 and nucleoside diphosphates: tales of a suicidal enzyme. *Biochemistry* 31, 9744–9751.
47. Mao, S. S., Holler, T. P., Yu, G. X., Bollinger, J. M., Booker, S., Johnston, M. I., and Stubbe, J. (1992) A model for the role of multiple cysteine residues involved in ribonucleotide reduction: amazing and still confusing. *Biochemistry* 31, 9733–9743.
48. Lin, A. N. I., Ashley, G. W., and Stubbe, J. (1987) Location of the redox-active thiols of ribonucleotide reductase: sequence similarity between the *Escherichia coli* and *Lactobacillus leichmannii* enzymes. *Biochemistry* 26, 6905–6909.
49. Eriksson, H. K. (2001) Kinetics in the pre-steady state of the formation of cystines in ribonucleoside diphosphate reductase: evidence for an asymmetric complex. *Biochemistry* 40, 9631–9637.
50. Zhang, Z., Yang, K., Chen, C., Feser, J., and Huang, M. (2007) Role of the C terminus of the ribonucleotide reductase large subunit in enzyme regeneration and its inhibition by Sml1. *Proc. Natl. Acad. Sci. U.S.A.* 104, 2217–2222.
51. Ge, J., Yu, G. X., Ator, M. A., and Stubbe, J. (2003) Pre-steady-state and steady-state kinetic analysis of *E. coli* class I ribonucleotide reductase. *Biochemistry* 42, 10071–10083.
52. Uppsten, M., Färnegårdh, M., Domkin, V., and Uhlin, U. (2006) The first holocomplex structure of ribonucleotide reductase gives new insight into its mechanism of action. *J. Mol. Biol.* 359, 365–377.