

Enthalpy–Entropy Contribution to Carcinogen-Induced DNA Conformational Heterogeneity[†]

Fengting Liang and Bongsup P. Cho*

Department of Biomedical and Pharmaceutical Sciences, University of Rhode Island, Kingston, Rhode Island 02881

Received September 17, 2009; Revised Manuscript Received November 3, 2009

ABSTRACT: DNA damage by adduct formation is a critical step for the initiation of carcinogenesis. Aromatic amines are strong inducers of environmental carcinogenesis. Their DNA adducts are known to exist in an equilibrium between the major groove (B) and base-displaced stacked (S) conformations. However, the factors governing such heterogeneity remain unclear. Here we conducted extensive calorimetry/NMR/CD studies on the model DNA lesions caused by fluorinated 2-aminofluorene (FAF) and 4-aminobiphenyl (FABP) in order to gain thermodynamic and kinetic insights into the S/B conformational equilibrium. We demonstrate that there are large differences in enthalpy–entropy compensations for FABP and FAF. The small and flexible FABP exclusively adopts the less perturbed B conformer with small enthalpy ($\Delta\Delta H$ -2.7 kcal/mol)/entropy ($\Delta\Delta S$ -0.7 eu) change. In contrast, FAF stacks better and exists as a mixture of B and S conformers with large enthalpy ($\Delta\Delta H$ -13.4 kcal/mol)/entropy ($\Delta\Delta S$ -34.2 eu) compensation. van't Hoff analysis of dynamic ¹⁹F NMR data indicated $\Delta H_{B\leftrightarrow S}$ = 4.1 kcal/mol in favor of the B conformer and $\Delta S_{B\leftrightarrow S}$ = 15.6 cal mol⁻¹ K⁻¹ in favor of the intercalated S conformer. These findings demonstrate that the favorable entropy of the S conformer over B conformer determines the S/B population ratios at physiological temperatures.

The human genome is under constant assault from various sources of endogenous and exogenous chemical and radiation insult. Adduct formation is an important type of DNA damage that has been implicated in the etiology of various human cancers. Although occurring at a low frequency, DNA adducts in certain functionally active regions of the genome can elicit devastating molecular and cellular consequences (1). For example, mutations on the p53 tumor suppressor gene that alters transcriptional activity are responsible for approximately 50% of all cancers (2, 3). Elucidating the mechanisms of adduct-induced mutagenesis is an essential first step to developing effective strategies for risk assessment and novel chemotherapy targets (4–6).

We are particularly interested in the structure–function relationship of arylamine–DNA adducts, as they display a unique sequence-dependent conformational heterogeneity (7, 8). Depending on the location of the hydrophobic carcinogen on the DNA, the prototype arylamine carcinogen 2-aminofluorene (AF) adduct can exist as a mixture of two well-defined conformations: base-displaced stacked (S) or major groove B-type (B) (Figure 1) (9, 10). The S/B population ratio is dependent upon the 3'-flanking sequence; that is, 3'-purine promotes S-conformation (11, 12). The intrinsic fluorescence of 2-aminopurine positioned on the adducted strand flanking the AF on its 5'-side has been utilized to emphasize the important role of base stacking and hydrophobic interactions for sequence-dependent S/B equilibrium (13). Quantum mechanics calculations have further indicated the involvement of electrostatic interaction (12). In addition, S/B equilibrium is dependent on the size, coplanarity, and topology of the arylamine carcinogen. For example, B conformation

is predominant for the flexible one-ring aniline (14) and two-ring aminobiphenyl (ABP) adducts (15), whereas the larger amine adducts such as 1-aminopyrene (AP) (16) and isoquinoline (IQ) exist predominantly as S and/or wedge-type (W) conformers (17–19), the latter resulting from the carcinogen being pressed further into the narrow minor groove. However, the food-borne heterocyclic amine imidazopyridine (PhIP) is similar in size to AF but differs in topology so that the freely rotatable phenyl at C6 can be accommodated by the minor groove (20). As a result, a significant shift in the equilibrium toward the S conformer occurs in a temperature-dependent manner (20).

The AF-induced S/B heterogeneity may permit specific macromolecular interactions, thereby influencing binding affinity and substrate selectivity in repair and translesion synthesis (TLS) pathways, respectively. Recent crystallographic studies showed low electron densities around a weakly bound AF moiety in the active site, giving credence to the solution/solid comparability of adduct structures (21, 22). Likewise, we have found that S conformeric AF adducts are more efficiently excised in the *Escherichia coli* UvrABC nucleotide excision repair (NER) system (12). This study represents the first reported case of conformation-specific repair and provides molecular insights into the mechanisms of protein–DNA interactions involved in NER.

Despite the wealth of structural knowledge accumulated thus far, the factors defining carcinogen-induced conformational heterogeneity remain uncharacterized and its biological implications unclear. The objective of the present study was to gain thermodynamic and kinetic insights into the arylamine-induced S/B equilibrium using dynamic ¹⁹F NMR/CD¹ and DSC measurements.

[†]We are grateful to the NIH (R01CA098296) for financial support of this work. This research was also made possible in part by the use of the Research Core Facility supported by the NCRR/NIH (P20 RR016457) and NSF EPSCoR Proteomics (0554548).

*To whom correspondence should be addressed. E-mail: bcho@uri.edu. Telephone: 401-874-5024. Fax: 401-874-5766.

¹Abbreviations: FAF, *N*-(2'-deoxyguanosin-8-yl)-7-fluoro-2-aminofluorene; FABP, *N*-(2'-deoxyguanosin-8-yl)-4'-fluoro-4-aminobiphenyl; CD, circular dichroism; DSC, differential scanning calorimetry; ¹⁹F NMR, fluorine nuclear magnetic resonance spectroscopy; ICD_{290–350nm}, induced circular dichroism at 290–350 nm.




FIGURE 1: (a) Chemical structures of FAFP- and FAF-modified DNA. (b) AF-induced S/B-conformational heterogeneity. B and S refers to B-type and stacked conformers, respectively.

Our results indicate that enthalpy–entropy compensation is important for modulating bulky adduct-induced conformational heterogeneity.

EXPERIMENTAL PROCEDURES

Caution: ABP and AF derivatives are suspected human mutagens/carcinogens and therefore must be handled with caution.

Adduct Synthesis. Following the published procedures (12, 23), we prepared FAFP- and FAF-modified 11-mer oligonucleotides d(5'-CCATCG*CAACC-3'), in which G* is either an FAFP- or FAF-modified dG adduct (Figure 1a). The HPLC system used for purification of the oligos consisted of a Hitachi EZChrom Elite unit (Ibaraki-ken, Japan) with a L2450 diode array as a detector and employed a Phenomenex Luna C18 column (10 × 150 mm, 2.5 μ m) (Torrance, CA) with a 40 min gradient system involving 3–15% acetonitrile in pH 7.0 ammonium acetate buffer (0.10 M) with a flow rate of 2.0 mL/min. The modified oligos were characterized by ESI-Linear ion trap/FTMS analysis as reported previously (12) (Supporting Information Figures S1–S2) and annealed individually with the appropriate complementary strand to produce the desired duplexes. An identical set of unmodified duplexes was also prepared as controls.

UV-Melting and Circular Dichroism (CD). UV-melting data were obtained using a Beckman DU 800 UV/vis spectrophotometer using the general procedures reported previously (12, 24). A typical melting experiment consisted of forward/reverse scans and was repeated three times. Thermodynamic parameters were calculated using the program MELTWIN version 3.5. CD measurements were conducted on a Jasco J-810 spectropolarimeter using the published procedures (12, 25). Details on UV and CD are described in Supporting Information.

Dynamic ^{19}F NMR. Duplex samples (20–40 ODS) were dissolved in 300 μ L of a pH 7.0 buffer (100 mM NaCl, 10 mM Na_3PO_4 , and 100 μ M EDTA in 10% D_2O /90% H_2O) and filtered into through a Shigemi tube using a 0.2 μ m membrane filter. All ^1H and ^{19}F NMR results were recorded using a dedicated 5 mm $^{19}\text{F}/^1\text{H}$ dual probe on a Bruker DPX400 Avance spectrometer operating at 400.0 and 376.5 MHz, respectively, as described previously (24, 26). Details on NMR are found in Supporting Information.

DSC Measurements. Measurements were performed using a VP-DSC microcalorimeter (Microcal Inc., Northampton, MA).

Prior to temperature scanning, samples were degassed for at least 10 min under house vacuum in a closed vessel. Sample and reference solutions were loaded into the respective cells with a blunt-tip syringe. The cells were capped, and external pressure (~ 3 atm) was applied to prevent evaporation of the sample solution.

Raw data were collected as microwatts vs temperature. Typically, a 0.12 mM duplex sample solution was scanned against a control buffer from 15 to 85 $^\circ\text{C}$ at a rate of 0.75 $^\circ\text{C}/\text{min}$ with at least five repetitions. Usually, the first scan showed deviations due to the conditioning of the DSC cells to the temperature and to contact of solution and thus was excluded from further analysis. A buffer to buffer scan was subtracted from the sample scan, which was then normalized for the heating rate; i.e., each data point was divided by the corresponding heating rate. This results in base-corrected ΔC_p^{ex} vs temperature curves. Each transition showed negligible changes in heat capacity between the initial and final states; thus it was difficult to detect an appreciable $\Delta\Delta C_p^{\text{ex}}$ for the melting transitions; subsequently, $\Delta\Delta C_p^{\text{ex}} = 0$ was assumed. The area of the resulting curve is proportional to the transition heat, which, when normalized for the number of the moles of the sample, is equal to the transition enthalpy, ΔH :

$$\Delta H = \int \Delta C_p^{\text{ex}} dT$$

T_m was the temperature at half the peak area.

RESULTS

Model Systems. *In vivo* activation of arylamines and heterocyclic amines followed by reactions with cellular DNA results in N-deacetylated dG adduct formation as the major and the most persistent lesion (27). Due to the flexibility around the glycosidyl bond, these adducts can exist in an equilibrium between B and S conformations, as illustrated by the AF example in Figure 1b (7). As model systems, we prepared fully paired fluorine-labeled 7-fluoro-2-aminofluorene- (FAF-) and 4'-fluoro-4-aminobiphenyl- (FAFP-) modified 11-mer DNA duplexes [d(5'-CCATCG*CAACC-3')·d(5'-GGTTGCGATGG-3')] (G* = FAF and FAFP) (Figure 1a). The modified oligonucleotides were characterized by mass spectrometry (Supporting Information Figures S1 and S2). This duplex sequence has been used previously for structural investigations of various bulky adducts derived from arylamines




FIGURE 2: (a) ^{19}F NMR spectra of FABP- and FAF-modified 11-mer (b) CD and (c) UV spectra for FABP- and FAF-modified duplexes.

and polycyclic aromatic hydrocarbons (8, 25, 28). FAF and FABP are versatile fluorinated conformational probes for the potent carcinogens 2-aminofluorene (AF) and 4-aminobiphenyl (ABP), respectively (23, 26, 29–31).

UV Melting. The modified duplexes were significantly destabilized thermally and thermodynamically relative to the unmodified control in a wide range of concentrations, 0.3–10 μM (Supporting Information Figure S3). Adduct formation induced a major reduction in T_m of FAF and FABP at 0.1 mM by -8.5 and -10.7 $^{\circ}\text{C}$, respectively (Supporting Information Table 1). Free energy differences ($\Delta G_{37^{\circ}\text{C}}$) were similarly decreased by -3.2 and -3.3 kcal/mol, respectively. The more robust thermal destabilization exhibited by FABP was unexpected because of its conformational flexibility and lesser impact on the duplex structure (detailed below).

^{19}F NMR and $\text{ICD}_{290-350\text{nm}}$. The conformation difference between FABP- and FAF-modified duplexes is clearly evidenced by their ^{19}F NMR spectra obtained at 20 $^{\circ}\text{C}$ (Figure 2a). The FABP duplex exhibited a single homogeneous ^{19}F signal at -116.8 ppm. This signal was assigned to a B conformer because of its relatively large H/D isotope effect (~ 0.2 ppm) that indicated full exposure of the carcinogen to the solvent (23). The FAF duplex, on the other hand, yielded two signals at an approximate ratio of 34:66, favoring the S conformer. NOESY data (Supporting Information Figure S4) indicated that the two signals are gradually exchanged at ambient temperature. The peak at -117.4 ppm indicated a large H/D solvent effect (~ 0.2 ppm), whereas the upfield signal at -118.6 ppm showed a minimal effect (< 0.05 ppm) (Supporting Information Figures S5–S8). Consequently, these signals were designated as representing B and S conformers, respectively. The signal assignment was also verified by analyses of induced CD at 290 – 350 nm ($\text{ICD}_{290-350\text{nm}}$) (Figure 2b). We have previously shown that $\text{ICD}_{290-350\text{nm}}$ of a B conformer is mostly negative, while the W and S conformers are characterized by positive $\text{ICD}_{290-350\text{nm}}$ values (25). Figure 2b indicates a clear conformational difference between the FABP and FAF adducts, where a negative $\text{ICD}_{290-350\text{nm}}$ was observed for FABP and FAF exhibited a strong positive $\text{ICD}_{290-350\text{nm}}$. This observation is in contrast to the unmodified control which showed no noticeable ellipticities in the same region. For comparison, the UV spectra of FAF and FABP duplexes are shown in Figure 2c.

Microcalorimetry Data. To further investigate the thermodynamic impact of different carcinogens, the melting enthalpy of the duplexes with and without lesions was measured via differential scanning calorimetry (DSC). In contrast to UV melting, DSC procedures are model-independent, thus yielding more accurate measurements of enthalpy and T_m (31, 32). Figure 3 shows the DSC plots of excess heat capacity (C_p^{ex}) vs temperatures at a constant oligo concentration of 0.12 mM. The T_m of both FAF and FABP duplexes was reduced by -9.1 and -10.2 $^{\circ}\text{C}$, whereas $\Delta G_{37^{\circ}\text{C}}$ was decreased by -2.4 and -2.8 kcal/mol, respectively. Consistent with the UV-melting results (Supporting Information Figure 3 and Table 1), both FABP and FAF lesions produced significantly lower thermal (T_m) and thermodynamic (ΔG) stabilities of the duplex (Table 1).

Furthermore, the DSC results indicated that FABP modification produces a much smaller effect on the melting enthalpy, $\Delta\Delta H$ (-2.7 kcal/mol), compared to FAF (-13.4 kcal/mol) (Figure 3, Table 1). Given similar overall duplex thermodynamic stability ($\Delta\Delta G$), the large difference of enthalpy reduction between the FABP and FAF lesions must have originated from their differences in adduct conformation. For the FABP duplex, the carcinogen is bound in a major groove external conformation to a minimally perturbed B-DNA helix with all base pairs essentially intact and the modified dG in the normal *anti* glycosidic configuration (i.e., B conformer). Unlike FAF, FABP lacks a methylene group, thus allowing greater conformational flexibility. The flexible FABP in the major groove maintains its minimal interaction with the duplex, resulting in a small change in melting enthalpy ($\Delta\Delta H$ -2.7 kcal/mol) and entropy ($\Delta\Delta S$ -0.7 eu) (Table 1). Consequently, a small enthalpy–entropy compensation was obtained. For example, the enthalpy decreases ($\Delta\Delta H$ -2.7 kcal/mol) observed for the FABP adduct represent 3.4% loss of total enthalpy; nearly all enthalpy loss contributes to the free energy loss ($\Delta\Delta G_{37^{\circ}\text{C}}$ -2.4 kcal/mol), which represents almost 20% of the total free energy. The large thermal and thermodynamic destabilization observed for the FABP adduct is due to the mild interaction that lacks entropy compensation through adoption of a major groove binding B conformation.

In contrast, the flat FAF interacts strongly with the helical duplex, resulting in significant enthalpy reduction ($\Delta\Delta H$ 13.4 kcal/mol) with large entropy compensation (Figure 3). At 20 $^{\circ}\text{C}$, 66% of the FAF duplex adopts the S conformation with

both the modified dG and dC opposite the lesion being displaced in the major groove, while the FAF intercalated into the helix via a base displacement. The large enthalpy reduction observed is a reflection of the ruptured Watson–Crick base pairs and loss of normal stacking in the helix. In addition, the effect of the lesion on the duplex is not limited to the lesion site; it also propagates into the immediately flanking and next nearest neighbor base pairs. As shown in Figure 3, the larger entropy compensation, $T\Delta\Delta S$, causes $\Delta\Delta G$ to be similar to ABP modification (-2.8 vs -2.4 kcal/mol) at 37°C .

Dynamic ^{19}F NMR. Although DSC is a powerful technique, the conformational heterogeneity at or near the duplex melting temperature (T_m) makes it difficult to separate the respective contributions of B and S conformers to the thermodynamic destabilization of the duplex. Dynamic ^{19}F NMR, however,

provides a unique way to investigate the conformational dynamic and thermodynamic properties of a lesion-containing duplex by providing simplified NMR spectra in the lower temperature range ($5\text{--}40^\circ\text{C}$) (12, 23).

As shown in Figure 4a, the FAPB duplex exclusively exhibited B conformer signal throughout the entire temperature range ($5\text{--}60^\circ\text{C}$), with little change among the chemical shifts. These results make sense because the fully exposed characteristic of FAPB would protect against substantial effects resulting from B conformer to random single strand conformational fluctuations (12). For FAF (Figure 4b), however, the S conformer population was favorably interchanged in a gradual manner until it coalesced around 45°C and eventually stabilized at near complete melting at 60°C (see Supporting Information Figure S9 for complete spectral series).

The excellent separation of the B and S ^{19}F NMR signals allowed application of van't Hoff analysis, an efficient approach to estimate the relative energy difference between the B and S adduct conformers. The lower temperature range ($0\text{--}30^\circ\text{C}$) was used in order to minimize the interference of duplex melting on the conformational analysis (Figure 5). A matched NMR sample concentration (0.12 mM) from the DSC experiments was used. The van't Hoff plot ($\ln S/B$ vs $1/T$) indicated $\Delta H_{B\leftrightarrow S} = 4.1$ kcal/mol enthalpy difference in favor of the major groove binding B conformer, and $\Delta S_{B\leftrightarrow S} = 15.6$ cal mol $^{-1}$ K $^{-1}$ in favor of the intercalated S conformer (Figure 5a). Therefore, the S conformer exhibited greater stability at a physiologic temperature due to the favorable entropy (Figure 5b). It can be concluded that the increased stability exhibited by the S conformer in response to increasing temperatures is an entropy-driven event.

The van't Hoff data indicate that the free energy differences $\Delta G_{B\leftrightarrow S}(20^\circ\text{C}) = -0.39$ kcal/mol favor S over B conformers at 20°C . Previous studies of FAF have indicated stacking and van der Waal dispersive forces (collectively known as “hydrophobic effect”) as well as electrostatic effects as responsible for shifting the balance from B to S. Our present data indicate that these factors may not be sufficiently responsible for the preference of S conformers at a physiological temperature. In regard to the S conformer, the possible gain in enthalpy from carcinogen stacking was offset by rupture of hydrogen bonds, unfavorable steric interactions, and loss of base stacking induced by base displacement as well as misplaced nucleotides at the lesion site (12). B conformer is favored enthalpically and would be




FIGURE 3: (a) DSC and (b) thermodynamic plots of the FAPB- and FAF-modified duplexes and the control DNA duplex.

Table 1: Thermodynamic Parameters Derived from DSC Experiments

CCATCGCAACC GGTAGCGTTGG				
	ΔH kcal/mol	ΔS eu	ΔG_{37} kcal/mol	T_m °C
Control	79.1	214.9	12.45	64.5
FAPB	76.4	214.1	10.02	54.3
FAF	65.7	180.7	9.66	55.4
	$-\Delta\Delta H$ kcal/mol	$-\Delta\Delta S$ eu	$-\Delta\Delta G_{37}$ kcal/mol	$-\Delta T_m$ °C
FAPB	2.7	0.7	2.43	10.2
FAF	13.4	34.2	2.79	9.1




FIGURE 4: Dynamic ^{19}F NMR of the FABP- and FAF-modified DNA duplexes as a function of temperatures.

expected to dominate the balance at lower temperatures. However, S-related entropy compensation would lead to dominant S conformers at higher temperatures. The larger entropy in S conformers was likely an effect of the increased conformational freedom of the modified dG and opposite dC that are displaced into the major groove and the reorientation of water molecules in the hydration shell of the hydrophobic carcinogen moiety.

Combining the conformational thermodynamic results derived from DSC melting and ^{19}F NMR measurements, the destabilization effect of respective B and S conformers could be estimated. The ratio of B to S was approximately 25:75 at 30 °C (Figure 4b) at which the duplex melting process begins. The melting enthalpy ΔH of the integration DSC profile included contributions from both conformers, and their respective melting thermodynamic parameters are as follows:

$$\begin{aligned} \text{B conformer : } \Delta H^{\text{B}} &= 68.8 \text{ kcal/mol,} \\ \Delta S^{\text{B}} &= 192.0 \text{ cal/(mol K),} \\ \Delta G^{\text{B}}_{37^\circ\text{C}} &= 9.3 \text{ kcal/mol} \\ \text{S conformer : } \Delta H^{\text{S}} &= 64.7 \text{ kcal/mol,} \\ \Delta S^{\text{S}} &= 176.4 \text{ cal/(mol K),} \\ \Delta G^{\text{S}}_{37^\circ\text{C}} &= 10.0 \text{ kcal/mol} \end{aligned}$$

These results indicated that the FAF in a B conformer also interacts considerably with the duplex, resulting in reduction of melting enthalpy ($\Delta\Delta H = -10.3$ kcal/mol) and gain of entropy ($\Delta\Delta S = -22.9$ cal/(mol K)), although to a lesser degree than that in an S conformer ($\Delta\Delta H = -14.4$ kcal/mol, $\Delta\Delta S = -38.5$ cal/(mol K)). As shown above, FABP produces minimal disturbances with duplex structures by adopting the B conformer




FIGURE 5: (a) van't Hoff plot of the FAF-induced B/S conformational equilibrium as a function of $1/T$ (0–30 °C). (b) Conformational thermodynamics of B–S equilibrium derived from van't Hoff parameters.

state, resulting in nominal reduction of enthalpy and entropy compensation. The stronger interaction of the FAF B conformer, relative to FABP existing exclusively in B conformer state, may be a reflection of its larger ring size and more rigid nature. It is noteworthy that the interaction among the carcinogen, DNA duplex, and solvent system may produce a more complex situation. The differences between B and S conformer heat capacity, ΔC_p , might be significant because of considerable structural differences. Thus, the conformational enthalpy difference, $\Delta H_{\text{B/S}}$, and entropy difference, $\Delta S_{\text{B/S}}$, might be temperature-dependent.

Line Shape Analysis. A two-site dynamic exchange line shape was applied to calculate the kinetic parameters for the FAF duplex (Table 2) (12, 23). The S/B interconversion energy barrier (ΔG^\ddagger) obtained at 20 °C and using the Eyring equation,

Table 2: Kinetic and Thermodynamic Parameters Derived from Dynamic ^{19}F NMR Experiments for the FAF-modified DNA Duplex

B:S ratio, 20 °C	34%:66%
$\Delta G^\circ_{20^\circ\text{C}}$	-0.39 kcal/mol
ΔH°	4.04 kcal/mol
$k_{30^\circ\text{C}}$	300 s^{-1}
τ (1/ k)	3.33 ms
$k_{\text{C}} = 2.22\Delta\nu$	921 s^{-1}
$\Delta G^\ddagger_{20^\circ\text{C}}$	14.3 kcal/mol
ΔH^\ddagger	13.6 kcal/mol

$\Delta G^\ddagger_{\text{C}} = 4.58T_{\text{C}}(10.32 + \log T_{\text{C}}/k_{\text{C}})$ cal/mol, was 13.6 kcal/mol. The exchange rate constant at T_{C} ($k_{\text{C}} = 2.22\Delta\nu$) was determined to be 921 s^{-1} . The correlation time ($\tau = 1/k$) of the FAF duplex, which signifies the amount of time the lesion spends in the B conformer state before changing to S, or vice versa, was 3.3 ms at 20 °C. This value is comparable to the lifespan in G-tract G·C pairs but is about 10-fold smaller than those for regular G·C Watson–Crick base pairs (33, 34).

DISCUSSION

Our results showed a typical carcinogen-induced thermal (ΔT_{m}) and thermodynamic ($\Delta\Delta G$) destabilization upon FAFB and FAF modification (24, 35). $\Delta\Delta G$ is considered to be a more reliable predictor for lesion-induced destabilization because of its focus on the behavior of the DNA duplex at physiologically relevant temperatures (25 or 37 °C) (32). The exclusive B conformer FAFB interacts with the helix and causes minimal disturbance at the lesion site resulting in a small change in both enthalpy and entropy, both of which are compensated. The coplanar FAF, however, interacts significantly with the duplex, yielding an S/B conformational heterogeneity. Consequently, FAF duplex formation resulted in a major reduction in enthalpy changes, the bulk of which was compensated for by favorable entropy gained from the conformational freedom of the displaced base and the rearranged water molecules in the hydration shell of the hydrophobic carcinogen at the lesion site. This enthalpy–entropy compensation appears to minimize sequence-dependent variations in duplex stability ($\Delta G_{37^\circ\text{C}}$) (12, 24). The compensation effect could also explain why the B conformer is dominant at low temperatures, while the S conformer becomes favored at higher temperatures (12). Thus, favorability of the B conformer is enthalpic in origin at lower temperatures, but the equilibrium becomes entropically driven at higher temperatures and favors the S conformer. This appears to be the case regardless of the nature of sequence contexts flanking the FAF-modified DNA duplexes (12, 24). A greater stacking ability of the hydrophobic FAF, in comparison to FAFB, should play a role in shifting equilibrium toward the S conformer, but the stacking alone may not be sufficient to compensate for the loss of enthalpy resulting from ruptured hydrogen bonding, loss of base stacking, and unfavorable steric strain at the lesion site (12). It should be noted that the larger amine adducts such as aminopyrene (16) and isoquinoline (19) favor the S conformation. It is the favorable entropy of the S over B conformer that defines the relative S/B population ratio at a given temperature.

Pronounced enthalpy–entropy compensation has widely been observed for various weakly interacting macromolecular settings, including conformational fluctuations occurring during the melting of the DNA helix and the binding of small ligands or drugs to DNA and/or proteins (34, 36, 37). It is difficult to dissect the myriad of forces that govern the kinetic and thermodynamic

parameters involved in damaged DNA and protein phenomena. The molecular contacts between lesion-bound DNA and nearby amino acid residues, such as those in the active site of a polymerase or a repair protein, can also alter the conformational dynamics at the lesion site (38, 39). Thus, protein enzymatic actions can magnify the enthalpic effects of the lesion while minimizing the entropy compensation; indeed, the enthalpy differences induced by adduct formation may serve as a better indicator of how enzymes perform replication and repair proteins recognize DNA damage (40). Other, more subtle contributory factors, such as changes in solvation characteristics around and within the protein–DNA assembly complex, can further complicate the analysis (36). For instance, the enhanced free energy differences which guide the competition between correct and incorrect base pairing during DNA replication could be large when considered as enthalpy differences, presumably accounting for the fidelity of a polymerase (31). These scenarios appear to be generally valid for many entropy-driven biological systems.

The AF-induced S/B equilibrium is analogous to the so-called “base flipping”, a mostly enzyme-induced mechanism that is frequently involved in precarious epigenetic control of gene expression as well as base excision repair (BER) of a damaged DNA (33). In the latter, upon recognition of DNA damage a glycosylase utilizes base flipping to form a catalytic package in the active site of the enzyme, and then a glycosidic bond is broken to form an abasic site followed by insertion of the correct base. It has been shown that the enthalpy cost of removing a base from its helical stack is compensated for by the entropy gain from the increased conformational freedom gained upon base flipping (41, 42). Usually, the voided helix space created by base flipping is transiently occupied by nearby amino acids in the active site of the enzyme (43, 44). The AF-induced S/B equilibrium is different in that the duplex cavity created by the rotation of the glycosidyl bond of the modified dG at the lesion site is occupied by either the hydrophobic carcinogen (as with S conformer) or modified dG (as with B conformer) (12). The compensation effect observed for the S/B equilibrium, therefore, appears to be a consequence of the intrinsic ability of the carcinogen to influence its interaction with the helical DNA structure. The correlation time ($\tau = 1/k$) of the FAF duplex was found to be 3.3 ms at 20 °C. This value is comparable to the lifespan in G-tract G·C pairs but is about 10-fold smaller than those for regular G·C Watson–Crick base pairs (33, 34). The amount of time the lesion spends in the B state before changing to “S” includes an anti to syn conformational change of the glycosidic bond, which is not normal in unmodified DNA. Perhaps G tracts have some unusual properties, and the low correlation time for the adduct compared to normal G·C pairs may reflect the syn/anti transition. This situation is reminiscent of a model study conducted by Jiang et al. (45), in which the aromatic surrogate pyrene in a DNA strand opposite the uracil nucleotide acts both as a wedge to push the uracil away from the base stack in the free DNA and as a plug to hinder its reinsertion after base flipping, restoring the catalytic activity of mutant uracil DNA glycosylase. Using difluorophenyl as a model uracil, the same group has shown that the hydrophobic moiety is stabilized within the helix despite a rapid exchange rate ($k_{\text{C}} > 1200\text{ s}^{-1}$) between stacked and unstacked states, a range comparable to that observed for the S/B equilibrium ($k_{\text{C}} > 921\text{ s}^{-1}$) (45).

Base flipping is also observed in nucleotide excision repair (NER), the major pathway for repair of bulky DNA lesions (46).

During prokaryotic NER, lesions are excised through an elaborate two-tier damage recognition process involving initial binding to UvrA and verification of the damage by UvrB (47, 48). The tandem detection by UvrAB mediates the subsequent formation of a stable preincision complex using a padlock-like mode of binding in which a β -hairpin is inserted between the two strands of the DNA duplex in the vicinity of the DNA damage. Finally, a dual incision of the strand containing the lesion is carried out by UvrC. Correspondingly, Malta et al. (49) have shown that several bases around a cholesterol adduct, including the base immediately adjacent to the lesion on the 3'-side as well as its partner base on the complementary strand, are flipped out of the DNA duplex in NER. It has been shown that the yeast XPC orthologue Rad4 inserts a β -hairpin through the DNA duplex, causing the cyclobutane pyrimidine dimer (CPD) lesion to flip out of the double helix. The expelled nucleotides of the undamaged strand are recognized by Rad4, whereas the two CPD-linked nucleotides become disordered (50). Tubbs et al. (51) have recently shown that alkyl-transferase-like proteins (ATLs) generate a stable complex involving a nonenzymatic flipping of the endogenous alkyl-guanine lesions into a bulky lesion for the NER pathway. Flipping patterns involving NER proteins appear to be dependent on the nature of lesions.

We have shown previously that the S-conformeric FAF lesion is more susceptible to repair than the B conformer in the *E. coli* UvrABC system (11). The molecular basis of this conformation-specific NER is not well understood, although the presence of Watson–Crick hydrogen base pairs at the lesion site and groove surface characteristics could be important (12). The influence of sequence context and the quality of Watson–Crick base pairs at or in the vicinity of the lesion are also important. For example, the AF adduct in the TG**T* sequence context is incised by UvrABC proteins with consistently greater efficiency than in the CG**C* context (52–54). Similar results have been obtained for nucleotide excision repair of oxidative DNA damage (8-oxo-dG and FAPY) (55, 56) and DNA modified by the 10(S)-(+)-*trans*-anti-[BP]-N2-dG adduct (57), a major bulky adduct derived from a prominent environmental carcinogen. Results indicate a weakening of local base stacking interactions and poor quality of Watson–Crick base pairing at and in the vicinity of the lesion in the TG**T* sequence context. A similar sequence effect has been observed in human NER for the BP lesion; i.e., the ...CG*GC... sequence context elicited NER efficiency 4 times greater than did the ...CG**C*... context, and a destabilized Watson–Crick hydrogen bonding in the CG*GC elicited the most robust NER responses (58). Taken together, these results indicate that the local distortion of base pairing interactions constitutes a recognition element for the NER repair apparatus.

In summary, combined calorimetric/CD/¹⁹F NMR studies provide valuable mechanistic insights into how arylamine carcinogens interact with the duplex at the lesion site. Our results also demonstrate how thermodynamic parameters contribute to conformational equilibrium of the AF–DNA adduct. The AF-induced S/B equilibrium strongly depends on not only the nature of the carcinogen's structure and flanking sequences but also its entropy compensation effect. A more complete understanding of these types of entropy-driven conformational heterogeneity requires additional investigations of the adducts in other DNA sequences, alone or in combination with appropriate repair proteins or polymerases. These investigations are in progress in our laboratory.

SUPPORTING INFORMATION AVAILABLE

Experimental details, ESI-MS spectra (Figures S1 and S2) and UV melting curves (Figure S3) of FAFP- and FAF-modified oligonucleotides, NOESY/EXSY spectra of the FAF duplex (Figure S4), H/D isotope effects (Figures S5 and S7), imino proton spectra (Figures S6 and S7), full dynamic ¹⁹F NMR spectra (Figure S9), and UV melting thermodynamics (Table S1). This material is available free of charge via the Internet at <http://pubs.acs.org>.

REFERENCES

1. Vogelstein, B., and Kinzler, K. W. (1992) p53 function and dysfunction. *Cell* 70, 523–526.
2. Menendez, D., Ling, A., and Resnick, M. A. (2009) The expanding universe of p53 targets. *Nat. Rev. Cancer* 9, 724–737.
3. Greenblatt, M. S., Bennett, W. P., Hollstein, M. H., and Harris, C. C. (1994) Mutations in the p53 tumor suppressor gene: clues to cancer etiology and molecular pathogenesis. *Cancer Res.* 54, 4855–4878.
4. Ljungman, M. (2009) Targeting the DNA damage response in cancer. *Chem. Rev.* 109, 2929–2950.
5. Berdis, A. J. (2008) DNA Polymerases as therapeutic targets. *Biochemistry* 47, 8253–8260.
6. Hussain, S. P., Hollstein, M. H., and Harris, C. C. (2000) p53 tumor suppressor gene: at the crossroads of molecular carcinogenesis, molecular epidemiology, and human risk assessment. *Ann. N.Y. Acad. Sci.* 919, 79–85.
7. Cho, B. P. (2004) Dynamic conformational heterogeneities of carcinogen-DNA adducts and their mutagenic relevance. *J. Environ. Sci. Health, Part C: Environ. Carcinog. Ecotoxicol. Rev.* 22, 57–90.
8. Patel, D. J., Mao, B., Gu, Z., Hingerty, B. E., Gorin, A., Basu, A. K., and Broyde, S. (1998) Nuclear magnetic resonance solution structures of covalent aromatic amine-DNA adducts and their mutagenic relevance. *Chem. Res. Toxicol.* 11, 391–407.
9. Cho, B. P., Beland, F. A., and Marques, M. M. (1994) NMR structural studies of a 15-mer DNA duplex from a ras protooncogene modified with the carcinogen 2-aminofluorene: conformational heterogeneity. *Biochemistry* 33, 1373–1384.
10. Eckel, L. M., and Krugh, T. R. (1994) 2-Aminofluorene modified DNA duplex exists in two interchangeable conformations. *Nat. Struct. Biol.* 1, 89–94.
11. Meneni, S., Shell, S. M., Zou, Y., and Cho, B. P. (2007) Conformation-specific recognition of carcinogen-DNA adduct in *Escherichia coli* nucleotide excision repair. *Chem. Res. Toxicol.* 20, 6–10.
12. Meneni, S. R., Shell, S. M., Gao, L., Jurecka, P., Lee, W., Sponer, J., Zou, Y., Chiarelli, M. P., and Cho, B. P. (2007) Spectroscopic and theoretical insights into sequence effects of aminofluorene-induced conformational heterogeneity and nucleotide excision repair. *Biochemistry* 46, 11263–11278.
13. Jain, N., Reshetnyak, Y. K., Gao, L., Chiarelli, M. P., and Cho, B. P. (2008) Fluorescence probing of aminofluorene-induced conformational heterogeneity in DNA duplexes. *Chem. Res. Toxicol.* 21, 445–452.
14. Shapiro, R., Ellis, S., Hingerty, B. E., and Broyde, S. (1998) Effect of ring size on conformations of aromatic amine-DNA adducts: the aniline-C8 guanine adduct resides in the B-DNA major groove. *Chem. Res. Toxicol.* 11, 335–341.
15. Cho, B. P., Beland, F. A., and Marques, M. M. (1992) NMR structural studies of a 15-mer DNA sequence from a ras protooncogene, modified at the first base of codon 61 with the carcinogen 4-aminobiphenyl. *Biochemistry* 31, 9587–9602.
16. Mao, B., Vyas, R. R., Hingerty, B. E., Broyde, S., Basu, A. K., and Patel, D. J. (1996) Solution conformation of the N-(deoxyguanosin-8-yl)-1-aminopyrene ([AP]dG) adduct opposite dC in a DNA duplex. *Biochemistry* 35, 12659–12670.
17. Wang, F., DeMuro, N. E., Elmquist, C. E., Stover, J. S., Rizzo, C. J., and Stone, M. P. (2006) Base-displaced intercalated structure of the food mutagen 2-amino-3-methylimidazo[4,5-f]quinoline in the recognition sequence of the NarI restriction enzyme, a hotspot for –2 bp deletions. *J. Am. Chem. Soc.* 128, 10085–10095.
18. Elmquist, C. E., Wang, F., Stover, J. S., Stone, M. P., and Rizzo, C. J. (2007) Conformational differences of the C8-deoxyguanosine adduct of 2-amino-3-methylimidazo[4,5-f]quinoline (IQ) within the NarI recognition sequence. *Chem. Res. Toxicol.* 20, 445–454.
19. Wang, F., Elmquist, C. E., Stover, J. S., Rizzo, C. J., and Stone, M. P. (2007) DNA sequence modulates the conformation of the food

- mutagen 2-amino-3-methylimidazo[4,5-f]quinoline in the recognition sequence of the NarI restriction enzyme. *Biochemistry* 46, 8498–8516.
20. Brown, K., Hingerty, B. E., Guenther, E. A., Krishnan, V. V., Broyde, S., Turteltaub, K. W., and Cosman, M. (2001) Solution structure of the 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine C8-deoxyguanosine adduct in duplex DNA. *Proc. Natl. Acad. Sci. U.S.A.* 98, 8507–8512.
 21. Dutta, S., Li, Y., Johnson, D., Dzantiev, L., Richardson, C. C., Romano, L. J., and Ellenberger, T. (2004) Crystal structures of 2-acetylaminofluorene and 2-aminofluorene in complex with T7 DNA polymerase reveal mechanisms of mutagenesis. *Proc. Natl. Acad. Sci. U.S.A.* 101, 16186–16191.
 22. Hsu, G. W., Kiefer, J. R., Burnouf, D., Becherel, O. J., Fuchs, R. P. P., and Beese, L. S. (2004) Observing translesion synthesis of an aromatic amine DNA adduct by a high-fidelity DNA polymerase. *J. Biol. Chem.* 279, 50280–50285.
 23. Zhou, L., Rajabzadeh, M., Traficante, D. D., and Cho, B. P. (1997) Conformational heterogeneity of arylamine-modified DNA: ^{19}F NMR evidence. *J. Am. Chem. Soc.* 119, 5384–5389.
 24. Meneni, S. R., D'Mello, R., Norigian, G., Baker, G., Gao, L., Chiarelli, M. P., and Cho, B. P. (2006) Sequence effects of amino-fluorene-modified DNA duplexes: thermodynamic and circular dichroism properties. *Nucleic Acids Res.* 34, 755–763.
 25. Liang, F., Meneni, S., and Cho, B. P. (2006) Induced circular dichroism characteristics as conformational probes for carcinogenic amino-fluorene-DNA adducts. *Chem. Res. Toxicol.* 19, 1040–1043.
 26. Meneni, S., Liang, F., and Cho, B. P. (2007) Examination of the long-range effects of amino-fluorene-induced conformational heterogeneity and its relevance to the mechanism of translesional DNA synthesis. *J. Mol. Biol.* 366, 1387–1400.
 27. Beland, F. A., and Kadlubar, F. F. (1990) *Handbook of Experimental Pharmacology*, Springer-Verlag, Heidelberg.
 28. Geacintov, N. E., Cosman, M., Hingerty, B. E., Amin, S., Broyde, S., and Patel, D. J. (1997) NMR solution structures of stereoisomeric covalent polycyclic aromatic carcinogen-DNA adduct: principles, patterns, and diversity. *Chem. Res. Toxicol.* 10, 111–146.
 29. Jain, N., Li, Y., Zhang, L., Meneni, S. R., and Cho, B. P. (2007) Probing the sequence effects on NarI-induced -2 frameshift mutagenesis by dynamic ^{19}F NMR, UV, and CD spectroscopy. *Biochemistry* 46, 13310–13321.
 30. Cho, B. P., and Zhou, L. (1999) Probing the conformational heterogeneity of the acetylaminofluorene-modified 2'-deoxyguanosine and DNA by ^{19}F NMR spectroscopy. *Biochemistry* 38, 7572–7583.
 31. Liang, F., and Cho, B. P. (2007) Probing the thermodynamics of amino-fluorene-induced translesion DNA synthesis by differential scanning calorimetry. *J. Am. Chem. Soc.* 129, 12108–12109.
 32. Plum, G. E., Breslauer, K. J., and Roberts, R. W. (2002) *Thermodynamics and Kinetics of Nucleic Acid Association/Dissociation and Folding Processes*, Pergamon, Amsterdam.
 33. Priyakumar, U. D., and MacKerell, A. D., Jr. (2006) Computational approaches for investigating base flipping in oligonucleotides. *Chem. Rev.* 106, 489–505.
 34. Chen, C., and Russu, I. M. (2004) Sequence-dependence of the energetics of opening of at basepairs in DNA. *Biophys. J.* 87, 2545–2551.
 35. Geacintov, N. E., Broyde, S., Buterin, T., Naegeli, H., Wu, M., Yan, S., and Patel, D. J. (2002) Thermodynamic and structural factors in the removal of bulky DNA adducts by the nucleotide excision repair machinery. *Biopolymers* 65, 202–210.
 36. Petruska, J., and Goodman, M. F. (1995) Enthalpy-entropy compensation in DNA melting thermodynamics. *J. Biol. Chem.* 270, 746–750.
 37. Qu, X., Ren, J., Riccelli, P. V., Benight, A. S., and Chaires, J. B. (2003) Enthalpy/entropy compensation: influence of DNA flanking sequence on the binding of 7-amino actinomycin D to its primary binding site in short DNA duplexes. *Biochemistry* 42, 11960–11967.
 38. Broyde, S., Wang, L., Zhang, L., Rechkobl, O., Geacintov, N. E., and Patel, D. J. (2008) DNA adduct structure-function relationships: comparing solution with polymerase structures. *Chem. Res. Toxicol.* 21, 45–52.
 39. Guengerich, F. P. (2006) Interactions of carcinogen-bound DNA with individual DNA polymerases. *Chem. Rev.* 106, 420–452.
 40. Brabec, V., Stehlikova, K., Malina, J., Vojtiskova, M., and Kasparkova, J. (2006) Thermodynamic properties of damaged DNA and its recognition by xeroderma pigmentosum group A protein and replication protein A. *Arch. Biochem. Biophys.* 446, 1–10.
 41. O'Neil, L. L., and Wiest, O. (2008) Structures and energetics of base flipping of the thymine dimer depend on DNA sequence. *J. Phys. Chem. B* 112, 4113–4122.
 42. Klimasauskas, S., Szyperski, T., Serva, S., and Wuthrich, K. (1998) Dynamic modes of the flipped-out cytosine during HhaI methyltransferase-DNA interactions in solution. *EMBO J.* 17, 317–324.
 43. Huffman, J. L., Sundheim, O., and Tainer, J. A. (2005) DNA base damage recognition and removal: new twists and grooves. *Mutat. Res.* 577, 55–76.
 44. Beuck, C., Singh, I., Bhattacharya, A., Hecker, W., Parmar, V. S., Seitz, O., and Weinhold, E. (2003) Polycyclic aromatic DNA-base surrogates: high-affinity binding to an adenine-specific base-flipping DNA methyltransferase. *Angew. Chem., Int. Ed. Engl.* 42, 3958–3960.
 45. Jiang, Y. L., McDowell, L. M., Poliks, B., Studelska, D. R., Cao, C., Potter, G. S., Schaefer, J., Song, F., and Stivers, J. T. (2004) Recognition of an unnatural difluorophenyl nucleotide by uracil DNA glycosylase. *Biochemistry* 43, 15429–15438.
 46. Gillet, L. C. J., and Scharer, O. D. (2005) Molecular mechanisms of mammalian global genome nucleotide excision repair. *Chem. Rev.* 106, 253–276.
 47. Truglio, J. J., Croteau, D. L., Van Houten, B., and Kisker, C. (2006) Prokaryotic nucleotide excision repair: the UvrABC system. *Chem. Rev.* 106, 233–252.
 48. Pakotiprapha, D., Liu, Y., Verdine, G. L., and Jeruzalmi, D. (2009) A structural model for the damage-sensing complex in bacterial nucleotide excision repair. *J. Biol. Chem.* 284, 12837–12844.
 49. Malta, E., Moolenaar, G. F., and Goosen, N. (2006) Base flipping in nucleotide excision repair. *J. Biol. Chem.* 281, 2184–2194.
 50. Min, J. H., and Pavletich, N. P. (2007) Recognition of DNA damage by the Rad4 nucleotide excision repair protein. *Nature* 449, 570–575.
 51. Tubbs, J. L., Latypov, V., Kanugula, S., Butt, A., Melikishvili, M., Kraehenbuehl, R., Fleck, O., Marriott, A., Watson, A. J., Verbeek, B., McGown, G., Thorncroft, M., Santibanez-Koref, M. F., Millington, C., Arvai, A. S., Kroeger, M. D., Peterson, L. A., Williams, D. M., Fried, M. G., Margison, G. P., Pegg, A. E., and Tainer, J. A. (2009) Flipping of alkylated DNA damage bridges base and nucleotide excision repair. *Nature* 459, 808–813.
 52. Zou, Y., Shell, S. M., Utzat, C. D., Luo, C., Yang, Z., Geacintov, N. E., and Basu, A. K. (2003) Effects of DNA adduct structure and sequence context on strand opening of repair intermediates and incision by UvrABC nuclease. *Biochemistry* 42, 12654–12661.
 53. Jain, N., Meneni, S., Jain, V., and Cho, B. P. (2009) Influence of flanking sequence context on the conformational flexibility of amino-fluorene-modified dG adduct in dA mismatch DNA duplexes. *Nucleic Acids Res.* 37, 1628–1637.
 54. Ruan, Q., Liu, T., Kolbanovskiy, A., Liu, Y., Ren, J., Skovvaga, M., Zou, Y., Lader, J., Malkani, B., Amin, S., Van Houten, B., and Geacintov, N. E. (2007) Sequence context- and temperature-dependent nucleotide excision repair of a benzo[a]pyrene diol epoxide-guanine DNA adduct catalyzed by thermophilic UvrABC proteins. *Biochemistry* 46, 7006–7015.
 55. Kalam, M. A., and Basu, A. K. (2005) Mutagenesis of 8-oxoguanine adjacent to an abasic site in simian kidney cells: tandem mutations and enhancement of G→T transversions. *Chem. Res. Toxicol.* 18, 1187–1192.
 56. Kalam, M. A., Haraguchi, K., Chandani, S., Loechler, E. L., Moriya, M., Greenberg, M. M., and Basu, A. K. (2006) Genetic effects of oxidative DNA damages: comparative mutagenesis of the imidazole ring-opened formamidopyrimidines (Fapy lesions) and 8-oxo-purines in simian kidney cells. *Nucleic Acids Res.* 34, 2305–2315.
 57. Cai, Y., Patel, D. J., Geacintov, N. E., and Broyde, S. (2007) Dynamics of a benzo[a]pyrene-derived guanine DNA lesion in TGT and CGC sequence contexts: enhanced mobility in TGT explains conformational heterogeneity, flexible bending, and greater susceptibility to nucleotide excision repair. *J. Mol. Biol.* 374, 292–305.
 58. Kropachev, K., Kolbanovskii, M., Cai, Y., Rodriguez, F., Kolbanovskii, A., Liu, Y., Zhang, L., Amin, S., Patel, D., Broyde, S., and Geacintov, N. E. (2009) The sequence dependence of human nucleotide excision repair efficiencies of benzo[a]pyrene-derived DNA lesions: insights into the structural factors that favor dual incisions. *J. Mol. Biol.* 386, 1193–1203.