

# A Hearing Loss-Associated myo1c Mutation (R156W) Decreases the Myosin Duty Ratio and Force Sensitivity<sup>†</sup>

Tianming Lin,<sup>‡,||</sup> Michael J. Greenberg,<sup>‡,||,§</sup> Jeffrey R. Moore,<sup>§</sup> and E. Michael Ostap<sup>\*,‡</sup>

<sup>‡</sup>*Pennsylvania Muscle Institute and Department of Physiology, University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania 19104, United States, and* <sup>§</sup>*Department of Physiology and Biophysics, Boston University School of Medicine, Boston, Massachusetts 02118, United States. These authors contributed equally to this work.*

*Received October 18, 2010; Revised Manuscript Received January 20, 2011*

**ABSTRACT:** myo1c is a member of the myosin superfamily that has been proposed to function as the adaptation motor in vestibular and auditory hair cells. A recent study identified a myo1c point mutation (R156W) in a person with bilateral sensorineural hearing loss. This mutated residue is located at the start of the highly conserved switch 1 region, which is a crucial element for the binding of nucleotide. We characterized the key steps on the ATPase pathway at 37 °C using recombinant wild-type (myo1c<sup>31Q</sup>) and mutant myo1c (R156W-myo1c<sup>31Q</sup>) constructs that consist of the motor domain and three IQ motifs. The R156W mutation only moderately affects the rates of ATP binding, ATP-induced actomyosin dissociation, and ADP release. The actin-activated ATPase rate of the mutant is inhibited >4-fold, which is likely due to a decrease in the rate of phosphate release. The rate of actin gliding, as measured by the *in vitro* motility assay, is unaffected by the mutation at high myosin surface densities, but the rate of actin gliding is substantially reduced at low surface densities of R156W-myo1c<sup>31Q</sup>. We used a frictional loading assay to measure the effect of resisting forces on the rate of actin gliding and found that R156W-myo1c<sup>31Q</sup> is less force-sensitive than myo1c<sup>31Q</sup>. Taken together, these results indicate that myo1c with the R156W mutation has a lower duty ratio than the wild-type protein and motile properties that are less sensitive to resisting forces.

Myo1c is the single-headed member of the myosin superfamily that plays roles in trafficking of GLUT4-containing vesicles to the plasma membrane in response to insulin stimulation (1, 2) and in compensatory endocytosis following regulated exocytosis (3). Myo1c has also been proposed to play a key role in the process of adaptation in specialized sensory cells, where it is thought to dynamically adjust tension on mechanosensitive ion channels via its interaction with actin (4, 5).

Cell biological studies have shown that myo1c localizes and fractionates with cell membranes, and biochemical experiments have shown myosin I isoforms bind directly to phosphoinositides (6, 7). Additionally, mechanical experiments have shown that a related myosin isoform (myo1b) acts as a tension sensor by responding to small resisting loads by increasing its actin attachment lifetime, allowing it to generate and sustain tension for extended periods of time (8). Thus, evidence points to myo1c acting as a tension-sensing motor protein that links cellular membranes to the underlying actin cytoskeleton.

Consistent with the proposed role of myo1c acting as a tension-sensing adaptation motor in sensory cells, a myo1c point mutation (R156W) has recently been identified in an individual with bilateral sensorineural hearing loss (Figure 1) (9). R156 is highly conserved among the members of the myosin superfamily.

It is located at the start of the switch 1 region in the motor domain, which is a crucial structural element involved in nucleotide binding, although R156 is not proposed to interact directly with the nucleotide (10). Point mutations in switch 1 of myosins II and V affect nucleotide binding, ATP hydrolysis, phosphate release, and ADP release (11–14). Thus, the R156W mutation likely affects the motile and tension-sensing properties of myo1c. Indeed, the role of myo1c in adaptation may be particularly sensitive to even minor alterations in the ATPase kinetics and load-dependent mechanochemistry, because the processes of hearing and balance have stringent force and kinetic requirements (15, 16).

In this study, we determined the effect of the R156W mutation on key steps in the myo1c ATPase cycle (Scheme 1) using transient and steady-state kinetic techniques. We also determined the effect of this mutation on the motile properties of the motor. Experiments were performed with wild-type (myo1c<sup>31Q</sup>)<sup>1</sup> and mutated (R156W-myo1c<sup>31Q</sup>) myo1c constructs that consist of the motor domain, regulatory domain (contains three calmodulin-binding IQ motifs), and a C-terminal biotinylation tag for site-specific attachment of myosin for motility assays (Figure 1). We found that the R156W mutation causes a reduction in the myosin duty cycle, likely by reducing the rate of phosphate release. Furthermore, the mutation appears to cause a reduction in the

<sup>†</sup>This work was supported by grants from the National Institutes of Health to E.M.O. (GM57247), J.R.M. (HL077280), and M.J.G. (AR053461), and by the American Heart Association to M.G. (0815704D).

\*To whom correspondence should be addressed: Department of Physiology, University of Pennsylvania School of Medicine, B400 Richards Building, Philadelphia, PA 19104-6085. Phone: (215) 573-9758. Fax: (215) 898-2653. E-mail: ostap@mail.med.upenn.edu.

<sup>1</sup>Abbreviations: myo1c<sup>31Q</sup>, recombinant myosin IC consisting of the motor domain and three IQ motifs; R156W-myo1c<sup>31Q</sup>, myo1c<sup>31Q</sup> containing the R156W mutation; pyrene-actin, actin labeled at Cys-374 with pyrenyl iodoacetamide.





FIGURE 1: (A) Schematic of the expressed myo1b<sup>3IQ</sup> construct showing the relationship of the motor domain (large rectangle) to the IQ motifs (smaller numbered rectangles). The inset shows the positional relationship of the mutated residue (underlined) to switch 1 (bold). (B) Sodium dodecyl sulfate–polyacrylamide gel electrophoresis showing purified myo1c<sup>3IQ</sup> and R156W-myo1c<sup>3IQ</sup>. (C) Steady-state fluorescence emission spectra of 1  $\mu$ M (—) myo1c<sup>3IQ</sup> and (···) R156W-myo1c<sup>3IQ</sup> ( $\lambda_{\text{ex}}$  = 280 nm). The peak of the R156W-myo1c<sup>3IQ</sup> spectrum (335 nm) has a 1.3-fold higher intensity and is red-shifted compared to that of myo1c<sup>3IQ</sup> (332 nm).

#### Scheme 1



sensitivity of myo1c to resisting loads, possibly explaining the basis of the observed mutant phenotype.

## EXPERIMENTAL PROCEDURES

**Reagents, Proteins, and Buffers.** Rabbit skeletal muscle actin was prepared and gel filtered (17). Actin concentrations were determined by absorbance at 290 nm ( $\epsilon_{290}$  = 26600 M<sup>-1</sup> cm<sup>-1</sup>). Actin was labeled with pyrenyl iodoacetamide (pyrene-actin) and gel filtered (18). All actin was stabilized with a molar equivalent of phalloidin (Sigma). Calmodulin was expressed in *Escherichia coli* and purified as described previously (19). ADP and ATP concentrations were determined before each experiment by absorbance ( $\epsilon_{259}$  = 15400 M<sup>-1</sup> cm<sup>-1</sup>). Unless stated otherwise, all experiments were performed in KMg25 buffer (10 mM Mops, 25 mM KCl, 1 mM MgCl<sub>2</sub>, 1 mM EGTA, and 1 mM DTT). Unless otherwise stated, all experiments were conducted in the presence of 1  $\mu$ M free calmodulin (i.e., in excess of the calmodulin already bound to the myosin from the purification).

**Myosin I Expression and Purification.** Constructs consisting of the motor domain and three IQ motifs of mouse myo1c

were constructed, expressed in *Sf9* cells, and purified as described previously (20). The R156W-myo1c<sup>3IQ</sup> construct was made via QuickChange (Stratagene), and the sequence was verified by sequencing.

**Steady-State ATPase Measurements.** Steady-state ATPase activities were measured in KMg25 at 37 °C using the NADH-coupled assay as described previously (21, 22). The final protein concentrations after mixing were 100 nM myo1c<sup>3IQ</sup>, 0–175  $\mu$ M actin, and 1  $\mu$ M free calmodulin. Steady-state fluorescence measurements were taken with a Photon Technology International fluorometer.

**Transient Kinetic Measurements.** Transient kinetic measurements were taken with an Applied Photophysics (Surrey, U. K.) SX.18MV stopped-flow instrument (22). A 400 nm long-pass filter was used to monitor pyrene fluorescence ( $\lambda_{\text{ex}}$  = 365 nm), and a 320 nm long-pass filter was used to monitor intrinsic tryptophan fluorescence ( $\lambda_{\text{ex}}$  = 295 nm). Transients were fit to exponential functions using the software supplied with the stopped-flow instrument. All concentrations are given as final after mixing. Single-turnover ATPase measurements were fit to kinetic simulations of Scheme 2 in Berkley Madonna (<http://www.berkeleymadonna.com/>) to determine the rate constants. For the single-turnover measurements, the ratio of myosin to free calmodulin was maintained at 1:1.

**In Vitro Motility Assays.** In vitro motility assays using biotinylated myo1b<sup>3IQ</sup> and R156W-myo1b<sup>3IQ</sup> were performed in standard motility chambers at 37 °C as described previously (23). The temperature was controlled using an objective heater (Biopetechs). Nitrocellulose-coated coverslips were exposed to 0.1 mg/mL streptavidin and then blocked with 1 mg/mL BSA. Biotinylated myo1c<sup>3IQ</sup> (50–250 nM) was added to the coverslip and allowed to bind to the immobilized streptavidin. The rate of actin filament gliding was determined using Metamorph (Universal Imaging).

Frictional loading motility assays were performed using a procedure identical to that of the unloaded motility assays except that the desired amount of  $\alpha$ -actinin (Cytoskeleton Inc., Denver, CO) diluted in KMg25 buffer was added to the flow cell and allowed to incubate for 1 min before the addition of streptavidin. The myosin concentration added to the flow cell was 250 nM. The average gliding rate of 15–35 filaments was drift corrected and measured over five frames, captured at rates varying from 10 to 45 s/frame, using the freeware tracking software Retrac (<http://mc11.mcri.ac.uk/Retrac>). Single-exponential curves were fit to the data for visualization.

## RESULTS

**Protein Expression and Steady-State ATPase Activity.** We expressed wild-type and R156W recombinant myo1c protein constructs that contain the motor domain and three IQ motifs, including a C-terminal sequence for site-specific biotinylation (Figure 1). The presence of the additional tryptophan is apparent in the intrinsic fluorescence of the protein, as the steady-state emission spectrum ( $\lambda_{\text{ex}}$  = 280 nm) of R156W-myo1c<sup>3IQ</sup> has a peak intensity that is 1.3-fold greater than that of myo1c<sup>3IQ</sup> (Figure 1). There is also a 3 nm red shift in the emission peak of the R156W-myo1c<sup>3IQ</sup> protein.

The steady-state ATPase activity of myo1c<sup>3IQ</sup> is linearly activated by actin with an apparent second-order rate constant of  $v_a$  [ $v_a$  = 0.041 ( $\pm$ 0.0015)  $\mu$ M<sup>-1</sup> s<sup>-1</sup> (Figure 2)]. We were not able to achieve actin concentrations higher than 175  $\mu$ M because

of mixing artifacts resulting from high viscosities. This linear actin activation suggests that the maximal steady-state ATPase rate ( $V_{\max}$ ) is faster than  $7 \text{ s}^{-1}$ . It also indicates that the affinity of myo1c<sup>31Q</sup> for actin in the pre-power stroke states is very weak, as found for other myosin I isoforms (24, 25). The ATPase activity of R156W-myo1c<sup>31Q</sup> is also linearly related to the actin concentration [ $v_a = 0.010 (\pm 0.0004) \mu\text{M}^{-1} \text{ s}^{-1}$  (Figure 2)], but the rates





FIGURE 2: (A) Steady-state ATPase activity of myo1c. Actin concentration dependence of the steady-state ATPase rate of (●) myo1c<sup>31Q</sup> and (■) R156W-myo1c<sup>31Q</sup> measured using the NADH-coupled assay at 37 °C. Data for filled symbols were acquired in the presence of KMg25, and data for empty symbols were acquired in the same buffer in the absence of KCl. Solid lines are linear fits to the data acquired in KMg25 with slopes of  $0.041 \mu\text{M}^{-1} \text{ s}^{-1}$  for myo1c<sup>31Q</sup> and  $0.010 \mu\text{M}^{-1} \text{ s}^{-1}$  for R156W-myo1c<sup>31Q</sup>. (B) Single-turnover measurements of myo1c ATPase activity acquired in KMg25. Pyrene fluorescence transients were obtained by mixing  $0.75 \mu\text{M}$  ATP with a pre-equilibrated mixture of  $2.5 \mu\text{M}$  myosin and  $7.5 \mu\text{M}$  pyrene-actin at 37 °C. Five traces were averaged and normalized. Transients were fit to kinetic simulations of Scheme 2 (smooth lines) to obtain effective rate constants for detachment ( $k_{\text{off}} = 0.1 \mu\text{M}^{-1} \text{ s}^{-1}$  for myo1c<sup>31Q</sup>, and  $k_{\text{off}} = 0.28 \mu\text{M}^{-1} \text{ s}^{-1}$  for R156W-myo1c<sup>31Q</sup>) and for the strong binding of myosin to pyrene-actin ( $k_{\text{on}} = 0.05 \mu\text{M}^{-1} \text{ s}^{-1}$  for myo1c<sup>31Q</sup>, and  $k_{\text{on}} = 0.008 \mu\text{M}^{-1} \text{ s}^{-1}$  for R156W-myo1c<sup>31Q</sup>).

are 4-fold lower than that of myo1c<sup>31Q</sup>. ATPase rates obtained in the absence of added salt were only slightly faster [ $v_a = 0.046 (\pm 0.0052) \mu\text{M}^{-1} \text{ s}^{-1}$  for myo1c<sup>31Q</sup>, and  $v_a = 0.016 (\pm 0.0004) \mu\text{M}^{-1} \text{ s}^{-1}$  for R156W-myo1c<sup>31Q</sup>], indicating that the affinity of myo1c<sup>31Q</sup> for actin is less sensitive to ionic strength than other characterized myosins (26, 27).

To ensure that the measured steady-state ATPase activity of the mutant was not decreased because of the presence of inactive protein, we performed single-turnover ATPase measurements (Figure 2). ATP ( $0.75 \mu\text{M}$ ) was mixed with a pre-equilibrated complex of  $7.5 \mu\text{M}$  pyrene-labeled actin and  $2.5 \mu\text{M}$  myosin. Upon mixing, the intensity of fluorescence transients increased rapidly because of ATP-induced dissociation of myosin from pyrene-actin, which was followed by slow quenching because of repopulation of the strong binding state. Experiments performed with R156W-myo1c<sup>31Q</sup> had a quenching time course that was substantially slower than that of myo1c<sup>31Q</sup> (Figure 2). Thus, the rate of the kinetic step that limits entry into the quenched pyrene state is decreased for R156W-myo1c<sup>31Q</sup>. Data were fit via simulation to the following pathway:

Scheme 2



where  $A^*$  is the quenched state of actin,  $A^{**}$  is the high-fluorescence state of actin,  $k_{\text{off}}$  is the effective rate constant for ATP-induced dissociation of the pyrene-actomyosin complex, and  $k_{\text{on}}$  is the effective rate constant for myosin binding to actin and formation of a strongly bound state. Fits of the data yielded a  $k_{\text{on}}$  of  $0.05 \mu\text{M}^{-1} \text{ s}^{-1}$  for myo1c<sup>31Q</sup> and a  $k_{\text{on}}$  of  $0.008 \mu\text{M}^{-1} \text{ s}^{-1}$  for R156W-myo1c<sup>31Q</sup>, which are similar to those measured for the steady-state ATPase rates (Figure 2 and Table 1). These results confirm that the observed reduction of the ATPase activity of the mutant protein is due to kinetic changes in the myosin and also suggest that the rate-limiting step of the ATPase cycle is a transition that precedes entry into the strong binding state.

**ATP-Induced Population of the Weakly Bound States.** Pyrene-actin fluorescence was used to measure the rate of ATP binding and population of the weakly bound states at 37 °C (22, 28). Mixing actomyo1c with ATP resulted in transient increases in pyrene-actin fluorescence that were best fit to the sum of two exponential rates (Figure 3). The rates were

Table 1: Rates and Equilibrium Constants for the myo1c<sup>31Q</sup> and R156W-myo1c<sup>31Q</sup> ATPase<sup>a</sup>

	myo1c <sup>31Q</sup>	R156W-myo1c <sup>31Q</sup>
steady-state ATPase activity		
actin concentration dependence of the ATPase rate	$0.041 (\pm 0.0015) \mu\text{M}^{-1} \text{ s}^{-1}$	$0.010 (\pm 0.0004) \mu\text{M}^{-1} \text{ s}^{-1}$
$V_{\max}$	$> 7 \text{ s}^{-1}$	$> 2 \text{ s}^{-1}$
ATP binding		
$1/K_1'$	$220 (\pm 60) \mu\text{M}$	$250 (\pm 40) \mu\text{M}$
$k_2'$	$150 (\pm 15) \text{ s}^{-1}$	$230 (\pm 14) \text{ s}^{-1}$
$K_\alpha$	$1.2 (\pm 0.31)$	$4.2 (\pm 1.6)$
$k_{+\alpha}$	$37 (\pm 5.4) \text{ s}^{-1}$	$26 (\pm 3.1) \text{ s}^{-1}$
$k_{-\alpha}$	$31 (\pm 9.1) \text{ s}^{-1}$	$6.2 (\pm 2.5) \text{ s}^{-1}$
ATP hydrolysis		
$k_3^{\text{app}}$	$140 (\pm 9.0) \text{ s}^{-1}$	$160 (\pm 14) \text{ s}^{-1}$
ADP release		
$K_5'$	$1.8 (\pm 0.54) \mu\text{M}$	$1.6 (\pm 0.27) \mu\text{M}$
$k_{+5}'$	$24 (\pm 0.50) \text{ s}^{-1}$	$19 (\pm 0.087) \text{ s}^{-1}$
$k_{-5}'$	$13 (\pm 4.0) \mu\text{M}^{-1} \text{ s}^{-1}$	$12 (\pm 2.0) \mu\text{M}^{-1} \text{ s}^{-1}$

<sup>a</sup>Experiments performed in KMg25 [10 mM MOPS (pH 7.0), 25 mM KCl, 1 mM EGTA, 1 mM DTT, and 1 mM MgCl<sub>2</sub>] at 37 °C.





FIGURE 3: ATP-induced population of weakly bound actomyo1c states. Pyrene fluorescence transients obtained by mixing 1  $\mu\text{M}$  (●) myo1c<sup>31Q</sup>-pyrene-actin or (■) R156W-myo1c<sup>31Q</sup>-pyrene-actin with 16–1400  $\mu\text{M}$  ATP at 37 °C were fitted to a double-exponential function [ $k_{\text{obs}} = A_{\text{fast}}(1 - e^{-k_{\text{fast}}t}) + A_{\text{slow}}(1 - e^{-k_{\text{slow}}t})$ ]. The rates (A)  $k_{\text{fast}}$  and (B)  $k_{\text{slow}}$  are plotted as a function of ATP concentration. The fractional amplitude of the slow phase [ $A_{\text{slow}}/(A_{\text{slow}} + A_{\text{fast}})$ ] is also plotted (C) as a function of ATP concentration. Solid lines are the best fits of the data to eq 1.

hyperbolically related to the ATP concentration, with R156W-myo1c<sup>31Q</sup> having faster rates than myo1c<sup>31Q</sup>. The fast phase ( $k_{\text{fast}}$ ) of the increase in pyrene-actin fluorescence was modeled as ATP binding to the AM state and subsequent population of the AM. ATP state ( $K_1'k_{+2}'$ ), and the slow phase ( $k_{\text{slow}}$ ) was modeled as a transition ( $k_{+\alpha}$ ) from a nucleotide-insensitive state (AM) to a state that can bind ATP (AM') as proposed by Geeves (29):

Scheme 3



where A\* represents the unquenched fluorescent state of pyrene-actin. We analyzed the ATP dependence of the fast phase as

$$k_{\text{fast}} = \frac{k_{+2}'[\text{ATP}]}{\frac{1}{K_1'} + [\text{ATP}]} \quad (1)$$

where  $K_1'$  is a rapid equilibrium and  $k_{+2}'$  is a rate-limiting isomerization to the high-fluorescence AM.ATP state. Values for  $K_1'$  and  $k_{+2}'$  for the wild-type and mutant proteins are listed in Table 1.

At high ATP concentrations, the rate of the slow phase reports the isomerization of AM to AM' ( $k_{+\alpha}$ ), and the ratio of the amplitudes of the fast phase to the slow phase reports the equilibrium constant between AM and AM' (28, 29). Values for  $K_{\alpha}$ ,  $k_{+\alpha}$ , and  $k_{-\alpha}$  were determined by averaging points acquired at ATP concentrations of >300  $\mu\text{M}$ . Interestingly, R156W-myo1c<sup>31Q</sup> has a larger  $K_{\alpha}$  than the wild-type protein






FIGURE 4: Rate of ATP hydrolysis as measured by intrinsic tryptophan fluorescence. ATP concentration dependence of the rate of change in tryptophan fluorescence of (●) myo1c<sup>31Q</sup> and (■) R156W-myo1c<sup>31Q</sup> at 37 °C. Solid lines are hyperbolic fits, yielding maximal rates:  $k_3^{\text{app}} = 140 (\pm 9.0) \text{ s}^{-1}$  for myo1c<sup>31Q</sup>, and  $k_3^{\text{app}} = 160 (\pm 14) \text{ s}^{-1}$  for R156W-myo1c<sup>31Q</sup>. The inset shows time courses of a fluorescence increase after mixing proteins with 73  $\mu\text{M}$  ATP. The smooth lines are the best fits of the data to a single-exponential function with the following rates:  $k_{\text{obs}} = 110 (\pm 4.1) \text{ s}^{-1}$  for myo1c<sup>31Q</sup>, and  $k_{\text{obs}} = 120 (\pm 2.7) \text{ s}^{-1}$  for R156W-myo1c<sup>31Q</sup>.

Scheme 4



Scheme 5



(Table 1), which can be seen in a plot of the ATP dependence of the amplitude of the slow phase, which reports the mole fraction of the AM state (Figure 3 and Scheme 3).

**Intrinsic Tryptophan Fluorescence Changes in myo1c<sup>31Q</sup> and R156W-myo1c<sup>31Q</sup>.** Changes in the intrinsic tryptophan fluorescence of myo1c<sup>31Q</sup> and R156W-myo1c<sup>31Q</sup> upon mixing with ATP were determined by the stopped-flow method at 37 °C (Figure 4). Fluorescence time courses were best fit to a single-exponential function, and the rates were found to increase hyperbolically with ATP concentration (Figure 4). Previous kinetic reports associated this myo1c fluorescence change with the structural transition that accompanies ATP hydrolysis, as shown in Scheme 4 where M\* represents the enhanced fluorescence state (30).

Thus, the maximal rate of the fluorescence change reports the effective rate of ATP hydrolysis ( $k_3^{\text{app}} = k_{+3} + k_{-3}$ ) for the wild-type [ $k_3^{\text{app}} = 140 (\pm 9.0) \text{ s}^{-1}$ ] and mutant proteins [ $k_3^{\text{app}} = 160 (\pm 14) \text{ s}^{-1}$ ]. The amplitudes and rates of the fluorescent transients for the two proteins are similar, so it appears that the fluorescence of the tryptophan introduced via the point mutation does not change upon ATP binding or hydrolysis, despite its proximity to the nucleotide binding site.

**ADP Release.** The rate of ADP release [ $k_{+5}'$  (Scheme 1)] was determined by ATP-induced dissociation of the myo1c<sup>31Q</sup> constructs from pyrene-actin (22). When the active sites of myo1c<sup>31Q</sup> and R156W-myo1c<sup>31Q</sup> are saturated with ADP, ATP binding is rate-limited by the slow dissociation of ADP [ $k_5'$  (Scheme 5)].

ADP (10  $\mu\text{M}$ ) was incubated with 250 nM pyrene-actomyo1c and mixed with 500  $\mu\text{M}$  ATP, and transients were best fit by a






FIGURE 5: (A) Release of ADP from actomyosin<sup>31Q</sup>. The fractional amplitudes of the slow phase were obtained by fitting pyrene-actomyosin dissociation transients from experiments with (●) myo1c<sup>31Q</sup> or (■) R156W-myo1c<sup>31Q</sup> to a double-exponential function as a function of ADP concentration at 37 °C. Values were normalized to the total change in amplitude. The lines are fits of the (—) myo1c<sup>31Q</sup> and (---) R156W-myo1c<sup>31Q</sup> data to a hyperbolic function. The inset shows time courses of pyrene-actin fluorescence after mixing 500 μM ATP with 0.25 μM myo1c<sup>31Q</sup>-pyrene-actin and R156W-myo1c<sup>31Q</sup>-pyrene-actin equilibrated with 10 μM ADP. The smooth lines are best fits of the data to a single-exponential function. (B) Calculated duty ratio as a function of actin concentration for (—) myo1c<sup>31Q</sup> and (---) R156W-myo1c<sup>31Q</sup>. Duty ratios were calculated as defined in eq 2.

single-exponential function (Figure 5, inset). The rate of release of ADP from myo1c<sup>31Q</sup> [ $k_5' = 24 (\pm 0.50) \text{ s}^{-1}$ ] is slightly faster than the rate of release from R156W-myo1c<sup>31Q</sup> [ $k_5' = 19 (\pm 0.090) \text{ s}^{-1}$  (Table 1)].

ATP-induced pyrene-actin fluorescence transients were acquired as a function of ADP concentration to determine the affinity of ADP ( $K_5'$ ) for actin-bound myo1c<sup>31Q</sup> proteins. Time courses were fit to two-exponential functions, with the slow component reporting ADP release ( $k_5'$ ) and the rate of the fast phase reporting ATP binding ( $k_1'/k_{+2}'$ ). The affinity of the actomyosin<sup>31Q</sup> for ADP was determined by monitoring the change in the amplitude of the slow phase as a function of ADP concentration (Figure 5). Hyperbolic fits to the data yield similar affinities for myo1c<sup>31Q</sup> [ $K_5' = 1.8 (\pm 0.54) \mu\text{M}$ ] and R156W-myo1c<sup>31Q</sup> [ $K_5' = 1.6 (\pm 0.27) \mu\text{M}$  (Table 1)].

**In Vitro Motility Assays.** We determined the motile activity of myo1c<sup>31Q</sup> and R156W-myo1c<sup>31Q</sup> at 37 °C using the in vitro motility assay (23, 31). The myosin proteins contain a C-terminal biotin that allows site-specific attachment of the myosin to the motility surface (see Experimental Procedures). Incubation of 250 nM myosin in streptavidin-coated chambers resulted in actin gliding rates of myo1c<sup>31Q</sup> that are the same as that of R156W-myo1c<sup>31Q</sup> (Figure 6). This finding is consistent with the two proteins having similar rates of ADP release (Figure 5), as this step is expected to limit the rate of unloaded sliding at high ATP concentrations of several different myosin isoforms (e.g., refs 8 and 32–34).

Actin gliding velocity increased ~1.5-fold at lower myo1c<sup>31Q</sup> concentrations, which is likely due to a reduced drag on actin at lower surface densities. However, R156W-myo1c<sup>31Q</sup> did not




FIGURE 6: Velocity of actin filament gliding measured by the in vitro motility assay. Actin gliding rates were measured at five different surface densities of (●) myo1c<sup>31Q</sup> and (■) R156W-myo1c<sup>31Q</sup> at 37 °C. Myosin concentrations indicate the quantity of protein incubated in the flow chamber before washing (see Experimental Procedures). Gliding rates were determined over a 5 min time period as a function of actin filament length. Actin filaments with rates of 0 nm/s were not immobile but rather exhibited nondirectional diffusion. Points are the averages of 10 filaments, and error bars are standard deviations.




FIGURE 7: Frictional loading assays. Actin filament sliding velocity was measured as a function of α-actinin concentration for both (●) myo1c<sup>31Q</sup> and (■) R156W-myo1c<sup>31Q</sup> at 37 °C. Two-parameter exponential decays were fit to the data for visualization. Points are the average sliding velocity of 15–35 filaments averaged over five frames, and the error bars are standard errors of the mean.

exhibit increased motility rates at lower surface densities. Interestingly, when the concentration of R156W-myo1c<sup>31Q</sup> in the motility chambers was decreased, we found that shorter actin filaments did not undergo directed motility but rather appeared to be detached from the coverslip. This inability of R156W-myo1c<sup>31Q</sup> to power motility of short actin filaments is likely due to a decrease in the motor's duty ratio (see below).

We tested the ability of myosin to propel actin filaments in the presence of a load by performing frictional loading assays (35, 36). α-Actinin transiently binds to sliding actin filaments, providing a frictional load that opposes the driving force of the bed of myosin. The reduction in actin sliding velocity with load is due to both the drag force of the α-actinin and the load-dependent kinetics of the myosin. The rate of actin gliding was measured in the presence of a range of α-actinin surface densities (Figure 7). The motility rates of myo1c<sup>31Q</sup> decreased by 2-fold with the addition of 60 nM α-actinin, while R156W-myo1c<sup>31Q</sup> required approximately 100 nM α-actinin to cause the same

decrease in velocity. Thus, although sensitive to resisting loads, the sliding velocity of R156W-myolc<sup>31Q</sup> appears to be substantially less force-sensitive than that of the wild-type protein.

## DISCUSSION

**ATP-Induced Dissociation of Actomyolc<sup>31Q</sup> and ATP Hydrolysis.** The R156W mutation increases the rate of ATP-induced dissociation of myolc<sup>31Q</sup> from actin ( $k_2'$ ) ~1.5-fold. This elevated rate is not expected to affect the population of the actin-bound states, because it is substantially faster than the rates that limit the entry and exit from the force-bearing states (below). Consistent with this notion, the unloaded motility of R156W mutant myosin is unchanged compared to that of the wild type.

It has been proposed that the time course of ATP-induced changes in the intrinsic tryptophan fluorescence of myolc reports the rate of ATP hydrolysis (30), as shown for other myosins (e.g., refs 37 and 38). We found the rates of the fluorescence transients from the wild-type and mutant proteins to be fast and not significantly different (Table 1). Thus, it appears that the R156W mutation does not alter ATP hydrolysis kinetics, in contrast to other switch 1 mutations (11–13). It is also interesting to note that, in the absence of actin, conformational changes that have been reported to occur near switch 1 (39) do not affect the fluorescence emission of W156 in the mutant protein.

**Steady-State ATP Hydrolysis and Phosphate Release.** The R156W point mutation inhibits the actin-activated ATPase activity of myolc<sup>31Q</sup> ~4-fold. This inhibition most likely occurs at the phosphate release step [ $k_4'$  (Scheme 1)] for the following reasons. (a) The other key steps on the ATPase pathway ( $k_1'$ ,  $k_2'$ ,  $k_3$ , and  $k_5'$ ) are substantially faster than the rate that limits the steady-state ATPase activity (Table 1). (b) Single-turnover experiments (Figure 2) show that the rate of isomerization into a strongly bound state (a transition associated with phosphate release) is the same as the rate that limits the steady-state ATPase activity. (c) Phosphate release has been shown to be the rate-limiting step for other myosin I isoforms (24, 25, 28, 40). (d) The lack of an effect of the mutation on the velocity of actin gliding indicates that the inhibited step occurs while myosin is in a weak binding state (33).

We were not able to determine the rate constant for phosphate release ( $k_4'$ ) directly, so it is possible that the R156W mutation weakens actin affinity, which would lead to a decreased  $V_a$ . Switch 1 is not at the actin-binding interface (41), so the mutation is unlikely to affect actin binding via a direct disruption of the actin-binding site. However, the conformational state of switch 1 impacts actin affinity via modulation of the conformation of myosin's actin-binding cleft (14, 42). Therefore, this mutation may affect affinity by modulating the ability of the actin-binding cleft to close, which ultimately affects phosphate release. Nevertheless, the functional consequence of the R156W mutation is a decreased myosin duty ratio, which results in decreased force from groups of myosins (below).

**ADP Release, in Vitro Motility, and Duty Ratio.** The R156W mutation only slightly affects the rate of release of ADP ( $k_{+5}'$ ) from actomyolc (Table 1). Because ADP release limits the rate of detachment from the strong binding states, it is not surprising that the maximal actin gliding velocities are only slightly affected by the R156W mutation. This is true despite the fact that the steady-state ATPase activity of the mutant is inhibited 4-fold. This result is a strong indication that the steady-state ATPase rate is inhibited at a kinetic step that occurs

when myosin is detached from (or weakly bound to) actin filaments (33).

The duty ratio is the fraction of the ATPase cycle during which myosin is bound to actin in the strong binding (force-bearing) states. We can calculate the duty ratios of the wild-type and mutant myosins by assuming that the steady-state ATPase assay reports the rate that limits entry into the strong binding states ( $k_9k_4'$ ), and ADP release ( $k_5'$ ) limits exit from these states:

$$\text{duty ratio} = \frac{[A]v_A}{[A]v_A + k_{+5}'} \quad (2)$$

where  $v_A$  is the actin-dependent ATPase rate and  $[A]$  is the actin concentration (Figure 5) (43). At the maximal experimental actin concentration (175  $\mu\text{M}$ ), the duty ratio of R156W-myolc<sup>31Q</sup> is ~2.4-fold lower than that of the wild-type protein. This finding is consistent with the in vitro motility assays that show that lower surface densities of the mutant protein are less likely to propel short actin filaments (Figure 6); i.e., short actin filaments have fewer myosin binding sites, further reducing the probability of actomyosin interactions necessary for movement.

The frictional loading assays (Figure 7) indicate that load-induced inhibition of actin gliding is less pronounced in the mutant myosin. The ADP release step ( $k_5'$ ) has been shown to be the most load-dependent step for other myosin isoforms (8, 44–46), so it is likely that the load dependence of this step is affected by the R156W mutation. It is interesting to note that the R156W mutation increases the equilibrium constant ( $K_a$ ) that defines the transition between the nucleotide-sensitive (AM') and nucleotide-insensitive (AM) states (Figure 3 and Table 1). It has been proposed that myosins that undergo the AM-to-AM' transition are highly force-sensitive (29) and that this transition may be similar to the force-sensitive transition (AM·ADP to AM') during ATP cycling. Thus, an increase in the stabilization in  $K_a$  (i.e., a stabilization of the nucleotide-accessible AM' state) may be an indication of decreased tension sensitivity. Single-molecule measurements are required to address this speculation directly (8, 44).

**Comparison with Previous Studies.** A kinetic analysis comparing wild-type and R156W myolc was recently published by Adamek et al. (47) using myosin constructs in which the native light chain binding domain was replaced with a single  $\alpha$ -helix domain from myo10. The ATPase rate constants obtained by Adamek et al. were substantially slower than those measured in this study, and they did not provide information about the effect of R156W on the myolc duty ratio. Furthermore, the maximal steady-state ATPase rate measured with the construct containing the single  $\alpha$ -helix domain (0.66  $\text{s}^{-1}$ ) is more than 10-fold slower than the maximal rate measured here ( $>7 \text{ s}^{-1}$ ). It is possible that the proteins used by Adamek et al. had altered kinetics because of the absence of an intact regulatory domain (20). In a separate study by Adamek et al., utilizing myolc with a single IQ motif, the maximal steady-state ATPase rate measured for the wild-type myosin (1.8  $\text{s}^{-1}$ ) was only 25% of the rate measured here ( $>7 \text{ s}^{-1}$ ) (30). The reason for this discrepancy is unclear.

**Physiological Impact of the R156W Mutation.** Despite the normal sliding velocity of R156W-myolc<sup>31Q</sup> at low loads and high myosin densities, we propose that ensemble force generation by mutant myosins is impaired. The amount of force generated by a group of myosins is proportional to the number of motors that are strongly bound to actin. Because R156W-myolc<sup>31Q</sup> has a decreased duty ratio, an ensemble of mutant motors will generate

less force than wild-type motors. We also propose that the rate at which R156W-myolc<sup>31Q</sup> detaches from actin is less sensitive to resisting forces (Figure 7), resulting in an impaired ability to alter the duty ratio in response to tension (8). If myolc is in fact the "adaptation motor", this altered tension sensing may result in improper gating of mechanosensitive channels. Future mechanical experiments will better address the effect of force on ensembles and single myolc molecules.

A missense mutation associated with human deafness (E385D) has been identified in the switch II region of the myosin I family member myo1a (48). Like the R156W mutation in myolc, this mutation has a decreased actin-activated ATPase rate (49). However, this protein is unable to propel actin in gliding filament assays. Interestingly, this switch II mutation also results in the improper localization of the protein. Given the importance of the motor domain in myosin I localization (50, 51), it will be important to determine if the R156W mutation in myolc also has an effect on the subcellular localization of this molecular motor.

## REFERENCES

- Bose, A., Robida, S., Fuccinitti, P. S., Chawla, A., Fogarty, K., Corvera, S., and Czech, M. P. (2004) Unconventional myosin Myo1c promotes membrane fusion in a regulated exocytic pathway. *Mol. Cell. Biol.* 24, 5447–5458.
- Bose, A., Guilherme, A., Robida, S. I., Nicoloso, S. M., Zhou, Q. L., Jiang, Z. Y., Pomerleau, D. P., and Czech, M. P. (2002) Glucose transporter recycling in response to insulin is facilitated by myosin Myo1c. *Nature* 420, 821–824.
- Sokac, A. M., Schietroma, C., Gundersen, C. B., and Bement, W. M. (2006) Myosin-Ic couples assembling actin to membranes to drive compensatory endocytosis. *Dev. Cell* 11, 629–640.
- Holt, J. R., Gillespie, S. K., Provance, D. W., Shah, K., Shokat, K. M., Corey, D. P., Mercer, J. A., and Gillespie, P. G. (2002) A chemical-genetic strategy implicates myosin-Ic in adaptation by hair cells. *Cell* 108, 371–381.
- Stauffer, E. A., Scarborough, J. D., Hirono, M., Miller, E. D., Shah, K., Mercer, J. A., Holt, J. R., and Gillespie, P. G. (2005) Fast adaptation in vestibular hair cells requires myosin-Ic activity. *Neuron* 47, 541–553.
- Hokanson, D. E., and Ostap, E. M. (2006) Myo1c binds tightly and specifically to phosphatidylinositol 4,5-bisphosphate and inositol 1,4,5-trisphosphate. *Proc. Natl. Acad. Sci. U.S.A.* 103, 3118–3123.
- Hokanson, D. E., Laakso, J. M., Lin, T., Sept, D., and Ostap, E. M. (2006) Myo1c Binds Phosphoinositides through a Putative Pleckstrin Homology Domain. *Mol. Biol. Cell* 17, 4856–4865.
- Laakso, J. M., Lewis, J. H., Shuman, H., and Ostap, E. M. (2008) Myosin I can act as a molecular force sensor. *Science* 321, 133–136.
- Zadro, C., Alemanno, M. S., Bellacchio, E., Ficarella, R., Donaudy, F., Melchionda, S., Zelante, L., Rabionet, R., Hilgert, N., Estivill, X., Van Camp, G., Gasparini, P., and Carella, M. (2009) Are MYO1C and MYO1F associated with hearing loss? *Biochim. Biophys. Acta* 1792, 27–32.
- Smith, C. A., and Rayment, I. (1996) X-ray structure of the magnesium(II)-ADP-vanadate complex of the *Dictyostelium discoideum* myosin motor domain to 1.9 Å resolution. *Biochemistry* 35, 5404–5417.
- Shimada, T., Sasaki, N., Ohkura, R., and Sutoh, K. (1997) Alanine scanning mutagenesis of the switch I region in the ATPase site of *Dictyostelium discoideum* myosin II. *Biochemistry* 36, 14037–14043.
- Li, X. D., Rhodes, T. E., Ikebe, R., Kambara, T., White, H. D., and Ikebe, M. (1998) Effects of mutations in the  $\gamma$ -phosphate binding site of myosin on its motor function. *J. Biol. Chem.* 273, 27404–27411.
- Forgacs, E., Sakamoto, T., Cartwright, S., Belknap, B., Kovacs, M., Toth, J., Webb, M. R., Sellers, J. R., and White, H. D. (2009) Switch I mutation S217A converts myosin V into a low duty ratio motor. *J. Biol. Chem.* 284, 2138–2149.
- Kintsjes, B., Gyimesi, M., Pearson, D. S., Geeves, M. A., Zeng, W., Bagshaw, C. R., and Malnasi-Csizmadia, A. (2007) Reversible movement of switch I loop of myosin determines actin interaction. *EMBO J.* 26, 265–274.
- Gillespie, P. G. (2004) Myosin I and adaptation of mechanical transduction by the inner ear. *Philos. Trans. R. Soc. London, Ser. B* 359, 1945–1951.
- Batters, C., Wallace, M. I., Coluccio, L. M., and Molloy, J. E. (2004) A model of stereocilia adaptation based on single molecule mechanical studies of myosin I. *Philos. Trans. R. Soc. London, Ser. B* 359, 1895–1905.
- Spudich, J. A., and Watt, S. (1971) The regulation of rabbit skeletal muscle contraction. I. Biochemical studies of the interaction of the tropomyosin-troponin complex with actin and the proteolytic fragments of myosin. *J. Biol. Chem.* 246, 4866–4871.
- Kouyama, T., and Mihashi, K. (1981) Fluorimetry study of N-(1-pyrenyl)iodoacetamide-labelled F-actin. Local structural change of actin protomer both on polymerization and on binding of heavy meromyosin. *Eur. J. Biochem.* 114, 33–38.
- Putkey, J. A., Slaughter, G. R., and Means, A. R. (1985) Bacterial expression and characterization of proteins derived from the chicken calmodulin cDNA and a calmodulin processed gene. *J. Biol. Chem.* 260, 4704–4712.
- Manceva, S., Lin, T., Pham, H., Lewis, J. H., Goldman, Y. E., and Ostap, E. M. (2007) Calcium regulation of calmodulin binding to and dissociation from the myo1c regulatory domain. *Biochemistry* 46, 11718–11726.
- De La Cruz, E. M., Sweeney, H. L., and Ostap, E. M. (2000) ADP inhibition of myosin V ATPase activity. *Biophys. J.* 79, 1524–1529.
- De La Cruz, E. M., and Ostap, E. M. (2009) Kinetic and equilibrium analysis of the myosin ATPase. *Methods Enzymol.* 455, 157–192.
- Lin, T., Tang, N., and Ostap, E. M. (2005) Biochemical and motile properties of Myo1b splice isoforms. *J. Biol. Chem.* 280, 41562–41567.
- Ostap, E. M., and Pollard, T. D. (1996) Biochemical kinetic characterization of the *Acanthamoeba* myosin-I ATPase. *J. Cell Biol.* 132, 1053–1060.
- El Mezgueldi, M., Tang, N., Rosenfeld, S. S., and Ostap, E. M. (2002) The kinetic mechanism of Myo1e (human myosin-IC). *J. Biol. Chem.* 277, 21514–21521.
- Furch, M., Rammel, B., Geeves, M. A., and Manstein, D. J. (2000) Stabilization of the actomyosin complex by negative charges on myosin. *Biochemistry* 39, 11602–11608.
- Furch, M., Geeves, M. A., and Manstein, D. J. (1998) Modulation of actin affinity and actomyosin adenosine triphosphatase by charge changes in the myosin motor domain. *Biochemistry* 37, 6317–6326.
- Lewis, J. H., Lin, T., Hokanson, D. E., and Ostap, E. M. (2006) Temperature dependence of nucleotide association and kinetic characterization of myo1b. *Biochemistry* 45, 11589–11597.
- Geeves, M. A., Perreault-Micale, C., and Coluccio, L. M. (2000) Kinetic analyses of a truncated mammalian myosin I suggest a novel isomerization event preceding nucleotide binding. *J. Biol. Chem.* 275, 21624–21630.
- Adamek, N., Coluccio, L. M., and Geeves, M. A. (2008) Calcium sensitivity of the cross-bridge cycle of Myo1c, the adaptation motor in the inner ear. *Proc. Natl. Acad. Sci. U.S.A.* 105, 5710–5715.
- Kron, S. J., and Spudich, J. A. (1986) Fluorescent actin filaments move on myosin fixed to a glass surface. *Proc. Natl. Acad. Sci. U.S.A.* 83, 6272–6276.
- Rock, R. S., Rice, S. E., Wells, A. L., Purcell, T. J., Spudich, J. A., and Sweeney, H. L. (2001) Myosin VI is a processive motor with a large step size. *Proc. Natl. Acad. Sci. U.S.A.* 98, 13655–13659.
- Siemankowski, R. F., Wiseman, M. O., and White, H. D. (1985) ADP dissociation from actomyosin subfragment 1 is sufficiently slow to limit the unloaded shortening velocity in vertebrate muscle. *Proc. Natl. Acad. Sci. U.S.A.* 82, 658–662.
- Rief, M., Rock, R. S., Mehta, A. D., Mooseker, M. S., Cheney, R. E., and Spudich, J. A. (2000) Myosin-V stepping kinetics: A molecular model for processivity. *Proc. Natl. Acad. Sci. U.S.A.* 97, 9482–9486.
- Bing, W., Knott, A., and Marston, S. B. (2000) A simple method for measuring the relative force exerted by myosin on actin filaments in the in vitro motility assay: Evidence that tropomyosin and troponin increase force in single thin filaments. *Biochem. J.* 350 (Part 3), 693–699.
- Greenberg, M. J., and Moore, J. R. (2010) The molecular basis of frictional loads in the in vitro motility assay with applications to the study of the loaded mechanochemistry of molecular motors. *Cytoskeleton* 67, 273–285.
- De La Cruz, E. M., Wells, A. L., Rosenfeld, S. S., Ostap, E. M., and Sweeney, H. L. (1999) The kinetic mechanism of myosin V. *Proc. Natl. Acad. Sci. U.S.A.* 96, 13726–13731.
- Johnson, K. A., and Taylor, E. W. (1978) Intermediate states of subfragment 1 and actosubfragment 1 ATPase: Reevaluation of the mechanism. *Biochemistry* 17, 3432–3442.

39. Malnasi-Csizmadia, A., Dickens, J. L., Zeng, W., and Bagshaw, C. R. (2005) Switch movements and the myosin crossbridge stroke. *J. Muscle Res. Cell Motil.* 26, 31–37.
40. Jontes, J. D., Milligan, R. A., Pollard, T. D., and Ostap, E. M. (1997) Kinetic characterization of brush border myosin-I ATPase. *Proc. Natl. Acad. Sci. U.S.A.* 94, 14332–14337.
41. Rayment, I., Holden, H. M., Whittaker, M., Yohn, C. B., Lorenz, M., Holmes, K. C., and Milligan, R. A. (1993) Structure of the actin-myosin complex and its implications for muscle contraction. *Science* 261, 58–65.
42. Furch, M., Fujita-Becker, S., Geeves, M. A., Holmes, K. C., and Manstein, D. J. (1999) Role of the salt-bridge between switch-1 and switch-2 of *Dictyostelium* myosin. *J. Mol. Biol.* 290, 797–809.
43. De La Cruz, E. M., Wells, A. L., Sweeney, H. L., and Ostap, E. M. (2000) Actin and light chain isoform dependence of myosin V kinetics. *Biochemistry* 39, 14196–14202.
44. Laakso, J. M., Lewis, J. H., Shuman, H., and Ostap, E. M. (2010) Control of myosin-I force sensing by alternative splicing. *Proc. Natl. Acad. Sci. U.S.A.* 107, 698–702.
45. Veigel, C., Schmitz, S., Wang, F., and Sellers, J. R. (2005) Load-dependent kinetics of myosin-V can explain its high processivity. *Nat. Cell Biol.* 7, 861–869.
46. Veigel, C., Molloy, J. E., Schmitz, S., and Kendrick-Jones, J. (2003) Load-dependent kinetics of force production by smooth muscle myosin measured with optical tweezers. *Nat. Cell Biol.* 5, 980–986.
47. Adamek, N., Geeves, M. A., and Coluccio, L. M. (2011) Myo1c mutations associated with hearing loss cause defects in the interaction with nucleotide and actin. *Cell. Mol. Life Sci.* 68, 139–150.
48. Donaudy, F., Ferrara, A., Esposito, L., Hertzano, R., Ben-David, O., Bell, R. E., Melchionda, S., Zelante, L., Avraham, K. B., and Gasparini, P. (2003) Multiple mutations of MYO1A, a cochlear-expressed gene, in sensorineural hearing loss. *Am. J. Hum. Genet.* 72, 1571–1577.
49. Yengo, C. M., Ananthanarayanan, S. K., Brosey, C. A., Mao, S., and Tyska, M. J. (2008) Human deafness mutation E385D disrupts the mechanochemical coupling and subcellular targeting of myosin-1a. *Biophys. J.* 94, L5–L7.
50. Ruppert, C., Godel, J., Muller, R. T., Kroschewski, R., Reinhard, J., and Bahler, M. (1995) Localization of the rat myosin I molecules myr 1 and myr 2 and in vivo targeting of their tail domains. *J. Cell Sci.* 108 (Part 12), 3775–3786.
51. Tang, N., and Ostap, E. M. (2001) Motor domain-dependent localization of myo1b (myr-1). *Curr. Biol.* 11, 1131–1135.