


Crystal Structure of the Central Coiled-Coil Domain from Human Liprin- β 2

Ryan L. Stafford, Ming-Yun Tang, Michael R. Sawaya, Martin L. Phillips, and James U. Bowie*

Department of Chemistry and Biochemistry, UCLA-DOE Institute of Genomics and Proteomics, Molecular Biology Institute, University of California, Los Angeles, Boyer Hall 611 Charles E. Young Dr. E., Los Angeles, California 90095-1570, United States

S Supporting Information

ABSTRACT: Liprins are a conserved family of scaffolding proteins important for the proper regulation and development of neuronal synapses. Humans have four liprin- α s and two liprin- β s which all contain long coiled-coil domains followed by three tandem SAM domains. Complex interactions between the coiled-coil and SAM domains are thought to create liprin scaffolds, but the structural and biochemical properties of these domains remain largely uncharacterized. In this study we find that the human liprin- β 2 coiled-coil forms an extended dimer. Several protease-resistant subdomains within the liprin- β 1 and liprin- β 2 coiled-coils were also identified. A 2.0 Å crystal structure of the central, protease-resistant core of the liprin- β 2 coiled-coil reveals a parallel helix orientation. These studies represent an initial step toward determining the overall architecture of liprin scaffolds and understanding the molecular basis for their synaptic functions.



LAR family transmembrane protein–tyrosine phosphatase interacting proteins, or liprins, are a conserved family of proteins involved in the formation of the presynaptic cytomatrix of the active zone.^{1,2} The human genome encodes four liprin- α s and two liprin- β s, but only liprin- α s bind to LAR.² Liprin- α 1 has also been found to interact with numerous other proteins including CASK,³ ERC,⁴ GIT,⁵ KIF1A,⁶ RIM,⁷ and RSY-1.⁸ Liprin- α 1 was first directly linked to the neural synapse with the discovery that a mutation in the *C. elegans* liprin- α 1 homologue, *syd-2*, led to the development of abnormally large synaptic active zones and impaired synaptic transmission.⁹ Subsequently, liprin- α 1 has been shown to be important for photoreceptor targeting and selection in *Drosophila*,^{10,11} axon guidance in *C. elegans*,¹² and synapse development in rat hippocampal neurons.¹³ At postsynaptic sites, liprin- α 1 is involved in the trafficking of AMPA receptors through its interaction with GIT.⁵ Recently, the human protein kazrin, previously known for its involvement in the regulation of desmosome assembly, was found to be a member of the liprin family.¹⁴ Specifically, the kazrinE isoform has sequence similarity to liprins and also binds to LAR. In *Drosophila*, the kazrinE homologue, termed liprin- γ , was found to work in concert with liprin- α and liprin- β to regulate the development of photoreceptor and neuromuscular synapses.¹⁵

All liprins possess the same overall domain composition comprised of an N-terminal predicted coiled-coil region followed by three tandem SAM domains. The coiled-coil domains of liprin- β s are significantly shorter than liprin- α s (Figure 1A).² Immunoprecipitation experiments have shown that liprin- α s interact with other liprin- α s through their coiled-coil domains, but not liprin- β s.² Similarly, liprin- β coiled-coils interact with other liprin- β coiled-coil domains, but not liprin- α s. The C-terminal SAM domains appear to mediate interactions between liprin- α s and liprin- β s. The extensive interactions between liprins, coupled with their high expression in brain tissue, suggested early




Figure 1. Domain organization of liprin- β 1 and - β 2. (A) The diagram shows the relative sizes and domain composition of liprin- α s (blue) to liprin- β s (orange). (B) A COILS¹⁶ prediction using a 24-residue window for the coiled-coil domains from human liprin- β 1 (dashed line) and liprin- β 2 (solid line).

on that they possessed the ability to make complex neural scaffolding structures.²

The overall architectures of the presumed liprin scaffolds are unknown since there is little structural information for the coiled-coil or SAM domains from any of the liprin proteins. Though there are many reliable algorithms to determine the presence of

Received: January 27, 2011
Revised: March 16, 2011
Published: April 04, 2011

coiled-coil domains in proteins directly from the primary amino acid sequence,^{16–18} it remains challenging to determine the number and orientations of the coiled-coils in each oligomer.^{19–22} SAM domains are also readily identified by their amino acid sequence and usually function as protein–protein interaction motifs, but they can have many different functions, including binding to RNA, lipids, and forming homo- and heteropolymers.²³ In the case of the liprin SAM domains, they have been predicted to form polymers, but this has not been verified experimentally.²⁴ Understanding the individual structures of these coiled-coil and SAM domains from liprins should help explain how they form the scaffolds necessary to execute their function in synapse development.

In this study we have focused on determining the structure of the coiled-coil domains of human liprin- β s. Specifically, we cloned the complete coiled-coil domains of human liprin- β 1 and - β 2 and focused the majority of our efforts studying liprin- β 2. The coiled-coil domain of liprin- β 2 was found to be predominantly dimeric by equilibrium sedimentation and chemical cross-linking experiments. Three protease-resistant subdomains of the liprin- β 1 and - β 2 coiled-coils were also identified including the central third of the liprin- β 2 coiled-coil, for which we were able to determine a crystal structure.

MATERIALS AND METHODS

Cloning and Mutagenesis. The template DNA for cloning all human liprin- β 1 (Image ID 5276255; Accession # BC050281) and liprin- β 2 (Image ID 4826610; Accession # BC021714) constructs was obtained from Open Biosystems. The complete coiled-coil construct for liprin- β 1 was created by PCR amplifying DNA corresponding to residues G97 through T342 and inserting the fragment into a modified pET-3a vector between MluI and SalI restriction sites containing an N-terminal sequence (MEKTR) for efficient expression and a C-terminal His-tag (RRHHHHHH) for purification. The complete coiled-coil construct for liprin- β 2 was similarly cloned by inserting residues S100 through E315 into the same pET-3a plasmid restriction sites. The 21 kDa liprin- β 1 protease-resistant fragment from E110 to K290 was cloned into a pQE2 vector (Qiagen) between restriction sites SphI and SalI containing a cleavable N-terminal His-tag (MKHHHHHHHMHAK). The 19 and 9 kDa liprin- β 2 protease-resistant fragment from A102 to A265 and K185 to A265, respectively, were cloned into the same pQE2 vector restriction sites. Note that an additional lysine residue (underlined) was inserted between the His-tag and the first protein residue to provide a stopping point for the exoprotease. This additional lysine was unnecessary for the 9 kDa construct which already contained an initial lysine. The C145S mutant of the liprin- β 2 construct was created by QuickChange site-directed mutagenesis (Stratagene). All plasmid constructs were verified by DNA sequencing (Genewiz).

Protein Expression and Purification. Complete liprin- β 1 and liprin- β 2 coiled-coils were expressed in BL-21 (DE3) cells. Two separate 2 L cultures of LB were incubated at 37 °C until OD₆₀₀ \approx 0.6 and induced with 0.2 μ g/mL (final concentration) IPTG for 3–4 h at 37 °C. Cells were harvested by centrifugation, flash frozen in liquid nitrogen, and stored at –80 °C. Cells (7.22 g for liprin- β 1 wild-type, 8.23 g for liprin- β 2 wild-type, and 10.4 g liprin- β 2 C145S) were lysed by sonication in 40–50 mL of lysis buffer (50 mM HEPES-NaOH, 500 mM NaCl, 1 mM TCEP or 5 mM BME, 0.5–1 mg/mL lysozyme, 0.1 mM PMSF, and

DNase, pH 7.5). The lysate was then centrifuged for 10 min at 15 krpm in a Sorvall SS-34 rotor at 4 °C. The desired protein was found in the soluble and insoluble fractions, but the soluble protein did not bind efficiently to the Ni column under native conditions, perhaps due to proteolysis of the affinity tag. The protein could be purified from the insoluble fraction, however. The insoluble pellet was dissolved in 40 mL of denaturing binding/wash buffer (20 mM Tris-HCl, 300 mM NaCl, 6 M urea, 10 mM BME, and 10 mM imidazole pH 8.0) and incubated with turning at 4 °C for 1 h. Insoluble material was removed by centrifuging for 10 min at 15 krpm as above, and the supernatant was incubated with gentle stirring with 5 mL (bed volume) Ni-NTA resin for 1 h at 4 °C. The Ni beads were collected over a column and extensively washed with more than 50 mL of binding/wash buffer. The protein was eluted with 25 mL of denaturing elution buffer (20 mM Tris-HCl, 300 mM NaCl, 200 mM imidazole, 6 M urea, pH 8.0). After diluting the protein to 50 mL with 8 M urea, the protein was refolded by dialyzing into refolding buffer at 4 °C (20 mM HEPES-NaOH, 500 mM NaCl, 1 mM EDTA, 1 mM TCEP, pH 7.5). The C145S mutant was refolded into the same buffer containing 1 M ammonium sulfate. Solid ammonium sulfate was added to the refolded wild-type proteins (1 M final concentration) which were then purified over a 5 mL HiTrap HP phenylsepharose hydrophobic interaction column, washing first with 45 mL of buffer A (0.5 M ammonium sulfate, 20 mM Tris-HCl, pH 7.5) and eluting using a gradient to 100% buffer B (20 mM Tris-HCl, pH 7.5) over 45 mL. All proteins were further purified on a Superdex 200 column in 20 mM HEPES-NaOH, 500 mM NaCl, pH 7.5. Wild-type proteins were first reduced with excess DTT before gel filtration.

The 21, 19, and 9 kDa liprin β 1 and β 2 fragments were expressed and stored in frozen M15 cells (Qiagen) as described above for BL21(DE3). Cells were lysed by sonication in lysis buffer as above, and the protein was purified directly from the soluble fraction which was bound directly to Ni resin for 1 h at 4 °C. Unbound protein was washed from the resin using nondenaturing wash buffer (20 mM HEPES-NaOH, 500 mM NaCl, 5 mM imidazole, pH 7.5), and the bound protein was eluted with nondenaturing elution buffer (20 mM HEPES-NaOH, 500 mM NaCl, 200 mM imidazole, pH 7.5). Proteins were dialyzed overnight at 4 °C into phosphate buffer (20 mM PBS, 500 mM NaCl, pH 7.2). To remove the N-terminal His-tag, 15 mL aliquots of 21, 19, and 9 kDa fragments (1, 1, and 3 mg/mL) were treated with 50 μ L of DAPase (10 U/mL, Qiagen) preactivated with 50 μ L of 20 mM cystamine-HCl for 5 min at room temperature. The cleavage reaction was allowed to proceed at 37 °C for 3 h, and the DAPase and uncleaved protein were then removed using subtractive Ni affinity chromatography.

Equilibrium Sedimentation. Equilibrium sedimentation was performed at 20 °C using a Beckman Optima XL-A analytical centrifuge. From the amino acid sequence of liprin- β 2 coiled-coil C145S mutant the following were calculated: a monomeric molecular weight of 26 992.4, an extinction coefficient of 9970 M^{–1} cm^{–1}, and a partial specific volume of 0.7327 at 20 °C.^{25–27} Initial solutions of liprin- β 2 coiled-coil C145S were 0.18, 0.12, and 0.059 mg/mL, which were centrifuged at 10K, 12K, 14K, and 17K rpm. The equilibrium profiles were measured at 228 nm. A cell with a 1.2 mm path length was used for all experiments, and the buffer for all samples was 20 mM Tris-HCl, 500 mM NaCl, 100 mM ammonium sulfate, 1 mM EDTA, pH 7.5 (1.0269 calculated density at 20 °C). Samples were centrifuged at each

speed until the absorbance profile difference did not change significantly over ~ 4 h. Baseline measurements were obtained to perform experimental background correction by spinning at high speed (40K rpm) to completely pellet the protein. The data were globally fit using a nonlinear least-squares exponential fitting algorithm for a single nonassociating species with Beckman Origin software. Attempts to fit the data to a monomer–dimer equilibrium model yielded a large association constant suggesting the system effectively behaved as a dimer at the concentrations investigated. Sedimentation experiments using higher protein concentrations (0.5–2.0 mg/mL) were consistent with a mixture of low molecular species (i.e., mostly dimer and some tetramer) with a small amount of higher molecular weight oligomers that complicated fitting to a simple dimer–tetramer equilibrium model (GOF > 14). Size exclusion chromatography experiments in the buffer used for equilibrium sedimentation were compared to a lower salt buffer (20 mM Tris-HCl, 200 mM NaCl, 1 mM EDTA, pH 7.5) which yielded essentially the same elution volumes, indicating no significant effects of salt concentration on the oligomer size.

Chemical Cross-Linking. The liprin- $\beta 2$ coiled-coil mutant C145S was dialyzed into 20 mM phosphate, 500 mM NaCl, pH 7.2, to give a stock solution of 21 μ M. The BS3 cross-linking agent was dissolved in 20 mM phosphate, pH 7.2, to give a stock solution of 1 mM. Protein dilutions were performed in the same phosphate buffer containing 500 mM NaCl, and BS3 dilutions were made in straight phosphate buffer. For the BS3 titration and time series, cross-linking was initiated by mixing 5 μ L of protein solution with 5 μ L of cross-linker solution and allowed to proceed at room temperature. Reactions were quenched at different times with 10 μ L of 100 mM Tris-HCl, pH 8.0, and 20 μ L of SDS loading buffer was added to each sample before boiling 10 min and running on an SDS-PAGE gel. For the protein dilution series, cross-linking was initiated by mixing 18 μ L of protein solution with 2 μ L of BS3 stock solution and allowed to proceed at room temperature for 5 min. Reactions were quenched with 20 μ L of 100 mM Tris-HCl, pH 8.4. A total of 40 μ L of 2X SDS loading buffer was added to each reaction before boiling for 10 min and running on an SDS-PAGE gel.

Limited Proteolysis. Wild-type liprin- $\beta 1$ and liprin- $\beta 2$ coiled-coils were concentrated using Amicon Ultra 10k MWCO centrifugal concentrators to 2 mg/mL in 20 mM HEPES-NaOH, 500 mM NaCl, 2 mM DTT, pH 7.5. Lyophilized proteases α -chymotrypsin (type II from bovine pancreas), proteinase K (from *Tritirachium album*), and trypsin (TPCK treated from bovine pancreas) were obtained from Sigma and dissolved in water to give 1 mg/mL stock solutions that were used immediately. Dilutions of proteases were also made in water. A master reaction mix was first made containing 187.5 μ L of the 2.5X reaction buffer (500 mM Tris-HCl, 50 μ M CaCl_2 , pH 7.5), 120 μ L of protein stocks, and 67.5 μ L of fresh 10 mM DTT, from which 16 μ L aliquots were mixed with 4 μ L of protease solutions. Final concentrations of proteases in the reactions were 100 μ g/mL, 20 μ g/mL, 4 μ g/mL, 800 ng/mL, and 160 ng/mL. Reactions were initiated by addition of protease and quenched after exactly 20 min with 20 μ L 2X stop buffer (125 mM Tris-HCl, 5 mM PMSF, 4% (w/v) SDS, 5% (v/v) BME, 20% (v/v) glycerol, pH 6.75, and trace bromophenol blue). All reactions were boiled immediately for 10 min before running on an SDS-PAGE gel.

Large-scale reactions with proteinase K were scaled accordingly, quenched with PMSF, and purified by size-exclusion chromatography using a Superdex 200 10/300 GL column

with 20 mM HEPES-NaOH, 500 mM NaCl, pH 7.5. Aliquots from fractions containing protein fragments were desalted and analyzed by mass spectrometry (Applied Biosystems/MDS Sciex QSTAR XL mass spectrometer). All fractions were also run on an SDS-PAGE gel with 0.1 mM thioglycolate in the upper running buffer and blotted for 2 h at 120 mA onto a Sequi-Blot PVDF membrane (Bio-Rad) using 10 mM CAPS, pH 11, 10% methanol as the transfer buffer. The PVDF membrane was stained with 0.025% Coomassie blue in 40% methanol and destained with 50% methanol, i.e., in the absence of acetic acid to avoid N-terminal acetylation. Bands corresponding to each protein fragment were cut from the membrane and analyzed by three cycles of Edman degradation. The software PAWS (i.e., Protein Analysis Work Sheet, Genomic Solutions Inc.) was used to identify sequences within each coiled-coil domain consistent with the observed masses and N-termini.

Crystal Structure Determination. The recombinant 9 kDa liprin- $\beta 2$ coiled-coil fragment was dialyzed into 20 mM Tris, 5% glycerol, pH 8.0, and concentrated to 14.9 mg/mL using a 5000 MWCO Amicon Ultra centrifugal concentrator. Crystals were grown by the hanging drop method using 190 mM ammonium iodide, 23% PEG 3350 as the well solution and mixing 1 μ L of the protein sample with 0.5 μ L of the well solution. Crystals grew overnight at room temperature ($\sim 21^\circ\text{C}$) and were cryoprotected using the well solution supplemented with 30% glycerol before data collection. Three data sets, collected from different sections of a single large crystal at Advanced Photon Source (APS, Argonne National Laboratories), were processed and merged with Denzo/Scalepack.²⁸ The crystals belonged to space group $C222_1$ with cell dimensions of $a = 29.965 \text{ \AA}$, $b = 162.653 \text{ \AA}$, $c = 34.387 \text{ \AA}$, and $\alpha = \beta = \gamma = 90^\circ$ with a single molecule in the asymmetric unit. The molecular replacement search model was generated by submitting the complete sequence of the crystalized fragment from K185 through A265 of liprin- $\beta 2$ to the Phyre²⁹ (Protein Homology/analogueY Recognition Engine) Web server which generated a model corresponding to residues L212 through T264 of liprin- $\beta 2$ based on the coiled-coil structure of tropomyosin. An initial molecular replacement solution was found by Phaser³⁰ using this homology model. The amino acid register was set automatically using ARP/wARP³¹ building into a side-chain omit map. The initial solution from Phaser was refined by iterative cycles of manual building with COOT³² and REFMACS.³³ During refinement the anisotropy was corrected by performing ellipsoidal truncation and anisotropic scaling using the UCLA MBI Diffraction Anisotropy Server (<http://services.mbi.ucla.edu/anisotropy/>).³⁴ TLS refinement was used toward the end of refinement using a single TLS group for the entire protein molecule in the asymmetric unit. Residues with clear density for more than one side-chain conformation were refined with partial occupancy in dual conformations.

Structure Analysis. The packing of the liprin- $\beta 2$ coiled-coil dimer was analyzed using the SOCKET³⁵ Web server, and the coiled-coil parameters were analyzed using TWISTER.³⁶ In order to be analyzed by both programs, a dimeric coiled-coil structure was first generated by applying a crystallographic 2-fold symmetry operation to create a coordinate file for the dimer model. As alternate conformations are not allowed by these programs, a single conformation was assigned to each residue. Graphics were prepared in PyMOL (Schrödinger, LLC).




Figure 2. Liprin- β 2 coiled-coil forms an extended dimer. (A) The liprin- β 2 coiled-coil domain concentration distributions after sedimentation to equilibrium at various speeds and protein concentrations. The three different concentrations (light blue, 0.18 mg/mL; orange, 0.12 mg/mL; gray, 0.059 mg/mL) were equilibrated at four different speeds (10K, 12K, 14K, and 17K rpm, bottom to top) and globally fit to a single exponential (shown as lines) with a variable molecular weight. The data were best fit with the molecular weight equal to the dimer (52 598 Da measured from fit; 53 984.8 Da calculated from sequence). (B) An SDS-PAGE gel showing the results of a liprin- β 2 coiled-coil domain chemical cross-linking time-series experiment performed using a constant concentration of protein (5 μ M) and BS3 cross-linker (50 μ M). (C) An SDS-PAGE gel showing the results of a chemical cross-linking of the liprin- β 2 coiled-coil domain at different protein concentrations, at a constant BS3 cross-linker concentration (100 μ M) for a total time of 5 min. The ladder used for the gels in (B) and (C) contains standard proteins of 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 120, 160, and 220 kDa (BenchMark Protein Ladder, Invitrogen). For reference the 20 and 50 kDa bands of the ladder are indicated on each gel. (D) Superdex S-200 gel filtration chromatography of the liprin- β 2 coiled-coil domain C145S. The elution volume of a set of globular protein standards is also indicated. The protein standards from largest to smallest are ferritin, aldolase, ovalbumin, ribonuclease A, and aprotinin (GE Healthcare). The void volume of the column was determined using blue dextran 2000 to be \sim 8.3 mL. The liprin- β 2 coiled-coil (concentration <0.5 mg/mL) elutes at 12.6 mL (\sim 525 kDa) or \sim 20-fold higher than its monomeric molecular weight.

RESULTS

Design of β 1 and β 2 Coiled-Coil Expression Constructs. To define the extent of the coiled-coil domains for both liprin β 1 and β 2 (1011 and 876 amino acids, respectively), we employed the COILS¹⁶ coiled-coil prediction server. As shown in Figure 1B, residues 97–342 of liprin β 1 and 100–315 of liprin β 2 were strongly predicted to be coiled-coils. Both regions were expressed in *E. coli* and purified.

Both β 1 and β 2 coiled-coil constructs were prone to intermolecular oxidative disulfide bond formation as evidenced by peak shifts in gel filtration experiments in the presence and absence of a reducing agent (data not shown). To eliminate complications due to disulfide bond formation during further biochemical analysis of the liprin β 2 construct, a C145S mutant was made by site-directed mutagenesis to remove its single Cys. The C145S mutant eluted from the gel filtration column at the same volume as the reduced wild-type construct (data not shown), indicating that this mutant was the same size and shape as the reduced wild-type protein, but not prone to oxidation.

Liprin β 2 Coiled-Coil Forms a Dimer. As the coiled-coil domains were expected to have an unusually extended architecture,

we employed equilibrium sedimentation to determine the molecular weight of the oligomer because the method measures mass independent of shape (Figure 2A). Sedimentation experiments were performed at three concentrations (6.8, 4.5, and 2.2 μ M) and four different speeds (10K, 12K, 14K, and 17K rpm) using the liprin β 2 C145S coiled-coil. As shown in Figure 2A, the data were globally fit well by a single exponential for a simple dimeric species (GOF = 1.17).

Chemical cross-linking using BS3 was performed to further establish the oligomer size of the β 2 coiled-coil (Figure 2B,C). BS3 contains two amine-reactive succinimidyl ester moieties \sim 11 Å apart that should covalently couple exposed primary amines (e.g., from lysine residues) in close proximity. First, the C145S β 2 coiled-coil concentration was held constant at 5 μ M in the presence of a 10-fold stoichiometric excess of BS3 (50 μ M) (Figure 2B). The cross-linking reaction was quenched at different time points to monitor the extent of cross-linking as a function of time. The reaction is essentially complete within 1 min, and only monomeric and dimeric species are observed despite the large excess of cross-linker. A set of bands centered at the dimer molecular weight, rather than a single band, is




Figure 3. Limited proteolysis of liprin- β 1 and - β 2 coiled-coils. (A) An SDS-PAGE gel showing the results of limited proteolysis of the liprin- β 1 coiled-coil (30.8 kDa) by several proteases at a series of concentrations (μ g/mL). (B) An SDS-PAGE gel showing the results of limited proteolysis of the liprin- β 2 coiled-coil (27.0 kDa) by several proteases at a series of concentrations (μ g/mL). The 21 kDa (*), 19 kDa (x), and 9 kDa (o) bands were identified by mass spectrometry and Edman degradation. The ladder is the same as in Figure 2. (C) ClustalW⁴¹ sequence alignment of human liprin- β 1 and - β 2 coiled-coils. The regions cloned in the full-length coiled-coil constructs are highlighted in yellow. The 21 kDa liprin- β 1 fragment is indicated by a solid line above the alignment, and the 19 and 9 kDa fragments of liprin- β 2 are indicated by solid lines below the alignment. The extent of sequence conservation is indicated below the alignment.

observed, presumably due to different cross-linking points that affect migration through the gel. The cross-linking experiment was then performed for 5 min at a constant BS3 concentration (100 μ M) using a dilution series of the C14SS β 2 coiled-coil (Figure 2C). This experiment shows that the relative ratio of dimer to monomer remains constant regardless of the protein concentration. These experiments suggest that the β 2 coiled-coil is primarily dimeric.

The size of the liprin- β 2 coiled-coil domain was also estimated by size exclusion chromatography. As shown in Figure 2D, it migrates at a position corresponding to a 525 kDa globular protein or the size of a globular 20-mer. The large hydrodynamic radius suggests that the protein is highly extended as might be expected for a coiled-coil structure.

Identification of Stable, Protease-Resistant β 1 and β 2 Coiled-Coil Subdomains. Despite extensive efforts, we were unable to obtain high-quality crystals of the full length coiled-coil domains. We therefore searched for smaller, stable subdomains. To identify smaller and presumably more rigid subdomains, the wild-type liprin β 1 and β 2 coiled-coil constructs were subjected to limited proteolysis using α -chymotrypsin, proteinase K, and trypsin under reducing conditions (Figure 3A,B). Limited proteolysis of the liprin β 1 coiled-coil yielded a single, ~21 kDa fragment in the presence of all three enzymes as assessed by SDS-PAGE analysis. Limited proteolysis of the liprin β 2 coiled-coil

yielded two fragments, approximately 9 and 19 kDa, as assessed by SDS-PAGE analysis. The consistent size of these fragments regardless of the different sequence specificities of these proteases suggests that these fragments are likely to represent common, stable subdomains within the entire coiled-coil domains.

To identify the sequences of the protease stable subdomains, limited proteolysis experiments of both liprin β 1 and β 2 coiled-coils using proteinase K were scaled up, and the resulting components were purified by size-exclusion chromatography. Fractions containing the purified fragments were analyzed by ESI mass spectrometry and N-terminal sequencing by Edman degradation (Figure 3C). A single consistent sequence was identified for the 21 kDa liprin β 1 coiled-coil subdomain (calculated mass 21 385.5 kDa; measured mass 21 389.0 kDa; N-terminal sequence: END) from E110 through K290. A single consistent sequence was also identified for the 19 kDa liprin β 2 coiled-coil subdomain (calculated mass 19 120.6 kDa; measured mass 19 121.0 kDa; N-terminal sequence: ASN) from A102 through A265. The smaller 9 kDa subdomain of the liprin β 2 coiled-coil appeared to be heterogeneous by both mass spectrometry and Edman degradation analysis. Nevertheless, a likely component of this mixture was identified from the largest species (calculated mass 9 723.1 kDa; measured mass 9 722.2 kDa; N-terminal sequence: KLK) that encompassed amino acids K185 through

Table 1. X-ray Diffraction Data and Refinement Statistics

data collection	liprin- β 2 (K185 to A265) (PDB code 3QH9)
wavelength (Å)	1.4938
space group	C222 ₁
cell dimensions	
<i>a</i> , <i>b</i> , <i>c</i> (Å)	30.0, 162.7, 34.4
α , β , γ (deg)	90, 90, 90
Matthews coefficient (Å ³ /Da)	2.15
% solvent	42.95
resolution (Å) (highest shell)	80.00–2.00 (2.07–2.00)
<i>R</i> _{sym}	0.099 (0.348)
<i>I</i> / <i>I</i> σ	20.7 (5.0)
% completeness	97.0 (84.1)
redundancy	10.2 (7.2)
refinement	
no. reflections	5774
<i>R</i> _{work}	0.232
<i>R</i> _{free} ^a	0.298
no. atoms	638
<i>B</i> -factor (Å ²)	
all atoms	37.0
protein	38.0
water	19.6
rmsd bond lengths (Å)	0.0234
rmsd bond angle (deg)	2.0
Ramachandran statistics (%)	
most favored	100
additionally allowed	0
generously allowed	0
disallowed	0

^aCalculated from 5% random sampling of data excluded from refinement.

A265. The common C-terminus of this 9 kDa and the 19 kDa fragment further suggests that K185-A265 is the dominant component of this mixture.

Three constructs encompassing these protease-resistant cores of the liprin β 1 and β 2 coiled-coils were cloned into a pQE2 bacterial expression vector containing a cleavable, N-terminal His-tag for purification purposes. All constructs expressed into the soluble fraction and were purified under native conditions. The N-terminal His-tag was removed using an exopeptidase and the resulting tag-free constructs were purified by a subtractive Ni affinity column. The 21 kDa β 1 or 19 kDa β 2 fragments did not yield crystals, but we were able to obtain crystals of the 9 kDa fragment of liprin- β 2.

X-ray Crystallography of the Central β 2 Coiled-Coil Core. The 9 kDa β 2 coiled-coil crystals belonged to space group C222₁ with a unit cell of *a* = 29.965 Å, *b* = 162.653 Å, *c* = 34.387 Å, and $\alpha = \beta = \gamma = 90^\circ$ (Table 1). The crystals diffracted to ~ 2.0 Å, but the diffraction was strongly anisotropic with significantly reduced diffraction intensity along the *a*^{*} reciprocal axis (Supporting Information Figure 1). The structure was solved by molecular replacement using a homology model as the search model. Remarkably, the homology model generated by the Phyre server²⁹ was based on a tropomyosin structure that bore only

16% identity to the liprin β 2 construct. The search model corresponded to residues L212 through T264, but electron density for residues preceding L212 was clearly visible in the initial molecular replacement solution, lending confidence that the correct solution had been obtained. To further validate this initial solution, all residues were converted to alanine to create an omit map that clearly showed positive side-chain electron density throughout the entire structure. The amino acid register was determined automatically by allowing ARP/wARP³¹ to build into this omit map. It became clear during refinement that the strongly anisotropic data needed to be corrected by performing ellipsoidal truncation and scaling.³⁴ Before this anisotropy correction the refinement stalled (*R*/*R*_{free} \approx 0.35–0.40), and the electron density was less well-defined than would be expected from a 2.0 Å electron density map (Supporting Information Figure 2). Immediately following the anisotropy correction, the electron density became much better defined and the *R*/*R*_{free} dropped accordingly (final *R*/*R*_{free} of 0.232/0.298). Attempts to refine the structure in a lower symmetry space group did not improve the *R*-factor or the quality of the electron density map.

Analysis of the β 2 Coiled-Coil Structure. Although the construct used for crystallization included residues K185 through A265 of liprin- β 2, the electron density was only interpretable for residues Q198 through R263. A single protein molecule was found in the asymmetric unit that formed a single, continuous α -helix which interacted with a symmetry related copy to form a parallel, homodimeric coiled-coil ~ 100 Å long (Figure 4A,B). Throughout the crystal lattice the dimeric coiled-coils pack with their long axes parallel to the long *b*-axis of the unit cell. Analysis by SOCKET³⁵ reveals typical knobs-into-holes packing throughout most of the dimer with typical Leu and Val residues occupying the majority of the “a” and “d” residues of the heptad repeat involved in packing between the two molecules of the coiled-coil (Figure 4C). A notable exception are residues R226 and Y229 which occupy successive “a” and “d” residues near the center of the coiled-coil but do not exhibit classical knobs-into-holes type packing. Analysis by TWISTER³⁶ reveals a sharp increase in the local pitch of the coiled-coil surrounding these residues, indicating that the coiled-coil unwinds slightly at this location, but the register of the heptad pattern does not change (Figure 4D,E). Overall, the two-stranded coiled-coil forms a typical left-handed supercoil with an average coiled-coil pitch of 161 ± 38 Å.

DISCUSSION

Synaptic vesicles are organized by a complex protein assembly that ensures the rapid release of neurotransmitters in response to incoming action potentials.^{1,37} The liprin protein family plays an important part in regulating and maintaining this synaptic architecture,³⁸ but the molecular basis for these functions is unknown. Previous immunoprecipitation experiments have shown that liprin- β coiled-coils interact with themselves,² but the basic stoichiometry and orientation of these interactions have not been defined. The equilibrium sedimentation and cross-linking experiments presented here demonstrate that the coiled-coil of liprin- β 2 predominantly forms a dimer. The sequence similarity and similar size-exclusion retention times suggest that liprin- β 1 and liprin- β 2 are approximately the same size and shape, so it seems probable that liprin- β 1 is also dimeric. The orientation of the liprin- β 2 coiled-coil oligomer was established from the crystal structure of the central, protease-resistant core.




Figure 5. Possible organization of liprin scaffolds. (A) A parallel homodimer model of the complete liprin- β 2. The complete coiled-coil (~ 300 Å) was modeled using a combination of BEAMMOTIF and MAKECCSC.⁴² The model is merely intended to convey the orientation and relative sizes of the coiled-coil domains. (B) A speculative scaffold organization is shown consisting of liprin- α 1 and liprin- β 2 molecules. For simplicity, the entire liprin- α 1 coiled-coil (~ 900 Å) was also assumed to form a two-stranded parallel structure which was constructed using MAKECCSC with no attempt to adjust for stutters, stammers, or gaps. The three tandem SAM domains of liprin- α s have been shown in other studies to interact with liprin- β s,² which we show here in a “closed dimer” model of α and β dimers. (C) An alternative “open scaffold” model is shown in which multiple α - and β -liprins stack indefinitely. In both “closed dimer” and “open scaffold” models the coiled-coil domains are predicted to radiate from the central SAM domain interactions.

more than one dimer as shown in Figure 5C. Another more complex possibility is that α and β SAMs form a copolymer which has been observed before with the SAM domains from Scm and polyhomeotic.⁴⁰ Individual liprin SAM domains have also been computationally predicted to form polymeric structures, which could be an important aspect of their scaffolding architecture.²⁴

Obviously, concepts of the overall scaffolding structure of liprins remain highly speculative since little is known about the interactions between the SAM domains and the structure of the liprin- α coiled-coils. Regardless, in all of the scaffolding models, the coiled-coil domains are predicted to radiate out from the central axis of SAM domains to interact with other synaptic proteins. At present, the distinct functional roles of the two different liprin- β proteins and four different liprin- α proteins in humans are unknown, but it is known that they can associate. It seems plausible that the coiled-coil domains may form distinct hetero-oligomers that may elicit different functions. Much work remains before we can visualize how liprins organize the complex set of partner proteins that contribute to the construction of the cytomatrix of the active zone.

■ ASSOCIATED CONTENT

S Supporting Information. Two figures showing an analysis of the anisotropic diffraction and electron density maps. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Accession Codes

Coordinates have been deposited in the RCSB Protein Data Bank: PDB code 3QH9 (liprin- β 2 residues K185 to A265).

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: bowie@mbi.ucla.edu. Phone: (310)206-4747. Fax: (310)206-4749.

Funding Sources

This work was supported by NIH Grant R01GM093393 to J.U.B. and a NIH Ruth L. Kirschstein Postdoctoral Fellowship F32GM084615 to R.L.S.

■ ACKNOWLEDGMENT

We thank Julian Whitelegge and Sara Bassilian (UCLA Pasarow Mass Spectrometry Laboratory) for assistance with mass spectrometry experiments and Kathy Schegg (UNR INBRE Nevada Proteomics Center; NIH Grant Number P20 RR-016464) for Edman sequencing.

■ ABBREVIATIONS

SAM, sterile alpha motif; PCR, polymerase chain reaction; BME, β -mercaptoethanol; HEPES, *N*-(2-hydroxyethyl)piperazine-*N'*-2-ethanesulfonic acid; TCEP, (tris(2-carboxyethyl)phosphine); PMSF, phenylmethylsulfonyl fluoride; LB, Luria broth; IPTG, isopropyl- β -D-1-thiogalactopyranoside; EDTA, ethylenediaminetetraacetic acid; DTT, DL-dithiothreitol; PBS, phosphate buffered saline; BS3, bis(sulfosuccinimidyl) suberate; PVDF, poly(vinylidene fluoride); SDS, sodium dodecyl sulfate; PAGE, polyacrylamide gel electrophoresis; PEG, poly(ethylene glycol); ESI, electrospray ionization; ML, midloop; EH, end-helix.

■ REFERENCES

- (1) Schoch, S., and Gundelfinger, E. D. (2006) Molecular organization of the presynaptic active zone. *Cell Tissue Res.* 326, 379–391.
- (2) Serra-Pagès, C., Medley, Q. G., Tang, M., Hart, A., and Streuli, M. (1998) Liprins, a family of LAR transmembrane protein-tyrosine phosphatase-interacting proteins. *J. Biol. Chem.* 273, 15611–15620.
- (3) Olsen, O., Moore, K. A., Fukata, M., Kazuta, T., Trinidad, J. C., Kauer, F. W., Streuli, M., Misawa, H., Burlingame, A. L., Nicoll, R. A., and Bredt, D. S. (2005) Neurotransmitter release regulated by a MALSLiprin- α presynaptic complex. *J. Cell Biol.* 170, 1127–1134.
- (4) Ko, J., Na, M., Kim, S., Lee, J., and Kim, E. (2003) Interaction of the ERC family of RIM-binding proteins with the liprin- α family of multidomain proteins. *J. Biol. Chem.* 278, 42377–42385.
- (5) Ko, J., Kim, S., Valtschanoff, J. G., Shin, H., Lee, J., Sheng, M., Premont, R. T., Weinberg, R. J., and Kim, E. (2003) Interaction between liprin- α and GIT1 is required for AMPA receptor targeting. *J. Neurosci.* 23, 1667–1677.
- (6) Shin, H., Wyszynski, M., Huh, K., Valtschanoff, J. G., Lee, J., Ko, J., Streuli, M., Weinberg, R. J., Sheng, M., and Kim, E. (2003) Association of the kinesin motor KIF1A with the multimodular protein liprin- α . *J. Biol. Chem.* 278, 11393–11401.
- (7) Schoch, S., Castillo, P. E., Jo, T., Mukherjee, K., Geppert, M., Wang, Y., Schmitz, F., Malenka, R. C., and Südhof, T. C. (2002) RIM1 α forms a protein scaffold for regulating neurotransmitter release at the active zone. *Nature* 415, 321–326.
- (8) Patel, M. R., and Shen, K. (2009) RSY-1 is a local inhibitor of presynaptic assembly in *C. elegans*. *Science* 323, 1500–1503.
- (9) Zhen, M., and Jin, Y. (1999) The liprin protein SYD-2 regulates the differentiation of presynaptic termini in *C. elegans*. *Nature* 401, 371–375.
- (10) Hofmeyer, K., Maurel-Zaffran, C., Sink, H., and Treisman, J. E. (2006) Liprin- α has LAR-independent functions in R7 photoreceptor axon targeting. *Proc. Natl. Acad. Sci. U.S.A.* 103, 11595–11600.

- (11) Choe, K., Prakash, S., Bright, A., and Clandinin, T. R. (2006) Liprin-alpha is required for photoreceptor target selection in *Drosophila*. *Proc. Natl. Acad. Sci. U.S.A.* 103, 11601–11606.
- (12) Ackley, B. D., Harrington, R. J., Hudson, M. L., Williams, L., Kenyon, C. J., Chisholm, A. D., and Jin, Y. (2005) The two isoforms of the *Caenorhabditis elegans* leukocyte-common antigen related receptor tyrosine phosphatase PTP-3 function independently in axon guidance and synapse formation. *J. Neurosci.* 25, 7517–7528.
- (13) Dunah, A. W., Hueske, E., Wyszynski, M., Hoogenraad, C. C., Jaworski, J., Pak, D. T., Simonetta, A., Liu, G., and Sheng, M. (2005) LAR receptor protein tyrosine phosphatases in the development and maintenance of excitatory synapses. *Nature Neurosci.* 8, 458–467.
- (14) Nachat, R., Cipolat, S., Sevilla, L. M., Chhatrivala, M., Groot, K. R., and Watt, F. M. (2009) KazrinE is a desmosome-associated liprin that colocalises with acetylated microtubules. *J. Cell. Sci.* 122, 4035–4041.
- (15) Astigarraga, S., Hofmeyer, K., Farajian, R., and Treisman, J. E. (2010) Three *Drosophila* liprins interact to control synapse formation. *J. Neurosci.* 30, 15358–15368.
- (16) Lupas, A., Van Dyke, M., and Stock, J. (1991) Predicting coiled coils from protein sequences. *Science* 252, 1162–1164.
- (17) Wolf, E., Kim, P. S., and Berger, B. (1997) MultiCoil: a program for predicting two- and three-stranded coiled coils. *Protein Sci.* 6, 1179–1189.
- (18) McDonnell, A. V., Jiang, T., Keating, A. E., and Berger, B. (2006) Paircoil2: improved prediction of coiled coils from sequence. *Bioinformatics* 22, 356–358.
- (19) Mason, J. M., and Arndt, K. M. (2004) Coiled coil domains: stability, specificity, and biological implications. *ChemBioChem* 5, 170–176.
- (20) Grigoryan, G., and Keating, A. E. (2008) Structural specificity in coiled-coil interactions. *Curr. Opin. Struct. Biol.* 18, 477–483.
- (21) Yadav, M. K., Leman, L. J., Price, D. J., Brooks, C. L., Stout, C. D., and Ghadiri, M. R. (2006) Coiled coils at the edge of configurational heterogeneity. Structural analyses of parallel and antiparallel homotetrameric coiled coils reveal configurational sensitivity to a single solvent-exposed amino acid substitution. *Biochemistry* 45, 4463–4473.
- (22) Apgar, J. R., Gutwin, K. N., and Keating, A. E. (2008) Predicting helix orientation for coiled-coil dimers. *Proteins* 72, 1048–1065.
- (23) Qiao, F., and Bowie, J. U. (2005) The many faces of SAM. *Sci. STKE* 2005, re7.
- (24) Meruelo, A. D., and Bowie, J. U. (2009) Identifying polymer-forming SAM domains. *Proteins* 74, 1–5.
- (25) Pace, C. N., Vajdos, F., Fee, L., Grimsley, G., and Gray, T. (1995) How to measure and predict the molar absorption coefficient of a protein. *Protein Sci.* 4, 2411–2423.
- (26) Cohn, E. J., and Edsall, J. T. (1943) *Proteins, Amino Acids and Peptides as Ions and Dipolar Ions*, pp 370–381, Reinhold, New York.
- (27) Laue, T. M., Shah, B. D., Ridgeway, S. L., and Pelletier, S. L. (1992) *Analytical Ultracentrifugation in Biochemistry and Polymer Science*, pp 90–125, The Royal Society of Chemistry, Cambridge.
- (28) Minor, W., and Otwinowski, Z. (1997) Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol.* 276, 307–326.
- (29) Kelley, L. A., and Sternberg, M. J. E. (2009) Protein structure prediction on the Web: a case study using the Phyre server. *Nature Protoc.* 4, 363–371.
- (30) McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C., and Read, R. J. (2007) Phaser crystallographic software. *J. Appl. Crystallogr.* 40, 658–674.
- (31) Langer, G., Cohen, S. X., Lamzin, V. S., and Perrakis, A. (2008) Automated macromolecular model building for X-ray crystallography using ARP/wARP version 7. *Nature Protoc.* 3, 1171–1179.
- (32) Emsley, P., Lohkamp, B., Scott, W. G., and Cowtan, K. (2010) Features and development of Coot. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* 66, 486–501.
- (33) Vagin, A. A., Steiner, R. A., Lebedev, A. A., Potterton, L., McNicholas, S., Long, F., and Murshudov, G. N. (2004) REFMAC5 dictionary: organization of prior chemical knowledge and guidelines for its use. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* 60, 2184–2195.
- (34) Strong, M., Sawaya, M. R., Wang, S., Phillips, M., Cascio, D., and Eisenberg, D. (2006) Crystal structure of a PE/PPE protein complex from *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci. U.S.A.* 103, 8060–8065.
- (35) Walshaw, J., and Woolfson, D. N. (2001) Socket: a program for identifying and analysing coiled-coil motifs within protein structures. *J. Mol. Biol.* 307, 1427–1450.
- (36) Strelkov, S. V., and Burkhard, P. (2002) Analysis of alpha-helical coiled coils with the program TWISTER reveals a structural mechanism for stutter compensation. *J. Struct. Biol.* 137, 54–64.
- (37) Sudhof, T. C. (2004) The synaptic vesicle cycle. *Annu. Rev. Neurosci.* 27, 509–547.
- (38) Spangler, S. A., and Hoogenraad, C. C. (2007) Liprin-alpha proteins: scaffold molecules for synapse maturation. *Biochem. Soc. Trans.* 35, 1278–1282.
- (39) Kurabi, A., Brener, S., Mobli, M., Kwan, J. J., and Donaldson, L. W. (2009) A nuclear localization signal at the SAM-SAM domain interface of AIDA-1 suggests a requirement for domain uncoupling prior to nuclear import. *J. Mol. Biol.* 392, 1168–1177.
- (40) Kim, C. A., Sawaya, M. R., Cascio, D., Kim, W., and Bowie, J. U. (2005) Structural organization of a Sex-comb-on-midleg/polyhomeotic copolymer. *J. Biol. Chem.* 280, 27769–27775.
- (41) Chenna, R., Sugawara, H., Koike, T., Lopez, R., Gibson, T. J., Higgins, D. G., and Thompson, J. D. (2003) Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res.* 31, 3497–3500.
- (42) Offer, G., Hicks, M. R., and Woolfson, D. N. (2002) Generalized Crick equations for modeling noncanonical coiled coils. *J. Struct. Biol.* 137, 41–53.