


Structure of Even/Odd Trinucleotide Repeat Sequences Modulates Persistence of Non-B Conformations and Conversion to Duplex

Amalia Ávila Figueroa,[†] Douglas Cattie,[†] and Sarah Delaney*

Department of Chemistry, Brown University, Providence, Rhode Island 02912, United States

 Supporting Information

ABSTRACT: Expansion of trinucleotide repeats (TNR) has been implicated in the emergence of neurodegenerative diseases. Formation of non-B conformations such as hairpins by these repeat sequences during DNA replication and/or repair has been proposed as a contributing factor to expansion. In this work we employed a combination of fluorescence, chemical probing, optical melting, and gel shift assays to characterize the structure of a series of (CTG)_n sequences and the kinetic parameters describing their interaction with a complementary sequence. Our structure-based experiments using chemical probing reveal that sequences containing an even or odd number of CTG repeats adopt stem-loop hairpins that differ from one another by the absence or presence of a stem overhang. Furthermore, we find that this structural difference dictates the rate at which the TNR hairpins convert to duplex with a complementary CAG sequence. Indeed, the rate constant describing conversion to (CAG)₁₀/(CTG)_n duplex is slower for sequences containing an even number of CTG repeats than for sequences containing an odd number of repeats. Thus, when both the CAG and CTG hairpins have an even number of the repeats, they display a longer lifetime relative to when the CTG hairpin has an odd number of repeats. The difference in lifetimes observed for these TNR hairpins has implications toward their persistence during DNA replication or repair events and could influence their predisposition toward expansion. Taken together, these results contribute to our understanding of trinucleotide repeats and the factors that regulate persistence of hairpins in these repetitive sequences and conversion to canonical duplex.



The expansion of trinucleotide repeat (TNR) sequences has been implicated in an array of neurodegenerative disorders.^{1–4} The length of these repeat polymorphisms predicts whether an individual will be healthy or will be afflicted by disease. The number of TNR repeats falls into one of three categories: (1) a small number of repeats in which case the individual will be healthy and the repeat tract will be stably transmitted to offspring; (2) an intermediate or premutation number of repeats in which case individuals typically do not present symptoms of the disease, but are capable of transmitting an expanded TNR tract to their offspring; or (3) a large number of repeats in which case the individual will develop the disorder. Notably, from the disease state the number of repeats can expand further in subsequent generations. As a result, a distinct trait of TNR disorders is the phenomenon of anticipation, in which the offspring of individuals that are afflicted by the disorder will have TNR tracts that are expanded further.⁵ As an example, in Huntington's disease (HD) which is caused by the expansion of a (CAG)_n/(CTG)_n TNR sequence, healthy individuals have 5–35 repeats, individuals with 36–39 repeats are in the premutation range, and individuals with 40 or more repeats will develop the disorder.⁶

While the molecular basis for TNR disorders is known to be expansion of the repetitive DNA sequence, the mechanism of the expansion is not fully understood. However, a common element of TNR sequences that are prone to expansion is their ability to

adopt non-B DNA conformations such as triplexes, quadruplexes, and hairpins.^{7–15} For example, (CAG)_n and (CTG)_n sequences have been shown to form hairpins both in vitro¹⁶ and in vivo.¹⁷ Additionally, TNR sequences have been observed to form slipped-strand structures where the repetitive nature of the sequence allows the complementary strands to slip out of register and fold into hairpins.^{18–20} The propensity to form such non-B DNA structures has been shown to increase with length of the TNR sequence.²¹ Indeed, formation and persistence of non-B conformations have been proposed as contributing factors in the prevailing models for expansion, which involve polymerase nonprocessivity caused by slippage of the nascent strand during DNA synthesis or aberrant repair events following DNA damage.^{1,3,4,22,23}

Given the proposed importance of non-B conformations and repeat length in expansion, in this work we define the structural properties of TNR sequences of varying length and delineate the role of structure in modulating the persistence of non-B TNR hairpins. Specifically, using a (CAG)₁₀ sequence, we examined the role of the length of the complementary (CTG)_n sequence (when *n* = 6–14) in influencing the structure and lifetime of the

Received: March 17, 2011

Revised: April 26, 2011

Published: April 28, 2011

TNR DNA. Surprisingly, using sequences that correspond to the healthy allele in HD, we find that incremental changes of one repeat unit affect the structure and persistence of the non-B conformations. While all the sequences adopt hairpin structures with a loop size of four nucleobases, hairpins with an even number of repeats have blunt-ended stems whereas those with an odd number of repeats have one repeat overhanging the stem. Importantly, we also find that the presence of an overhanging repeat decreases the lifetime of the non-B conformations by facilitating conversion to (CAG)₁₀/(CTG)_n duplex. Understanding structure-based reactivity and lifetime of even- and odd-repeat hairpins can provide insight into how TNR sequences expand and, thus, shed light on the molecular mechanism of several diseases.

EXPERIMENTAL PROCEDURES

Synthesis and Purification of Oligonucleotides. Oligonucleotides were synthesized using standard phosphoramidite chemistry.²⁴ All phosphoramidites, including 5'-fluorescein (FL) and 3'-(*E*)-N-(3-(dihydroxymethoxy)propyl)-4-((4-(dimethylamino)phenyl)diazetyl)benzamide (DCL), were purchased from Glen Research. Oligonucleotides were purified using a Dynamax Microsorb C18 reverse phase HPLC column (10 × 250 mm) as described previously.²⁵ Quantification was performed at 90 °C using ϵ_{260} values estimated for single-stranded DNA.²⁶

Fluorescence Measurements. The (CTG)_n oligonucleotides were prepared in 20 mM sodium phosphate, 10 mM NaCl, pH 7.0, at a concentration of 3 μ M. In order to obtain the thermodynamically favored DNA conformations, oligonucleotides were incubated for 5 min at 90 °C and cooled to room temperature. The FL-(CAG)₁₀-DCL oligonucleotide was prepared in the same fashion with all sample manipulations performed in the dark. The FL-(CAG)₁₀-DCL and (CTG)_n oligonucleotides were chilled on ice for 1 h prior to mixing in a 1:1 ratio for a final sample (25 μ L) that contained 1.5 μ M FL-(CAG)₁₀-DCL and 1.5 μ M (CTG)_n in 20 mM sodium phosphate, 10 mM NaCl, pH 7.0. After mixing, the samples were subjected to the following temperature cycle on a MX4000 real-time PCR instrument (Stratagene): 10 °C for 10 min, 10 to 90 °C at 1 °C/min, 5 min at 90 °C, and 90 to 10 °C at 1 °C/min. Over the course of this temperature cycle the samples were excited at 492 nm, and the emission was monitored at 520 nm. The melting temperature (T_m) of FL-(CAG)₁₀-DCL was taken as the maximum in the first derivative of the fluorescence profile. The experiments were performed in triplicate.

Characterization of Oligonucleotides Using the Chemical Probe DEPC. Oligonucleotides were 5'-[³²P] end-labeled using T4 polynucleotide kinase according to the manufacturer's protocol. Radiolabeled oligonucleotide was supplemented with the unlabeled version to obtain a final DNA concentration of 5 μ M in 20 mM sodium phosphate, 10 mM NaCl, pH 7.0. The oligonucleotides were incubated for 5 min at 90 °C and slowly cooled to room temperature. The DNA samples were incubated with 4% diethyl pyrocarbonate (DEPC, v/v) for 30 min at 37 °C with vortexing at the start of the reaction and halfway through incubation. Following incubation with DEPC, the samples were dried under vacuum, treated with 10% piperidine (v/v) for 60 min at 90 °C, and again dried under vacuum. Samples were resuspended in denaturing loading buffer (80% formamide, 0.1 mM EDTA, 0.1% xylene cyanol, and 0.1% bromophenol blue), electrophoresed through an 18% denaturing PAGE gel,

and visualized using phosphorimager. Control samples were prepared similarly but excluded DEPC and/or piperidine treatment. To obtain values of percent cleavage, the amount of strand cleavage at a given nucleotide was determined relative to total intensity in the lane. Prior to analysis, each lane was corrected for background using the rolling disk lane background correction in Quantity One software. Experiments were performed in triplicate.

Optical Melting Analysis. Melting temperatures for the structures adopted by each oligonucleotide were obtained using a Beckman Coulter DU800 UV-vis spectrophotometer equipped with a Peltier thermoelectric device. Oligonucleotides were diluted to 325 μ L in 20 mM sodium phosphate, 10 mM NaCl, pH 7.0, such that at 90 °C they would have an absorbance at 260 nm of 1.0 ± 0.1 . Prior to optical analysis samples were incubated for 5 min at 90 °C and slowly cooled to room temperature. The samples were then heated at a rate of 1 °C/min from 25 to 90 °C while monitoring absorbance at 260 nm, held at 90 °C for 5 min, and returned to the starting temperature at a rate of 1 °C/min. The T_m was taken as the maximum in the first-derivative plot of the absorbance profile generated during heating from 25 to 90 °C. Experiments were performed in triplicate.

In order to extract van't Hoff enthalpies (ΔH_{VH}) from the data obtained by optical melting analysis, the melting curves were baseline adjusted and fit to a sigmoidal equation.²⁷ The van't Hoff enthalpies were then calculated from the following equation as reported by Breslauer,²⁷ where α represents the fraction of oligonucleotide present in the hairpin conformation:

$$\Delta H_{VH} = 4RT_m^2 \left(\frac{\partial \alpha}{\partial T} \right)_{T=T_m}$$

Determination of Rate Constants for Hairpin to Duplex Conversion. The (CAG)₁₀ strand was 5'-[³²P] end-labeled. Radiolabeled DNA was supplemented with unlabeled DNA to obtain the desired concentrations. The (CAG)₁₀ or (CTG)_n hairpins were obtained by incubating the appropriate oligonucleotide at 90 °C for 5 min followed by slow cooling to room temperature. Prior to mixing, each hairpin was incubated at 37 °C for 30 min; the hairpins were then combined and incubated at 37 °C. Experiments were performed under pseudo-first-order conditions using (CAG)₁₀ and (CTG)_n ($n = 10$ or 11) in 20 mM sodium phosphate, 10 mM NaCl, pH 7.0, in a 10 μ L total sample volume. DNA concentrations were 0.04 μ M for (CAG)₁₀ and 0.4 μ M for (CTG)₁₀ or (CTG)₁₁. The longest time point was mixed first. After the desired incubation time, 5 μ L of nondenaturing loading buffer (15% Ficoll, 0.25% xylene cyanol, and 0.25% bromophenol blue) was added, and the sample was immediately loaded onto a 12% nondenaturing PAGE gel and electrophoresed at 50 V at 4 °C. Mixing of the remaining samples was timed such that there was a 15 min delay between the loading of each sample. After the final sample was loaded onto the gel, electrophoresis at 50 V proceeded for 6 h at 4 °C. Products were visualized using phosphorimager. The amount of duplex product was quantified and plotted as a function of time. The data were fitted in either Origin or Kaleidagraph to a monoexponential curve using the equation $D = H_0(1 - e^{-k't})$, where D is percentage of duplex at time t (in seconds), H_0 is percentage of hairpin at $t = 0$ which was set to 100% during data fitting,²⁸ and k' is the observed rate. The reported second-order rate constant




Figure 1. Overview of the fluorescence quenching experiment showing schematic representations of the nonfluorescent and fluorescent conformations of the FL-(CAG)₁₀-DCL probe.

was obtained by dividing k' by the concentration of the sequence that was in excess. Half-lives were obtained using the equation $t_{1/2} = 1/(kH_0)$, where k is the second-order rate constant and H_0 is the concentration of the (CTG)_{*n*} hairpin at $t = 0$. Experiments were performed in triplicate.

RESULTS

Reactivity of a (CAG)₁₀ Fluorescent Probe in the Presence of (CTG)_{*n*} Sequences of Variable Length. We have previously described a fluorescence-based strategy to assess the propensity of a CAG hairpin to convert to duplex in the presence of a complementary CTG hairpin.²⁵ The fluorescent probe, FL-(CAG)₁₀-DCL, is a (CAG)₁₀ oligonucleotide with fluorescein covalently attached at the 5'-end and a dabcyI quencher at the 3'-end. When the probe is in a hairpin conformation, the fluorophore and quencher are in close proximity and fluorescence quenching occurs (Figure 1). In contrast, when the probe is unstructured or is part of a CAG/CTG duplex the fluorophore and quencher are spatially separated and fluorescence is observed. Using this fluorescent probe, we reported previously the mechanism by which the (CAG)₁₀ and (CTG)₁₀ hairpins convert to the (CAG)₁₀/(CTG)₁₀ duplex.²⁵ The goal of the work described here was to determine how length of the CTG sequence modulates reactivity with the FL-(CAG)₁₀-DCL hairpin. The fluorescence profile of FL-(CAG)₁₀-DCL was obtained by monitoring fluorescence intensity as a function of temperature. As a control, the fluorescence profile of FL-(CAG)₁₀-DCL in the absence of other DNA is shown in Figure 2A (open circles). While at low temperatures the fluorescence is quenched, as the temperature is increased from 10 to 90 °C there is an increase in fluorescence intensity. This increase is consistent with FL-(CAG)₁₀-DCL forming a hairpin at low temperatures which melts as the temperature is raised; a melting temperature (T_m) of 47.0 ± 0.8 °C for FL-(CAG)₁₀-DCL is obtained from this fluorescence profile.




Figure 2. (A) Fluorescence intensity at 520 nm of 1.5 μ M FL-(CAG)₁₀-DCL alone (open circles), 1.5 μ M FL-(CAG)₁₀-DCL in the presence of 1.5 μ M (CTG)₁₀ (closed circles), and 1.5 μ M FL-(CAG)₁₀-DCL in the presence of 1.5 μ M (CTG)₁₁ (open squares). (B) Melting temperature of FL-(CAG)₁₀-DCL in presence of (CTG)_{*n*} sequences. All samples were prepared in 20 mM sodium phosphate, 10 mM NaCl, pH 7.0.

The fluorescent probe was then combined with an equimolar amount of (CTG)_{*n*} where $n = 6 - 14$. Notably, this series of CTG sequences contains 5 even-repeat sequences ($n = 6, 8, 10, 12, 14$) and 4 odd-repeat sequences ($n = 7, 9, 11, 13$). As seen in Figure 2A, addition of an equimolar amount of (CTG)₁₀ results in a shift of the transition in the fluorescence profile to a lower temperature (38.7 ± 0.2 °C; closed circles). Addition of (CTG)₁₁ (open squares) shifts the transition lower still (32.8 ± 0.5 °C). A pattern emerges when comparing the fluorescence profiles of the FL-(CAG)₁₀-DCL probe in the presence of the series of (CTG)_{*n*} sequences (Figure 2B). The transition temperature is higher for the even-repeat (CTG)_{*n*} sequences than for the odd-repeat sequences. In an attempt to reveal the source of this behavior, we next examined the structure adopted by the series of even-repeat and odd-repeat CTG sequences.

Structural Characterization of Even-Repeat and Odd-Repeat (CTG)_{*n*} Sequences. In order to evaluate the structure adopted by the family of CTG sequences, (CTG)₆₋₁₄ were reacted with the chemical probe diethyl pyrocarbonate (DEPC).

DEPC is known to modify adenines and guanines.^{29,30} Furthermore, the level of reactivity of a nucleobase toward DEPC reflects the extent of solvent accessibility, and as a result, adenines and guanines in loops, bulges, or overhangs are more reactive than those in canonical base pairs.^{31,32} For the CTG sequences used in this work the differential reactivity of unpaired guanines was used to reveal loops, bulges, and overhangs and also to distinguish them from guanines in well-matched base pairs. We have previously characterized (CTG)₁₀ based on its reactivity toward DEPC.²⁵ The selective modification of two centrally located guanines (G5 and G6) indicates that the sequence adopts a hairpin conformation with a loop of four nucleobases and a blunt-ended stem with no overhangs (Figure 3A). It is of note that the reactivity at G6 is reduced in comparison to G5; thus, we propose

that G5 is in the loop while G6 is part of the loop-closing base pair with C5.

When examining the reactivity toward DEPC for the rest of the (CTG)_n sequences, there are two patterns that emerge (Figure 4). The first pattern is that there are two reactive guanines in the loop region of even-repeat sequences and three reactive guanines in the loop region of odd-repeat sequences (Figure 4A,C, dashed boxes). The second pattern is the oscillation in the level of reactivity of G1, which is the terminal 5'-G; G1 is significantly more reactive toward DEPC in odd-repeat sequences than in even-repeat sequences (Figure 4B,D). These reactivity patterns are likely due to the presence of a mixture of hairpins that have either 5' or 3' overhangs as in the proposed structures for (CTG)₁₁ (Figure 3B).

To determine whether the oscillating patterns of reactivity toward DEPC observed for the (CTG)_n sequences correspond to these proposed structures, we synthesized a series of standards that mimic the blunt-ended and overhanging stem regions proposed for (CTG)₁₀ and (CTG)₁₁, respectively (Figure 5A). By replacing the two apical nucleobases in the loop of the CTG hairpins with 5'-TT-3', the DNA hairpins can be locked into distinct conformations. First, in the case of (CTG)₁₀ a blunt-ended hairpin (No Overhang) is created. For (CTG)₁₁, substitution of these apical nucleobases creates hairpins with one CTG repeat overhanging the 5' end (5' Overhang), or one repeat overhanging the 3' end (3' Overhang). These standards were subjected to chemical probing with DEPC, and the results obtained were compared to those for (CTG)₁₀ and (CTG)₁₁.

The reactivity of G1 in 5' Overhang is the most intense of all the sequences, indicative of a hairpin conformation with a CTG repeat overhanging the 5' end (Figure 5B,C). On the other hand, 3' Overhang displayed reactivity toward DEPC that is




Figure 3. Schematic representation of the proposed structures for (A) (CTG)₁₀ and (B) (CTG)₁₁ sequences.




Figure 4. (A) Autoradiogram showing DEPC chemical probe reactivity profiles of (CTG)₆ through (CTG)₁₀. Each group contains two control lanes (N is DNA only and C is DNA treated with piperidine) and three replicates (lanes 1, 2, and 3) corresponding to reaction of the DNA with DEPC and piperidine. Conditions were 5 μ M DNA in 20 mM sodium phosphate, 10 mM NaCl, pH 7.0. (B) Percent cleavage at G1 for (CTG)₆₋₁₀. Error represents the standard deviation obtained for the three replicates. (C) Autoradiogram showing DEPC chemical probe reactivity profiles of (CTG)₁₀ through (CTG)₁₄. Each group contains one control lane C which is piperidine-treated DNA and three replicates (lanes 1, 2, and 3) corresponding to reaction with DEPC and piperidine. (D) Percent cleavage at G1 for (CTG)₁₀₋₁₄. Error represents the standard deviation obtained for the three replicates.




Figure 5. (A) Schematic representations of the standards No Overhang, 5' Overhang, 3' Overhang, and Mixed Overhang. The asterisk indicates the location of the 5'-[³²P]-radiolabel. (B) DEPC chemical probe reactivity profiles for standard sequences at 5 μM. Each group contains one control lane (C) which is piperidine-treated DNA and three replicates (lanes 1, 2, and 3) corresponding to reaction with DEPC and piperidine. All samples were prepared in 20 mM sodium phosphate, 10 mM NaCl, pH 7.0. (C) Percent cleavage at G1. Error represents the standard deviation obtained for the three replicates.

comparable to the blunt-ended hairpin No Overhang; this lack of reactivity at G1 is consistent with the absence of a 5' overhang. Lastly, an equimolar mixture of 5' Overhang and 3' Overhang (Mixed Overhang) shows a percent cleavage at G1 that is comparable to that of (CTG)₁₁ (Figure 5C), suggesting that a combination of 5' and 3' overhang-containing structures are the most likely conformations adopted by (CTG)₁₁.

Enthalpy of Melting of Even-Repeat and Odd-Repeat (CTG)_n Hairpins. Analysis by optical melting offers further information about the series of (CTG)_n hairpins by providing melting temperatures and thermodynamic parameters describing the transition from the structured to unstructured state. Optical melting profiles were obtained for the (CTG)_n hairpins in which the absorbance of the DNA was monitored as a function of temperature (Supporting Information). As shown in Table 1,

Table 1. Hairpin Melting Temperatures and van't Hoff Enthalpies of Melting^a

strand	<i>T</i> _m (°C)	Δ <i>H</i> _{VH} (kcal/mol)
(CTG) ₆	54.3 ± 0.8	38 ± 1
(CTG) ₇	54.6 ± 0.8	40 ± 2
(CTG) ₈	56.0 ± 0.2	46 ± 1
(CTG) ₉	55.5 ± 0.8	48 ± 2
(CTG) ₁₀	56.4 ± 0.3	56 ± 2
(CTG) ₁₁	56.3 ± 0.2	56 ± 5
(CTG) ₁₂	57.5 ± 0.3	57 ± 2
(CTG) ₁₃	58.3 ± 0.1	56 ± 1
(CTG) ₁₄	57.2 ± 0.1	65 ± 4
No Overhang	56.3 ± 0.1	82 ± 7
5' Overhang	55.3 ± 0.2	66 ± 2
3' Overhang	54.6 ± 0.1	64 ± 1

^a Error represents standard deviation derived from three experiments.




Figure 6. Enthalpies of melting for (CTG)_n sequences derived from optical melting data by van't Hoff analysis.

there is a general increase in the *T*_m across the series of hairpins from *n* = 6 to 14. To understand the thermodynamic impact of additional CTG repeats, we extracted the model-dependent van't Hoff enthalpy (Δ*H*_{VH}) from the optical melting profiles. As seen in Table 1, Δ*H*_{VH} increases across the series of hairpins. Instead of increasing linearly with the number of repeats an apparent stepping is observed (Figure 6). Starting with the hairpins (CTG)₆ and (CTG)₇, both show similar enthalpies of melting. Likewise, (CTG)₈ and (CTG)₉ have similar enthalpies of melting with the Δ*H*_{VH} being larger relative to (CTG)₆ and (CTG)₇. (CTG)₁₀, (CTG)₁₁, (CTG)₁₂, and (CTG)₁₃ display similar enthalpies of melting at the next "step". Finally, (CTG)₁₄ displays the largest enthalpy of melting in this series of hairpins.

Rate of Hairpin to Duplex Conversion for (CAG)₁₀ and (CTG)_n Hairpins. We have previously shown that (CAG)₁₀ and (CTG)₁₀ hairpins can convert to the (CAG)₁₀/(CTG)₁₀ duplex.²⁵ In order to quantitatively describe the (CAG)₁₀ and (CTG)_n hairpins converting to (CAG)₁₀/(CTG)_n duplex, we determined the rate constant for this process. These kinetic experiments were performed under pseudo-first-order conditions in which radiolabeled 5'-³²P-(CAG)₁₀ hairpin was combined with a 10-fold excess of (CTG)₁₀, which was used to represent the even-repeat hairpins, or (CTG)₁₁, which was




Figure 7. (A) Autoradiogram revealing conversion of (CAG)₁₀ and (CTG)₁₁ hairpins to the (CAG)₁₀/(CTG)₁₁ duplex and (B) plot of percent duplex as a function of time showing the fit to a monoexponential curve. (C) Autoradiogram revealing conversion of (CAG)₁₀ and (CTG)₁₀ hairpins to (CAG)₁₀/(CTG)₁₀ duplex and (D) plot of percent duplex as a function of time showing the fit to a monoexponential curve. Experiments were carried out at 37 °C in 20 mM sodium phosphate, 10 mM NaCl, pH 7.0.

used to represent the odd-repeat hairpins. After incubation of 5'-³²P-(CAG)₁₀ and (CTG)_n at 37 °C for a defined period of time the hairpin and duplex species were separated by non-denaturing PAGE. The amount of duplex product was quantitated and plotted as a function of time, and the data were fitted to a monoexponential curve in order to obtain the rate constant for duplex formation.

Using 0.04 μM (CAG)₁₀ and 0.4 μM (CTG)₁₁, a rate constant of $4625 \pm 728 \text{ M}^{-1} \text{ s}^{-1}$ was obtained for the hairpin to duplex conversion (Figure 7A,B). When comparable experiments were performed using 0.04 μM (CAG)₁₀ and 0.4 μM (CTG)₁₀, the rate was determined to be $1085 \pm 214 \text{ M}^{-1} \text{ s}^{-1}$ (Figure 7C,D), 4-fold slower than duplex conversion for (CAG)₁₀ and (CTG)₁₁. Thus, conversion to duplex for (CAG)₁₀ in presence of the odd-repeat hairpin (CTG)₁₁ occurs faster than in presence of the even-repeat hairpin (CTG)₁₀. On the basis of these rates, the upper-limit for the half-life of the (CTG)₁₁ hairpin is $9 \pm 1 \text{ min}$ and $40 \pm 8 \text{ min}$ for the (CTG)₁₀ hairpin.

DISCUSSION

(CTG)_n Repeat Number Modulates Reactivity with (CAG)₁₀ Complement. We have previously described the mechanism of hairpin to duplex conversion for the TNR hairpins (CAG)₁₀ and (CTG)₁₀.²⁵ In fact, we identified two distinct interactions that mediate conversion to duplex by the TNR hairpins. These are kissing interactions, which result from base pairing by the complementary nucleobases in the loop of the CAG and CTG hairpins, and cruciform interactions, which occur between the terminal nucleobases of the hairpin stems. In this previous report we focused on characterizing sequences containing 10 repeat units, and we sought here to expand our knowledge regarding the effect of repeat sequence length of the complementary (CTG)_n sequence on the (CAG)₁₀ hairpin.

Beginning with our fluorescence-based assay, we characterized the thermal stability of the FL-(CAG)₁₀-DCL hairpin probe in the presence of the family of (CTG)_n sequences. We observed an oscillating pattern in the thermal stability of the probe in presence of even-repeat sequences versus odd-repeat sequences. The CAG hairpin probe melts at a lower temperature in presence of a CTG hairpin containing an odd number of repeats as compared to an even number of repeats. This pattern revealed a higher degree of reactivity of the odd-repeat CTG sequences with the complementary (CAG)₁₀ hairpin, and we hypothesized that differences in structure between even- and odd-repeat sequences were the cause of the oscillating pattern observed by fluorescence.

Chemical Probes Reveal a Constant Loop Size for the (CTG)_n Hairpins and an Overhanging Repeat in the Odd-Repeat Hairpins. In order to characterize the structure(s) adopted by these sequences in solution, we conducted chemical probing experiments with DEPC. Interestingly, the oscillating pattern observed in the fluorescence profiles for the even- and odd-repeat sequences is echoed in the DEPC reactivity of the family of (CTG)_n sequences. Notably, even-repeat sequences show two reactive guanines in the loop region and low reactivity at G1, whereas odd-repeat sequences show three reactive guanines in the loop region and high reactivity at G1.

There are two scenarios that could explain these observed DEPC reactivity patterns. First, the size of the loop might be alternating between the even- and odd-repeat hairpins. For instance, even-repeat hairpins could have a loop of 4 nucleobases while odd-repeat hairpins could have a loop of 7 nucleobases. If this were the case, the stem of both the even- and odd-repeat hairpins would be blunt-ended, and there would be no overhanging bases.

On the basis of NMR studies using (CTG)_n sequences where $n = 4-10$, Chi and Lam concluded that loop size alternates

between 3 and 4 nucleobases for odd- and even-repeat sequences, respectively.³³ However, in order to suppress the formation of homoduplex at the relatively high DNA concentrations required for NMR, the repeat hairpins also included a well-matched and nonrepetitive stem extension of 4 base pairs. While this stem extension provides a kinetic advantage to the formation of an intramolecular hairpin over an intermolecular homoduplex, it also prevents overhangs since this extension forms the end of the stem. Thus, we cannot compare our results directly to those obtained by Chi and Lam.

Further support for loop size alternating between 3 and 4 nucleobases for odd- and even-repeat sequences came from Mariappan and co-workers, who used NMR to examine the conformation adopted by (CTG)₅ and (CTG)₆.³⁴ They reported that (CTG)₅ forms a 3 nucleobase loop while (CTG)₆ forms a 4 nucleobase loop. The stem of both hairpins is blunt-ended. Additionally, on the basis of cleavage of the DNA by single-strand specific nucleases, Amrane and co-workers proposed that the loop size alternates between 3 and 4 nucleobases for (CTG)₁₅ and (CTG)₁₆; notably, features such as 5' bulges and 3' overhangs were also proposed.³⁵

If the odd-repeat hairpins had smaller loops than the even-repeat hairpins, we would expect a trend opposite to that seen in Figure 4A,C, with even-repeat hairpins having more reactive guanines in the loop region. In contrast, we observe that the even-repeat hairpins have two reactive guanines while the odd-repeat hairpins have three. While it may be tempting to suggest, based on the number of guanines in the loop region that are modified by DEPC, that the odd-repeat hairpins have a larger loop than even-repeat hairpins, this conclusion is not consistent with the reactivity observed in the stem region.

A second scenario that can explain the DEPC pattern observed for the family of (CTG)_n hairpins would be for the loop size to remain constant rather than alternate between even- and odd-repeat hairpins. If the loop size remains the same across the family of (CTG)_n hairpins, it would be most likely comprised of 4 nucleobases. In order to accommodate this constant loop size, one CTG repeat would overhang the 5' or 3' end of the odd-repeat hairpins. Since the same type of nucleobases (C, T, G) would be present in either a 5' or 3' overhang neither is expected to form preferentially. In fact, a mixture of both species of overhangs is likely. It is noteworthy that if the loop size alternates between even- and odd-repeat hairpins, the reactivity toward DEPC would represent the reactivity of a *single* DNA conformation, whereas if the loop size was constant across the series, the reactivity pattern would represent a superimposition of those generated by *two* species in solution.

If there are no overhangs present and the hairpins have blunt ends, we would expect a similar amount of modification of G1 across the series of hairpins. If odd-repeat hairpins have a mixture of 5' and 3' overhangs, we would expect the species with the 5'-overhang to be more reactive toward DEPC at G1 than the even-repeat hairpins. Indeed, we observe more reactivity at G1 in the case of the odd-repeat hairpins (Figure 4B,D).

The results we obtained for modification of the (CTG)_n (*n* = 6–14) hairpins by DEPC are consistent with this series of hairpins maintaining the same loop size. We came to this conclusion by examining the DEPC reactivity patterns in both the loop and stem regions of the hairpins. Therefore, we propose that both the even-repeat and odd-repeat hairpins have a loop of 4 nucleobases and that the stem is blunt-ended when there is an even number of CTG repeats and one repeat overhanging the

stem in odd-repeat hairpins. It is noteworthy that our proposed structure for (CTG)₆ is consistent with that reported by Mariappan and co-workers based on NMR characterization of the same sequence.³⁴

If all the hairpins have the same size loop, why is the number of reactive guanines in the loop region not the same across the series? Examination of the two structures proposed for (CTG)₁₁ depicted in Figure 3B provides the rationale: the 5' and 3' overhang species have different guanines in the loop and loop-closing base pair. In the (CTG)₁₁ hairpin with a 5' overhang, G6 is in the loop and G7 is in the loop-closing base pair. For the (CTG)₁₁ hairpin with a 3' overhang, G5 is in the loop and G6 is in the loop-closing base pair. When the DEPC reactivity patterns of these two hairpins are overlaid, one would expect to see three reactive guanines: G5, G6, and G7. Moreover, the extent of reactivity for each guanine will depend on its position in the two hairpins. G6 is expected to have the highest reactivity, as it either is in the loop or is part of a loop-closing base pair, G5 would have less reactivity toward DEPC than G6 because it is either in the loop or in the stem, and last, G7 will be the least reactive toward DEPC since it is in either the loop-closing base pair or the stem. Indeed, consistent with the predicted structures for (CTG)₁₁, the observed level of reactivity toward DEPC is G6 > G5 > G7.

To confirm these predictions, we constructed a series of standards that, due to substitution of various nucleobases in the loop, adopt locked structures differing only in presence or absence of 5' and 3' overhangs. Several predictions can be made about the expected reactivity of these standards toward DEPC. First, the No Overhang hairpin has a blunt end, and therefore, reactivity of guanines in the stem should mimic those in the (CTG)₁₀ stem. In contrast, the 5' Overhang and 3' Overhang hairpins were designed to mimic the proposed (CTG)₁₁ structures that have either a 5' or 3' repeat overhanging the stem, respectively. Therefore, the 5' Overhang standard should display more reactivity at G1 than (CTG)₁₁ because (CTG)₁₁ is predicted to have half as many G1 nucleobases in a 5' overhang as compared to the 5' Overhang standard. The 3' Overhang standard has only a 3' overhang, and not a 5' overhang, and should not have heightened reactivity at G1 relative to (CTG)₁₁. The Mixed Overhang sample is an equimolar mixture of the 5' and 3' Overhang hairpin standards, and therefore, if the proposed (CTG)₁₁ structures are correct, the reactivity profile at G1 for Mixed Overhang should resemble that observed for (CTG)₁₁.

Indeed, the observed reactivity of these standards toward DEPC is consistent with our proposed (CTG)₁₁ structures and confirms that (CTG)₁₁ adopts two distinct structures with a 4 nucleobase loop and either a 5' or 3' overhang. The use of these standards supports our conclusions that loop size is constant across the series of (CTG)_n hairpins (*n* = 6–14) and that it is the absence or presence of an overhang that varies in the even- and odd-repeat hairpins.

Enthalpy of Melting Supports a Constant Loop Size in Even-Repeat and Odd-Repeat Hairpins. Van't Hoff analysis of thermal melting data revealed that (CTG)₆ and (CTG)₇ have a similar ΔH_{VH} . Our proposed structures for these (CTG)_{even} and (CTG)_{odd} hairpins have the same loop size and the same number of well-matched and mismatched base pairs in their stems. Assuming that the base stacking contributed by the nucleobases in the overhang is minimal, we would predict they would have a similar ΔH_{VH} . Similarly, on the basis of our proposed structures, we would predict that (CTG)₈ and (CTG)₉ have similar enthalpies of melting with ΔH_{VH} being greater than that observed when

$n = 6$ or 7 due to the base stacking and hydrogen-bonding interactions of three additional base pairs in the stem. At the next “step” in ΔH_{VH} we observe (CTG)₁₀, (CTG)₁₁, (CTG)₁₂, and (CTG)₁₃. Notably, a plateau in ΔH has been reported previously for TNR hairpins.³⁵ However, we observe that ΔH_{VH} of (CTG)₁₄ is larger than that observed for (CTG)₁₀, (CTG)₁₁, (CTG)₁₂, and (CTG)₁₃. It should be noted that van’t Hoff analysis assumes a two-state transition, where the hairpin proceeds directly from the structured to the unstructured state. If intermediates populate this transition, ΔH_{VH} will be an underestimation of ΔH .²⁷ Using calorimetric methods, which provide values of ΔH that are independent of the nature of the melting transition (ΔH_{cal}), it was observed that when there are 12 or more CTG repeats, $\Delta H_{\text{VH}} \neq \Delta H_{\text{cal}}$.³⁵ Thus, the ΔH_{VH} we obtained for sequences containing 12, 13, and 14 CTG repeats is likely an underestimate of ΔH and does not reflect a true plateau.

With an understanding of the structure and thermodynamics of the (CTG)_{*n*} sequences, the oscillating pattern of fluorescence transitions observed for even- and odd-repeat sequences can be rationalized. The transitions observed in the presence of the odd-repeat hairpins are at lower temperatures than those for the even-repeat hairpins (Figure 2B). The transition temperature reflects FL-(CAG)₁₀-DCL melting as a result of conversion to duplex to form FL-(CAG)₁₀-DCL/(CTG)_{*n*}. While the FL-(CAG)₁₀-DCL/(CTG)_{*n*} duplexes formed with even- and odd-repeat CTG sequences are expected to have the same thermodynamic stability, the fluorescence data suggest that odd-repeat hairpins, which have one repeat overhanging the stem, convert more readily to duplex than the even-repeat hairpins that lack these overhangs. Thus, we determined the rate constant describing the hairpin to duplex conversion for even- and odd-repeat hairpins.

Kinetic Experiments Reveal That Structural Differences Alter the Rate of Hairpin to Duplex Conversion. We reported previously that the complementary (CAG)₁₀ and (CTG)₁₀ hairpins can convert to duplex via two independent mechanisms: kissing or cruciform interactions in which the hairpins interact by the loop or stem regions, respectively.²⁵ Furthermore, we have shown that kissing and cruciform interactions contribute equally to the rate of duplex formation by (CAG)₁₀ and (CTG)₁₀. In this work, our kinetic data reveal that hairpin to duplex conversion occurs faster for the (CAG)₁₀ hairpin in the presence of the odd-repeat hairpin (CTG)₁₁ than in the presence of (CTG)₁₀. In fact, we find that the rate of formation of (CAG)₁₀/(CTG)₁₁ duplex is 4-fold faster than formation of the (CAG)₁₀/(CTG)₁₀ duplex. These results indicate that, when present together, the lifetimes of the (CAG)₁₀ and (CTG)₁₀ hairpins are longer than the lifetimes of (CAG)₁₀ and (CTG)₁₁. We postulate that the unpaired nucleobases of the overhanging repeat on either the 3' or 5' end of the (CTG)₁₁ hairpin facilitate cruciform interactions and are responsible for the faster conversion to duplex and shorter lifetimes of the (CAG)₁₀ and (CTG)₁₁ hairpins.

Taken together, the results obtained by chemical probing with DEPC, optical melting analysis, and kinetic analysis by non-denaturing PAGE indicate that the structural differences in the series of (CTG)_{*n*} modulate their ability to form (CAG)₁₀/(CTG)_{*n*} duplex and that these differences are responsible for the oscillating even/odd patterns seen in our fluorescence experiments using the probe FL-(CAG)₁₀-DCL.

Lifetime of Hairpins as a Contributing Factor to TNR Expansion. It has been proposed that the presence of hairpins formed by TNR sequences induces expansion. If hairpin formation were to occur during DNA replication, slippage of polymerase

during DNA synthesis could promote expansion. Additionally, during the gap-filling step of long-patch base excision repair (LP-BER), formation of a structured hairpin flap could inhibit cleavage by the flap endonuclease FEN1 and lead to expansion.³ We can therefore envision that lifetime of these hairpins during replication and/or repair may dictate whether or not expansion occurs.

In this work we have shown that the lifetime of (CTG)_{*n*} TNR hairpins is modulated by whether the number of repeats is even or odd. Indeed, we have shown that when $n = 10$ the lifetime of the hairpin is significantly longer than when $n = 11$. Thus, the lifetime of hairpins formed during replication or from the single-stranded flap generated in LP-BER may be controlled by the number of repeats in the hairpin.

It is of note that an oscillating pattern in the behavior of even- and odd-repeat TNR sequences was also reported by Darlow and Leach, who evaluated the size of viral plaques in bacteria infected with phage containing an even or odd number of CAG/CTG repeats.³⁶ The repeats were embedded in a palindromic sequence which is capable of forming a hairpin. Using bacteriophages that contained 1–5 CAG/CTG repeats, plaque size was smaller for bacteria infected with phages containing an even number of repeats than for those containing an odd number of repeats.³⁶ Earlier work from the same laboratory³⁷ demonstrated a correlation between plaque size, phage infectivity, and palindromic hairpin stability; phages containing more stable hairpins are less infective and yield smaller plaques. Thus, Darlow and Leach postulated that since phages containing even-repeat sequences yield smaller plaques, even-repeat sequences form more stable hairpins than odd-repeat hairpins.

While admittedly quite different systems, our results can shed light on the viral infectivities observed for phage DNA containing an even or odd number of TNR repeats. In the phage system used by Darlow and Leach the even-repeat hairpins may have longer lifetimes than the odd-repeat hairpins. When considering DNA constrained in a circular phage genome, as opposed to the unconstrained oligonucleotides used in the work described here, cruciform interactions, as opposed to kissing interactions, are the likely mechanism for hairpin to duplex conversion since the terminal stem bases of the complementary strands can interact and unzip the hairpins to convert to duplex. While overhangs are not possible in circular DNA, the structure of the even versus odd repeats may still modulate conversion to duplex.

Our findings pose an interesting question regarding TNR expansion: Is expansion more likely in alleles that contain an even number of repeats than in alleles that have an odd number of repeats? Clinical data gathered to examine the phenomenon of anticipation in HD patients display no noticeable oscillation in expansion propensity between an even or odd number of repeats. Offspring of HD patients possess TNR regions that are dependent on the total number of repeats in the parental allele regardless of whether the parental allele had an even or odd number of repeats.³⁸ This observation suggests that in vivo the structural differences in sequences containing an even or odd number of repeats are not as clearly differentiated as the ones studied in this work. However, it is noteworthy that our studies characterized sequence lengths that would be found in healthy individuals; repeat tracts of these lengths are not prone to expansion. Interestingly, our findings suggest that even though sequences of this length are not prone to expansion the ability for hairpins to form and persist in repeat lengths that are considered healthy may not be uniform across the entire genome.

■ ASSOCIATED CONTENT

S Supporting Information. Optical melting curves for (CTG)_{6–14} and standard sequences. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Tel: (401) 863-3590. Fax: (401) 863-9368. E-mail: sarah_delaney@brown.edu

Author Contributions

[†]These authors contributed equally to this work.

Funding Sources

This work was supported by the National Institute of Environmental Health Sciences (R01ES019296). A.A.F. was supported by a National Science Foundation Graduate Research Fellowship.

■ ACKNOWLEDGMENT

We thank Prof. David Cane for use of phosphorimaging equipment, Ms. Diana Lizarazo for conducting initial experiments, and Mr. Daniel Jarem and Mr. Craig Yennie for helpful discussions. The real-time PCR instrument used in this work is located in the Molecular Pathology Core of the COBRE Center for Cancer Research Development, NIH P20 RR17695, awarded by the National Center for Research Resources, Institutional Development Award (IdeA) Program.

■ ABBREVIATIONS

TNR, trinucleotide repeat; FL-(CAG)₁₀-DCL, fluorescein-5'-(CAG)₁₀-3'-(E)-N-(3-(dihydroxymethoxy)propyl)-4-((4-(dimethylamino)phenyl)diazanyl)benzamide; DEPC, diethyl pyrocarbonate; PAGE, polyacrylamide gel electrophoresis.

■ REFERENCES

- (1) Mirkin, S. M. (2007) Expandable DNA repeats and human disease. *Nature* 447, 932–940.
- (2) Kovtun, I. V., and McMurray, C. T. (2008) Features of trinucleotide repeat instability in vivo. *Cell Res.* 18, 198–213.
- (3) McMurray, C. T. (2010) Mechanisms of trinucleotide repeat instability during human development. *Nat. Rev. Genet.* 11, 786–799.
- (4) Castel, A. L., Cleary, J. D., and Pearson, C. E. (2010) Repeat instability as the basis for human diseases and as a potential target for therapy. *Nat. Rev. Mol. Cell Biol.* 11, 165–170.
- (5) Orr, H. T., and Zoghbi, H. Y. (2007) Trinucleotide repeat disorders. *Annu. Rev. Neurosci.* 30, 575–621.
- (6) MacDonald, M. E. et al. (1993) A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's-disease chromosomes. *Cell* 72, 971–983.
- (7) Mitas, M. (1997) Trinucleotide repeats associated with human disease. *Nucleic Acids Res.* 25, 2245–2253.
- (8) Paiva, A. M., and Sheardy, R. D. (2004) Influence of sequence context and length on the structure and stability of triplet repeat DNA oligomers. *Biochemistry* 43, 14218–14227.
- (9) Paiva, A. M., and Sheardy, R. D. (2005) The influence of sequence context and length on the kinetics of DNA duplex formation from complementary hairpins possessing (CNG) repeats. *J. Am. Chem. Soc.* 127, 5581–5585.
- (10) Mirkin, S. M. (2006) DNA structures, repeat expansions and human hereditary disorders. *Curr. Opin. Struc. Biol.* 16, 351–358.

- (11) Wells, R. D. (2007) Non-B DNA conformations, mutagenesis and disease. *Trends Biochem. Sci.* 32, 271–278.
- (12) Degtyareva, N. N., Reddish, M. J., Sengupta, B., and Petty, J. T. (2009) Structural Studies of a trinucleotide repeat sequence using 2-aminopurine. *Biochemistry* 48, 2340–2346.
- (13) Degtyareva, N. N., Barber, C. A., Sengupta, B., and Petty, J. T. (2010) Context dependence of trinucleotide repeat structures. *Biochemistry* 49, 3024–3030.
- (14) Jarem, D. A., Huckaby, L. V., and Delaney, S. (2010) AGG interruptions in (CGG)_n DNA repeat tracts modulate the structure and thermodynamics of non-B conformations in vitro. *Biochemistry* 49, 6826–6837.
- (15) Degtyareva, N. N., Barber, C. A., Reddish, M. J., and Petty, J. T. (2011) Sequence length dictates repeated CAG folding in three-way junctions. *Biochemistry* 50, 458–465.
- (16) Gacy, A. M., Goellner, G., Juranic, N., Macura, S., and McMurray, C. T. (1995) Trinucleotide repeats that expand in human-disease form hairpin structures in-vitro. *Cell* 81, 533–540.
- (17) Liu, G. Q., Chen, X. M., Bissler, J. J., Sinden, R. R., and Leffak, M. (2010) Replication-dependent instability at (CTG)·(CAG) repeat hairpins in human cells. *Nat. Chem. Biol.* 6, 652–659.
- (18) Pearson, C. E., and Sinden, R. R. (1996) Alternative structures in duplex DNA formed within the trinucleotide repeats of the myotonic dystrophy and fragile X loci. *Biochemistry* 35, 5041–5053.
- (19) Pearson, C. E., Wang, Y. H., Griffith, J. D., and Sinden, R. R. (1998) Structural analysis of slipped-strand DNA (S-DNA) formed in (CTG)_n·(CAG)_n repeats from the myotonic dystrophy locus. *Nucleic Acids Res.* 26, 816–823.
- (20) Pearson, C. E., Tam, M., Wang, Y. H., Montgomery, S. E., Dar, A. C., Cleary, J. D., and Nichol, K. (2002) Slipped-strand DNAs formed by long (CAG)·(CTG) repeats: slipped-out repeats and slip-out junctions. *Nucleic Acids Res.* 30, 4534–4547.
- (21) Pearson, C. E., and Sinden, R. R. (1998) Trinucleotide repeat DNA structures: dynamic mutations from dynamic DNA. *Curr. Opin. Struc. Biol.* 8, 321–330.
- (22) Kovtun, I. V., and McMurray, C. T. (2007) Crosstalk of DNA glycosylases with pathways other than base excision repair. *DNA Repair* 6, 517–529.
- (23) Zhao, J., Bacolla, A., Wang, G., and Vasquez, K. M. (2010) Non-B DNA structure-induced genetic instability and evolution. *Cell. Mol. Life Sci.* 67, 43–62.
- (24) Beaucage, S. L., and Caruthers, M. H. (2000) Synthetic strategies and parameters involved in the synthesis of oligodeoxyribonucleotides according to the phosphoramidite method. *Curr. Protoc. Nucleic Acid Chem.* 3.3.1–3.3.20.
- (25) Ávila-Figueroa, A., and Delaney, S. (2010) Mechanistic studies of hairpin to duplex conversion for trinucleotide repeat sequences. *J. Biol. Chem.* 285, 14648–14657.
- (26) Warshaw, M. M., and Tinoco, I., Jr. (1966) Optical properties of sixteen dinucleoside phosphates. *J. Mol. Biol.* 20, 29–38.
- (27) Marky, L. A., and Breslauer, K. J. (1987) Calculating thermodynamic data for transitions of any molecularity from equilibrium melting curves. *Biopolymers* 26, 1601–1620.
- (28) Vo, M. N., Barany, G., Rouzina, I., and Musier-Forsyth, K. (2006) Mechanistic studies of mini-TAR RNA/DNA annealing in the absence and presence of HIV-1 nucleocapsid protein. *J. Mol. Biol.* 363, 244–261.
- (29) Leonard, N., McDonald, J., Henderson, R., and Reichmann, M. (1971) Reaction of diethyl pyrocarbonate with nucleic acid components. Adenosine. *Biochemistry* 10, 3335–3342.
- (30) Vincze, A., Henderson, R., McDonald, J., and Leonard, N. (1973) Reaction of diethyl pyrocarbonate with nucleic acid components. Bases and nucleosides derived from guanine, cytosine and uracil. *J. Am. Chem. Soc.* 95, 2677–2682.
- (31) Herr, W. (1985) Diethyl pyrocarbonate: a chemical probe for secondary structure in negatively supercoiled DNA. *Proc. Natl. Acad. Sci. U.S.A.* 82, 8009–8013.
- (32) Huertas, D., Bellsollé, L., Casasnovas, J., Coll, M., and Azorín, F. (1993) Alternating d(GA)_n DNA sequences form antiparallel

stranded homoduplexes stabilized by the formation of G•A base pairs. *EMBO J.* 12, 4029–4038.

(33) Chi, L. M., and Lam, S. L. (2005) Structural roles of CTG repeats in slippage expansion during DNA replication. *Nucleic Acids Res.* 33, 1604–1617.

(34) Mariappan, S. V. S., Garcia, A. E., and Gupta, G. (1996) Structure and dynamics of the DNA hairpins formed by tandemly repeated CTG triplets associated with myotonic dystrophy. *Nucleic Acids Res.* 24, 775–783.

(35) Amrane, S., Sacca, B., Mills, M., Chauhan, M., Klump, H. H., and Mergny, J. L. (2005) Length-dependent energetics of (CTG)_(n) and (CAG)_(n) trinucleotide repeats. *Nucleic Acids Res.* 33, 4065–4077.

(36) Darlow, J. M., and Leach, D. R. F. (1995) The effects of trinucleotide repeats found in human inherited disorders on palindrome inviability in *Escherichia coli* suggest hairpin folding preferences in vivo. *Genetics* 141, 825–832.

(37) Davison, A., and Leach, D. R. F. (1994) Two-base DNA hairpin-loop structures in vivo. *Nucleic Acids Res.* 22, 4361–4363.

(38) Wheeler, V. C., Persichetti, F., McNeil, S. M., Mysore, J. S., Mysore, S. S., MacDonald, M. E., Myers, R. H., Gusella, J. F., Wexler, N. S., and The US-Venezuela Collaborative Research Group. (2007) Factors associated with HD CAG repeat instability in Huntington disease. *J. Med. Genet.* 44, 695–701.