


Retinal Degeneration 3 (RD3) Protein Inhibits Catalytic Activity of Retinal Membrane Guanylyl Cyclase (RetGC) and Its Stimulation by Activating Proteins

Igor V. Peshenko,[‡] Elena V. Olshevskaya,[‡] Seifollah Azadi,[§] Laurie L. Molday,[§] Robert S. Molday,[§] and Alexander M. Dizhoor^{*,‡}

[‡]Department of Basic Sciences and Pennsylvania College of Optometry, Salus University, Elkins Park, Pennsylvania 19027, United States

[§]Department of Biochemistry and Molecular Biology, University of British Columbia, Vancouver, BC V6T 1Z3, Canada

ABSTRACT: Retinal membrane guanylyl cyclase (RetGC) in the outer segments of vertebrate photoreceptors is controlled by guanylyl cyclase activating proteins (GCAPs), responding to light-dependent changes of the intracellular Ca^{2+} concentrations. We present evidence that a different RetGC binding protein, retinal degeneration 3 protein (RD3), is a high-affinity allosteric modulator of the cyclase which inhibits RetGC activity at submicromolar concentrations. It suppresses the basal activity of RetGC in the absence of GCAPs in a noncompetitive manner, and it inhibits the GCAP-stimulated RetGC at low intracellular Ca^{2+} levels. RD3 opposes the allosteric activation of the cyclase by GCAP but does not significantly change Ca^{2+} sensitivity of the GCAP-dependent regulation. We have tested a number of mutations in RD3 implicated in human retinal degenerative disorders and have found that several mutations prevent the stable expression of RD3 in HEK293 cells and decrease the affinity of RD3 for RetGC1. The RD3 mutant lacking the carboxy-terminal half of the protein and associated with Leber congenital amaurosis type 12 (LCA12) is unable to suppress the activity of the RetGC1/GCAP complex. Furthermore, the inhibitory activity of the G57V mutant implicated in cone-rod degeneration is strongly reduced. Our results suggest that inhibition of RetGC by RD3 may be utilized by photoreceptors to block RetGC activity during its maturation and/or incorporation into the photoreceptor outer segment rather than participate in dynamic regulation of the cyclase by Ca^{2+} and GCAPs.



Cyclic GMP in vertebrate photoreceptors couples rhodopsin in disk membranes to cGMP-gated cation channels in the plasma membrane of rod outer segments (ROS) through a phototransduction cascade (reviewed in ref 1). Following photoexcitation, cGMP in photoreceptors is first hydrolyzed by a light-activated phosphodiesterase (PDE6) and then replenished by a retinal membrane guanylyl cyclase (RetGC).^{2–4} In addition to its role as a key enzyme in photoreceptor physiology, RetGC has been linked to multiple cases of congenital blinding disorders in human patients including Leber congenital amaurosis type 1 (LCA1) and dominant cone-rod dystrophy.^{5–7} RetGC activity in rods and cones is controlled by Ca^{2+} -sensitive guanylyl cyclase activating proteins, GCAPs,^{8–11} via a Ca^{2+} feedback mechanism. When cGMP-gated channels close in response to light, the reduced influx of Ca^{2+} through the channels causes GCAP to switch from a Ca^{2+} -bound (inhibitor) state to a Mg^{2+} -bound (RetGC activator) state (reviewed in¹²). This accelerates the resynthesis of cGMP for timely recovery of photoreceptors to their dark state or establishing light adaptation.^{13,14}

More recently, it has been shown that RD3, a 23-kDa protein unrelated to GCAPs and linked to Leber congenital amaurosis type12 (LCA12) and rapid retinal degeneration in the *rd3*

mouse,¹⁵ coimmunoprecipitates with RetGC and is essential for the normal expression of RetGC in rod and cone photoreceptor cells.¹⁶ The RD3 gene transcripts are highly abundant in the retina and concentrated in photoreceptors but undetectable in other mouse tissues.¹⁵ An anti-RD3 polyclonal antibody was observed to stain the inner and outer segments of mouse rod and cone photoreceptor cells by immunofluorescence microscopy.¹⁶ RD3 also colocalized with RetGC in intracellular vesicles when both proteins were coexpressed in HEK293 cells,¹⁶ thus leading to a hypothesis that at least one function of RD3 in photoreceptor cells is to participate in the intracellular trafficking of the cyclase.

In the present study, we have investigated whether the association of RetGC with RD3 can affect the catalytic function of RetGC. We find that RD3 acts as a high-affinity allosteric inhibitor of RetGC, capable of both effectively competing with GCAPs and suppressing the catalytic activity of the cyclase. Some mutations in the human RD3 gene found in patients with

Received: August 24, 2011

Revised: September 16, 2011

Published: September 19, 2011

congenital retinal disorders strongly affect the inhibitory activity of the human recombinant RD3 protein in vitro.

EXPERIMENTAL PROCEDURES

Antibodies. Rabbit polyclonal antibody RD3 497 was raised against the isolated full-length human recombinant RD3 expressed in *E. coli* as described below and purified from the serum by immunoaffinity chromatography on RD3 coupled to CNBr-activated Sepharose CL-4B (GE Heath Sciences). The Rho-1D4 mouse monoclonal antibody¹⁷ was against the TETSQVAPA peptide used as a C-terminal tag in some RD3 constructs, and the anti-RD3 mouse monoclonal RD3-9D12 antibody was produced against the C-terminal peptide of RD3.¹⁶ The rabbit polyclonal anti-RetGC1 antibody was produced against the catalytic domain of human RetGC1,¹⁸ and the mouse monoclonal GC-8A5 antibody was raised against the C-terminus of mouse RetGC1.

Human Recombinant RD3 Expression in *E. coli*. Human RD3 cDNA was amplified from a pCMV-SPORT6/MHS1010–9206149 cDNA clone (Open Biosystems/Thermo Scientific) using high-fidelity Phusion Flash DNA polymerase (Finnzymes), subcloned into the *NcoI*/*BamHI* sites of pET11d vector (Novagen/Calbiochem), sequenced, and expressed in a BL21(DE3)CodonPlus (Agilent Technologies) *E. coli* strain in the presence of isopropyl β -D-1-thiogalactopyranoside for 2 h. The protein which accumulated in inclusion bodies was purified by series of sonication and centrifugation cycles described for GCAP purification,¹⁹ solubilized in 10 mM Tris-HCl (pH 7.5) buffer containing 2 mM EDTA, 8 M urea, and 14 mM 2-mercaptoethanol, and dialyzed against 2 \times 300 volumes of 10 mM Tris-HCl (pH 7.5) buffer containing 0.1 mM EDTA and 14 mM 2-mercaptoethanol at 4 °C. Insoluble protein was removed by centrifugation at 10000g for 10 min at 4 °C, and the supernatant containing RD3 (typically 80–90% purity by SDS PAGE) was collected and used either immediately or after storage at –70 °C with 50% v/v glycerol. RD3 has a tendency to precipitate under normal storage conditions. For expression of RD3 mutants¹⁵ in *E. coli*, the corresponding mutations were introduced into the RD3 cDNA using Phusion Flash DNA polymerase (Finnzymes) by a conventional “splicing by overlap extension” method and the mutated cDNA was verified by sequencing the entire coding region of the resulting plasmid.

Expression of a Full-Length Human RD3 in HEK293 Cells Used in RetGC Activity Assays. The RD3 cDNA was inserted into the *HindIII*/*XbaI* sites of a modified pRCCMV vector (Invitrogen), transfected into a 50–80% confluent cell culture (ca. 20 μ g of DNA per 100-mm dish) using the calcium phosphate DNA precipitation protocol, and expressed for 24–36 h. The soluble fraction containing RD3 was extracted from the harvested cells which were subjected to a hypotonic shock on ice in a buffer solution containing 10 mM Tris-HCl (pH 7.5), 0.1 mM EGTA, 14 mM 2-mercaptoethanol, 60 μ g/mL aprotinin, and 30 μ g/mL leupeptin and further disrupted by 10–15 strokes in a Dounce glass-to-glass homogenizer. The soluble fraction was clarified by centrifugation at 12000g for 30 min, 4 °C, and concentrated in a Millipore Amicon Ultra-4 MW10000 cartridge. A 10- μ L sample was subjected to SDS PAGE on a 15% gel. Expression of RD3 was confirmed by immunoblotting probed with the rabbit polyclonal anti-RD3 antibody 497 and visualized using a SuperSignal chemiluminescence peroxidase substrate kit (Pierce/ThermoFisher) in a FotoDyne Luminous FX imaging system.

For comparative testing of the RD3 mutant expression in cultured HEK293T cells, wild type and the mutant RD3 cDNA constructs generated using the QuikChange Mutagenesis kit (Agilent Technologies) all contained an extension that coded for the C-terminal Rho-1D4 peptide, TETSQVAPA. HEK293T cells were transfected with 10 μ g of the RD3 plasmid DNA per a 100-mm culture dish using the calcium phosphate precipitation protocol. Forty-eight hours after transfection, the cells from two plates were solubilized in 0.8 mL of 1% Triton X-100 in PBS containing Complete Protease Inhibitor cocktail (Roche) for 15 min at 4 °C, and the insoluble material was removed by centrifugation. Each sample was resolved on a 10% SDS polyacrylamide gel and analyzed on immunoblots sequentially probed with the Rho-1D4 mouse monoclonal antibody and secondary goat antimouse antibody conjugated to IR dye 680 for imaging on a LI-COR Odyssey infrared imaging system.

GCAP1 and GCAP2 Expression. GCAP1 and GCAP2 were expressed from pET11d vector in a BLR(DE3) *E. coli* strain harboring yeast *N*-myristoyl transferase as previously described.^{19,20}

RetGC Activity. RetGC cyclase activity in ROS fraction isolated from mouse retinas using Optiprep density gradient centrifugation as described in ref 21 was assayed in the dark using infrared viewers as described previously.^{14,22,23} The activity of the recombinant human RetGC1 expressed from pRCCMV vector in HEK293 cells transfected by calcium phosphate precipitation method was assayed under normal illumination as described previously.²² The assay mixture (25 μ L) contained (unless indicated otherwise) 30 mM MOPS–KOH (pH 7.2), 60 mM KCl, 4 mM NaCl, 1 mM DTT, 2 mM Ca^{2+} /EGTA buffer, 1 mM free Mg^{2+} , 0.3 mM ATP, 4 mM cGMP, 1 mM GTP, 10 mM creatine phosphate, 0.5 unit of creatine phosphokinase, 1 μ Ci of [α -³²P]GTP, 0.1 μ Ci of [³-H]cGMP (Perkin-Elmer), PDE6 inhibitors zaprinast and dipyrindamole, and variable concentrations of GCAPs, RD3, and membranes containing RetGC1. The reaction continued at a linear time-course for 12 min at 30 °C (40 min for the recombinant RetGC1) and subsequently heat-inactivated at 95 °C for 2 min. The [³²P]cGMP was quantified using a thin-layer chromatography on polyethylenimine cellulose plates. Statistical analysis of the data, where applicable, was performed using KaleidaGraph (Synergy Software).

Ca^{2+} Buffers. Free Ca^{2+} and Mg^{2+} concentrations in the RetGC assay were maintained using a series of Ca^{2+} /EGTA mixtures prepared using the Tsien and Pozzan method,²⁴ and their free Ca^{2+} and Mg^{2+} concentrations were calculated using the Marks and Maxfield algorithm in Bound and Determined and MaxChelator software which corrects for the effects of other components of the assay mixture. The calculated Ca^{2+} /EGTA buffering was verified by fluorescent indicator dyes as previously described in detail.^{20,25}

Co-immunoprecipitation (co-IP). For the co-IP studies, the HEK293T cell cultures were transfected with 10 μ g of the RD3 plasmid and 10 μ g of human RetGC1 plasmid DNA per a 100-mm diameter culture dish. Twenty-four hours after the transfection, the cells were solubilized in 800 μ L 1% Triton X-100 in PBS containing Complete Protease Inhibitor cocktail (Roche). After removal of the insoluble material by centrifugation, the supernatant was applied to an Rd3-9D12-Sepharose immunoaffinity column. After 1 h the column was washed several times with 0.1% Triton X-100 in PBS and the




Figure 1. RD3 inhibits RetGC activity at submicromolar concentrations. (A) Wild-type mouse ROS were titrated with human recombinant RD3 in the presence of 1 mM free Mg^{2+} and 2 mM EGTA. Inset: Coomassie Brilliant Blue R250-stained gel of human RD3 isolated from *E. coli* (left lane) and molecular weight standards (right lane). (B) Human recombinant RetGC1 activated by 1.5 μM GCAP1 in the presence of 1 mM free Mg^{2+} and 2 mM EGTA was assayed at different concentrations of the human recombinant RD3 expressed and purified from *E. coli*. (C) Human recombinant RetGC1 was activated by 1.5 μM bovine GCAP1 in the presence of 1 mM free Mg^{2+} and 2 mM EGTA and titrated with protein extracts from HEK293 cells either expressing or not expressing RD3. To exclude the possibility of a nonspecific effect of different total protein concentrations, the total amount of protein was equilibrated with a control protein extract in every assay mixture. Inset: immunoblotting of HEK293 cell extracts probed with anti-RD3 antibody 497 (left lane, nontransfected cells; right lane, RD3 plasmid-transfected cells) (notice that there is no endogenous RD3 expression in the nontransfected cells). The data in A–D are fitted by the equation $a = (a_{max} - a_{min}) / (1 + [RD3] / IC_{50}) + a_{min}$, where a_{max} and a_{min} are the maximal and minimal activity of guanylyl cyclase in the experiment, respectively, and the IC_{50} is the concentration of RD3 producing 50% inhibition. (D–F) The effect of RD3 on guanylyl cyclase catalytic activity in ROS fractions measured in the absence of GCAPs. The RetGC activity in ROS fraction from GCAP1,2 $^{-/-}$ mouse retinas²⁴ was titrated with the purified *E. coli*-expressed RD3 in the presence of saturating 10 mM Mg^{2+} and 2 mM EGTA; GTP concentration in D was 1 mM and in E–F varied as indicated. (E) Michaelis plot of the nonstimulated RetGC catalytic activity in the absence (●) or in the presence of 30 nM (■) or 60 nM (▲) purified recombinant RD3, representative from three similar independent experiments. (F) Lineweaver–Burke plot for data from panel E illustrates a respective ~2.5-fold and 5-fold suppression of the V_{max} by 30 nM and 60 nM RD3 without a major effect on K_{mGTP} . The K_i calculated using the equation for a noncompetitive inhibition, $V_i = V_{max}(1 + [RD3] / K_i)^{-1}$, from three independent experiments was 19 nM \pm 7 SD. The activity in assays containing retinal membranes is presented per rhodopsin content, in membranes expressing recombinant RetGC it is normalized by the maximal activity for each series of the membrane preparations.

fraction containing bound proteins was eluted with 80 μL of 3% SDS.

RESULTS

Inhibition of the Basal and the GCAP-Stimulated RetGC Activity by RD3. Recombinant RD3 strongly inhibits the activity of the native RetGC1/GCAP1 complex in ROS membranes and the recombinant RetGC1 expressed in HEK293 membranes and reconstituted with purified GCAP1 (Figure 1A–C). The suppression of the RetGC activity was observed by RD3 expressed and purified from *E. coli* (Figure 1A,B) and RD3 present in a soluble extract from the HEK293 cells expressing recombinant protein (Figure 1C). The experiment presented in Figure 1D also demonstrates that RD3 suppresses the catalytic activity of RetGC in the absence of GCAPs. Two isoforms of GCAPs, GCAP1 and GCAP2, are the products of neighboring genes, which are both eliminated by a single knockout construct in GCAP1,2 $^{-/-}$ mice.²⁶ GCAP1 and GCAP 2 are the only two GCAP isoforms present in the mouse genome. Therefore, RetGC in the GCAP1,2 $^{-/-}$ remains

insensitive to Ca^{2+} ,^{13,26} and its basal cyclase activity is much lower than the GCAP-stimulated RetGC,²¹ even though the level of RetGC expression in the double knockout retinas remains normal.^{21,26} Since RD3 inhibits RetGC activity in ROS membranes from the GCAP1,2 $^{-/-}$ retinas (Figure 1D), it must directly inhibit the catalytic activity of the cyclase per se.

Classical inhibitory analysis of Michaelis kinetics is presented in Figure 1E,F for basal RetGC activity. There is no significant difference between the K_{mGTP} values (mean \pm SD from three independent measurements) in the absence (1.31 ± 0.17 mM) or in the presence of 30 nM (1.57 ± 0.25 mM) and 60 nM (0.94 ± 0.41 mM) RD3, whereas the V_{max} was reduced nearly 2.5-fold and 5-fold, from 3.5 ± 0.7 to 1.4 ± 0.4 and 0.7 ± 0.4 nmol min $^{-1}$ mg Rh $^{-1}$, respectively ($P < 0.05$). This argues that the inhibition of the basal catalytic activity of RetGC by RD3 in the absence of GCAPs is mostly noncompetitive ($K_i \sim 20$ nM). On this basis, it is unlikely that RD3 directly binds to the catalytic pocket of the enzyme and affects substrate binding. Since the RetGC is active only as a homodimer,^{7,27–29} one possible mechanism for the RD3 affecting RetGC catalytic




Figure 2. RD3 inhibits RetGC activation through competition with GCAPs. (A) The guanylyl cyclase activity in HEK293 homogenates containing RetGC1 expressed alone (●) or coexpressed with human RD3 (▲) was assayed in the presence of added recombinant GCAP1. The data were fitted assuming Michaelis hyperbolic function, $a = a_{\max}[\text{GCAP}]/(K_{1/2} + [\text{GCAP}])$; equalized by RetGC1 content in both samples. (B) Immunoblotting. The cells expressing RetGC1, either alone or cotransfected with RD3, from panel A were probed with anti-RetGC1 (left) or anti-RD3 polyclonal antibody 497 (right). Nontransfected HEK293 cells (leftmost in each panel) were used as a specificity control. (C–F) Competition of RD3 with GCAP1 and GCAP2 in RetGC assay. (C, D) RetGC1 expressed in HEK293 cells was activated by purified GCAP1 (C) or GCAP2 (D) in the absence (●) or in the presence of 3 nM RD3 (▲) or 9 nM RD3 (■). The data were fitted by Michaelis hyperbolic function. Maximal RetGC1 activity (a_{\max} , mean \pm SD) at 0 nM, 3 nM or 9 nM RD3 was 4.8 ± 0.13 , 4.8 ± 0.11 , and 4.6 ± 0.16 nmol/min/mg protein, respectively, when activated by GCAP1 and 2.1 ± 0.05 , 2.6 ± 0.2 , and 2.7 ± 0.1 nmol/min/mg when activated by GCAP2. The respective concentrations of GCAP producing half-maximal activation ($K_{1/2}$) were 1.1 ± 0.12 , 4.1 ± 0.6 , and $7.5 \pm 0.5 \mu M$ (GCAP1) and 5.9 ± 0.6 , 19 ± 2.6 , and $36 \pm 1.5 \mu M$ (GCAP2). (E, F) Double-reciprocal plots related to panels C and D, respectively.

activity is by altering the conformation of the cyclase dimer. This possible mechanism requires further study.

In addition to the basal activity, RD3 also inhibits the activity of RetGC reconstituted with GCAP as well as the RetGC activity in ROS fraction isolated from wild type mice (Figure 1A). This suggests that the GCAP-stimulated activity of RetGC can also be affected by RD3, perhaps through its competition with GCAPs. Indeed, the data presented in Figure 2 directly support the possibility that RD3 strongly competes with GCAP1 for the recombinant human RetGC1 when both regulator proteins are present in the assay. When RetGC1 is coexpressed with RD3, the concentration-dependence of its activation by GCAP measured in HEK293 cell homogenates shifts to a much higher range compared to the RetGC1 expressed in the absence of RD3 (Figure 2A). Likewise, the competition between RD3 and GCAPs for the cyclase is also evident from the experiments when RetGC1 expressed in

HEK293 cells is activated by purified GCAP1 (Figure 2C) or GCAP2 (Figure 2D) in the presence of different concentrations of purified recombinant RD3. While the maximal level of the cyclase activity stimulated by GCAPs at saturation is only slightly affected by RD3, the concentration-dependence of the activation by each GCAP shifts toward much higher EC_{50} values. The latter can only be interpreted as a competitive blocking of the GCAP-dependent stimulation of RetGC by RD3. We emphasize that in these experiments RD3 competes with the GCAPs regardless of what expression system is used to produce the RD3 protein, i.e., HEK 293 cells (Figure 2A) or *E. coli* (Figure 2C,D). Therefore, the effect we observe cannot be attributed to a nonspecific artifact(s) of a particular RD3 expression system.

RD3 Affects the Activation of RetGC Rather than Ca²⁺ Sensitivity of the GCAP-Dependent Regulation. The effect of RD3 on Ca²⁺ and Mg²⁺ dependence of RetGC




Figure 3. RetGC activation by GCAP suppressed by RD3 retains near normal sensitivity to Ca^{2+} and Mg^{2+} . (A–C) ROS fraction isolated from C57B6 mouse retinas was assayed for RetGC activity (open symbols) at various free Ca^{2+} concentrations and either 1 mM (A) or 6 mM (B) free Mg^{2+} concentrations in the presence or in the absence of 0.1 μM or 0.5 μM recombinant RD3 as indicated; in panel C, the activities normalized to the maximal RetGC activity in each assay series are shown as a function of free Ca^{2+} concentrations. (D–F) Membrane fraction from HEK 293 cells expressing human RetGC1 (filled symbols) was reconstituted with 5 μM recombinant GCAP1 in the absence or in the presence of 10 nM and 30 nM RD3 as indicated and assayed as described in the Experimental Procedures.

regulation by GCAPs was tested using two different preparations, the native ROS fraction obtained from dark-adapted mouse retinas by density gradient centrifugation (Figure 3A–C) and a membrane fraction from the HEK293 cells expressing RetGC1 and reconstituted with the recombinant GCAP1 (Figure 3D–F). When RetGC activity partially suppressed by RD3 is measured as a function of Ca^{2+} concentration, its sensitivity to Ca^{2+} remains similar to that in the absence of RD3. The sensitivity of cyclase activity to Mg^{2+} is directly relevant to its sensitivity to Ca^{2+} . This is because Mg^{2+} is required for maintaining the activator conformation of GCAP1 in the light^{12,20,26,30} and competes with Ca^{2+} binding in the EF hand domains of GCAP,²⁰ thus shifting the Ca^{2+} sensitivity of the cyclase regulation by GCAPs.²³ This shift appears in the presence of RD3 when the free Mg^{2+} level in the assay is increased from a near-physiological 1 mM to 6 mM and is similar to the control samples assayed in the absence of RD3 (Figure 3C,F).

Mutations in Human RD3 Affect Its Expression in HEK293 Cells and RetGC Activity in Vitro. A nonsense

mutation in *rd3* mice that terminates RD3 synthesis after Glu¹⁰⁶ and a mutation in LCA12 patients that alters a splicing signal and also generates a stop codon after Arg⁹⁹ both delete the C-terminal half of the protein.¹⁵ In addition to the deletion mutation directly linked to LCA12, a number of missense mutations were detected in patients with other retinal disorders,¹⁵ although their roles in development of the disease remain to be established. We have tested the effect of several such mutations on human RD3 expression in HEK293 cells and their ability to associate with RetGC in co-IP experiments (Figure 4). The F100ter mutation mimicking the LCA-linked truncation of RD3 as well as the G57V mutant failed to accumulate in transfected HEK293T cells at detectable levels (Figure 4A,B). This is in agreement with the very low expression of the related mouse RD3 truncated protein observed in transfected COS-1 culture cells.¹⁵ Other tested RD3 mutants were expressed in HEK293 cells and coimmunoprecipitated with RetGC1, indicating that they retain their general ability to interact with the cyclase (Figure 4C). We then evaluated the inhibitory activity of different mutants




Figure 4. Mutations found in human patients with retinal diseases¹⁵ affect RD3 expression in HEK293 cells. (A) Expression of RD3 mutants in HEK293 cells. Human wild type and mutant RD3 tagged with the 1D4 peptide at the C-terminus were detected by immunoblotting as described in the Experimental Procedures. Actin immunostaining (upper panel) was used as a protein load control. (B) The average band intensities of the mutant RD3 proteins relative to WT RD3 band were determined from 3 independent experiments described in panel A (mean \pm SD). (C) Co-IP experiment. RetGC1 and RD3 were coexpressed as described in Experimental procedures and co-IP using Sepharose-coupled anti-RD3 9D12 antibody. The samples applied to the column (marked "I" for "input") and the eluted fractions (marked "B" for "bound") were analyzed by immunoblotting probed with anti-RetGC1 (GC-8A5; upper panel) and anti-RD3 (RD3-9D12; lower panel) antibodies. The G57V RD3 mutant that fails to express in HEK293 cells was used as a control to confirm that RetGC1 does not bind nonspecifically to the 9D12 antibody-coupled Sepharose. The lower band on the blot is specific for RD3-transfected cells and is most likely a partially proteolyzed RD3 polypeptide.

expressed and purified from *E. coli* (Figure 5). Although the expression of the F100ter and G57V mutants was undetectable in HEK293 cells, their expression in *E. coli* is fairly robust. All RD3 mutants expressed in bacteria were purified in quantities sufficient for the *in vitro* RetGC assay (Figure 5A). When reconstituted with the recombinant RetGC1 activated by GCAP1, the purified F100ter RD3 did not compete with the GCAP and was unable to suppress the RetGC activity (Figure 5B). All other mutants are able to inhibit the activity of the GCAP1/RetGC1 complex, but to varying degrees. The W6R/E23D and the G35R mutants both inhibited RetGC1 activity in a manner nearly indistinguishable from the wild type protein, whereas the R68W, K130M, and G57V mutants were significantly less effective in suppressing cyclase activation by GCAP. This is particularly true for the G57V mutant, which has an IC_{50} value that is more than 10-fold higher than wild type RD3 (Figure 5B).




Figure 5. Effect of mutations in RD3 found in human patients with retinal diseases on RetGC activity *in vitro*. (A) Wild-type and mutant RD3¹⁵ expressed in *E. coli* were stained by Coomassie Brilliant Blue R-250 after SDS PAGE, 15% gel. For the use in subsequent RetGC assays, RD3 concentration in each case was that of the main band. (B) Inhibition of RetGC1 activity by the *E. coli*-expressed RD3 mutants. RetGC activity in independent assays was normalized per activity measured in the absence of RD3. Recombinant RetGC1 expressed in HEK293 cells was reconstituted with 1.5 μ M GCAP1 at indicated concentrations of RD3 (○) or its mutants: W6R/E23D (●), G35R (△), G57V (□), R68W (■), K130M (▲), and F100ter (◆). The IC_{50} values (mean \pm SE, *n*) for the inhibition were 4.6 \pm 1.3 nM, 5 (WT); 6.4 \pm 2 nM, 3 (W6R/E23D); 7.9 \pm 1 nM, 3 (G35R); 68 \pm 13 nM, 3 (G57V); 17.3 \pm 3.8 nM, 3 (R68W); and 14.3 \pm 2.4 nM, 3 (K130M); the IC_{50} for the F100ter was \gg 10 μ M. The difference in the IC_{50} values from the wild type was significant (*p* in unpaired Student *t* test assuming equal variance) for the G57V (<0.001), F100ter (<0.0001), R68W (0.012), and K130M (0.014) mutants.

DISCUSSION

It was recently found that a retina-specific protein RD3 linked to LCA12 blindness in humans and rapid retinal degeneration in *rd3* mouse line¹⁵ was expressed in rods and cones and was capable of associating with the photoreceptor-specific guanylyl cyclase RetGC.¹⁶ Our present data argue that this association has a strong functional consequence and that RD3 is a novel potent inhibitor of RetGC, capable of suppressing both the basal and GCAP-stimulated activity of the cyclase. A diagram summarizing the RD3 effects on the cyclase activity is presented in Figure 6. In the absence of GCAPs, RD3 suppresses the basal catalytic activity of the RetGC acting as a noncompetitive inhibitor (see also Figure 1E,F). It also counteracts the GCAP-dependent activation of the cyclase acting as a negative allosteric modulator (see also Figure 2). It




Figure 6. Diagram depicting the inhibition of RetGC by RD3. A RetGC homodimer forms an active site by combining the Mg^{2+} -coordinating center from one subunit with the GTP binding center from another subunit.^{27–29} There are two symmetrical active sites in the dimer, but only one such site is depicted here for simplicity. The low RetGC basal catalytic activity (left) becomes accelerated up to 20–100 fold^{21,31} when GCAPs bind the cytoplasmic portion of RetGC (middle). RD3 binding to the cyclase (right) inhibits the RetGC catalytic activity but does not prevent binding of the Mg^{2+} -GTP substrate in the active site; at the same time, RD3 acts as a negative high-affinity modulator of the RetGC/GCAP complex by displacing GCAP; this prevents dynamic activation by GCAPs via Ca^{2+} -feedback mechanism. The GCAP-stimulated activity is much higher than the basal RetGC activity;^{21,31} therefore, the strongest effect of RD3 as a negative allosteric modulator of the cyclase results from its competition with GCAP. Other explanations are provided in the Discussion.

should be emphasized that stimulation by GCAP increases the V_{max} for RetGC in mouse ROS up to 20-fold²¹ and up to 100-fold in HEK293 membranes (ref 31 and Figure 2). Therefore, compared to the noncompetitive inhibition of the basal RetGC activity in GCAPs^{−/−} ROS (Figure 1E,F), a direct competition with the stimulating effect of GCAPs clearly dominates the overall effect of RD3 on the cyclase regulation in a RetGC/GCAP complex (Figure 2). Indeed, once RD3 partially or completely displaces GCAP from the complex with the cyclase, the activity of the recombinant RetGC would fall as much as 100-fold since the cyclase is no longer activated by GCAP. The additional suppression of the basal cyclase activity by RD3 would also contribute to the overall inhibition, but relatively less than the displacement of GCAP from the cyclase. As a result, the noncompetitive inhibition of the catalytic activity evident in Figure 1D–F becomes masked in experiments presented in Figure 2 by the more robust allosteric modulation, i.e., RD3 competition with GCAPs. These results would be consistent with a model in which RD3 does not strongly inhibit the catalytic activity of the RetGC in complex with GCAP until after GCAP is displaced from the complex (Figure 6).

The exact binding sites for RD3 in RetGC1 have not yet been identified. Deletion of a 48-amino acid fragment from the C-terminus of RetGC hampered RD3 binding in coimmunoprecipitation experiments,¹⁶ but at this point it is unclear whether or not this is the only critically important region for RD3 binding and whether or not the binding sites for GCAP and RD3 in RetGC overlap. Multiple fragments of RetGC primary structure have been proposed by different groups to participate in binding of GCAP, including a portion of the cyclase catalytic domain.^{32–36} Yet, the fragments of the catalytic domain that may participate in GCAP binding (or at least be in a close proximity to the GCAP binding site, refs 33–36) are thus far located upstream from the C-terminal fragment whose

deletion affects coimmunoprecipitation with RD3. However, it is also important to note that although a competition for an overlapping epitope(s) in RetGC appears to be the simplest and the most likely possibility, RD3 and GCAP do not necessarily have to bind to exactly the same peptide epitopes; the effect of their mutual exclusion can even be potentially achieved through changing RetGC dimer conformation upon binding of the two allosteric protein regulators in different places.

At the same time, we see no evidence that RD3 is able to critically affect Ca^{2+} sensitivity of RetGC regulation by GCAPs either in the native ROS fraction or as a recombinant cyclase (Figure 3). This is generally consistent with the model presented in Figure 6, where RD3 mainly competes with GCAPs rather than alters the activity of the GCAP/RetGC complex. Two different physiological forms of a metal-bound GCAP1 have been characterized—a Mg^{2+} -bound activator form to which GCAP converts in the light and a Ca^{2+} -bound inhibitor form which suppresses the activity of RetGC in the dark when the free Ca^{2+} rises (reviewed in ref 12). Competition between Ca^{2+} and Mg^{2+} for EF-hands in GCAP1 and GCAP2 therefore strongly affects Ca^{2+} sensitivity of RetGC regulation by GCAPs resulting in a prominent right-shift of the Ca^{2+} sensitivity curve as Mg^{2+} concentrations increase.^{23,31} This same shift remains in the presence of RD3 (Figure 3). These results argue that RD3 does not significantly affect metal binding to GCAP and is unlikely to alter the dynamic regulation of the cyclase in response to light induced changes in Ca^{2+} concentrations in photoreceptor cells.^{12,13}

What then is the possible role for the inhibitory activity of RD3 on RetGC activity in a photoreceptor cell? Evidently, the biochemical mechanism of RetGC inhibition by RD3 itself does not require the environment of the photoreceptor cell (Figures 1 and 2), yet it should be specific for the photoreceptors since both RD3 and RetGC¹⁶ are both highly expressed in rod and cone cells and largely absent in other cells of the body.¹⁵ RD3 displays a much stronger apparent affinity than GCAP for RetGC since it inhibits the cyclase in a nanomolar range, whereas GCAPs activate RetGC at micromolar concentrations (Figure 2). Furthermore, unlike GCAP/RetGC complex, the RD3/RetGC complex is sufficiently stable in detergent to sustain the co-IP procedures (Figure 4 and ref 16). The precise binding stoichiometry of RD3 and RetGC1 remains unknown at this point, but the concentration of RD3 could hamper the GCAP-dependent stimulation based on the following estimate. Since nearly half of RetGC monomers may be coimmunoprecipitated with RD3 from the detergent-solubilized retinal membranes as an irreversible complex¹⁶ and the concentration of RetGC dimer in mammalian ROS lies in a micromolar range^{36,37} reaching $\sim 3.6 \mu M$ in mouse ROS (or $7.2 \mu M$ RetGC monomer),²¹ the total concentration of RD3 could approach $\sim 3.6 \mu M$ (assuming for simplicity that one RD3 binds per RetGC monomer). The total concentration of GCAPs can reach as high as $\sim 17 \mu M$ in the mouse ROS cytosol, nearly five times higher than RD3 (calculated on the basis of refs 21 and 36). However, RD3 exerts its allosteric modulation of the cyclase at concentrations nearly 100 times lower than GCAPs (Figures 2, 3, and 5). Therefore, even by a conservative estimate, RD3 has the potential for suppressing a large fraction of the cyclase activity. Even though we do not know at this time the fine mechanism of RD3/RetGC complex formation and quantification of RD3 in photoreceptors needs further investigation, we would not anticipate that RD3-dependent

inhibition of the cyclase activity or its high-affinity competition with GCAP contributes to the dynamic regulation of RetGC by Ca^{2+} and GCAPs. Furthermore, based on our data it would also very likely require additional mechanisms to reduce or prevent the RD3 inhibitory effects on RetGC at the cellular level to allow the dynamic regulation of RetGC in the outer segment.

A possible functional role of RD3 in photoreceptor physiology is to suppress the basal and GCAP activated RetGC activity in the inner segment during its transport to the outer segment. Another possibility is that RD3 itself can actively participate in the process of the transport¹⁶ and/or assembly of the cyclase complex into the proper structure of the outer segment, during which its ability to strongly compete with GCAPs can prevent GCAP from prematurely interfering with this photoreceptor-specific process.

An apparent involvement of mutated RD3 in congenital retinal diseases corroborates its possible physiological role in photoreceptors. The 195-amino acid product of *Rd3* (*C1of36*) gene originally identified by Friedman and co-workers¹⁵ has been linked to a recessive retinal degeneration in the *rd3* mice^{38,39} and LCA12 patients.¹⁵ Although RD3 effectively competes with GCAPs for the target enzyme, it fails to do so when truncated by the F100ter mutation linked to LCA12 (Figure 5). A similar shortening of RD3 in *rd3* mice¹⁵ also leads to the loss of interaction between RD3 and RetGC and dramatically suppresses the levels of RetGC1 and RetGC2 expression in mouse photoreceptors *in vivo*.¹⁶ A variety of other mutations in the *RD3* gene were reported in patients with retinal disorders.¹⁵ While the importance of those mutations for the development of congenital diseases remains to be determined, we find that some of these mutations can affect RD3 expression in human cultured cells and reduce the inhibitor activity of RetGC (Figures 4 and 5).

One could expect that a possible consequence of mutations hampering RD3 binding to the cyclase would be to prevent RetGC from reaching the outer segment resulting in its accumulation in the inner segment. Evidently, this is not the case in *rd3* mice.¹⁶ When RD3 is truncated similar to the F100ter mutant found in LCA12 patients and is unable to bind RetGC, the photoreceptors suppress production of RetGC rather than allow it to accumulate in the inner segment. It is likely that the photoreceptors possess a safety check capability that prevents the accumulation of RetGC in the absence of RD3. It is therefore tempting to speculate that failure of the mutant RD3 to properly bind the cyclase could trigger the mislocalization and rapid destruction of the cyclase as a protection against unsuitable activation of cGMP synthesis. Depletion of RetGC in ROS would inevitably suppress cGMP production resulting in a drop in intracellular Ca^{2+} to a level which could cause photoreceptor degeneration.¹⁶ The consequence of the altered activity of several RD3 mutants shown in this study requires further in-depth analysis of the mutants *in vivo*.

AUTHOR INFORMATION

Corresponding Author

*Phone: 215-780-1468. Fax: 215-780-1464. E-mail: adizhoor@salus.edu.

Funding

This work was supported by NIH grants EY11522 (AMD) and EY02422 (RSM) and Formula Grant from Pennsylvania Department of Health (AMD).

ACKNOWLEDGMENTS

A.M.D. is a Martin and Florence Hafter Chair Professor of Pharmacology, and R.S.M. is a Canada Research Chair in Vision and Macular Degeneration.

ABBREVIATIONS

BSA, bovine serum albumin; EGTA, ethylene glycol-bis(2-aminoethylether)-*N,N,N',N'*-tetraacetic acid; DTT, dithiothreitol; GCAP, guanylyl cyclase activating protein; co-IP, coimmunoprecipitation; IPTG, isopropyl- β -D-thiogalactoside; KHD, kinase homology domain; LCA, Leber congenital amaurosis; LDAO, lauryldimethylamine oxide; MOPS, 3-(*N*-morpholino) propanesulfonic acid; RD3, retinal degeneration 3 protein; RetGC, retinal guanylyl cyclase; ROS, outer segment; SDS, sodium dodecyl sulfate

REFERENCES

- (1) Fu, Y., and Yau, K. W. (2007) Phototransduction in mouse rods and cones. *Pflügers Arch.* 454, 805–819.
- (2) Lowe, D. G., Dizhoor, A. M., Liu, K., Gu, Q., Spencer, M., Laura, R., Lu, L., and Hurley, J. B. (1995) Cloning and expression of a second photoreceptor-specific membrane retina guanylyl cyclase (RetGC), RetGC-2. *Proc. Natl. Acad. Sci. U.S.A.* 92, 5535–5539.
- (3) Yang, R. B., Foster, D. C., Garbers, D. L., and Fulle, H. J. (1995) Two membrane forms of guanylyl cyclase found in the eye. *Proc. Natl. Acad. Sci. U.S.A.* 92, 602–606.
- (4) Pugh, E. N. Jr., Duda, T., Sitaramayya, A., and Sharma, R. K. (1997) Photoreceptor guanylate cyclases: a review. *Biosci. Rep.* 17, 429–473.
- (5) Perrault, I., Rozet, J. M., Gerber, S., Ghazi, I., Ducroq, D., Souied, E., Leowski, C., Bonnemaïson, M., Dufier, J. L., Munnich, A., and Kaplan, J. (2000) Spectrum of retGC1 mutations in Leber's congenital amaurosis. *Eur. J. Hum. Genet.* 8, 578–582.
- (6) Perrault, I., Rozet, J. M., Gerber, S., Kelsell, R. E., Souied, E., Cabot, A., Hunt, D. M., Munnich, A., and Kaplan, J. (1998) A retGC-1 mutation in autosomal dominant cone-rod dystrophy. *Am. J. Hum. Genet.* 63, 651–654.
- (7) Ramamurthy, V., Tucker, C., Wilkie, S. E., Daggett, V., Hunt, D. M., and Hurley, J. B. (2001) Interactions within the coiled-coil domain of RetGC-1 guanylyl cyclase are optimized for regulation rather than for high affinity. *J. Biol. Chem.* 276, 26218–26229.
- (8) Gorczyca, W. A., Gray-Keller, M. P., Detwiler, P. B., and Palczewski, K. (1994) Purification and physiological evaluation of a guanylate cyclase activating protein from retinal rods. *Proc. Natl. Acad. Sci. U.S.A.* 91, 4014–4018.
- (9) Palczewski, K., Subbaraya, I., Gorczyca, W. A., Helekar, B. S., Ruiz, C. C., Ohguro, H., Huang, J., Zhao, X., Crabb, J. W., Johnson, R. S., Walsh, K. A., Gray-Keller, M. P., Detwiler, P. B., and Baehr, W. (1994) Molecular cloning and characterization of retinal photoreceptor guanylyl cyclase-activating protein. *Neuron* 13, 395–404.
- (10) Dizhoor, A. M., Lowe, D. G., Olshevskaya, E. V., Laura, R. P., and Hurley, J. B. (1994) The human photoreceptor membrane guanylyl cyclase, RetGC, is present in outer segments and is regulated by calcium and a soluble activator. *Neuron* 12, 1345–1352.
- (11) Dizhoor, A. M., Olshevskaya, E. V., Henzel, W. J., Wong, S. C., Stults, J. T., Ankoudinova, I., and Hurley, J. B. (1995) Cloning, sequencing, and expression of a 24-kDa Ca^{2+} -binding protein activating photoreceptor guanylyl cyclase. *J. Biol. Chem.* 270, 25200–25206.
- (12) Dizhoor, A. M., Olshevskaya, E. V., and Peshenko, I. V. (2010) $\text{Mg}^{2+}/\text{Ca}^{2+}$ cation binding cycle of guanylyl cyclase activating proteins (GCAPs): role in regulation of photoreceptor guanylyl cyclase. *Mol. Cell. Biochem.* 334, 117–124.
- (13) Burns, M. E., Mendez, A., Chen, J., and Baylor, D. A. (2002) Dynamics of cyclic GMP synthesis in retinal rods. *Neuron* 36, 81–91.
- (14) Makino, C. L., Peshenko, I. V., Wen, X. H., Olshevskaya, E. V., Barrett, R., and Dizhoor, A. M. (2008) A role for GCAP2 in regulating

the photoresponse. Guanylyl cyclase activation and rod electrophysiology in GUCA1B knock-out mice. *J. Biol. Chem.* 283, 29135–43.

(15) Friedman, J. S., Chang, B., Kannabiran, C., Chakarova, C., Singh, H. P., Jalali, S., Hawes, N. L., Branham, K., Othman, M., Filippova, E., Thompson, D. A., Webster, A. R., Andreasson, S., Jacobson, S. G., Bhattacharya, S. S., Heckenlively, J. R., and Swaroop, A. (2006) Premature truncation of a novel protein, RD3, exhibiting subnuclear localization is associated with retinal degeneration. *Am. J. Hum. Genet.* 79, 1059–70.

(16) Azadi, S., Molday, L. L., and Molday, R. S. (2011) RD3, the protein associated with Leber congenital amaurosis type 12, is required for guanylate cyclase trafficking in photoreceptor cells. *Proc. Natl. Acad. Sci. U.S.A.* 107, 21158–21163.

(17) Hodges, R. S., Heaton, R. J., Parker, J. M., Molday, L., and Molday, R. S. (1988) Antigen-antibody interaction. Synthetic peptides define linear antigenic determinants recognized by monoclonal antibodies directed to the cytoplasmic carboxyl terminus of rhodopsin. *J. Biol. Chem.* 263, 11768–11775.

(18) Laura, R. P., Dizhoor, A. M., and Hurley, J. B. (1996) The membrane guanylyl cyclase, retinal guanylyl cyclase-1, is activated through its intracellular domain. *J. Biol. Chem.* 271, 11646–11651.

(19) Olshevskaya, E. V., Hughes, R. E., Hurley, J. B., and Dizhoor, A. M. (1997) Calcium binding, but not a calcium-myristoyl switch, controls the ability of guanylyl cyclase-activating protein GCAP-2 to regulate photoreceptor guanylyl cyclase. *J. Biol. Chem.* 272, 14327–14333.

(20) Peshenko, I. V., and Dizhoor, A. M. (2006) Ca^{2+} and Mg^{2+} binding properties of GCAP-1. Evidence that Mg^{2+} -bound form is the physiological activator of photoreceptor guanylyl cyclase. *J. Biol. Chem.* 281, 23830–23841.

(21) Peshenko, I. V., Olshevskaya, E. V., Savchenko, A. B., Karan, S., Palczewski, K., Baehr, W., and Dizhoor, A. M. (2011) Enzymatic properties and regulation of the native isozymes of retinal membrane guanylyl cyclase (RetGC) from mouse photoreceptors. *Biochemistry* 50, 5590–600.

(22) Peshenko, I. V., Olshevskaya, E. V., and Dizhoor, A. M. (2008) Binding of guanylyl cyclase activating protein 1 (GCAP1) to retinal guanylyl cyclase (RetGC1). The role of individual EF-hands. *J. Biol. Chem.* 283, 21747–21757.

(23) Peshenko, I. V., and Dizhoor, A. M. (2004) Guanylyl cyclase-activating proteins (GCAPs) are $\text{Ca}^{2+}/\text{Mg}^{2+}$ sensors: implications for photoreceptor guanylyl cyclase (RetGC) regulation in mammalian photoreceptors. *J. Biol. Chem.* 279, 16903–16906.

(24) Tsien, R., and Pozzan, T. (1989) Measurement of cytosolic free Ca^{2+} with quin2. *Methods Enzymol.* 172, 230–62.

(25) Peshenko, I. V., and Dizhoor, A. M. (2007) Activation and inhibition of photoreceptor guanylyl cyclase by guanylyl cyclase activating protein 1 (GCAP-1): the functional role of $\text{Mg}^{2+}/\text{Ca}^{2+}$ exchange in EF-hand domains. *J. Biol. Chem.* 282, 21645–21652.

(26) Mendez, A., Burns, M. E., Sokal, I., Dizhoor, A. M., Baehr, W., Palczewski, K., Baylor, D. A., and Chen, J. (2001) Role of guanylate cyclase-activating proteins (GCAPs) in setting the flash sensitivity of rod photoreceptors. *Proc. Natl. Acad. Sci. U.S.A.* 98, 9948–9953.

(27) Liu, Y., Ruoho, A. E., Rao, V. D., and Hurley, J. H. (1997) Catalytic mechanism of the adenylyl and guanylyl cyclases: modeling and mutational analysis. *Proc. Natl. Acad. Sci. U.S.A.* 94, 13414–13419.

(28) Tucker, C. L., Hurley, J. H., Miller, T. R., and Hurley, J. B. (1998) Two amino acid substitutions convert a guanylyl cyclase, RetGC-1, into an adenylyl cyclase. *Proc. Natl. Acad. Sci. U.S.A.* 95, 5993–5997.

(29) Ramamurthy, V., Tucker, C., Wilkie, S. E., Daggett, V., Hunt, D. M., and Hurley, J. B. (2001) Interactions within the coiled-coil domain of RetGC-1 guanylyl cyclase are optimized for regulation rather than for high affinity. *J. Biol. Chem.* 276, 26218–26229.

(30) Lim, S., Peshenko, I., Dizhoor, A., and Ames, J. B. (2009) Effects of Ca^{2+} , Mg^{2+} , and myristoylation on guanylyl cyclase activating protein 1 structure and stability. *Biochemistry* 48, 850–862. *Biochemistry* 48, 850–862.

(31) Peshenko, I. V., Moiseyev, G. P., Olshevskaya, E. V., and Dizhoor, A. M. (2004) Factors that determine Ca^{2+} sensitivity of photoreceptor guanylyl cyclase. Kinetic analysis of the interaction between the Ca^{2+} -bound and the Ca^{2+} -free guanylyl cyclase activating proteins (GCAPs) and recombinant photoreceptor guanylyl cyclase 1 (RetGC-1). *Biochemistry* 43, 13796–13804.

(32) Lange, C., Duda, T., Beyermann, M., Sharma, R. K., and Koch, K. W. (1999) Regions in vertebrate photoreceptor guanylyl cyclase ROS-GC1 involved in Ca^{2+} -dependent regulation by guanylyl cyclase-activating protein GCAP-1. *FEBS Lett.* 460, 27–31.

(33) Sokal, I., Haeseleer, F., Arendt, A., Adman, E. T., Hargrave, P. A., and Palczewski, K. (2001) Identification of a guanylyl cyclase-activating protein-binding site within the catalytic domain of retinal guanylyl cyclase 1. *Biochemistry* 38, 1387–1393.

(34) Krylov, D. M., and Hurley, J. B. (1999) Identification of proximate regions in a complex of retinal guanylyl cyclase 1 and guanylyl cyclase-activating protein-1 by a novel mass spectrometry-based method. *J. Biol. Chem.* 276, 30648–30654.

(35) Duda, T., Fik-Rymarkiewicz, E., Venkataraman, V., Krishnan, R., Koch, K. W., and Sharma, R. K. (2005) The calcium-sensor guanylate cyclase activating protein type 2 specific site in rod outer segment membrane guanylate cyclase type 1. *Biochemistry* 44, 7336–7345.

(36) Hwang, J. Y., Lange, C., Helten, A., Hoppner-Heitmann, D., Duda, T., Sharma, R. K., and Koch, K. W. (2003) Regulatory modes of rod outer segment membrane guanylate cyclase differ in catalytic efficiency and Ca^{2+} -sensitivity. *Eur. J. Biochem.* 270, 3814–3821.

(37) Helten, A., Saftel, W., and Koch, K. W. (2007) Expression level and activity profile of membrane bound guanylate cyclase type 2 in rod outer segments. *J. Neurochem.* 103, 1439–46.

(38) Chang, B., Heckenlively, J. R., Hawes, N. L., and Roderick, T. H. (1993) New mouse primary retinal degeneration (rd-3). *Genomics* 16, 45–49.

(39) Danciger, M., Ogando, D., Yang, H., Matthes, M. T., Yu, N., Ahern, K., Yasumura, D., Williams, R. W., and Lavail, M. M. (2008) Genetic modifiers of retinal degeneration in the rd3 mouse. *Invest. Ophthalmol. Vis. Sci.* 49, 2863–2869.