


# P3–P3' Residues Flanking Scissile Bonds in Factor VIII Modulate Rates of Substrate Cleavage and Procofactor Activation by Thrombin

Jennifer L. Newell-Caito,<sup>†</sup> Amy E. Griffiths, and Philip J. Fay\*

Department of Biochemistry and Biophysics, University of Rochester School of Medicine, 601 Elmwood Avenue, Rochester, New York 14642, United States

**ABSTRACT:** Thrombin-catalyzed activation of factor VIII (FVIII) occurs through proteolysis at three P1 Arg residues: Arg<sup>372</sup> and Arg<sup>740</sup> in the FVIII heavy chain and Arg<sup>1689</sup> in the FVIII light chain. Cleavage at the latter two sites is relatively fast compared with cleavage at Arg<sup>372</sup>, which appears to be rate-limiting. Examination of the P3–P3' residues flanking each P1 site revealed that those sequences at Arg<sup>740</sup> and Arg<sup>1689</sup> are more optimal for thrombin cleavage than at Arg<sup>372</sup>, suggesting these sequences may impact reaction rates. Recombinant FVIII variants were prepared with mutations swapping scissile bond flanking sequences in the heavy chain individually and in combination with a second swap or with a P1 point mutation. Rates of generation of A1 and A3-C1-C2 subunits were determined by Western blotting and correlated with rates of cleavage at Arg<sup>372</sup> and Arg<sup>1689</sup>, respectively. Rates of thrombin cleavage at Arg<sup>372</sup> were increased ~10- and ~3-fold compared with that of wild-type FVIII when it was replaced with P3–P3' residues flanking Arg<sup>740</sup> and Arg<sup>1689</sup>, respectively, and these values paralleled increased rates of A2 subunit generation and procofactor activation. Positioning of more optimal residues flanking Arg<sup>372</sup> abrogated the need for initial cleavage at Arg<sup>740</sup> to facilitate this step. These results show marked changes in cleavage rates correlate with the extent of cleavage-optimal residues flanking the scissile bond and modulate the mechanism for procofactor activation.



Factor VIII (FVIII) is an inactive procofactor that is essential in blood coagulation as mutations or splicing defects in the FVIII gene lead to hemophilia A. FVIII is synthesized as an ~300 kDa single-chain molecule corresponding to 2332 amino acids in six distinct domains (NH<sub>2</sub>-A1-A2-B-A3-C1-C2-COOH).<sup>1,2</sup> Segments rich in acidic residues border the A domains of FVIII and are designated a1 (residues 337–372) and a2 (residues 711–740), which directly follow the A1 and A2 domains, respectively, and a3 (residues 1649–1689), preceding the A3 domain. FVIII circulates primarily as a heterodimer composed of a heavy chain (A1-a1-A2-a2-B domains) and a light chain (a3-A3-C1-C2 domains) as a result of proteolytic processing at the B–a3 junction.<sup>3</sup>

The activated cofactor, factor VIIIa (FVIIIa), binds to factor IXa (FIXa) on a membrane surface, and this complex activates factor X (FX). FVIIIa increases the catalytic efficiency of FIXa by several orders of magnitude in a primarily *k*<sub>cat</sub>-dependent manner.<sup>4</sup> Conversion of the procofactor to activated FVIIIa occurs through limited proteolysis by either thrombin or FXa<sup>5,6</sup> via cleavage of three P1 residues at Arg<sup>740</sup> (a2–B junction), Arg<sup>372</sup> (a1–A2 junction), and Arg<sup>1689</sup> (a3–A3 junction).<sup>5</sup> Thrombin is considered the physiological activator of FVIII inasmuch as von Willebrand factor inhibits the membrane-dependent procofactor activation catalyzed by FXa.<sup>5,7</sup> An essential step in activation of the procofactor is cleavage at Arg<sup>372</sup>, as this cleavage allows exposure of FIXa-interactive sites in the A2 domain that are otherwise masked.<sup>8</sup> Cleavage at Arg<sup>1689</sup> is also critical as it not only increases cofactor-specific activity but also releases FVIIIa from von Willebrand factor.<sup>9–11</sup> Inactivation of FVIIIa resulting in downregulation of the

intrinsic factor Xase complex<sup>3</sup> occurs through two mechanisms: spontaneous inactivation via dissociation of the weak electrostatic interaction between the A1/A3-C1-C2 dimer and the A2 subunit<sup>12,13</sup> and proteolytic inactivation via cleavage at Arg<sup>336</sup> and Arg<sup>562</sup> by activated protein C<sup>5,12,14</sup> and FXa.<sup>5,15</sup>

Substrate recognition by thrombin occurs through interaction with one or both anion binding exosites (ABE1 and ABE2) followed by active site docking.<sup>16</sup> This docking primarily affects the *V*<sub>max</sub> for substrate activation by thrombin, while it is believed that exosite binding exclusively determines substrate affinity.<sup>17</sup> Examination of the P3–P3' region of thrombin substrates reveals a variety of residues that can be accommodated by thrombin, and it is thought that P1–P3' flanking sites primarily affect substrate docking in a manner independent of binding affinity.<sup>16,18,19</sup>

Previous studies of the proteolytic activation of FVIII by thrombin have indicated that both ABE1 and ABE2 contribute to the activation mechanism.<sup>20</sup> Furthermore, there appears to be a competition between the heavy chain and the light chain for this exosite-dependent binding, with binding to and cleavage of FVIII heavy chain being preferred.<sup>21</sup> Additionally, this mechanism is further complicated by the presence of three P1 cleavage sites that differ in P3–P3' sequences flanking the P1 Arg<sup>372</sup>, Arg<sup>740</sup>, and Arg<sup>1689</sup>, which are represented by <sup>370</sup>QIR<sup>1</sup>SVA<sup>375</sup>, <sup>738</sup>EPR<sup>1</sup>SFS<sup>743</sup>, and <sup>1687</sup>SPR<sup>1</sup>SFQ<sup>1692</sup>, respectively. Residues flanking Arg<sup>372</sup> are considered non-

Received: January 17, 2012

Revised: March 23, 2012

Published: March 28, 2012



optimal for thrombin cleavage, with only two residues being optimal (bold) for cleavage in the P3–P3' sequence, while residues flanking the two other P1 sites are considered near-optimal, with four of six residues being optimal (bold).<sup>22</sup> Inasmuch as the rate-limiting cleavage occurs at Arg<sup>372</sup>,<sup>3,23</sup> there is a possible role for residues flanking the P1 site in the overall reaction mechanism. In this study, the effects of P3–P3' residues on thrombin-catalyzed cleavage and activation of FVIII were investigated using recombinant FVIII variants possessing P3–P3' sequence swaps of residues flanking P1 residues Arg<sup>372</sup>, Arg<sup>740</sup>, and Arg<sup>1689</sup>. Results from this study show marked changes in site-specific cleavage rates that correlate with the degree of cleavage-optimal residues present flanking the scissile bond.

## ■ EXPERIMENTAL PROCEDURES

**Reagents.** Coagulation proteins FIXa, FXa, FX, and human  $\alpha$ -thrombin were obtained from Enzyme Research Laboratories. Chromogenic FXa substrate Pefachrome FXa (CH<sub>3</sub>OCO-D-CHA-Gly-Arg-pNA) (Pentapharm), horseradish peroxidase-labeled streptavidin (Calbiochem), and recombinant non-sulfated hirudin-(55–65) (Calbiochem) were purchased from the indicated suppliers. P. Lollar and J. Healey (Emory University, Atlanta, GA) generously provided the Bluescript FVIII vector (pBS FVIII) and B-domainless FVIII expression construct RENeo FVIII. Monoclonal antibodies C5 (recognizing the A1 region) and 2D2 (recognizing the A3 domain) were gifts from Z. Ruggeri and L. Regan (Bayer), respectively. Monoclonal FVIII antibodies R8B12 (GMA-012) (recognizing the A2 domain)<sup>12</sup> and GMA-8003 (recognizing the C2 domain) were obtained from Green Mountain Antibodies.

**Phospholipid Vesicles.** N-Octyl glucoside was used to prepare phospholipid vesicles containing 20% phosphatidylserine, 40% phosphatidylcholine, and 40% phosphatidylethanolamine (Avanti Polar Lipids).<sup>24</sup>

**FVIII-Deficient Plasma.** FVIII-deficient plasma was prepared by incubating pooled human plasma with 25 mM EDTA for ~18 h at 4 °C followed by addition of 25 mM CaCl<sub>2</sub>.<sup>25</sup> Aliquots were snap-frozen in liquid N<sub>2</sub> and stored at –80 °C.

**Construction, Expression, and Purification of Recombinant Proteins.** The FVIII expression construct FVIIIHSQ-MSAB-NotI-RENeo was restricted using endonucleases *Xho*I and *Not*I to create B-domainless FVIII cDNA and cloned into the pBluescript II K/S vector. The B-domainless FVIII cDNA was then further restricted using endonucleases *Sac*II and *Apa*I to create the Arg<sup>1689</sup> and Arg<sup>740</sup> P3–P3' mutants and endonucleases *Spe*I and *Sac*II to create the Arg<sup>372</sup> P3–P3' mutants. All P3–P3' mutants were subcloned into the pBluescript II K/S vector and introduced into the construct using the Stratagene QuikChange site-directed mutagenesis kit as previously described.<sup>26,27</sup> Dideoxy sequencing was used to confirm the presence of only the desired mutation. The mutated FVIII cDNA was then ligated back into the FVIII expression shuttle construct and subjected to a second round of dideoxy sequencing, again to confirm that only the desired mutation was present. The FVIII expression construct was transfected to BHK cells using FuGENE6 (Roche). Geneticin antibiotic selection, subcloning, and cloning of stable transfectants were performed by standard methods, and the cloned cells were cultured in roller bottles for protein expression.<sup>27</sup> Approximately 300 mL of conditioned medium was collected daily for 4 days, and the expressed proteins were purified by SP-Sepharose (Amersham Biosciences) column chromatography as

previously described.<sup>27</sup> Fractions were assayed using FVIII-deficient plasma in a one-stage clotting assay to identify fractions containing FVIII. Yields of purified wild-type (WT) and variant FVIII ranged from ~0.2 to 1.0 mg/L of conditioned medium. Resultant FVIII was >90% pure as judged by SDS–PAGE, with the main contaminant being albumin. FVIII samples were snap-frozen in liquid N<sub>2</sub> and stored at –80 °C.

**One-Stage Clotting Assay.** FVIII-deficient pooled human plasma was incubated with activated partial thromboplastin reagent (Trinity Biotech) for 7 min at 37 °C. A dilution of FVIII was added to the cuvette, and after incubation at 37 °C for 1 min, CaCl<sub>2</sub> (final concentration of 6 mM) was added and the time to form a clot measured using an ST4 coagulation analyzer (Diagnostica Stago) and compared with that of a pooled normal plasma standard (George King Biomedical, Inc.).

**Enzyme-Linked Immunosorbent Assay (ELISA).** A sandwich ELISA was used to determine the final concentration of purified FVIII proteins as previously described.<sup>26</sup> Briefly, GMA-8003 was used as the capture antibody and biotinylated R8B12 was used as the detection antibody. A chromogenic assay was used to determine the amount of FVIII bound to the plate utilizing streptavidin-linked horseradish peroxidase (Calbiochem) with the chromogen *o*-phenylenediamine dihydrochloride (Sigma). Purified commercial recombinant FVIII (Kogenate, Bayer) was used as the standard. FVIII specific activity values were determined using one-stage clotting and an ELISA.<sup>26</sup>

**Thrombin Cleavage of FVIII.** WT or mutant FVIII (100 nM) was incubated at 22 °C with indicated concentrations of thrombin (see the figure legends) in a buffer containing 20 mM HEPES, 0.14 M NaCl, 5 mM CaCl<sub>2</sub>, and 0.01% Tween 20 (pH 7.2). Samples were removed at the indicated points during the time course; reactions were terminated by addition of SDS–PAGE sample buffer [0.04 M Tris-HCl, 2% (w/v) SDS, 4% (v/v) glycerol, and 0.05% bromophenol blue (pH 6.8)] containing 3% (v/v) 2-mercaptoethanol and mixtures boiled for 3 min. Cleavage rates were assessed by rates of A1, A2, and/or A3–C1–C2 subunit generation.

**Electrophoresis and Western Blotting.** Thrombin cleavage of WT and mutant FVIII was evaluated by SDS–PAGE<sup>28</sup> on 8% polyacrylamide gels using a Bio-Rad mini-gel apparatus at 175 V for 1 h. Protein transfer occurred on a polyvinylidene fluoride membrane for 1 h at 100 V in ice-cold transfer buffer [10 mM 3-(cyclohexylamino)propane-1-sulfonic acid and 10% (v/v) methanol (pH 11)]. Western blotting was performed by probing the membranes with anti-FVIII monoclonal antibodies as indicated in the figure legends, followed by addition of an alkaline phosphatase-linked goat anti-mouse antibody (Sigma) as the secondary antibody. An enhanced chemifluorescence (ECF) system (Amersham Biosciences) was used to detect the signal, and the blots were scanned over a range of 650–740 nm with a Storm 860 instrument (GE Healthcare). Densitometry was used to quantitate the linear density regions of the blots using ImageQuant (GE Healthcare).

**Determination of Rates of Subunit Generation.** Rates of subunit generation following thrombin cleavage of the WT and mutants were calculated from the blots on the basis of summation of all FVIII band density values. The intensity of each band was determined by volume integration and corrected for variability in background staining by using densities in the areas immediately adjacent to each band. The concentrations of

**Table 1. Specific Activities and Sequences of FVIII P3–P3' Mutants**

factor VIII	372(P3–P3') sequence <sup>a</sup> (residues 370–375)	740(P3–P3') sequence <sup>a</sup> (residues 738–743)	1689(P3–P3') sequence <sup>a</sup> (residues 1687–1692)	specific activity <sup>b</sup> (%)
WT	QIR <sup>1</sup> SVA	EPR <sup>1</sup> SFS	SPR <sup>1</sup> SFQ	100 ± 6
372(P3–P3')740	EPR <sup>1</sup> SFS			98 ± 5
372(P3–P3')1689	SPR <sup>1</sup> SFQ			14 ± 2
372(P3–P3')740/R740Q	EPR <sup>1</sup> SFS	EPQ <sup>1</sup> SFS		38 ± 4
372(P3–P3')740/740(P3–P3')372	EPR <sup>1</sup> SFS	QIR <sup>1</sup> SVA		122 ± 18
740(P3–P3')372		QIR <sup>1</sup> SVA		73 ± 2

<sup>a</sup>Sequences for the P3–P3' residues flanking Arg<sup>372</sup>, Arg<sup>740</sup>, and Arg<sup>1689</sup> are shown using single-letter amino acid code, and specific mutations are shown in bold. Arrows indicate the position of the scissile bond. <sup>b</sup>Specific activities are presented as a percentage ± the standard deviation from the WT value.

A1, A2, and A3-C1-C2 subunits were normalized to the total density in each lane to correct for any differences in loading. Initial time points were fit using nonlinear least-squares regression to the single-exponential equation

$$A = A_0(1 - e^{k_{\text{obs}}t}) \quad (1)$$

where  $A_0$  is the total nanomolar concentration of the A1, A2, or A3-C1-C2 subunit and  $t$  is the time in minutes. The absolute value of  $k_{\text{obs}}$  represents the rate of A1, A2, or A3-C1-C2 subunit generation that was normalized by the thrombin concentration and expressed in nanomoles of A1, A2, or A3-C1-C2 subunit per minute per nanomole of thrombin, respectively. All experiments were performed at least three separate times, and the average values with standard deviations are shown. The quality of the fits was acceptable with  $R^2$  values ranging from 0.946 to 0.995. Data were also subjected to analysis using a quadratic equation<sup>21</sup> and yielded rates that were typically within 30% of those obtained with the single-exponential equation (data not shown). Rate values are presented using the single exponential because the overall quality of the fitted curves was superior with this method.

**FXa Generation Assay.** A purified protein system was used to measure the rate of conversion of FX to FXa.<sup>29</sup> To assess thrombin activation of FVIII over time, we reacted FVIII (100 nM) with 5 or 0.5 nM thrombin in a buffer containing 20 mM HEPES, 0.14 M NaCl, 5 mM CaCl<sub>2</sub>, and 0.01% Tween 20 (pH 7.2). Samples were removed at the indicated times, and thrombin activity was inhibited by the addition of hirudin (2 units/mL). The dilute FVIIIa solution (final concentration of 2 nM) was added to FIXa (20 nM) in the presence of phospholipid vesicles (10 μM). FXa generation was initiated by addition of FX (300 nM). The reactions were terminated with EDTA (50 mM), and rates of FXa generation were determined by the addition of the chromogenic substrate Pefachrome FXa (final concentration of 0.46 mM). Reactions were read at 405 nm for 5 min using a  $V_{\text{max}}$  microtiter plate reader (Molecular Devices).

## RESULTS


### Specific Activity of Recombinant FVIII Protein.

Thrombin cleaves the FVIII procofactor at Arg<sup>740</sup>, Arg<sup>372</sup>, and Arg<sup>1689</sup>. Recent studies have shown that cleavage at Arg<sup>740</sup> appears severalfold faster than that at Arg<sup>1689</sup>, and this effect likely results in part from exosite-driven competition between the heavy chain and the light chain of FVIII for thrombin binding.<sup>21</sup> Furthermore, cleavage at Arg<sup>740</sup> facilitates cleavage at Arg<sup>372</sup>,<sup>30</sup> which appears to represent the rate-limiting step in procofactor activation. This study investigates the effect of the P3–P3' residues of FVIII on thrombin proteolysis at the three

cleavage sites, and to this end, several recombinant B domain-deleted FVIII mutants were prepared and stably expressed. 372(P3–P3')740, 372(P3–P3')1689, and 740(P3–P3')372 designate variants in which the P3–P3' sequences surrounding one scissile bond in FVIII were swapped with others (see Table 1). For example, in the 372(P3–P3')740 variant, the P3–P3' sequence at Arg<sup>372</sup> was replaced with that from Arg<sup>740</sup>. In addition, a variant possessing the double swap 372(P3–P3')740/740(P3–P3')372 and one possessing the 740 for 372 swap with a point mutation of Arg<sup>740</sup> to Gln to create a noncleavable site [372(P3–P3')740/R740Q] were prepared.

Specific activities of WT and variant FVIII were measured using one-stage clotting and ELISAs (Table 1). The specific activities suggest an essentially normal phenotype for the 372(P3–P3')740, 372(P3–P3')740/740(P3–P3')372, and 740(P3–P3')372 variants. The 372(P3–P3')740/R740Q variant showed a modestly reduced specific activity (~40% of that of WT), whereas the value for the 372(P3–P3')1689 variant (~15% of that of WT) was consistent with a mild hemophilic phenotype. The observed reduction in the specific activity for FVIII 372(P3–P3')740/R740Q can be attributed solely to the Gln mutations at Arg<sup>740</sup> because this value was similar to the specific activity determined previously for the single-P1 site mutation alone.<sup>30</sup>


**Thrombin-Catalyzed Cleavage of 372(P3–P3') and 740(P3–P3') Recombinant FVIII Mutants.** Variants 372(P3–P3')740 and 372(P3–P3')1689 were used to assess whether the more thrombin-optimal P3–P3' residues that surround Arg<sup>740</sup> and Arg<sup>1689</sup> accelerate thrombin-catalyzed cleavage at Arg<sup>372</sup>. These cleavage sites are schematically represented in Figure 1, which also illustrates the FVIII single



**Figure 1.** Forms of B-domainless FVIII secreted from BHK cells. Secreted B-domainless FVIII single-chain (A) and heterodimer (B) forms. Thrombin cleavage sites are denoted with arrows, and a star denotes the furin cleavage site. The B domain in B-domainless FVIII represents a 14-amino acid remnant connecting the a2 and a3 acidic regions.

chain (A) and heterodimer (B). The B-domainless recombinant FVIII used in this study is expressed in nearly equal amounts of single-chain and heterodimer forms. Furthermore, a B domain





**Figure 2.** Thrombin cleavage of the single-swap P3–P3' FVIII mutants. WT and the 372(P3–P3')740 and 740(P3–P3')372 FVIII variants (100 nM) were reacted with thrombin over a 25 min time course, and samples were subjected to Western blotting using C5, R8B12, and 2D2 monoclonal antibodies for detection of A1, A2, and A3-C1-C2 subunits, respectively, as described in Experimental Procedures. (A) Representative blots for rates of A1 subunit generation for reactions using 0.75 nM thrombin for the WT and 740(P3–P3')372 and 0.075 nM thrombin for 372(P3–P3')740. (B) A2 subunit generation using 0.75 nM thrombin for the WT and 740(P3–P3')372 and 0.075 nM thrombin for 372(P3–P3')740. (C) A3-C1-C2 subunit generation using 0.13 nM thrombin for the WT and variants. SC refers to single chain and LC to light chain.


remnant of 14 amino acid residues remains and separates the a2 and a3 segments in the single-chain form or represents the C-terminal end of the heavy chain of the heterodimer.

Reactions were conducted over a 25 min time course, and results were visualized using SDS–PAGE and Western blotting to evaluate rates of FVIIIa subunit generation by thrombin as described in Experimental Procedures. Reaction of WT FVIII with thrombin showed efficient cleavage of the single chain and heavy and light chains, resulting in the generation of the A1, A2, and A3-C1-C2 FVIIIa subunits (Figure 2). These blots were quantitated by scanning densitometry and subjected to nonlinear least-squares regression analysis (Figure 3) to calculate rates of generation of each FVIIIa subunit (Table 2). Generation of the A1 subunit results from a single cleavage at Arg<sup>372</sup>; therefore, the rate of cleavage at that residue is equivalent to the rate of A1 subunit generation. The A2 subunit is derived from two cleavages: one between the A1 and A2 domains at Arg<sup>372</sup> and the other between the a2 segment and B domain remnant at Arg<sup>740</sup> (Figure 1). While the A2 subunit released from the heavy chain of the heterodimer requires only the single cleavage at Arg<sup>372</sup>, both cleavages at Arg<sup>372</sup> and Arg<sup>740</sup> are necessary to excise this subunit from single-chain FVIII. Because the presence or absence of the remnant cannot be discerned using SDS–PAGE, it is not possible to directly determine the rate of cleavage at Arg<sup>740</sup> in recombinant B-domainless FVIII. However, rates of A2 subunit generation in the variants are useful for estimation of this parameter relative to the WT FVIII value. A single cleavage at Arg<sup>1689</sup> in both the light chain and the single chain yields the A3-C1-C2 subunit; therefore, the rate of generation of this subunit represents the rate of cleavage at Arg<sup>1689</sup>.

Evaluation of reactions using 372(P3–P3')740 (Figure 2A) and 372(P3–P3')1689 FVIII (blots not shown) revealed

10- and 3-fold increases, respectively, in the rates of cleavage at Arg<sup>372</sup> as determined by rates of A1 subunit generation for the variants as compared with WT FVIII (Figure 3A and Table 2). Reactions for the variants used 10% of the thrombin concentration employed for WT. Control experiments showed that cleavage rates were linear over this thrombin concentration range. Although the rate of thrombin cleavage at Arg<sup>372</sup> increased for the variants, this increased rate did not appear to be matched by the rate of Arg<sup>740</sup> cleavage as suggested by the appearance of an A2-LC intermediate that could be detected in the blots using the anti-A2 domain and anti-A3 domain antibodies (Figure 2B,C). Thus, the 10- and 8-fold increases in A2 subunit generation for the 372(P3–P3')740 and 372(P3–P3')1689 mutants, respectively, most likely reflect the increase in the rate of thrombin cleavage at Arg<sup>372</sup> (Table 2). The rate of cleavage at Arg<sup>1689</sup> for both 372(P3–P3') mutants was similar to that of WT FVIII, suggesting little effect of accelerating cleavage at Arg<sup>372</sup> on cleavage at Arg<sup>1689</sup>. These results suggest that placing more optimal residues at Arg<sup>372</sup> increases the rate of thrombin cleavage at that site. Furthermore, the P3–P3' residues flanking Arg<sup>740</sup> appeared to be more optimal for Arg<sup>372</sup> cleavage than those flanking Arg<sup>1689</sup> (Table 2). The P3–P3' sequences flanking Arg<sup>740</sup> and Arg<sup>1689</sup> differ only in the P3 and P3' residues, suggesting the P3 Glu and/or P3' Ser may be preferred over Ser and/or Gln, respectively.

Examination of the effects of replacing P3–P3' residues flanking Arg<sup>740</sup> on thrombin-catalyzed cleavages of FVIII with those of the less optimal Arg<sup>372</sup> flanking sequence employed a similar Western blot approach as described above. Results obtained for the 740(P3–P3')372 variant revealed an ~2-fold increase in the rate of A1 subunit generation as well as an increased rate of A2 subunit generation (Figure 3A,B and Table 2). These results were surprising and, for reasons that are unclear, indicated an accelerated rate of cleavage at P1 Arg<sup>372</sup>.



**Figure 3.** Analysis of the Western blots of the single-swap P3-P3' FVIII mutants. Panels A–C plot rates derived from densitometry scans of the blots in Figure 2 and for Western blot data of 372(P3-P3')1689 (data not shown) for rates of generation of A1 (A), A2 (B), and A3-C1-C2 (C) subunits for WT (●), 372(P3-P3')740 (○), 372(P3-P3')1689 (▲), and 740(P3-P3')372 (△) FVIII. Lines are drawn representing the initial time points of the data fit to a single-exponential equation using nonlinear least-squares regression as described in Experimental Procedures. Experiments were performed at least three separate times, and average values are shown.

However, the persistence of the A2 light chain and A2-a3 intermediates (Figure 2B,C) over an extended time course suggested impairment of the rate of cleavage at Arg<sup>740</sup>, consistent with the less optimal cleavage sequence at this site.

**Thrombin-Catalyzed Cleavage of Combined (P3-P3') Recombinant FVIII Mutants.** To gain insights into the coordination of heavy chain cleavage during the activation process, we prepared a FVIII variant possessing two reciprocal swaps in the sequences flanking Arg<sup>372</sup> and Arg<sup>740</sup>. Thus, results

**Table 2.** Rates of A1, A2, and A3-C1-C2 Subunit Generation during WT and P3-P3' Mutant FVIII Activation by Thrombin


factor VIII	A1 subunit generation <sup>a</sup> [nmol of A1 min <sup>-1</sup> (nmol of IIa) <sup>-1</sup> ]	A2 subunit generation <sup>a</sup> [nmol of A2 min <sup>-1</sup> (nmol of IIa) <sup>-1</sup> ]	A3-C1-C2 subunit generation <sup>a</sup> [nmol of A3-C1-C2 min <sup>-1</sup> (nmol of IIa) <sup>-1</sup> ]
WT	63 ± 0.9	21 ± 1.1	199 ± 29
372(P3-P3')740	652 ± 42	236 ± 25	121 ± 7.1
372(P3-P3')1689	175 ± 12	164 ± 22	123 ± 10
740(P3-P3')372	147 ± 22	168 ± 16	124 ± 16
372(P3-P3')740/R740Q	665 ± 19	157 ± 15	157 ± 13
372(P3-P3')740/740(P3-P3')372	576 ± 33	216 ± 22	99 ± 7.4

<sup>a</sup>Rates of A1, A2, and A3-C1-C2 subunit generation by thrombin cleavage of WT and mutant FVIII were estimated by nonlinear least-squares regression analysis of the data shown in Figures 2–5. The data represent the average ± SD values of at least three separate experiments. Western blot assays were performed and data analyzed as described in Experimental Procedures.

obtained with this variant, 372(P3-P3')740/740(P3-P3')372, would examine compensatory effects on cleavage within the same chain following replacement of the slow site at Arg<sup>372</sup> with the fast site at Arg<sup>740</sup> and vice versa. In an earlier study,<sup>30</sup> variants showing slower cleavage rates at Arg<sup>740</sup> also demonstrated reduced rates of cleavage at the Arg<sup>372</sup> site. Results from the Western blotting time course of thrombin activation using the 372(P3-P3')740/740(P3-P3')372 variant (Figure 4A,B) showed a 9-fold enhancement in the rate of Arg<sup>372</sup> cleavage (A1 subunit generation (Figure 5A,B and Table 2). Both of these effects were nearly identical to results obtained with the single-swap 740(P3-P3')372 variant. The blotting also revealed two A2-LC and A2-a3 intermediates (Figure 4B,C), suggesting inefficient cleavage at Arg<sup>740</sup> was a result of the swap for the Arg<sup>372</sup> flanking sequence. Interestingly, there was a 2-fold decrease in the rate of cleavage at Arg<sup>1689</sup> as seen by a decreased rate of A3-C1-C2 subunit generation (Table 2).

The combination mutation variant used above placed the nonoptimal P3-P3' sequence from Arg<sup>372</sup> at Arg<sup>740</sup> in attempt to slow cleavage at the latter site. In the following experiment, we used an additional combined variant in which Arg<sup>740</sup> was replaced with Gln to eliminate this cleavage. This variant, designated 372(P3-P3')740/Arg740Gln, still retains the optimal flanking residues for cleavage at Arg<sup>372</sup> as in the variant described above but is now refractory to cleavage at the newly created Gln<sup>740</sup> residue. Earlier studies employing the single-point mutations, where Gln replaced Arg<sup>740</sup>,<sup>30</sup> showed a markedly reduced rate of cleavage at Arg<sup>372</sup> (~700-fold), reflecting a linkage of these catalytic events. Thus, the rationale for our experiment was to assess the influence of the optimal flanking sequence surrounding Arg<sup>372</sup> when the alternate site could not be cleaved.

Results using the 372(P3-P3')740/Arg740Gln FVIII mutant indicated that thrombin cleavage at Arg<sup>740</sup> was inhibited as shown by the persistence of the A2-LC and A2-a3 intermediates (Figure 4B,C). However, as judged by A1 subunit generation, the rate of cleavage at Arg<sup>372</sup> was increased by 11-fold compared with that of the WT to a similar value observed for the 372(P3-P3')740 variant (Figure 5A and



**Figure 4.** Thrombin cleavage of the combination P3–P3' FVIII mutants. The WT and the 372(P3–P3')740/R740Q and 372(P3–P3')740/740(P3–P3')372 FVIII variants (100 nM) were reacted with thrombin, and samples were subjected to Western blotting as described in the legend of Figure 2. (A) Representative blots for rates of A1 subunit generation for reactions using 0.75 nM thrombin for the WT and 0.075 nM thrombin for the variants. (B) A2 subunit generation using 0.75 nM thrombin for the WT and 0.075 nM thrombin for the variants. (C) A3-C1-C2 subunit generation using 0.13 nM thrombin for the WT and variants. SC refers to single chain and LC to light chain.

Table 2). These results suggest placing the more optimal P3–P3' residues at Arg<sup>372</sup> can abrogate the influence of initial cleavages at Arg<sup>740</sup> to yield (nearly) maximal rates of cleavage at Arg<sup>372</sup>.


**Thrombin Activation of FVIII Proteins.** Activation of the FVIII variants was performed to assess the effects of the P3–P3' mutations on the generation of FVIIIa activity. WT and mutant FVIII (100 nM) were reacted with thrombin over a 30 min time course, and activity was monitored using a FXa generation assay as described in Experimental Procedures. Two levels of thrombin (5 or 0.5 nM) were used to activate WT FVIII. The rate of activation was directly proportional to the level of thrombin used. At the higher concentration of thrombin (5 nM), WT FVIII demonstrated a peak activity (~40 nM/min) at ~1 min, after which cofactor activity steadily declined to ~10% of peak activity at 30 min (Figure 6). The loss of WT FVIIIa activity during the time course is independent of thrombin and results from the dissociation of the A2 subunit from the A1/A3-C1-C2 dimer, leading to inactivation of the cofactor and dampening of the intrinsic factor Xase.<sup>31</sup> When 0.5 nM thrombin was used to activate WT FVIII, the activation profile yielded a slower activation rate, an ~5-fold reduction in peak activity, and a broader plateau of activation (~5–10 min). The plateau of activity likely reflects a balance in the slower rate of activation coupled with the decay of FVIIIa.<sup>31</sup>

FVIII variants were activated using the lower thrombin concentration. In all cases, the generation of FVIIIa occurred faster than when this thrombin concentration was used with WT FVIII (Figure 6). Similar activity profiles were observed for the 740(P3–P3')372 and 372(P3–P3')740/740(P3–P3')372 mutants. These variants demonstrated a faster rate of procofactor activation than the WT (1–2 min vs 5–10 min)

with an ~2-fold increase in peak height. The 372(P3–P3')740 and 372(P3–P3')740/R740Q variants also exhibited faster rates of activation (peaking at 1–2 and 2–5 min, respectively) and had similar peak values of activation compared to WT FVIII. Lastly, the 372(P3–P3')1689 variant, although showing a faster rate of activation, had an ~8-fold reduction in the peak level of activation. This variant also demonstrated low specific activity that suggested substitution of residues affected cofactor function. However, in all cases, these faster activating variants all exhibited increased rates of cleavage at Arg<sup>372</sup>, consistent with this site representing the rate-limiting step in the activation mechanism.

## DISCUSSION


In this study we examined thrombin-catalyzed cleavage and activation of FVIII substrates possessing swaps for the P3–P3' sequences flanking the scissile bonds in the heavy chain. Results obtained from variants possessing single sequence swaps, a double sequence swap, or a single swap combined with point mutations to render the P1 Arg<sup>740</sup> noncleavable yielded several observations regarding the role of the P3–P3' sequences in the activation mechanism. First, the sequence flanking a particular P1 makes a variable and, in some cases, substantial contribution to the rate of attack at that site, with more optimal residues for thrombin yielding enhanced reaction rates. Second, P3–P3' sequences at the scissile bonds in the heavy chain appear to modulate the rate of catalysis at distal P1 sites. Third, comparing effects of P3–P3' sequences flanking residues Arg<sup>740</sup> and Arg<sup>1689</sup>, which possess differences in only the P3 and P3' positions, suggests changes in individual residues removed from the P1 site can yield marked effects on the catalytic rate. Finally, swaps that place more optimal residues



**Figure 5.** Analysis of the Western blots of the combination P3-P3' FVIII mutants. Rates derived from densitometry scans of the blots in Figure 4 for rates of generation of A1 (A), A2 (B), and A3-C1-C2 (C) subunits for 372(P3-P3')740/R740Q (●) and 372(P3-P3')740/740(P3-P3')372 (○) FVIII. Lines are drawn representing the initial time points of the data fit to a single-exponential equation using nonlinear least-squares regression as described in Experimental Procedures. Experiments were performed at least three separate times, and average values are shown.

flanking the rate-limiting Arg<sup>372</sup> site result in more rapid rates of thrombin-catalyzed cleavage and procofactor activation, independent of the sequences flanking the scissile bond at Arg<sup>740</sup> or whether that bond can be cleaved. This latter observation is of particular interest in that it reflects a disengagement of the linked pathway for heavy chain cleavage.

Examination of the P3-P3' sequences flanking the three scissile bonds in FVIII cleaved by thrombin revealed that two of the P1 residues, Arg<sup>740</sup> and Arg<sup>1689</sup>, are flanked by sequences containing four of six residues deemed optimal for thrombin cleavage, whereas the sequence flanking Arg<sup>372</sup> is optimal at only the P1 and P1' sites.<sup>22</sup> Results of this study show that replacement of the sequence flanking Arg<sup>372</sup> with those of Arg<sup>740</sup> and Arg<sup>1689</sup> yielded 10- and 3-fold increases, respectively, in the rates of cleavage at the P1 Arg. Thus, enhancement of the rates of catalysis at this site is clearly influenced by sequence.



**Figure 6.** Activation of WT and variant FVIII by thrombin. FXa generation assays were used to monitor rates of activation of WT and variant FVIII (100 nM) by 5 and 0.5 nM thrombin as described in Experimental Procedures. Solid lines represent data for the WT (□) with 5 nM thrombin and the WT (■) and variants 372(P3-P3')740/740(P3-P3')372 (●), 740(P3-P3')372 (○), 372(P3-P3')740 (▲), 372(P3-P3')740/R740Q (△), and 372(P3-P3')1689 (◆) with 0.5 nM thrombin. All experiments were performed at least three separate times, and average values are shown.

A primary cause for this rate enhancement is likely derived from the P2 Pro residue, which is a preferred residue at this position.<sup>16</sup> Because the P2-P2' sequences flanking Arg<sup>740</sup> and Arg<sup>1689</sup> are identical, the rate differences observed in comparing the two replacements reflect differences at the P3 and P3' positions. Studies using libraries of natural thrombin substrates have revealed little influence of the P3 and P3' positions on catalysis.<sup>32</sup> However, in the case of FVIII, the P3 Glu and/or P3' Ser present in the Arg<sup>740</sup> sequence appears to be somewhat more preferred for cleavage compared with the Ser and/or Gln in the Arg<sup>1689</sup> sequence as judged by the ~3-fold difference in cleavage rates at the substituted Arg<sup>372</sup> site.

Both of the Arg<sup>740</sup> and Arg<sup>1689</sup> sequences affected faster cleavage at Arg<sup>372</sup> and yielded faster activation kinetics, consistent with this site representing the apparent rate-limiting step in the activation mechanism.<sup>23</sup> While replacement with the Arg<sup>740</sup> site was benign with respect to specific activity, replacement with the Arg<sup>1689</sup> flanking sequence yielded a marked (>80%) reduction in this parameter. The reason for this is not clear, but it may reflect replacement of specific residues, specifically, the P3 Ser and/or P3' Gln that inhibits FVIII function. Examination of a hemophilia A database<sup>33</sup> does not reveal any missense mutations at position 370 (P3) or 375 (P3') affecting phenotype.

Earlier studies from our laboratory examining thrombin-catalyzed cleavage and activation of FVIII when individual P1 sites were rendered noncleavable by replacement of the P1 Arg with Gln showed mutation at one site had significant effects at unmodified P1 sites. For example, rates of A1 and A2 subunit generation were reduced ~8–10-fold using a FVIII variant possessing an Arg1689Gln mutation.<sup>21</sup> Furthermore, an Arg740Gln variant showed an ~140-fold reduced rate of cleavage at Arg<sup>1689</sup>.<sup>30</sup> Results from this study show an ~2-fold reduction in the rate of cleavage at Arg<sup>1689</sup> when the Arg740Gln mutation is coupled with the 372(P3-P3')740 mutation. The reason for this large disparity in relative reaction rates may reflect the markedly faster cleavage rate at the Arg<sup>372</sup> site in the latter variant that in turn modulates the reaction at the light chain.



We have speculated that the apparent cleavage order ( $\text{Arg}^{740} > \text{Arg}^{1689} > \text{Arg}^{372}$ ) leading to procofactor activation represents a competition for active site docking to sites in the two chains of the FVIII heterodimer, with an initial attack of the heavy chain at  $\text{Arg}^{740}$  (followed by subsequent cleavage at  $\text{Arg}^{372}$ ) representing the favored pathway over initial cleavage at  $\text{Arg}^{1689}$ .<sup>21</sup> Furthermore, the slow cleavage of  $\text{Arg}^{372}$  was thought to be a result of the requirement to first cleave  $\text{Arg}^{740}$ , possibly requiring a ratcheting mechanism<sup>17</sup> to permit subsequent active site docking. Interestingly, mutations that place optimal residues flanking  $\text{Arg}^{372}$  appear to obviate the need for initial cleavage at  $\text{Arg}^{740}$  to facilitate  $\text{Arg}^{372}$  cleavage. We observed that either placing the nonoptimal residues flanking  $\text{Arg}^{372}$  at  $\text{Arg}^{740}$  to slow cleavage at this site or replacing the P1  $\text{Arg}^{740}$  with Gln resulted in sustained high rates of cleavage at  $\text{Arg}^{372}$  (~10-fold greater than the WT rate) when this site was flanked with the P3–P3' sequence from  $\text{Arg}^{740}$ . This faster cleavage of  $\text{Arg}^{372}$  for these two variants was confirmed following FVIII activation at low thrombin levels, which yielded peak activity values during a time course shorter than that observed for the WT protein. Taken together, these results suggested that thrombin bound to FVIII may have access to all three scissile bonds in the FVIII substrate and catalysis at a given bond reflects in large part docking preferences based upon S and P site interactions.

Sequences flanking scissile bonds and their complementary interaction with binding pocket residues of the proteinase make marked contributions to catalytic rates in a number of coagulation proteinase systems. For example, exchanging the P2 and P3 binding pocket residues of FXa and activated protein C switched the enzyme specificities for chromogenic substrates and inhibitors.<sup>34</sup> In another study, replacing the P2 and P3 residues of prothrombin 2 (Asp–Gly) with those from FIXa (Leu–Thr) resulted in ~20- and ~40-fold reductions in  $k_{\text{cat}}$  for FXa and prothrombinase, respectively.<sup>18</sup> Additionally, this study also showed that  $K_{\text{m}}$  values for prothrombin were essentially unaffected by this and other residue replacements, consistent with the dependence for substrate recognition and affinity on exosite interactions.

This influence of P1 flanking sequences on the catalytic mechanism of thrombin activation of FVIII appears to represent a common theme for cleavage of FVIII substrates by exosite-driven enzymes. In an earlier study<sup>19</sup> examining the role of P4–P3' sequences flanking the two scissile bonds in FVIII cleaved by activated protein C, we observed that replacing the faster-reacting site at  $\text{Arg}^{336}$  in the A1 domain with the slower-reacting site at  $\text{Arg}^{562}$  in the A2 domain resulted in an approximately  $\leq 100$ -fold reduction in the rate of cleavage at the P1  $\text{Arg}^{336}$  in FVIIIa, whereas the reciprocal swap accelerated cleavage at  $\text{Arg}^{562}$  in the procofactor by ~25-fold. Inasmuch as little information is available regarding S4–S3' sites in this proteinase, discrimination of optimal and nonoptimal P residues could not be distinguished.

Overall, the results of this study provide insights into the mechanism of the specificity of thrombin for the P1 sites of FVIII. Faster rates of cleavage at  $\text{Arg}^{740}$  and  $\text{Arg}^{1689}$  can be attributed in part to more optimal residues in the P3–P3' region, while the relatively slower rate of cleavage at  $\text{Arg}^{372}$  can be accelerated by replacement with more optimal residues for thrombin cleavage. Furthermore, this resultant acceleration appears to reflect the uncoupling of the linkage between cleavage of the two scissile bonds in the FVIII heavy chain. Thus, the P3–P3' residues surrounding  $\text{Arg}^{740}$ ,  $\text{Arg}^{1689}$ , and  $\text{Arg}^{372}$  in FVIII impact rates of thrombin proteolysis at each site

and contribute to the mechanism for thrombin activation of the procofactor.

## AUTHOR INFORMATION

### Corresponding Author

\*Department of Biochemistry and Biophysics, P.O. Box 712, University of Rochester Medical Center, 601 Elmwood Ave., Rochester, NY 14642. Telephone: (585) 275-6576. Fax: (585) 275-6007. E-mail: philip\_fay@urmc.rochester.edu.

### Present Address

†Department of Pathology, Microbiology, and Immunology, Vanderbilt University School of Medicine, Nashville, TN 37232.

### Funding

This work was supported by Grants HL38199 and HL76213 from the National Institutes of Health (to P.J.F.).

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

We thank Pete Lollar and John Healey for the gift of the FVIII cloning and expression vectors and Zaverio Ruggeri, Bill Church, and Lisa Regan for the C5, GMA-8003, and 2D2 monoclonal antibodies, respectively.

## ABBREVIATIONS

BHK, baby hamster kidney; HEPES, *N*-(2-hydroxyethyl)-piperazine-*N'*-2-ethanesulfonic acid; SDS–PAGE, sodium dodecyl sulfate–polyacrylamide gel electrophoresis; ELISA, enzyme-linked immunosorbent assay; WT, wild-type; FVIII, factor VIII; FVIIIa, factor VIIIa; FIXa, factor IXa; FX, factor X; FXa, factor Xa; ABE, anion binding exosite.

## ADDITIONAL NOTE

<sup>a</sup>Residues flanking the scissile bonds are denoted using the nomenclature of Schechter and Berger.<sup>35</sup> Residues preceding the scissile bond are designated P1–P<sub>n</sub> extending toward the NH<sub>2</sub> terminus, and residues following the scissile bond are designated P1'–P<sub>n'</sub> extending toward the COOH terminus.

## REFERENCES

- (1) Vehar, G. A., Keyt, B., Eaton, D., Rodriguez, H., O'Brien, D. P., Rotblat, F., Oppermann, H., Keck, R., Wood, W. I., Harkins, R. N., et al. (1984) Structure of human factor VIII. *Nature* 312, 337–342.
- (2) Wood, W. I., Capon, D. J., Simonsen, C. C., Eaton, D. L., Gitschier, J., Keyt, B., Seeburg, P. H., Smith, D. H., Hollingshead, P., Wion, K. L., et al. (1984) Expression of active human factor VIII from recombinant DNA clones. *Nature* 312, 330–337.
- (3) Fay, P. J., Anderson, M. T., Chavin, S. I., and Marder, V. J. (1986) The size of human factor VIII heterodimers and the effects produced by thrombin. *Biochim. Biophys. Acta* 871, 268–278.
- (4) Mann, K. G., Nesheim, M. E., Church, W. R., Haley, P., and Krishnaswamy, S. (1990) Surface-dependent reactions of the vitamin K-dependent enzyme complexes. *Blood* 76, 1–16.
- (5) Eaton, D., Rodriguez, H., and Vehar, G. A. (1986) Proteolytic processing of human factor VIII. Correlation of specific cleavages by thrombin, factor Xa, and activated protein C with activation and inactivation of factor VIII coagulant activity. *Biochemistry* 25, 505–512.
- (6) Parker, E. T., Pohl, J., Blackburn, M. N., and Lollar, P. (1997) Subunit structure and function of porcine factor Xa-activated factor VIII. *Biochemistry* 36, 9365–9373.
- (7) Pieters, J., Lindhout, T., and Hemker, H. C. (1989) In situ-generated thrombin is the only enzyme that effectively activates factor



VIII and factor V in thromboplastin-activated plasma. *Blood* 74, 1021–1024.

(8) Fay, P. J., Matri, M., Koszelak, M. E., and Wakabayashi, H. (2001) Cleavage of factor VIII heavy chain is required for the functional interaction of A2 subunit with factor IXa. *J. Biol. Chem.* 276, 12434–12439.

(9) Donath, M. J., Lenting, P. J., Van Mourik, J. A., and Mertens, K. (1996) Kinetics of factor VIII light-chain cleavage by thrombin and factor Xa. A regulatory role of the factor VIII heavy-chain region Lys713-Arg740. *Eur. J. Biochem.* 240, 365–372.

(10) Lollar, P., Hill-Eubanks, D. C., and Parker, C. G. (1988) Association of the factor VIII light chain with von Willebrand factor. *J. Biol. Chem.* 263, 10451–10455.

(11) Regan, L. M., and Fay, P. J. (1995) Cleavage of factor VIII light chain is required for maximal generation of factor VIIIa activity. *J. Biol. Chem.* 270, 8546–8552.

(12) Fay, P. J., Haidaris, P. J., and Smudzin, T. M. (1991) Human factor VIIIa subunit structure. Reconstruction of factor VIIIa from the isolated A1/A3-C1-C2 dimer and A2 subunit. *J. Biol. Chem.* 266, 8957–8962.

(13) Lollar, P., and Parker, E. T. (1991) Structural basis for the decreased procoagulant activity of human factor VIII compared to the porcine homolog. *J. Biol. Chem.* 266, 12481–12486.

(14) Fay, P. J., Smudzin, T. M., and Walker, F. J. (1991) Activated protein C-catalyzed inactivation of human factor VIII and factor VIIIa. Identification of cleavage sites and correlation of proteolysis with cofactor activity. *J. Biol. Chem.* 266, 20139–20145.

(15) Plantier, J. L., Rolli, V., Ducasse, C., Dargaud, Y., Enjolras, N., Boukerche, H., and Negrier, C. (2010) Activated factor X cleaves factor VIII at arginine 562, limiting its cofactor efficiency. *J. Thromb. Haemostasis* 2, 286–293.

(16) Di Cera, E. (2003) Thrombin interactions. *Chest* 124, 11S–17S.

(17) Krishnaswamy, S. (2005) Exosite-driven substrate specificity and function in coagulation. *J. Thromb. Haemostasis* 3, 54–67.

(18) Orcutt, S. J., Pietropaolo, C., and Krishnaswamy, S. (2002) Extended interactions with prothrombinase enforce affinity and specificity for its macromolecular substrate. *J. Biol. Chem.* 277, 46191–46196.

(19) Varfaj, F., Wakabayashi, H., and Fay, P. J. (2007) Residues surrounding Arg336 and Arg562 contribute to the disparate rates of proteolysis of factor VIIIa catalyzed by activated protein C. *J. Biol. Chem.* 282, 20264–20272.

(20) Nogami, K., Zhou, Q., Myles, T., Leung, L. L., Wakabayashi, H., and Fay, P. J. (2005) Exosite-interactive regions in the A1 and A2 domains of factor VIII facilitate thrombin-catalyzed cleavage of heavy chain. *J. Biol. Chem.* 280, 18476–18487.

(21) Newell, J. L., and Fay, P. J. (2009) Cleavage at Arg-1689 influences heavy chain cleavages during thrombin-catalyzed activation of factor VIII. *J. Biol. Chem.* 284, 11080–11089.

(22) Bode, W., Brandstetter, H., Mather, T., and Stubbs, M. T. (1997) Comparative analysis of haemostatic proteinases: Structural aspects of thrombin, factor Xa, factor IXa and protein C. *Thromb. Haemostasis* 78, 501–511.

(23) Hill-Eubanks, D. C., and Lollar, P. (1990) von Willebrand factor is a cofactor for thrombin-catalyzed cleavage of the factor VIII light chain. *J. Biol. Chem.* 265, 17854–17858.

(24) Mimms, L. T., Zampighi, G., Nozaki, Y., Tanford, C., and Reynolds, J. A. (1981) Phospholipid vesicle formation and transmembrane protein incorporation using octyl glucoside. *Biochemistry* 20, 833–840.

(25) Casillas, G., Simonetti, C., and Pavlovsky, A. (1971) Artificial substrate for the assay of factors V and VIII. *Coagulation* 4, 107–111.

(26) Jenkins, P. V., Freas, J., Schmidt, K. M., Zhou, Q., and Fay, P. J. (2002) Mutations associated with hemophilia A in the 558–565 loop of the factor VIIIa A2 subunit alter the catalytic activity of the factor Xase complex. *Blood* 100, 501–508.

(27) Wakabayashi, H., Freas, J., Zhou, Q., and Fay, P. J. (2004) Residues 110–126 in the A1 domain of factor VIII contain a Ca<sup>2+</sup>

binding site required for cofactor activity. *J. Biol. Chem.* 279, 12677–12684.

(28) Laemmli, U. K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680–685.

(29) Lollar, P., Fay, P. J., and Fass, D. N. (1993) Factor VIII and factor VIIIa. *Methods Enzymol.* 222, 128–143.

(30) Newell, J. L., and Fay, P. J. (2007) Proteolysis at Arg740 facilitates subsequent bond cleavages during thrombin-catalyzed activation of factor VIII. *J. Biol. Chem.* 282, 25367–25375.

(31) Fay, P. J. (2004) Activation of factor VIII and mechanisms of cofactor action. *Blood Rev.* 18, 1–15.

(32) Edwards, P. D., Mauger, R. C., Cottrell, K. M., Morris, F. X., Pine, K. K., Sylvester, M. A., Scott, C. W., and Furlong, S. T. (2000) Synthesis and enzymatic evaluation of a P1 arginine aminocoumarin substrate library for trypsin-like serine proteases. *Bioorg. Med. Chem.* 10, 2291–2294.

(33) Kemball-Cook, G., Tuddenham, E. G., and Wacey, A. I. (1998) The Factor VIII Structure and Mutation Resource Site: HAMSTeRS version 4. *Nucleic Acids Res.* 26, 216–219.

(34) Rezaie, A. R. (1996) Role of residue 99 at the S2 subsite of factor Xa and activated protein C in enzyme specificity. *J. Biol. Chem.* 271, 23807–23814.

(35) Schechter, I., and Berger, A. (1967) On the size of the active site in proteases. I. Papain. *Biochem. Biophys. Res. Commun.* 27, 157–162.