


Base Pairing and Miscoding Properties of 1,N⁶-Ethenoadenine- and 3,N⁴-Ethenocytosine-Containing RNA Oligonucleotides

Alessandro Calabretta and Christian J. Leumann*

Department of Chemistry and Biochemistry, University of Bern, Freiestrasse 3, CH-3012 Bern, Switzerland

S Supporting Information

ABSTRACT: Two RNA phosphoramidites containing the bases 1,N⁶-ethenoadenine (ϵ A) and 3,N⁴-ethenocytosine (ϵ C) were synthesized. These building blocks were incorporated into two 12-mer oligoribonucleotides for evaluation of the base pairing properties of these base lesions by UV melting curve (T_m) and circular dichroism measurements. The T_m data of the resulting duplexes with the etheno modifications opposing all natural bases showed a substantial destabilization compared to the corresponding natural duplexes, confirming their inability to form base pairs. The coding properties of these lesions were further investigated by introducing them into 31-mer oligonucleotides and assessing their ability to serve as templates in primer extension reactions with HIV, AMV, and MMLV reverse transcriptases (RT). Primer extension reactions showed complete arrest of the incorporation process using MMLV RT and AMV RT, while HIV RT preferentially incorporates dAMP opposite ϵ A and dAMP as well as dTMP opposite ϵ C. The properties of these RNA lesions are discussed in the context of its putative biological role.



The accumulation of DNA nucleobase lesions has been correlated with the onset of various diseases, first and foremost most types of cancer.^{1–4} The formation of such lesions has been extensively investigated in the past and can be attributed to a large variety of endogenous or exogenous agents.¹ In particular, oxidative stress is considered to be one of the most important processes leading to damaged nucleobases, yielding different classes of chemical modifications that can accumulate on the same target sequence.⁴ Reactive oxygen species (ROS) are endogenous compounds such as hydrogen peroxide, superoxide, and hydroxyl radicals. These ROS are constantly produced during normal mitochondrial aerobic respiration, during inflammation induced by viral or bacterial infections, by cytochromes P-450, and by the peroxisome-mediated degradation of fatty acids.⁵ Furthermore, ROS can attack nucleic acids in a direct or indirect way: in the first case, they react directly with the nucleobases, while in the second case, they react first with cellular metabolites, forming intermediate species that introduce other types of lesions into the nucleobases.⁵ 8-Oxo-dA, 8-oxo-dG, 5-hydroxy-dU, and 5-hydroxy-dC⁶ are the main lesions that are produced via the first pathway, while cyclic adducts such as etheno-dA, etheno-dC, and etheno-dG (Figure 1) are mainly produced using the indirect path.⁷

Etheno (ϵ) adducts were first identified as products of the reaction of chloroacetaldehyde with natural nucleobases, nucleosides, and DNA and RNA strands.^{8–12} In living organisms, chloroacetaldehyde occurs as a metabolite produced from vinyl chloride as a xenobiotic.¹³ More recently, ϵ -bases have been shown to occur during lipid peroxidation (LPO) by the




Figure 1. Chemical structures of the main etheno base lesions.

interaction of the resulting reactive aldehydes and hydroxy-alkenals with DNA.^{14–16} Indeed, high concentrations of ϵ -bases have been identified in those cells in which large amounts of oxygen reactive species are produced indirectly, such as white blood cells during chronic inflammation.² The mutagenicity of these lesions when they are introduced into oligodeoxynucleotides has been fully investigated via in vitro and in vivo assays, and the lesions have been shown to be strongly mutagenic when tested in human kidney cell lines.¹⁷

Received: January 30, 2013

Revised: February 19, 2013

Published: February 20, 2013



In this case, A/G transition and A/T transversion mutations were predominant.^{18,19} In other cells, such as *Escherichia coli*, minor mutagenic properties were observed for this lesion.^{19,20} Similarly, when tested in vivo, ϵ C showed higher mutagenic properties with mammalian cells than with *E. coli*. Indeed, while mainly C/T transitions were observed in mammalian cells, only C/A transversions were produced in *E. coli*, albeit at a lower frequency.^{20–23}

Although the formation of DNA base damage and its biological consequences are fairly well understood, surprisingly little is known about the effect of base lesions in RNA biochemistry and biology. Indeed, RNAs play a major role not only in translation (mRNAs and tRNA) but also in cellular regulation (miRNAs and other noncoding RNAs). Moreover, RNA is more abundant than DNA, is located in the cytoplasm where most of the reactive species (in particular ROS) are present, and is less protected by proteins than histone-organized DNA. Base-damaged ribonucleosides such as 8-oxo-rA, 8-oxo-rG, 5-HOrU, and 5-HOrC have recently been incorporated into oligoribonucleotides by solid-phase synthesis, and their base pairing properties with complementary RNA as well as their coding preferences in templated primer extension reactions (PEX) by reverse transcriptases (RTs) have been assessed.^{24–27} At the level of translation, oxidatively damaged mRNA has been demonstrated to cause largely diminished translation efficiency and abnormal protein synthesis, both in vitro and in vivo.^{28,29}

Here, we report on the synthesis of short 12-mer oligoribonucleotides containing either single ϵ A or ϵ C nucleotides as well as their base recognition properties upon duplex formation with complementary DNA and RNA. Furthermore, 31-mer RNA templates modified with these units were synthesized and used in PEX with three different RTs to determine the coding properties of these lesions during reverse transcription.

MATERIALS AND METHODS

Oligonucleotide Synthesis and Purification. Unmodified DNA and RNA oligonucleotides were purchased from Microsynth (Balgach, Switzerland) and purified via high-performance liquid chromatography (HPLC) or polyacrylamide gel electrophoresis (PAGE) when necessary. Natural 2'-TBDMS-protected RNA phosphoramidites and CPG solid supports were obtained from GlenResearch or AZCO. The syntheses were performed on a 1.3 μ mol scale on a Pharmacia Gene Assembler Plus DNA synthesizer following standard phosphoramidite protocols with 5-(ethylthio)-1H-tetrazole (0.25 M in CH₃CN) as the activator and coupling times of 12 min. Oligoribonucleotides were deprotected using a 33% NH₃/EtOH mixture (3:1) for 16 h at 25 °C in a screw cap 1.5 mL tube. Solid supports were filtered off and washed with the 3:1 NH₃/EtOH mixture. The solutions were evaporated to dryness at room temperature in a Speedvac evaporator. The resulting pellets were further dried by addition of dry ethanol followed by evaporation. The pellets were then dissolved in anhydrous DMSO (100 μ L), and neat triethylamine trihydrofluoride (125 μ L) was added. After the mixture had been shaken at 25 °C for 24 h, an aqueous solution of NaOAc (3 M, pH 5.2, 25 μ L) and *n*-butanol (1 mL) was added, and the mixture was chilled on dry ice for 45 min. After centrifugation for 20 min in an Eppendorf tube, the liquid was discarded and the pellet washed twice with chilled ethanol (80%, 1 mL) and dried under high vacuum at room temperature, yielding the crude oligonucleotides. The 31-mer oligoribonucleotides were purified on a preparative 20% denaturing polyacrylamide gel

(PAGE), and the product band was electroeluted with an Elutrap electroelution system (Schleicher & Schuell) and desalted using Amicon ultra 0.5 filters (molecular mass cutoff of 3 kDa, Millipore). The 12-mer was purified via HPLC on a DNAPac-200 column (4 mm \times 250 mm, Dionex) with a gradient from 100% solvent A [25 mM Tris (pH 8.0)] to 50% solvent B [25 mM Tris and 1.25 M NaCl (pH 8.0)] in 30 min at room temperature and finally desalted on a Sep-Pak C18 column (Waters Corp.). Oligonucleotide concentrations were determined using a NanoDrop ND-100 UV-vis spectrophotometer (NanoDrop Technologies, Inc.). Stock solutions were made from DEPC-treated H₂O.

UV Melting Curves. Solutions were prepared in standard saline buffer [10 mM NaH₂PO₄ and 150 mM NaCl (pH 7.0)] with duplex concentrations of 2 μ M in a 1:1 strand ratio. Thermal melting experiments were conducted on a Varian Cary 100-Bio UV-vis spectrophotometer (Varian Inc.), equipped with a Peltier element at 260 nm with a heating and cooling rate of 0.5 °C/min. *T_m* values were obtained from the maxima of the first derivatives of the melting curves using WinUV.

Labeling of Oligonucleotides. The DNA primer d(AT-CCCTGAGCGACTGGTGCA) was 5'-phosphorylated with [γ -³²P]ATP (Hartmann Analytic) and T4 polynucleotide kinase (Fermentas) as follows. DNA primer (30 pmol), [γ -³²P]ATP (60 mCi), and T4 PNK (10 units) were incubated in a total volume of 20 μ L at 37 °C for 30 min in T4 PNK buffer [50 mM Tris-HCl (pH 7.6), 10 mM MgCl₂, 5 mM DTT, 0.1 mM spermidine, and 0.1 mM EDTA]. The T4 PNK was inactivated when the sample was heated to 80 °C for 2 min. The labeled primer was purified from unreacted free label by Amicon Ultra centrifugal filters, dried under vacuum, and redissolved in 20 μ L of DEPC water. Labeled DNA was then




Figure 2. Chemical structures of ϵ A and ϵ C RNA phosphoramidites 1 and 2.

Table 1. Natural and Modified Sequences Used in This Work

oligonucleotide	RNA sequence
ON 1	5'-AUGCUG ϵ AAGUCGA-3'
ON 2	5'-AUGCUG ϵ CAGUCGA-3'
ON 3	5'-AUGCUG ϵ AAGUCGA-3'
ON 4	5'-AUGCUG ϵ CAGUCGA-3'
ON 5	5'-UAGUCUGCAC ϵ AUGCACCAGUCGUCAGGGAU-3'
ON 6	5'-UAGUCUGCAC ϵ CUGCACCAGUCGUCAGGGAU-3'
ON 7	5'-UAGUCUGCAC ϵ AUGCACCAGUCGUCAGGGAU-3'
ON 8	5'-UAGUCUGCAC ϵ CUGCACCAGUCGUCAGGGAU-3'




Figure 3. HPLC of crude ON 1 (left) and ON 2 (right).

Table 2. T_m Data ($^{\circ}\text{C}$, 260 nm) of Duplexes of Oligoribonucleotides ON 1–ON 4 with r(UCGACUXAGCAU) or d(TCGACTXAGCAT) as the Complement^a

	RNA complement				DNA complement			
	rA	rC	rG	rU	dA	dC	dG	dT
ON 1 (ϵA)	43.9	40.6	44.9	42.0	28.6	23.6	29.3	25.4
ON 3 (A)	44.9	46.6	46.7	57.6	29.6	34.6	29.9	45.3
ON 2 (ϵC)	43.9	39.3	46.6	41.9	25.2	20.8	25.3	24.3
ON 4 (C)	46.9	43.0	64.6	43.9	27.2	24.5	48.6	27.2

^aDuplex concentration of 2 μM in 10 mM NaH_2PO_4 and 150 mM NaCl (pH 7.0).

annealed to the RNA template (60 pmol) in the reaction buffer of the respective enzyme (see the description of RT assays) when the mixture was heated to 60 $^{\circ}\text{C}$ for 5 min and slowly cooled to room temperature.

RT Assays. RT assays were performed with HIV-1 RT (Worthington Biochemical Corp.), AMV RT (Promega Corp.), and MMLV RT (USB Corp./Affymetrix Inc.) in the following buffers: 50 mM Tris (pH 8.3), 50 mM NaCl, 8 mM MgCl_2 , and 1 mM DTT for HIV-1 RT and AMV RT and 50 mM Tris (pH 8.3), 75 mM KCl, 3 mM MgCl_2 , and 10 mM DTT for MMLV RT. Final reaction mixtures contained RNA template (100 nM), DNA primer (50 nM), and dNTP (20 mM). After addition of the enzyme (2–40 units), the mixtures were incubated at 37 $^{\circ}\text{C}$ for 1 h. The reactions were quenched with stop solution [98% formamide, 0.05% xylene cyanol (FF), and 0.05% bromophenol blue], heated to 90 $^{\circ}\text{C}$ for 5 min, and resolved by 20% PAGE. Visualization and quantification were conducted on a Storm 820 phosphorimager with ImageQuant (GE Healthcare).

RESULTS

Synthesis of the Modified Phosphoramidites and Oligoribonucleotides. The RNA oligonucleotides containing ϵA and ϵC modifications were synthesized using the RNA phosphoramidites 1 and 2 depicted in Figure 2. The synthesis of these building blocks was performed via a modification of a literature procedure.³⁰ The experimental details and the analytical data of the synthesized compounds can be found in the Supporting Information.

The oligoribonucleotides prepared for this study are summarized in Table 1. The 12-mer oligonucleotides ON 1–4 contain one of the damaged nucleobases or the corresponding natural bases in the center of the sequence and were used for T_m and circular dichroism (CD) studies, while the 31-mer oligonucleotides ON 5–8 were used as templates in PEX.

Coupling yields of the modified building blocks 1 and 2 were >98% as determined by a trityl assay. Oligoribonucleotides were first base-deprotected, phosphate-deprotected, and finally 2'-O-deprotected. The 12-mer sequences were purified by IE-HPLC, while the 31-mer oligonucleotides were purified via PAGE. To limit the extent of ethenobase degradation in alkaline media,³¹ we conducted the cleavage and base deprotection in a 33% NH_3 /ethanol mixture (3:1) at 25 $^{\circ}\text{C}$ overnight, obtaining the desired products in acceptable isolated yields: 14.8% for oligonucleotide ON 1 and 10.5% for oligonucleotide ON 2. HPLC traces of crude oligonucleotides (Figure 3) and mass spectrometric data (see the Supporting Information) demonstrated the relative stabilities of the modified oligonucleotides during deprotection.

UV Melting Curves and CD Spectroscopy. To study the influence of ϵ -lesions on the hybridization properties of RNA, melting curves of the duplexes containing the modified oligonucleotides ON 1 or ON 2 and the corresponding RNA or DNA complements were measured and compared with those of the corresponding natural duplexes containing oligoribonucleotides ON 3 and ON 4 (Table 2 and Figure 4).

The T_m analysis further underscores the drastic influence that the exocyclic lesions have on the hybridization properties of nucleic acids. Indeed, the loss of their ability to form standard Watson–Crick hydrogen bonds^{32,33} causes a major destabilization of the RNA–RNA duplexes with the cognate complementary base opposite the lesion [–15.6 $^{\circ}\text{C}$ in the case of ϵA and –18.0 $^{\circ}\text{C}$ in the case of ϵC (Table 2)]. The same trend can be observed in the case of RNA–DNA duplexes. Moreover, the T_m values are not further reduced when all the other nucleobases face the lesion, clearly showing that the natural base–lesion pair constitutes a mismatch, regardless of the nature of the base. In all cases, the duplexes formed with a G opposite the lesion are slightly more stable than the others, most likely because of stronger base stacking interactions.




Figure 4. T_m curves for duplexes of oligonucleotides containing a lesion (ON 1 and ON 2) or the corresponding natural base (ON 3 and ON 4) with r(UCGACUXAGCAU) or d(TCGACTXAGCAT). The duplex concentration was 2 μ M in 10 mM NaH_2PO_4 and 150 mM NaCl (pH 7.0).

To study the influence of these exocyclic base lesions on the structure of the duplexes, CD measurements were performed. ON 1 and ON 2 were hybridized with complementary RNA sequences, and the CD spectra of the resulting duplexes all showed a standard A-conformation (Figure 5) and do not deviate substantially from the corresponding unmodified duplexes (data not shown).

Primer–Template Extension Experiments. Further investigations of the coding properties of the ϵ -bases were

performed using standing-start template–primer extension reactions in a setup similar to that reported previously for other RNA lesions.^{27,34} The 5'-³²P-radiolabeled DNA primer, dNTPs, 31-mer templates containing the lesions, and three different RTs having different transcription fidelities (HIV1 RT, AMV RT, and MMLV RT) were used, and the corresponding gels are shown in Figures 6 and 7. Quantitative data of dNMP incorporation opposite the lesion are given in the Supporting Information.




Figure 5. CD spectra of duplexes containing ON 1 or ON 2 with r(UCGACUXAGCAU). The duplex concentration was 2 μ M (1:1 strand ratio) in 10 mM NaH₂PO₄ and 150 mM NaCl (pH 7.0) at 25 °C.




Figure 6. Autoradiograms of standing-start primer extension reactions for the RTs indicated using ON 5 containing the ϵ A lesion as the template and the corresponding natural template (ON 7). Reactions were performed at 37 °C for 1 h. The following enzyme amounts were used: 2 units of HIV RT, 8 units of AMV RT, and 40 units of MMLV RT. Lanes: ref, primer without enzyme; A, T, G, and C, reactions in the presence of the respective dNTP; N, reactions in the presence of all four dNTPs; Nat, unmodified RNA template and all four dNTPs.

When transcriptases AMV RT and MMLV RT were used, complete inhibition of reverse transcription was observed for both the ϵ A and ϵ C lesions. Even in the presence of all the natural dNTPs, no bands or only minor traces corresponding to elongated primer could be observed (Figures 6 and 7). On the other hand, a different behavior was observed in the case of HIV RT. This error-prone polymerase is able to read through both damages. In the case of the ϵ A-containing RNA template (ON 5), dAMP (78.3%) was the most efficiently incorporated nucleotide, followed by dGMP (71.3%). No significant incorporation of the base thymine (dTTP) could be observed (Figure 6). Thus, the order of preferential incorporation opposite ϵ A is as follows: A > G \gg C and T. In the presence of all dNTPs, full-length synthesis was observed, although it was slightly less efficient than with the undamaged template.

The results obtained using the ϵ C-containing RNA template (ON 6) are shown in Figure 7. In contrast with what was observed in the case of the ϵ A-containing template, HIV1 RT was completely unable to incorporate dGMP opposite ϵ C. As observed for ON 5, the most efficiently incorporated nucleotide was dAMP (79.2%). Moreover, the incorporation of dTTP was rather efficient (61.8%) but not as good as that of dAMP. The analysis of these results led to the following preferential order of incorporation opposite ϵ C: A > T \gg C > G. Again, in the presence of all the dNTPs, synthesis of the full-length product is observed and the lesion seems to have no inhibitory effect.

DISCUSSION

We describe here the chemical synthesis of oligoribonucleotides containing 1,N⁶-ethenoadenosine and 3,N⁴-ethenocytidine using standard phosphoramidite chemistry. Although the synthesis of such RNA oligonucleotides was previously reported,³⁰ to the best of our knowledge the pairing and coding properties of oligoribonucleotides containing such lesions have never been reported. From UV melting curve analysis, it appears that both etheno modifications completely lack recognition capacities for all natural bases. The slightly more stable arrangements with purine bases opposite the etheno lesions may be explained by more favorable stacking interactions. These results are in agreement with earlier findings for the hybridization properties of oligodeoxynucleotides containing ϵ dA or ϵ dC lesions.^{35–37} Even there, strong duplex destabilization against any natural base was observed. Again, in these cases, the most stable duplex resulted when a G was placed opposite the lesion,^{35,36} confirming




Figure 7. Autoradiograms of standing-start primer extension reactions for the RTs indicated using ON 6 containing the ϵ C lesion as a template and the corresponding natural template (ON 8). Reactions were performed at 37 °C for 1 h. The following enzyme amounts were used: 2 units of HIV RT, 8 units of AMV RT, and 40 units of MMLV RT. Lanes: ref, primer without enzyme; A, T, G, and C, reactions in the presence of the respective dNTP; N, reactions in the presence of all four dNTPs; Nat, unmodified RNA template and all four dNTPs.

the importance of (presumably interstrand) stacking interactions with respect to duplex stability.

The structure of the duplexes containing these lesions was investigated by CD spectroscopy. The results indicate that their presence does not affect the overall A-helix conformation of the duplexes. Thus, the expected local structural perturbations at the site of the base lesion are (not unexpectedly) invisible when assessed by CD spectroscopy. Clearly, more high-resolution structural work by NMR or X-ray crystallography is needed to reveal the precise arrangement of the modified bases.

The primer extension experiments nicely show the mutagenic potential of ϵ A and ϵ C during reverse transcription. While neither of the two etheno nucleosides investigated can be read by AMV or MMLV RT, leading to abortion of DNA synthesis, both of them are accepted by the more permissive HIV RT. However, depending on the nature of the lesion, the

coding preferences are somewhat different. Indeed, ϵ A preferentially leads to the incorporation of purine opposite itself with a slight advantage for dAMP (A-rule). This behavior has been observed in a similar experiment with an abasic site as a noncoding lesion in an RNA template.³⁴

Primer extension experiments using ϵ dA inserted in a DNA template were reported previously.³⁸ Surprisingly, from these experiments, we found that the only enzyme completely inhibited by the presence of this lesion is HIV RT. Quite interestingly, MMLV RT could elongate the DNA primer in different sequence contexts, although not to full length. This again is contrary to what we observed with the RNA template. It therefore seems that HIV RT shows a higher degree of sensitivity toward lesions on DNA than on RNA. This may imply that retroviral mutations are more likely to occur during DNA synthesis on the viral RNA template than during complementary DNA strand synthesis on the reverse-transcribed DNA strand.

Also in the case of ethenocytidine, there is precedence in the literature for its coding properties when present in a DNA template.³⁹ It needs to be noted though that these PEX were performed with DNA polymerases and not with reverse transcriptases. Compared with our results with HIV RT, lesion bypass synthesis is more strongly slowed for all of the DNA polymerases investigated in this study. The nature of the bases preferentially introduced opposite the lesion is highly dependent on the polymerase used. Indeed, Pol α incorporates dAMP and dTMP opposite the lesion, pol β dAMP and dCMP, pol η dAMP and dGMP, and pol κ dTMP.³⁹ Thus, the preferences of HIV1 incorporation in our experiments compare well to those of pol α . Again, we need to point out that these conclusions do not take into consideration variations in base preferences as a function of sequence context.

The mutagenicity of base lesions in DNA is a major threat to the genome integrity of living systems. In part, this is also true for base-damaged RNA in organisms, the genome of which consists of RNA (e.g., RNA viruses). Moreover, base damage to RNA can have additional biological implications. For example, it can deeply affect the cellular metabolism if noncoding RNAs involved in regulatory processes are affected. In addition, if present on mRNAs, such base lesions can lead to misincorporations of amino acids during translation (translational mutations), or alternatively, it can lead to ribosome arrest, which may severely impair cellular vitality.

■ ASSOCIATED CONTENT

§ Supporting Information

Synthetic and analytical details for phosphoramidites **1** and **2**, mass spectra for etheno-modified oligonucleotides, and quantitative data for dNMP incorporation for primer extension reactions. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Telephone: +41 31 631 4355. Fax: +41 31 631 3422. E-mail: leumann@ioc.unibe.ch.

Funding

This work was supported by the Swiss National Science Foundation (Grant 200020-130373) and by the University of Bern.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank Dr. Marcel Hollenstein and Dr. Pascal Küpfer for their helpful advice and critical reading of the manuscript.

REFERENCES

- (1) Speed, N., and Blair, I. A. (2011) Cyclooxygenase- and lipoxygenase-mediated DNA damage. *Cancer Metastasis Rev.* 30, 437–447.
- (2) Dechakhamphu, S., Pinlaor, S., Sitthithaworn, P., Bartsch, H., and Yongvanit, P. (2010) Accumulation of miscoding etheno-DNA adducts and highly expressed DNA repair during liver fluke-induced cholangiocarcinogenesis in hamsters. *Mutat. Res.* 691, 9–16.
- (3) Delaney, J. C., and Essigmann, J. M. (2008) Biological Properties of Single Chemical-DNA Adducts: A Twenty Year Perspective. *Chem. Res. Toxicol.* 21, 232–252.
- (4) Martinez, G. R., Loureiro, A. P. M., Marques, S. A., Miyamoto, S., Yamaguchi, L. F., Onuki, J., Almeida, E. A., Garcia, C. C. M., Barbosa, L. I. F., Medeiros, M. H. G., and Di Mascio, P. (2003) Oxidative and alkylating damage in DNA. *Mutat. Res.* 544, 115–127.
- (5) Blair, I. A. (2008) DNA Adducts with Lipid Peroxidation Products. *J. Biol. Chem.* 283, 15545–15549.
- (6) Cooke, M. S., Evans, M. D., Dizdaroglu, M., and Lunec, J. (2003) Oxidative DNA damage: Mechanisms, mutation, and disease. *FASEB J.* 17, 1195–1214.
- (7) Tolentino, J. H., Burke, T. J., Mukhopadhyay, S., McGregor, W. G., and Basu, A. K. (2008) Inhibition of DNA replication fork progression and mutagenic potential of 1,N⁶-ethenoadenine and 8-oxoguanine in human cell extracts. *Nucleic Acids Res.* 36, 1300–1308.
- (8) Jacobsen, J. S., and Humayun, M. Z. (1990) Mechanisms of Mutagenesis by the Vinyl Chloride Metabolite Chloroacetaldehyde. Effect of Gene-Targeted in Vitro Adduction of M 13 DNA on DNA Template Activity in Vivo and in Vitro. *Biochemistry* 29, 496–504.
- (9) Kusmierek, J. T., and Singer, B. (1982) Chloroacetaldehyde-Treated Ribo- and Deoxyribopolynucleotides. 1. Reaction Products. *Biochemistry* 21, 5717–5722.
- (10) Barrio, J. R., Sattangi, P. D., Gruber, B. A., Dammann, L. G., and Leonard, N. J. (1976) Species Responsible for the Fluorescence of 3,N⁴-Ethenocytidine. *J. Am. Chem. Soc.* 98, 7408–7413.
- (11) Secrist, J. A., Barrio, J. R., Leonard, N. J., and Weber, G. (1972) Fluorescent Modification of Adenosine-Containing Coenzymes. Biological Activities and Spectroscopic Properties. *Biochemistry* 11, 3499–3506.
- (12) Barrio, J. R., Secrist, J. A., and Sciences, N. J. (1972) Fluorescent Adenosine and Cytidine Derivatives. *Biochem. Biophys. Res. Commun.* 46, 597–604.
- (13) Jacobsen, J. S., Perkins, C. P., Callahan, J. T., Sambamurti, K., and Humayun, M. Z. (1989) Mechanisms of Mutagenesis by Chloroacetaldehyde. *Genetics* 121, 213–222.
- (14) Chung, F. L., Chen, H. J. C., and Nath, R. G. (1996) Lipid peroxidation as a potential endogenous source for the formation of exocyclic DNA adducts. *Carcinogenesis* 17, 2105–2111.
- (15) Zhang, W., Johnson, F., Grollman, A. P., and Shibutani, S. (1995) Miscoding by the Exocyclic and Related DNA Adducts 3,N⁴-Etheno-2'-deoxycytidine, 3,N⁴-Ethano-2'-deoxycytidine, and 3-(2-Hydroxyethyl)-2-deoxyuridine. *Chem. Res. Toxicol.* 8, 157–163.
- (16) El Ghissassi, F., Barbin, A., Nair, J., and Bartsch, H. (1995) Formation of 1,N⁶-Ethenoadenine and 3,N⁴-Ethenocytosine by Lipid Peroxidation Products and Nucleic Acid Bases. *Chem. Res. Toxicol.* 8, 278–283.
- (17) Maciejewska, A. M., Beata, S., Nowicki, A., and Kusmierek, J. T. (2011) The role of AlkB protein in repair of 1,N⁶-ethenoadenine in *Escherichia coli* cells. *Mutagenesis* 26, 401–406.
- (18) Levine, R. L., Yang, I.-Y., Hossain, M., Pandya, G. A., Grollman, A. P., and Moriya, M. (2000) Mutagenesis Induced by a Single 1,N⁶-

Ethenodeoxyadenosine Adduct in Human Cells. *Cancer Res.* 60, 4098–4104.

(19) Pandya, G. A., and Moriya, M. (1996) 1,N⁶-Ethenodeoxyadenosine, a DNA Adduct Highly Mutagenic in Mammalian Cells. *Biochemistry* 35, 11487–11492.

(20) Basu, A. K., Wood, M. L., Niedernhofer, L. J., Ramos, L. A., and Essigmann, J. M. (1993) Mutagenic and Genotoxic Effects of Three Vinyl Chloride-Induced DNA Lesions: 1,N⁶-Ethenoadenine, 3,N⁴-Ethenocytosine, and 4-Amino-5-(imidazol-2-yl)imidazole. *Biochemistry* 32, 12793–12801.

(21) Palejwala, V. A., Pandya, G. A., Bhanot, O. S., Solomon, J. J., Murphy, H. S., Dunman, P. M., and Humayun, M. Z. (1994) UVM, an Ultraviolet-inducible RecA-independent Mutagenic Phenomenon in *Escherichia coli*. *J. Biol. Chem.* 269, 27433–27440.

(22) Moriya, M., Zhang, W., Johnson, F., and Grollman, A. P. (1994) Mutagenic potency of exocyclic DNA adducts: Marked differences between *Escherichia coli* and simian kidney cells. *Proc. Natl. Acad. Sci. U.S.A.* 91, 11899–11903.

(23) Gros, L., Ishchenko, A. A., and Saparbaev, M. (2003) Enzymology of repair of etheno-adducts. *Mutat. Res.* 531, 219–229.

(24) Cui, S., Kim, Y.-H., Jin, C.-H., Kim, S. K., Rhee, M.-h., Kwon, O.-S., and Moon, B. J. (2009) Synthesis and base pairing properties of DNA-RNA heteroduplex containing 5-hydroxyuridine. *BMB Rep.* 42, 373–379.

(25) Kim, S. K., Kim, J. Y., Baek, A. K., and Moon, B. J. (2002) Base pairing properties of 8-oxo-7,8-dihydroadenosine in cDNA synthesis by reverse transcriptases. *Bioorg. Med. Chem. Lett.* 12, 1977–1980.

(26) Kim, S. K., Lee, S. H., Kwon, O.-S., and Moon, B. J. (2004) DNA-RNA heteroduplex containing 8-oxo-7,8-dihydroguanosine: Base pairing, structures, and thermodynamic stability. *J. Biochem. Mol. Biol.* 37, 657–662.

(27) Küpfer, P. A., and Leumann, C. J. (2011) Synthesis, base pairing properties and trans-lesion synthesis by reverse transcriptases of oligoribonucleotides containing the oxidatively damaged base 5-hydroxycytidine. *Nucleic Acids Res.* 39, 9422–9432.

(28) Shan, X., Chang, Y., and Lin, C. L. (2007) Messenger RNA oxidation is an early event preceding cell death and causes reduced protein expression. *FASEB J.* 21, 2753–2764.

(29) Tanaka, M., Chock, P. B., and Stadtman, E. R. (2007) Oxidized messenger RNA induces translation errors. *Proc. Natl. Acad. Sci. U.S.A.* 104, 66–71.

(30) Srivastava, S. C., Raza, S. K., and Misra, R. (1994) 1,N⁶-Etheno deoxy and ribo adenosine and 3,N⁴-etheno deoxy and ribo cytidine phosphoramidites. Strongly fluorescent structures for selective introduction in defined sequence DNA and RNA molecules. *Nucleic Acids Res.* 22, 1296–1304.

(31) Speina, E., Ciesla, J. M., Wojcik, J., Bajek, M., Kusmierek, J. T., and Tudek, B. (2001) The Pyrimidine Ring-opened Derivative of 1,N⁶-Ethenoadenine Is Excised from DNA by the *Escherichia coli* Fpg and Nth Proteins. *J. Biol. Chem.* 276, 21821–21827.

(32) Freisinger, E., Fernandes, A., Grollman, A. P., and Kisker, C. (2003) Crystallographic Characterization of an Exocyclic DNA Adduct: 3,N⁴-Etheno-2'-deoxycytidine in the Dodecamer 5'-CGCG-AATT1CGCG-3'. *J. Mol. Biol.* 329, 685–697.

(33) Leonard, G. A., McAuley-Hecht, K. E., Gibson, N. J., Brown, T., Watson, W. P., and Hunter, W. N. (1994) Guanine-1,N⁶-Ethenoadenine Base Pairs in the Crystal Structure of d(CGCGAAT-T(edA)GCG). *Biochemistry* 33, 4755–4761.

(34) Küpfer, P. A., Crey-Desbiolles, C., and Leumann, C. J. (2007) Trans-lesion synthesis and RNaseH activity by reverse transcriptases on a true abasic RNA template. *Nucleic Acids Res.* 35, 6486–6485.

(35) Seela, F., Schweinberger, E., Xu, K., Sirivolu, V. R., Rosemeyer, H., and Beckera, E.-M. (2007) 1,N⁶-Etheno-2'-deoxytubercidin and pyrrolo-C: Synthesis, base pairing, and fluorescence properties of 7-deazapurine nucleosides and oligonucleotides. *Tetrahedron* 63, 3471–3482.

(36) Gelfand, C. A., Plum, G. E., Grollman, A. P., Johnson, F., and Breslauer, K. J. (1998) The impact of an Exocyclic Cytosine Adduct on a DNA Duplex Properties: Significant Thermodynamic Consequences

Despite Modest Lesion-Induced Structural Alterations. *Biochemistry* 37, 12507–12512.

(37) Basu, A. K., Niedernhofer, L. J., and Essigmann, J. M. (1987) Deoxyhexanucleotide Containing a Vinyl Chloride Induced DNA Lesion, 1,N⁶-Ethenoadenine: Synthesis, Physical Characterization, and Incorporation into a Duplex Bacteriophage M13 Genome as Part of an Amber Codon. *Biochemistry* 26, 5626–5635.

(38) Litinski, V., Chenna, A., Sagi, J., and Singer, B. (1997) Sequence context is an important determinant in the mutagenic potential of 1,N⁶-ethenodeoxyadenosine (ϵ A): Formation of ϵ A basepairs and elongation in defined templates. *Carcinogenesis* 18, 1609–1615.

(39) Singer, B., Medina, M., Zhang, Y., Wang, Z., Guliaev, A. B., and Hang, B. (2002) 8-(Hydroxymethyl)-3,N⁴-etheno-C, a Potential Carcinogenic Glycidaldehyde Product, Miscodes In Vitro Using Mammalian DNA Polymerases. *Biochemistry* 41, 1778–1785.