


Structural Consequences of Mutations to the α -Tocopherol Transfer Protein Associated with the Neurodegenerative Disease Ataxia with Vitamin E Deficiency

Dennis Bromley,[†] Peter C. Anderson,^{†,§} and Valerie Daggett^{*,†,‡}

[†]Division of Biomedical and Health Informatics, Department of Biomedical Informatics and Medical Education, University of Washington, Seattle, Washington 98195, United States

[‡]Department of Bioengineering, University of Washington, Seattle, Washington 98195, United States

ABSTRACT: The α -tocopherol transfer protein (α -TTP) is a liver protein that transfers α -tocopherol (vitamin E) to very-low-density lipoproteins (VLDLs). These VLDLs are then circulated throughout the body to maintain blood α -tocopherol levels. Mutations to the α -TTP gene are associated with ataxia with vitamin E deficiency, a disease characterized by peripheral nerve degeneration. In this study, molecular dynamics simulations of the E141K and R59W disease-associated mutants were performed. The mutants displayed disruptions in and around the ligand-binding pocket. Structural analysis and ligand docking to the mutant structures predicted a decreased affinity for α -tocopherol. To determine the detailed mechanism of the mutation-related changes, we developed a new tool called ContactWalker that analyzes contact differences between mutant and wild-type proteins and highlights pathways of altered contacts within the mutant proteins. Taken together, our findings are in agreement with experiment and suggest structural explanations for the weakened ability of the mutants to bind and carry α -tocopherol.



α -Tocopherol transfer protein (α -TTP) transfers α -tocopherol, a strong antioxidant and a form of vitamin E, to very-low-density lipoproteins (VLDLs). These lipoproteins, located in the liver, are then circulated throughout the body to maintain blood α -tocopherol levels. Vitamin E occurs in multiple forms, including α -, β -, γ -, and δ -tocopherol. Each form has a hydrophobic tail with three chiral centers, and α -TTP preferentially binds RRR- α -tocopherol (Figure 1a).¹

Patients with mutations in the α -TTP gene have a weakened ability to transfer α -tocopherol to VLDLs and, as a result, are unable to maintain blood α -tocopherol levels. This condition, known as ataxia with vitamin E deficiency (AVED), is associated with peripheral nerve degeneration. Although the link between α -tocopherol deficiency and AVED-associated nerve damage is not fully understood, it is likely that α -tocopherol protects tissues from oxidative stress. Increasing the level of dietary consumption of vitamin E can be an effective treatment for AVED.²

α -TTP belongs to the SEC14-like family of proteins and is composed of two domains: an N-terminal three-helix bundle and a C-terminal CRAL-TRIO domain (Figure 1b). The latter domain is used for transporting hydrophobic molecules through hydrophilic environments; other SEC14-like proteins include human supernatant protein factor and cellular retinaldehyde-binding protein (CRALBP). The CRAL-TRIO domain has a ligand binding pocket, entry to which is controlled by a swinging “lid”. Investigations of SEC14 and CRALBP indicate that the lid is “hinged” by two regions of

residues: residues 242–244 and 263–265 for CRALBP³ and residues 212 and 213 and 239, 240, and 242 for Sec14.⁴ On the basis of sequence alignment of these proteins,³ these regions fall in the vicinity of residues 194–201 and 220–224 for α -TTP. These regions are termed hinge 1 (H1) and hinge 2 (H2), respectively (Figure 1b). In α -TTP, the lid is formed by residues 198–221,¹ the central structure of which is helix α 14.

Meier et al. crystallized α -TTP in both the lid-open (PDB entry 1OIZ) (Figure 1b) and lid-closed (PDB entry 1OIP) (Figure 1c) forms.¹ When α -TTP is not bound to α -tocopherol, the lid is open and the ligand binding pocket is accessible. When α -TTP is bound to α -tocopherol, the lid is closed and the binding pocket is separated from the external environment. The open and closed forms are overlaid in Figure 1d.

In this study, we investigated two mutants of α -TTP: one replaces a Glu with a Lys at position 141 (E141K), and the other replaces an Arg with a Trp at position 59 (R59W). Both mutants alter strongly conserved residues⁵ and are associated with severe clinical effects.^{1,6,7} Although experimental binding, ligand transfer, and urea denaturation data have been reported for these mutants,⁷ experimentally determined structures of these mutants are not available. Consequently, we have used all-atom molecular dynamics (MD) simulations to investigate the

Received: January 28, 2013

Revised: May 3, 2013

Published: May 29, 2013




Figure 1. Molecular structures of vitamin E and the different forms of α -TTP. (a) RRR- α -tocopherol (vitamin E). (b) In the unbound structure (PDB entry 1OIZ), a lid formed by the α 14 region is open, allowing entrance to a hydrophobic binding pocket. The CRAL-TRIO domain is indicated in brackets. The black and gray filled circles indicate the E141K and R59W mutation regions, respectively. The two hinge regions are circled and labeled as H1 and H2. (c) In the bound conformation (PDB entry 1OIP), the lid is closed, enclosing the binding pocket. (d) 1OIZ (blue) and 1OIP (green) overlaid and aligned by C α rmsd. The C α rmsd between the two structures excluding α 14 was 0.5 Å.

structural effects of these mutations. This protein has not been the subject of previous MD simulations to the best of our knowledge, but Ryan et al.⁴ and Schaff et al.⁸ have performed MD simulations of related Sec14 and Sfh1 proteins. In addition to standard structural analyses of the α -TTP simulations, docking calculations were also performed to assess the impact of mutations on the protein's ability to bind RRR- α -tocopherol.

Finally, to characterize how effects are propagated between the mutation site and the binding site, we created a new tool called ContactWalker. Analysis of these pathways provides a molecular description of the mutation-associated structural changes resulting from the E141K and R59W mutations.

METHODS

Structural Models. The 1.88 Å resolution crystal structure of α -TTP was obtained from the PDB⁹ (entry 1OIZ)¹ and was used for the starting structure for the wild-type and mutant simulations. The unbound, apo structure was used so we could investigate mutation-induced changes in the intrinsic structural and dynamic behavior of the protein and to avoid ligand-induced structural bias. Starting structures for both mutants were constructed by making the appropriate amino acid substitutions to the wild-type protein. The energies of the resulting structures were minimized for 100 steps *in vacuo* using the ENCAD simulation package¹⁰ and the Levitt et al. force field.¹¹ A different 1.95 Å resolution crystal structure of α -TTP (PDB entry 1OIP)¹ crystallized with bound α -tocopherol and the lid “closed” was used as a ligand docking baseline. The model of RRR- α -tocopherol used for docking studies was extracted from the 1OIP crystal structure.

Molecular Dynamics Simulations and Analysis. MD simulations were performed using the *in lucem* molecular mechanics (ilmm) software package¹² using a previously published protocol and potential energy function.^{11,13–15} Starting structures were prepared for MD using 1000 steps of steepest descent minimization and then solvated with flexible F3C water molecules¹⁴ in a periodic box with walls located at least 10 Å from all protein atoms. The solvent density was set to the experimentally determined value for water at 37 °C, 0.993 g/mL.¹⁶ The solvent energy was minimized for at least 500 steps before the solvent was heated again for 1 ps. The solvent energy was then minimized for an additional 500 steps and followed by an energy minimization of the entire system for 500 steps. Atomic velocities were assigned from a Maxwellian distribution at low temperatures and then brought to the target temperature of 37 °C (310 K). Thereafter, we used the NVE microcanonical ensemble, in which the box volume, number of particles, and total energy are fixed. A force-shifted nonbonded cutoff range of 10 Å was used for nonbonded interactions,¹⁷ and the interaction list was updated every two steps. Simulation steps were 2 fs, and the structures were saved every 1 ps. The temperature was set to 37 °C, and the simulations were performed at neutral pH (Asp and Glu negatively charged, Arg and Lys positively charged, and His neutral). Three independent simulations of the wild type and each mutant were performed for at least 51 ns. These simulations were performed as part of the SNP (single-nucleotide polymorphism) thrust of our Dynameomics project,^{18,19} an ongoing project to determine the native state dynamics and unfolding pathways of all known protein folds. α -TTP's three-helix bundle N-terminal domain is represented in fold rank 89 and the C-terminal CRAL-TRIO domain in fold rank 1251 in our 2009 Consensus Domain Dictionary^{20,21} (CDD). Protein images were created using Pymol²² and UCSF Chimera.²³

Inter-Residue Contacts. Inter-residue contact occupancies were calculated at 1 ps granularity using heavy-atom contacts from time steps greater than 25 ns. Contact distance thresholds were 5.4 Å for carbon–carbon contacts and 4.6 Å for all other heavy-atom contacts; residues were considered to be in contact

if at least one interatomic contact was within the appropriate threshold. Contacts between neighboring residues were ignored. The “residue occupancy difference” was calculated as the difference in occupancy between each wild-type residue and the corresponding mutant residue. As a result, negative values indicated greater prevalence in the mutant (mutant stabilization) and positive values reflected greater prevalence in the wild type (mutant destabilization).

ContactWalker. Because of the dynamic nature of proteins, interatomic contact distances change over time and even stable contacts oscillate around some mean distance. As a result, searching for structurally important contact changes between wild-type and mutant proteins can be challenging. To address this challenge, we built ContactWalker, a tool that measures the difference in interatomic contact occupancies between wild-type and mutant proteins.

ContactWalker calculates contact occupancy differences between a set of one or more wild-type simulations and a set of one or more mutant simulations. It uses the minimal demonstrated occupancy change between the wild-type and mutant simulations to avoid overestimating the significance of the change in occupancy. For example, the R54:Q145 contact had occupancy values of 24, 53, and 61% for the three wild-type simulations and 100, 100, and 98% for the three E141K simulations. The minimal demonstrated occupancy change between the wild type and the E141K mutant was -37% ($61\% - 98\%$). Using the minimal demonstrated occupancy change provided the most conservative measure of significant changes in the strength of protein contacts.

Contact Networks. A residue whose contacts have been altered, either by stabilization or by destabilization, is considered to be a disrupted residue. Preliminary observations found that disrupted residues tended to exist in connected networks rather than in isolation. ContactWalker can be used to visualize these networks, highlighting pathways of disrupted contacts (disruption pathways) and building on discretized, noninteractive network depictions such as those reported by Schmidlin et al.²⁴

To do this, ContactWalker builds a connected graph to represent all the contact changes between the wild-type and mutant proteins. Graph nodes represent residues, and graph edges represent nonbonded contacts with edge weights proportional to the change in contact occupancy. Backbone peptide bonds are inserted with an occupancy change value of zero.

Following construction of the contact graph, ContactWalker identifies the target residues. These are residues among which ContactWalker looks for networks of disrupted contacts. Residues can be manually specified if a region of interest is already known, or they can be automatically specified by selecting those residues with changes in contact occupancy that lie outside of some threshold. If the automatic method is used, additional manual targets may be specified as well.

Once the target residues have been identified, the user specifies the search parameters. These include options such as maximal search depth (the maximal number of contacts to search before trying another path) and a minimal edge value to determine if an edge is eligible for search. A cutoff in terms of absolute occupancy may also be specified. Backbone peptide bonds are always available for traversal.

Next, ContactWalker performs the actual search. A depth-first search (DFS) is performed between every pair of target nodes to find pathways of significant occupancy change. A DFS

is a graph-traversal algorithm that searches a graph by moving from parent node to child node, often limited to a maximal number of steps, before backing out and trying a new path. In this scenario, this limits the search to a maximal number of inter-residue contacts between significant residues.

In the final step, the output graph is laid out and rendered. The output graph is a subset of the original connectivity graph containing only those edges and nodes that meet the user's significance criteria. Graph images are rendered using the Graphviz graph layout tool²⁵ and the built-in scalable force directed placement graph layout algorithm.²⁶ The graph layout is determined by the connectivity patterns of significantly changed edges, not necessarily the proteins' three-dimensional structure. Approximate structural regions of the protein are highlighted on the graph. Edge colors indicate the degree of occupancy change along a green (mutant stabilized) to orange (mutant destabilized) spectrum, and edge size indicates the magnitude of the occupancy change. Node color designates secondary structure per the legend. ContactWalker also outputs a Pymol²² script to visualize the contact change data mapped onto the protein structure.

Ligand Docking. To predict the effect of distortions of the binding site on the binding affinity of α -tocopherol, we performed docking calculations of RRR- α -tocopherol on MD structures taken between 25 and 51 ns. Docking energies and conformational pose calculations were made using AutoDock Vina.²⁷ Of the 26000 structures between 26 and 51 ns, 260 (1%) were selected using simple random sampling without replacement. The sample mean, the 95% confidence interval (CI), and the sample standard deviation were calculated using a two-tailed t test, $\alpha = 0.05$, and either 260 or 780 degrees of freedom for individual simulation data or aggregated data, respectively. Docking energies are reported as the sample mean \pm the standard deviation. Additionally, a baseline docking energy was established by docking RRR- α -tocopherol into the 1OIP crystal structure.

α rmsd and α rmsf. α rmsd values were calculated using only α -carbons and excluded both termini (residues 9–23 and 266–274) and the α 14 region (residues 201–213). α rmsf data were calculated using MD structures beginning at 25 ns, used only α -carbons, and also excluded both termini. The α 14 region was included in the α rmsf calculations because visual inspection of the trajectories showed that the lid had closed by approximately 10 ns.

α -Tocopherol Headgroup-Coordinating Residues. The α -tocopherol headgroup-coordinating residues were defined as those residues with at least one atom within 5.4 Å of the headgroup of the vitamin E molecule. Water-coordinating residues were defined as residues with a nitrogen or oxygen atom within 4.6 Å of a water molecule. Calculations were performed using Pymol²² and the structure of PDB entry 1OIP¹ of α -TTP.

RESULTS AND DISCUSSION

In general, all simulations maintained the overall binding pocket structure and did not undergo significant unfolding; this is consistent with the urea denaturation experiments reported by Morley et al.⁷ that showed a 0.4 M change in C_{50} for the R59W and E141K mutants relative to that of the wild type (WT). The α rmsd of all three sets of simulations indicated that the proteins were relatively stable between 2 and 3 Å α rmsd, with the exceptions that each mutant had one simulation with α rmsd values of 4.0 Å (simulation 1 in each case)

(Figure 2). α rmsf analysis of all three proteins showed a similar trend (Figure 3); each protein demonstrated α rmsf




Figure 2. α rmsd values over time for all three proteins. Values were calculated using α -carbons excluding both termini and $\alpha 14$. (a) Wild type. (b) E141K. (c) R59W.

values around 1 Å in regions of secondary structure, and in each case, the simulation with higher α rmsd values also showed higher α rmsf values, often in the vicinity of $\alpha 13$ and $\alpha 14$.

α -Tocopherol docking studies resulted in a range of docking energies from -6.5 to -10.2 kcal/mol (Figure 4). The 95% confidence intervals for the mean potential energies were 0.1 and 0.0 kcal/mol for the per-simulation and aggregated docking data, respectively. For reference, the potential energy of docking α -tocopherol into the 1OIP closed structure was -10.8 kcal/mol, and the α rmsd between the 1OIZ and 1OIP crystal structures excluding $\alpha 14$ was 0.5 Å. Thus, this control suggests that the method can correctly identify and recover both the experimentally determined structure of the complex and its correct binding interactions. Per-simulation α -tocopherol docking analyses demonstrated that both wild-type and mutant proteins were capable of a range of docking energies (Figure 4a–c). In aggregate, the wild type and the E141K mutant had the same mean docking energy (-8.5 ± 0.7 and -8.5 ± 0.6 kcal/mol, respectively), and the R59W mutant performed less favorably (-8.1 ± 0.6 kcal/mol) (Figure 4d). Converting these potential energies to dissociation constants (K_d) resulted in values of 1000 nM for the wild type and E141K mutant and 1900 nM for the R59W mutant. This is approximately in keeping with the findings of Morley et al.,⁷ who reported 2- and 4-fold increases in K_d for the E141K and R59W mutants, respectively.

Wild-Type Simulations. All three wild-type trajectories stabilized between 2.5 and 3 Å α rmsd (Figure 2a), and simulation 2 displayed more structural fluctuation than the




Figure 3. α rmsf calculated using α -carbons and excluding both termini. (a) Black and gray bands indicate α -helices and β -sheets, respectively; blue squares indicate mutation sites, and red bands indicate hinge regions 1 and 2 (labeled H1 and H2, respectively). (b) Wild type. (c) E141K mutant. (d) R59W mutant.

other two (Figures 3b and 5). In all three wild-type simulations, $\alpha 14$ underwent a swinging motion from the lid-open position to the lid-closed position; Ryan et al.⁴ reported a similar conformational change in MD simulations of the related Sec14 protein. The β -sheet wall of the binding pocket remained stable in all simulations, while the α -helix wall exhibited more dynamic behavior (Figures 3b and 5). Despite these similarities, the three wild-type simulations each behaved slightly differently; to better characterize the range of behaviors of the wild-type protein, we analyzed each simulation separately.

As shown in Figure 5a, structures from simulation 1 did not deviate far from the minimized crystal structure with the exceptions of the $\alpha 3$ – $\alpha 4$ loop, $\alpha 5$, $\alpha 7$, $\alpha 13$, N227, and the $\alpha 17$ – $\alpha 18$ loop. Simulation 1 also provided the most favorable docking conformations of the three wild-type simulations with a mean potential energy of -9 ± 0.5 kcal/mol (Figure 4a). Because of its stability and favorable docking performance, wild-type simulation 1 was used as a structural baseline for subsequent contact comparisons.

Simulation 2 was more dynamic than either simulation 1 or simulation 3, particularly near $\alpha 14$ (Figures 3b and 5b), and the mean docking energy (-8.4 ± 0.6 kcal/mol) was between the energies of the other two simulations (Figure 4a). Contact occupancy comparisons of simulations 1 and 2 indicated that the structures in simulation 2 were missing several binding




Figure 4. Histograms of α -tocopherol docking potential energies. AutoDock Vina was used to dock α -tocopherol into 260 randomly selected structures from each simulation. Mean potential energies and standard deviations are shown in the legends. (a–c) Per-simulation histograms of the wild type, E141K mutant, and R59W mutant, respectively. (d) Aggregated energies for each protein.

pocket contacts that were present in simulation 1, including E82:K178 (100% difference) and I83:K178 (66% difference) contacts (Figure 5b). These two contacts tethered $\alpha 5$ to $\alpha 13$ and helped maintain the structure of the α -helix wall of the binding pocket. Both E82 and I83 established alternative stabilizing contacts, including 100% occupancy contacts with R134. K178 did not establish significant alternative stabilizing contacts and remained destabilized relative to simulation 1 (1.5 Å α rmsf in simulation 2 vs 0.75 Å α rmsf in simulation 1). Structures from simulation 2 showed, relative to simulation 1, a general lack of contacts between $\alpha 12$ – $\alpha 14$ lid and hinge regions. Of the 19 contacts in this region with occupancy differences of >50%, six were stabilized and 13 were destabilized. This difference in occupancy is reflected in the increased mobility and fluctuation of $\alpha 14$ in simulation 2 (Figures 3b and 5b). The internal structure of $\alpha 14$ saw a similar destabilization; of the 13 contacts within $\alpha 14$ with occupancy differences of >50%, three contacts were stabilized and ten contacts were destabilized.




Figure 5. Representative wild-type structures rendered as tubes. The tube radius and color reflect the α rmsd (left) and α rmsf (right). E82, I83, and K178 are colored per the legend and rendered as spheres. Simulations 1 (a), 2 (b), and 3 (c). Note the separation of K178 from E82 and I83 in simulation 2.

Simulation 3 had relatively low α rmsf values, similar to those of simulation 1 (Figures 3b and 5c), but it also had the least favorable docking energy of the three wild-type simulations (-8.1 ± 0.5 kcal/mol) (Figure 4a). In total, 27 contacts had occupancy differences of >99% relative to simulation 1, the majority of which were in the vicinity of the hinge and lid region formed by $\alpha 12$ – $\alpha 14$. Also, like simulation 2, the I83:K178 contact (66% occupancy) was not present in simulation 3. However, unlike simulation 2, the E82:K178 contact was intact, retaining some of the $\alpha 13$ tethering. Furthermore, also unlike simulation 2, K178 formed alternative stabilizing contacts with L183 (93%). Thus, K178 was not completely disconnected from E82 and I83, nor was it left in a destabilized state, resulting in slightly lower α rmsf values (1.1 Å vs 1.5 Å in simulation 2).

E141K Simulations. Overall, the E141K simulations were fairly stable and did not undergo any significant unfolding. As indicated by the α rmsd calculations (Figure 2b), simulations 2 and 3 stabilized between 2 and 2.5 Å, while simulation 1 deviated from the starting structure by as much as 4 Å. Consistent with the α rmsd data, the α rmsf data indicated that simulation 1 of the E141K mutant was more dynamic than simulation 2 or 3 (Figures 3c and 6).




Figure 6. Representative E141K structures rendered as tubes. The tube radius and color reflect the Ca rmsd (left) and Ca rmsf (right). The K141 mutation site is rendered as magenta spheres. Simulations 1 (a), 2 (b), and 3 (c).

Comparing contact occupancies of all three E141K simulations against those of the wild-type simulation 1 baseline, we found 15 contacts with a minimal demonstrated occupancy change of >95%. Of these, three were stabilizations of the hinge 2 region (K217:R221, K217:E220, and E216:E220), five were destabilizations that included the lid (F213:K217, F213:E216, L214:K217, P212:E216, and K211:T215), one tethered $\alpha 13$ to $\alpha 14$ (T184:T215), and one untethered $\alpha 5$ from $\alpha 13$ (W76:S186). All 10 of these contact changes were also observed in the wild-type simulations. However, their presence in the E141K mutant was both more common (all three E141K simulations) and, in the case of the hinge 2 region and $\alpha 13$ – $\alpha 14$ tethering, larger in magnitude (Table 1).

To isolate the specific effects of the E141K mutation from the inherent propensities of the protein, we compared all three E141K simulations against all three wild-type simulations. Only nine contacts were observed that had a minimal demonstrated occupancy difference of >20% (Table 2). Of these, four of the nine contacts were in or near hinge 2. Disruption pathways between the mutation site and the hinge 2 region (Figure 7) were found by increasing the sensitivity of ContactWalker (significance threshold of 9%, allowable threshold of 6%, search depth of 2, absolute occupancy cutoff of 0.01). The primary contact change at the mutation site was the R54:K141 contact

Table 1. Occupancy Differences in Contacts Common to E141K and the Wild Type

residue 1	residue 2	WT1 vs WT2 ^a	WT1 vs WT3 ^a	WT1 vs E141K (all) ^a
W76	S186	68%	97%	97%
K211	T215	96%	97%	95%
P212	E216	99%	99%	98%
F213	E216	100%	100%	100%
F213	K217	100%	100%	100%
L214	K217	100%	98%	100%
T184	T215	–43%	–49%	–96%
K217	E220 ^b	–24%	–4%	–99%
K217	R221 ^b	–86%	–57%	–99%
E216	E220 ^b	–20%	(0% in both WT1 and WT3)	–100%

^aNegative numbers indicate stabilization in the compared structure (underlined). ^bHinge 2 region.

Table 2. E141K Occupancy Change Magnitudes of >20%

residue 1	residue 2	occupancy difference
E216	E220	–80%
K217	E220	–75%
A10	D60	–53%
T184	T215	–47%
R54	Q145	–37%
L183	I218	–37%
A10	F61	–27%
R77	L137	73%
R54	K141	92%

(92% loss). As indicated in the figure, the disruption pathway included $\alpha 5$, $\alpha 10$, $\alpha 13$, and the hinge 2 region.

Zhang et al. reported that membrane binding is dependent on several hydrophobic residues, including F165 and F169.²⁸ While these residues did not experience appreciable contact disruption, they did have average Ca rmsf and Ca rmsd values different from those of WT; the Ca rmsd values were 2.9 and 2.6 Å in WT and 1.9 and 1.9 Å in E141K for residues F165 and F169, respectively, and the Ca rmsf values for F169 were 1.4 Å in WT and 1.2 Å in E141K.

Several headgroup-coordinating residues were disrupted: F133, L137, V182, L183, F187, and L198 (Figure 7). Although docking studies with MD structures did not indicate significant disruption to α -tocopherol binding for this mutant, these contact changes may be indicative of binding pocket disruptions that take place over time scales longer than those currently available with molecular dynamics. Also, the 1OIP crystal structure shows four crystallographic water molecules involved with the α -tocopherol headgroup binding. Three of the disrupted headgroup-coordinating residues, F133, V182, and F187, also help to coordinate these waters. Data reported by Schaaf et al.⁸ suggest that water coordination within the analogous hydrophobic ligand binding pocket of the related Sec14 and Sfh1 proteins may be important to the functioning of those proteins.

In addition to possible binding site disruption, these results indicate that a structural effect of the E141K mutation was to disrupt the hinge 2 region. This is reflective of the *in silico* findings of Ryan et al. that the G266D mutant of the related Sec14 protein disrupts hinge function,⁴ although we did not observe significant disruption to the gating module reported in that same study. The ubiquity of these disruptions within the




Figure 7. Contact disruption pathways in the E141K mutant. Contact occupancy differences were calculated between all wild-type simulations and all E141K simulations with a significance threshold of 0.09, an allowable traversal threshold of 0.06, and a search depth of 2. (a) ContactWalker disruption pathways. The thickness of the line indicates the magnitude of the occupancy change; green lines indicate mutant-associated stabilization, and orange lines indicate mutant-associated destabilization. Nodes are colored per the legend. The black line highlights a pathway between the mutation site and the hinge 2 region. (b) The 51 ns structure of E141K simulation 1. The black line indicates the same pathway that is labeled in panel a. Pathway residues are rendered as sticks and labeled for orientation. The K141 mutation site is rendered as magenta spheres. The final residue of the pathway (E220) is rendered as orange spheres.

mutant simulations suggests an explanation for part of the increase in K_d relative to that of the wild type reported by Morley et al.;⁷ the destabilization caused by the E141K mutation was enough to disrupt the hinge and lid region of the protein, but not enough to severely disrupt the actual α -tocopherol binding pocket (although, as discussed above, some loss of binding ability may be attributable to pocket disruption). As a result, the E141K *in silico* docking energies were almost identical to those of the wild type, while the *in vitro* experiments, which were dependent on the wild-type-like gate and hinge behavior, demonstrated an increase in K_d .

R59W Simulations. The R59W mutant simulations, like the E141K simulations, did not undergo significant unfolding. Simulations 2 and 3 were relatively stable with $C\alpha$ rmsd values stabilizing between 2.5 and 3 Å (Figure 2c). Simulation 1, however, was considerably more dynamic with $C\alpha$ rmsd values as high as 4.5 Å and $C\alpha$ rmsf values often twice those of the other simulations (Figure 3d).

Comparison of all three R59W simulations against the wild-type simulation 1 baseline revealed four disrupted contacts in the lid region: K211:T215 (83%), P212:E216 (86%), F213:E216 (100%), and F213:K217 (100%). Four contacts that tethered $\alpha 13$ in place were also destabilized: W76:F187 (58%), Y73:P188 (83%), W59:T184 (87%), and W76:S186 (91%).

Comparison of contact occupancies between all three wild-type simulations and all three R59W simulations resulted in low minimal demonstrated occupancy change values. Only 18 contacts demonstrated minimal occupancy changes of >20%. Of these, only six contacts exhibited occupancy change values of >50%. Notably, the W59 mutation site did not see significant occupancy change; in the aggregate, the largest minimal demonstrated occupancy change for a W59 contact was a 6% contact occupancy increase with R54.

R59W simulations 1 and 3, considered on their own, exhibited less favorable α -tocopherol docking energies than simulation 2 (Figures 4 and 8). They were also structurally similar to wild-type simulation 2, displaying a characteristic separation of $\alpha 13$ from $\alpha 5$ and $\alpha 10$ (Figure 5b). Contact comparison of only these two simulations against wild-type simulation 1 revealed additional disruptions to T184:K217 (98% destabilization), L214:K217 (100% destabilization), and K217:R221 (99% stabilization) contacts. The T184:K217 and L214:K217 contact disruptions were also present in the poorer-docking wild-type simulations and in the E141K simulations. Contact analysis also revealed stabilization in the hinge 2 region, similar to the E141K hinge 2 stabilizations.

Given these data, we took R59W simulation 1 and 3 conformations to be representative of mutation-induced disruptions. To isolate the structural disruptions associated




Figure 8. Representative R59W structures rendered as tubes. The tube radius and color reflect Ca rmsd (left) and Ca rmsf (right). The W59 mutation site is rendered as magenta spheres. Simulations 1 (a), 2 (b), and 3 (c).

with weakened docking performance, R59W simulations 1 and 3 were compared against R59W simulation 2; 50 contacts with

>90% occupancy change were identified, of which 72% were destabilized (Figure 9). Contacts responsible for holding $\alpha 13$ in place were destabilized, including E82:K178 (100%), E141:F187 (99%), Y73:S186 (100%), N72:D185 (99%), W76:V182 (97%), N72:S186 (96%), and F133:L183 (91%) contacts. Of the stabilized contacts, three contacts strengthened the connection between the C-terminus of $\alpha 13$ and the N-terminal three-helix bundle: W59:F187 (93%), L65:S186 (93%), and A58:L189 (98%). Notably, one of these (W59:F187) included the mutated residue. Furthermore, the hinge 1 region contained three significant destabilizing interactions [E199:N227 (91%), P200:H225 (95%), and I197:P200 (94%)]; the hinge 2 region contained one significant stabilizing interaction [E216:E220 (94%) as well as others in the near vicinity], and the $\alpha 14$ lid contained one significant destabilization [K211:E216 (97%)]. The gating module reported by Ryan et al.⁴ was disrupted only in the H101:S136 contact.

The R59W mutation strengthened the tendency of the protein to adopt conformations that were unfavorable for ligand docking; relative to the wild-type simulation 1 baseline, the lid and hinge regions were disrupted and $\alpha 13$ became disrupted more often in the R59W simulations than in the wild-type simulations. Furthermore, the *in silico* docking was less favorable than that of the wild type, reflecting structural changes to the binding pocket in addition to disruptions to the lid and hinge regions. Like the E141K mutation, the R59W mutation disrupted several headgroup-coordinating residues, including F133, S136, V182, L183, F187, and L189 (Figure 9). All of these residues except L183 also help to coordinate crystallographic waters in the ligand binding pocket of the 1OIP (closed) crystal structure.

Finally, like the E141K mutant, the R59W mutant did not cause significant contact disruption to F165 and F169, the residues reported by Zhang et al.²⁸ to be important to membrane binding. These simulations did, however, show a lower average Ca rmsd relative to the starting structure than did the WT residues (2.9 and 2.6 Å in the WT vs 2.3 and 2.2 Å in R59W for residues F165 and F169, respectively). Also, F165




Figure 9. Contact disruption pathways in the R59W mutant. The contact occupancy difference was calculated between simulation 2 and simulations 1 and 3 with a significance threshold of 0.9, an allowable traversal threshold of 0.9, and a search depth of 1. (a) ContactWalker disruption pathways. The thickness of the line indicates the magnitude of the occupancy change; green lines indicate stabilization of the mutant, and orange lines indicate mutant destabilization. Nodes are colored per the legend. The dotted circles highlight destabilized $\alpha 13$ residues. (b) The 51 ns structure of R59W simulation 1. Significant nodes are rendered as sticks and colored on the basis of the overall stability (green) or instability (orange). Note the prevalence of destabilized residues among $\alpha 5$, $\alpha 10$, and $\alpha 13$ coincident with the expansion of the ligand binding pocket.

had a larger average $C\alpha$ rmsf in the R59W simulations (1.2 Å in WT vs 1.5 Å in R59W).

CONCLUSIONS

In aggregate, the mutants had very few large occupancy changes relative to the wild type, indicating that their observed behaviors were already present in the wild type. The primary effect of both mutations was to exacerbate preexisting tendencies; in the case of the E141K mutant, we observed changes in the hinge 2 region. In the case of the R59W mutant, we observed a widening of the ligand binding pocket. In both cases, the mutant proteins remained folded and relatively stable, indicating that these mutants' weakened α -tocopherol binding is due to a shift in the population with an increase in conformers with altered binding regions.

Genetic diseases can have severe clinical effects while causing only subtle changes to protein structure. As demonstrated by the E141K and R59W α -TTP mutations, the cause of these disruptions can be masked by the dynamic propensities of the protein. In the case of the E141K mutant, finding that the hinge was disrupted with only minor disruptions to the actual ligand binding pocket highlights the complexity and subtlety of protein dynamics. Occupancy comparison was a useful tool for characterizing these dynamic systems because we were able to establish a wild-type baseline against which we could compare mutation-associated structural changes. We were also able to characterize large-scale structural changes without losing single-contact data resolution. Visualization of these contact changes, together with docking studies and traditional analyses, facilitated the exploration of gigabytes of MD data and provided both quantitative and qualitative accounts of the changes that occurred as a result of mutations to α -TTP. These findings offer a structural description of the E141K and R59W mutants and suggest a molecular basis for their attenuated ability to transport α -tocopherol.

AUTHOR INFORMATION

Corresponding Author

*E-mail: daggett@uw.edu. Phone: (206) 543-9305.

Present Address

[§]P.C.A.: Sandia National Laboratories, Box 969, MS 9291, Livermore, CA 94551-0969.

Funding

We are grateful for computer resources provided through the U.S. Department of Energy Office of Biological and Environmental Research as provided by the National Energy Research Scientific Computing Center, which is supported by the Office of Science of the U.S. Department of Energy under Contract DE-AC02-05CH11231. Financial support was also provided by the National Institutes of Health (Grant GM50789 to V.D. and TG 3 T15 LM007442-04S1 to P.C.A. and D.B.).

Notes

The authors declare no competing financial interests.

ABBREVIATIONS

PDB, Protein Data Bank; rmsf, root-mean-square fluctuation; rmsd, root-mean-square deviation.

REFERENCES

(1) Meier, R., Tomizaki, T., Schulze-Bries, C., Baumann, U., and Stocker, A. (2003) The molecular basis of vitamin E retention:

Structure of human α -tocopherol transfer protein. *J. Mol. Biol.* 331, 725–734.

(2) Di Donato, I., Bianchi, S., and Federico, A. (2010) Ataxia with vitamin E deficiency: Update of molecular diagnosis. *Neurol. Sci.* 31, 511–515.

(3) Liu, T., Jenwitheesuk, E., Teller, D. C., and Samudrala, R. (2005) Structural insights into the cellular retinaldehyde binding protein (CRALBP). *Proteins* 61, 412–422.

(4) Ryan, M. M., Temple, B. R. S., Phillips, S. E., and Bankaitis, V. A. (2007) Conformational Dynamics of the Major Yeast Phosphatidylinositol Transfer Protein Sec14p: Insight into the Mechanisms of Phospholipid Exchange and Diseases of Sec14p-Like Protein Deficiencies. *Mol. Biol. Cell* 18, 1928–1942.

(5) Panagabko, C., Morley, S., Hernandez, M., Cassolato, P., Gordon, H., Parsons, R., Manor, D., and Atkinson, J. (2003) Ligand specificity in the CRAL-TRIO protein family. *Biochemistry* 42, 6467–6474.

(6) Cavalier, L., Ouahchi, K., Kayden, H. J., Di Donato, S., Reutenauer, L., Mandel, J.-L., and Koenig, M. (1998) Ataxia with Isolated Vitamin E Deficiency: Heterogeneity of Mutations and Phenotypic Variability in a Large Number of Families. *Am. J. Hum. Genet.* 62, 301–310.

(7) Morley, S., Panagabko, C., Shineman, D., Mani, B., Stocker, A., Atkinson, J., and Manor, D. (2004) Molecular Determinants of Heritable Vitamin E Deficiency. *Biochemistry* 43, 4143–4149.

(8) Schaaf, G., Dynowski, M., Mousley, C. J., Shah, S. D., Yuan, P., Winklbauer, E. M., de Campos, M. K. F., Trettin, K., Quinones, M.-C., Smirnova, T. I., Yanagisawa, L. L., Ortlund, E. A., and Bankaitis, V. A. (2011) Resurrection of a functional phosphatidylinositol transfer protein from a pseudo-Sec14 scaffold by directed evolution. *Mol. Biol. Cell* 22, 892–905.

(9) Bernstein, F. C., Koetzle, T. F., Williams, G. J., Meyer, E. F., Brice, M. D., Rodgers, J. R., Kennard, O., Shimanouchi, T., and Tasumi, M. (1977) The protein data bank: A computer-based archival file for macromolecular structures. *J. Mol. Biol.* 112, 535–542.

(10) Levitt, M. (1990) *ENCAD-Energy calculations and dynamics*, Molecular Applications Group, Stanford University, Palo Alto, CA, and Yeda, Rehovot, Israel.

(11) Levitt, M., Hirshberg, M., Sharon, R., and Daggett, V. (1995) Potential energy function and parameters for simulations of the molecular dynamics of proteins and nucleic acids in solution. *Comput. Phys. Commun.* 91, 215–231.

(12) Beck, D. A. C., Alonso, D. O. V., and Daggett, V. (2000–2013) *ilmm*, in *luem Molecular Mechanics*, University of Washington, Seattle.

(13) Beck, D. A. C., and Daggett, V. (2004) Methods for molecular dynamics simulations of protein folding/unfolding in solution. *Methods* 34, 112–120.

(14) Levitt, M., Hirshberg, M., Sharon, R., Laidig, K. E., and Daggett, V. (1997) Calibration and Testing of a Water Model for Simulation of the Molecular Dynamics of Proteins and Nucleic Acids in Solution. *J. Phys. Chem. B* 101, 5051–5061.

(15) Armen, R. S., Bernard, B. M., Day, R., Alonso, D. O., and Daggett, V. (2005) Characterization of a possible amyloidogenic precursor in glutamine-repeat neurodegenerative diseases. *Proc. Natl. Acad. Sci. U.S.A.* 102, 13433–13438.

(16) Kell, G. S. (1967) Precise representation of volume properties of water at one atmosphere. *J. Chem. Eng. Data* 12, 66–69.

(17) Beck, D. A., Armen, R. S., and Daggett, V. (2005) Cutoff size need not strongly influence molecular dynamics results for solvated polypeptides. *Biochemistry* 44, 609–616.

(18) Beck, D. A. C., Jonsson, A. L., Schaeffer, R. D., Scott, K. A., Day, R., Toofanny, R. D., Alonso, D. O., and Daggett, V. (2008) Dynameomics: Mass annotation of protein dynamics and unfolding in water by high-throughput atomistic molecular dynamics simulations. *Protein Eng., Des. Sel.* 21, 353–368.

(19) Van der Kamp, M. W., Schaeffer, R. D., Jonsson, A. L., Scouras, A. D., Simms, A. M., Toofanny, R. D., Benson, N. C., Anderson, P. C., Merkley, E. D., Rysavy, S., Bromley, D., Beck, D. A. C., and Daggett, V. (2010) Dynameomics: A comprehensive database of protein dynamics. *Structure* 18, 423–435.

- (20) Day, R., Beck, D. A. C., Armen, R. S., and Daggett, V. (2003) A consensus view of fold space: Combining SCOP, CATH, and the Dali Domain Dictionary. *Protein Sci.* 12, 2150–2160.
- (21) Schaeffer, R. D., Jonsson, A. L., Simms, A. M., and Daggett, V. (2011) Generation of a consensus protein domain dictionary. *Bioinformatics* 27, 46–54.
- (22) DeLano, W. L. (2002) *The PyMOL molecular graphics system*, DeLano Scientific, San Carlos, CA.
- (23) Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C., and Ferrin, T. E. (2004) UCSF Chimera: A visualization system for exploratory research and analysis. *J. Comput. Chem.* 25, 1605–1612.
- (24) Schmidlin, T., Kennedy, B. K., and Daggett, V. (2009) Structural changes to monomeric CuZn superoxide dismutase caused by the familial amyotrophic lateral sclerosis-associated mutation A4V. *Biophys. J.* 97, 1709–1718.
- (25) Gansner, E. R., and North, S. C. (2000) An open graph visualization system and its applications to software engineering. *Software—Practice and Experience* 30, 1203–1233.
- (26) Hu, Y. (2005) Efficient, high-quality force-directed graph drawing. *Mathematica Journal* 10, 37–71.
- (27) Trott, O., and Olson, A. J. (2010) AutoDock Vina: Improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *J. Comput. Chem.* 31, 455–461.
- (28) Zhang, W. X., Thakur, V., Lomize, A., Pogozheva, I., Panagabko, C., Cecchini, M., Baptist, M., Morley, S., Manor, D., and Atkinson, J. (2010) The contribution of surface residues to membrane binding and ligand transfer by the α -tocopherol transfer protein (α -TTP). *J. Mol. Biol.* 405, 972–988.