

α -Synuclein Misfolding Assessed with Single Molecule AFM Force Spectroscopy: Effect of Pathogenic Mutations

Alexey V. Krasnoslobodtsev,[†] Ivan L. Volkov,^{†,§} Josephat M. Asiago,[‡] Jagadish Hindupur,[‡] Jean-Christophe Rochet,[‡] and Yuri L. Lyubchenko^{*,†}


[†]Department of Pharmaceutical Sciences, University of Nebraska Medical Center, 986025 Nebraska Medical Center, Omaha, Nebraska 68198, United States

[‡]Department of Medicinal Chemistry and Molecular Pharmacology, Purdue University, West Lafayette, Indiana 47907-2091, United States

[§]Department of Physics, Saint-Petersburg State University, Saint-Petersburg 198504, Russia

S Supporting Information

ABSTRACT: Misfolding and subsequent aggregation of alpha-synuclein (α -Syn) protein are critically involved in the development of several neurodegenerative diseases, including Parkinson's disease (PD). Three familial single point mutations, A30P, E46K, and A53T, correlate with early onset PD; however, the molecular mechanism of the effects of these mutations on the structural properties of α -Syn and its propensity to misfold remains unclear. Here, we address this issue utilizing a single molecule AFM force spectroscopy approach in which structural details of dimers formed by all four variants of α -Syn are characterized. Analysis of the force spectroscopy data reflecting contour length distribution for α -Syn dimer dissociation suggests that multiple segments are involved in the assembly of the dimer. The interactions are not limited to the central nonamyloid-beta component (NAC) of the protein but rather expand beyond this segment. All three mutations alter the protein's folding and interaction patterns affecting interactions far beyond their immediate locations. Implementation of these findings to our understanding of α -Syn aggregation pathways is discussed.



The aggregation of α -synuclein (α -Syn) is a hallmark of several neurodegenerative diseases including Parkinson's disease. α -Syn is an abundant presynaptic protein that belongs to a group of natively unfolded proteins lacking secondary/tertiary structure in aqueous solutions.^{1,2} Although unstructured, some preferences for adopting α -helical structure have been observed in α -synuclein primarily within the N-terminal domain and the nonamyloid- β component (NAC) region when the protein binds to lipids of membranes and vesicles.³ Structural transitions of α -Syn are also involved in the formation of amyloid-like fibrils with a high content of β -sheet secondary structure *in vivo* and under a variety of conditions *in vitro*.^{4–7}

Various studies established that the aggregation rate of α -Syn depends on environmental conditions and is accelerated by various factors such as low pH and the presence of biogenic polyamines.^{8–14} Changes in primary sequence associated with early onset familial Parkinson's (A30P,¹⁰ E46K,¹² and A53T¹¹ single point mutants) also enhance the rate of protein aggregation compared to that of the wild-type. The pathway of aggregation, however, was found to differ for each mutant. Whereas, E46K and A53T aggregate more rapidly than WT with the formation of fibrils,^{15,16} A30P assembles primarily in oligomeric forms.^{17,18} Unraveling the effects of these factors on

the misfolding of α -Syn is important for understanding the molecular basis of disease development.

The main challenge in studying protein misfolding phenomena is that misfolded conformations that lead to the pathological aggregation of a protein are only transiently populated. The ensemble for some proteins consists of rapidly interconverting compact conformations indicating the highly dynamic nature of protein conformational space.¹⁹ It was recently demonstrated that the aggregation propensity of α -Syn depends on how fast the protein accesses aggregation-prone conformations, a process termed reconfiguration rate.^{20–22} A monomeric protein must undergo structural reorganization to adopt transient conformational states (misfolded conformations) in order to initiate aggregation.²³ Interactions between misfolded proteins facilitate the very first step in the aggregation process, dimerization. Subpopulation of misfolded states in the overall ensemble can be very small. More importantly, it is not obvious which of the many species in the conformational space lead to pathological aggregation. Increasing evidence suggests that α -Syn populates an ensemble

Received: July 31, 2013

Revised: September 24, 2013

Published: September 25, 2013

of conformers ranging from extended to highly collapsed forms in aqueous solutions.^{24,25} Conditions that favor more compact structures have been reported to significantly increase aggregation rates of α -Syn.^{26–28} Ensemble-based (bulk) methods, however, cannot access a wealth of information that a heterogeneous ensemble of α -synuclein conformations might offer with respect to the aggregation process.

Single-molecule techniques can characterize properties of these individual molecules without contribution from the rest of the ensemble. Recently, AFM single-molecule force spectroscopy (SMFS) has been applied to study the conformational diversity of monomeric α -Syn.^{27,28} The study identified three main distinct conformations that exist within monomeric α -synuclein based on their resistance to mechanical pulling. These conformations were classified as random coil, mechanically weak, and β -like structures. The latter were proposed to be directly related to the aggregation of α -Syn.²⁸ Interestingly, pathological conditions which are known to increase the propensity of α -synuclein to aggregate, such as high ionic strength and the presence of Cu^{2+} , as well as mutations A30P, A53T, and E46K,²⁷ shifted the conformational equilibrium toward these β -like structures. We have approached the misfolding problem differently. We have used a recently proposed novel method to characterize misfolded states in dimeric forms of amyloidogenic proteins rather than monomeric proteins (reviewed in ref 29). In this approach, interactions between proteins immobilized on an AFM tip and the probed surface are measured with force spectroscopy with significant rupture forces, suggesting that misfolded proteins form dimeric states. Therefore, instead of probing the whole conformational space of a protein we probe only conformers that are capable of forming dimers, the simplest aggregated species. The advantage of this methodology is, therefore, that it probes conformers that associated with one another at the single-molecule level. We have recently applied this methodology to characterize the effect of spermidine on α -Syn misfolding.¹⁴

Here, we report an AFM force spectroscopy study of α -Syn folding and the effect of single point mutations of α -Syn on the structure of dimeric contact. Pair-wise interactions were probed between individual α -Syn molecules at low pH conditions that induce conformational transitions associated with enhanced aggregation. We applied the rupture length analysis to identify segments of α -Syn involved in the formation of dimers. This analysis showed that only a limited span of the C-terminal region of wild-type α -Syn is involved in the stabilization of dimers. Importantly, the single-point mutations located within the N-terminal region of the protein dramatically change the pattern of protein misfolding by extending interactions into the C-terminal region. These results suggest that the single point mutations affect distantly located interacting segments of the misfolded α -Syn. A model of the protein assembly involving misfolded dimers is proposed.

MATERIALS AND METHODS

Materials. The 20 mM stock solution of β -mercaptoethanol was prepared in water. The 2.92 mM stock solution of maleimide-polyethylene glycol-silatrane (MAS)³⁰ was dissolved in water and stored at -20°C . The buffers used in force measurement experiments were pH 3.7 (10 mM sodium acetate/acetic acid) and pH 7.0 (PBS buffer). All buffers were adjusted by sodium chloride to a final ionic strength of 150 mM. All other reagents used were of analytical grade from

Sigma-Aldrich unless specified otherwise. Deionized water (18.2 M Ω , 0.22 μm pore size filter, APS Water Services Corp., Van Nuys, CA) was used for all experiments.

Preparation of Recombinant α -Syn Proteins. Wild type A140C α -synuclein in which the C-terminal alanine was replaced with a cysteine and the double mutant A30P-A140C were prepared as described in ref 14. A cDNA encoding human α -Syn E46K or A53T was amplified by PCR and subcloned as an Nde I - Hind III fragment into the vector pT7-7, yielding the constructs pT7-E46K and pT7-A53T, respectively. The double mutants E46K-A140C A53T-A140C were generated by replacing a 354-base pair fragment from pT7-7 A140C (excised by digesting with Nde I and BamH I) with the equivalent DNA fragment from pT7-E46K or pT7-A53T, respectively. The sequence of the α -Syn-encoding insert in each construct was verified using an Applied Biosystems (ABI 3730 XL) DNA sequencer. Each α -Syn variant was expressed in the *E. coli* strain BL21 (DE3) and purified as described in refs 14 and 31. To ensure that the α -synuclein variants were purified as full-length, intact proteins, each lot of protein was characterized using two methods.¹⁴ As one approach, each protein was analyzed by reacting with 5,5'-dithiobis-(2-nitrobenzoic acid) to determine levels of reduced cysteine. From measurements of the absorbance of protein-bound 2-nitro-5-thiobenzoate at 412 nm, it was determined that the ratio of cysteine residues to protein molecules was approximately 1:1. As a second approach, each α -Syn variant was further purified by reverse phase high performance liquid chromatography and analyzed by matrix assisted laser desorption ionization (MALDI) mass spectrometry. The mass-to-charge (m/z) values obtained from the MALDI analysis corresponded to the predicted values for full-length α -Syn.

Tip Modification. Silicon nitride (Si_3N_4) AFM tips (SNL-10, Bruker AFM Probes) were immersed in ethanol for 30 min, rinsed thoroughly with water and dried followed by UV treatment for 30 min (CL-1000 Ultraviolet Cross-linker, UVP, Upland, CA). After the UV treatment, AFM tips were immersed in an aqueous solution of 167 μM maleimide silatrane (MAS) for 3 h followed by multiple rinses in water. For covalent attachment of α -Syn to MAS functionalized tips, a 19 nM protein solution in PBS buffer, pH 7.0, was treated with 1 mM DTT (to reduce any disulfide bonds formed between α -Syn molecules) for 10 min and filtered through 10 kDa MWCO centrifuge filter devices (Amicon Ultra-0.5, Millipore) 3 times. MAS functionalized tips were immersed in a 19 nM solution of α -Syn for 1 h. After rinsing with PBS buffer, unreacted maleimide moieties were quenched with 10 mM β -mercaptoethanol in PBS for 10 min. These functionalized probes were washed with PBS and immediately used for force measurements.

Surface Modification. Mica sheets (Asheville-Schoonmaker Mica Co., Newport News, VA) were cut into $\sim 1.5\text{ cm} \times 1.5\text{ cm}$ squares and glued onto glass slides using epoxy glue EPO-TEK 353ND (Epoxy Technology, Inc., Billerica, MA). Freshly cleaved mica surfaces were treated with MAS for 3 h. MAS modified mica squares were rinsed with water several times to remove nonbound MAS, incubated with a 19 nM α -Syn solution (in PBS, prepared as described in the previous section) for 1 h, and rinsed with PBS buffer. Unreacted maleimide moieties were quenched with 10 mM β -mercaptoethanol in PBS buffer for 10 min. The functionalized mica surfaces were washed with PBS buffer and immediately used for force measurements. The low concentration of α -

synuclein (19 nM) for surface modification was previously identified to provide a sparse surface coverage of monomeric protein. AFM imaging was used to verify that there was only one pair of molecules interacting within the contact area in the force measurements.⁴⁰

Force Measurements. Force–distance measurements were performed in either PBS buffer or acetate buffer (pH 3.7) at room temperature using the Force Robot 300 (JPK Instruments, Berlin, Germany). Silicon nitride cantilevers (SNL-10, Bruker AFM probes) with nominal values of spring constants of ~ 0.1 N/m were used. Spring constants for each cantilever were obtained using a thermal method with the Force Robot 300 instrument. The ramp size was 200 nm with approach and retraction velocities kept constant at 600 nm/s. A low trigger force (200 pN) was applied. Force–distance curves were acquired systematically at different points on the surface set by a grid (5 by 5 μm), where points were separated from each other by 50 nm. The dwell time was set at 0.5 s.

Data Analysis. Thousands of force–distance curves were collected for each α -Syn variant. The curves were filtered using the Data Processing (DP) software of the JPK Force Robot 300 instrument selecting those that contain rupture events. The curves were then fitted with a Worm Like Chain (WLC) model incorporated into the DP software of JPK. The WLC model was chosen because it describes well the elasticity of a polypeptide chain.^{32–34} Rupture force value was measured at the rupture point. The contour length (L_C) was evaluated as a variable parameter for the best fit of the curve while keeping the persistence length constant at 0.4 nm. This value of persistence length was found to adequately describe a variety of proteins.^{32,35,36} Measured values of contour length were assembled into statistical histograms and fitted in MagicPlot Pro 2.0 software with Gaussian functions to determine maxima of L_C distributions ($L_{C,\text{MAX}} \pm \text{SD}$). The length of the α -synuclein segment that undergoes stretching upon applied force was estimated following an approach proposed earlier.³⁷ The length of this segment was calculated by subtracting the length of the MAS linker (3 nm) from the total measured contour length and taking into account that one amino acid residue contributes 0.34 nm to the total contour length.³⁸

RESULTS

Experimental System and Approach. α -Syn molecules were covalently attached to an AFM probe and the mica surface via maleimide–cysteine coupling for the force spectroscopy experiments. The wild-type α -Syn sequence and its mutants (A30P, A53T, and E46K) do not contain cysteine residues except the one we introduced at the very end of the C-terminus (A140C). The cysteine–maleimide coupling chemistry ensured very site-specific immobilization of the protein on the AFM substrate and tip surfaces required for quantitative analysis of the rupture lengths.^{14,29,30} The following structural properties of α -Syn were considered for using the C-terminus for immobilization. There are three distinct domains in the primary structure of α -Syn: N-terminus (1–60 aa), NAC region (61–95 aa), and C-terminus (96–140 aa). We chose the end of the C-terminus as the point of α -Syn covalent attachment because structural changes in the protein were suggested to occur only within the N-terminal part and the NAC region of the protein.³ The N-terminal and NAC regions are responsible for the transition to α -helical structure and association with lipids.³ The NAC region was also found in fibrils and accommodates β -sheet secondary structure.³⁹

However, the C-terminus of α -Syn is a highly acidic and proline-rich region that has been suggested to be in a fully random conformation in both free and lipid bound forms of α -Syn.³ The C-terminal region was expected to act as a flexible linker for probing interactions between α -Syn molecules in force measurements. Therefore, we have used only a short linker within the MAS tether (4 units of ethylene glycol) for covalent attachment of α -Syn molecules.

Interactions between covalently attached α -Syn molecules were probed in multiple approach–retraction cycles at various positions on the surface. If no stable dimeric contact is formed, the retraction of the tip does not detect any rupture event. However, a stable dimeric contact produces a specific rupture signature in the force–distance curve. The beginning of such a force curve usually contains an adhesion peak (1) due to short-range nonspecific interactions between the tip and the surface (illustrated in Figure 1). The adhesion peak is followed by a




Figure 1. Typical regions on the force–distance curves: (1) an adhesion peak due to short-range nonspecific interactions between the tip and the surface, (2) gradual increase in force characteristic of polymer stretching, (3) complete rupture, and (4) the region where the tip comes free from the surface.

complete rupture event of dimeric contact (3) where the tip comes free from the surface (4). The rupture event is preceded by a part characteristic of the stretching of polymer linkers (2).

The control experiments carried out with only one of the surfaces carrying protein (modified tip/unmodified surface or modified surface/unmodified tip) show no or only a few rupture events in force–distance curves (Figure S1, Supporting Information). These control experiments suggest that the ruptures we observed are not due to nonspecific interactions of α -Syn molecules with the surface or AFM tip but rather specific interactions which are probed only when a contact between α -Syn molecules is established. The formation of a dimer makes it possible to observe subtle features of α -Syn dimeric structure and conformational characteristics of misfolded proteins that normally would not be detectable using ensemble measurements. Notably, detectable interactions were also observed between α -Syn at neutral pH in the presence of spermidine.¹⁴

Effect of pH on Misfolding of α -Syn. We have shown recently^{14,40} that the interactions between α -Syn molecules are very weak at neutral pH, and this finding correlates with the low aggregation propensity of α -Syn at this pH.⁴¹ Several recent studies have utilized low pH to convert α -Syn to its aggregation-prone state.^{8,9,20,42–45} We also used low pH to stimulate misfolding of α -Syn in our experiments. The effect of pH is evident from Figure 2A, which shows a set of representative force–distance curves measured for wild type α -Syn at neutral pH. No rupture events were observed beyond the short-range adhesion peak. Next, we continued to probe interactions using the same tip and the same substrate but




Figure 2. Representative force–distance curves for wild type α -Syn. (A) Measured at neutral pH (scale bar 100 pN) and (B) measured at low pH = 3.7 (scale bar 100 pN): (i) with no rupture events (group NR), (ii) single ruptures (group SR), and (iii) multiple ruptures (group MR).

replaced the buffer in the cell to perform probing at a lower pH value of 3.7. Figure 2B demonstrates that at low pH specific interactions appear in the force–distance curves. In addition to curve i with a shape similar to the one detected at neutral pH, clear rupture events were detected (curves ii and iii). We assembled the rupture events into three major groups: NR, force curves with no rupture events (Figure 2B-i); SR, curves that feature a single rupture event (Figure 2B-ii); and MR, curves with multiple ruptures (Figure 2B-iii).

Force–Distance Curves with Single Rupture. Wild Type Protein. First, we analyzed force–distance curves that belong to group SR with a single rupture peak. We followed an approach recently developed in our lab that allows localizing segments of the protein involved in dimer formation.^{37,40,46} Using this approach, we characterized structural differences of misfolded α -Syn variants detected at low pH. Each force–distance curve with a specific rupture event was fitted with a worm-like chain model providing a contour length (L_C) that characterizes the length of stretchable parts of α -Syn molecules not involved in dimeric interaction. A statistical histogram of L_C values measured this way for SR force–distance curves of WT α -Syn is shown in Figure 3A. The distribution has a set of clear

features and was approximated with five peaks representing the most reproducible positions of the rupture events. A Gaussian fit of the histogram resulted in maxima of the distribution at 28 nm (peak 0), 34 nm (peak 1), 44 nm (peak 2), 54 nm (peak 3), and 68 nm (peak 4). The most prominent peak appears at 44 nm, which contributes 42% of all detected single rupture curves for WT α -Syn. The fact that the distribution reveals several peaks suggests that there is more than one conformation of WT α -Syn capable of forming the dimer.

Force–Distance Curves with Single Rupture. Mutants.

Next, we probed interactions involving pathological single point mutants, which correlate with familial early onset Parkinson's disease, A30P, A53T, and E46K.^{10–12} Again, similar to wild type α -Syn, probing interactions involving mutants at pH 7.0 did not reveal specific rupture events. However, specific rupture events appeared on the force–distance curves measured at pH 3.7. Similar to WT α -Syn, all three mutants had three major groups of force–distance curves: NR (no rupture), SR (single rupture), and MR (multiple ruptures). Results of force–distance curve analysis are shown in Figure 3B–D. Similar to the data for WT α -Syn, the distributions of contour lengths have specific features, although they appear simpler with fewer peaks present and slightly narrower than for WT α -Syn (Figures 3A–D). The distributions reveal different patterns for each α -Syn variant suggesting the existence of more than one conformation within the dimer. The major effect of the contour length measurements is a dramatic sensitivity of the contour length profiles upon introducing mutations that can be seen without any additional analysis (Figure 3). This effect reflects the change of the interprotein interaction, and we used fitting of the overall contour length distribution profiles with a set of peaks to facilitate the data interpretation. Four prominent peaks are clearly seen for the A30P mutant with maxima of Gaussians at 30, 36, 41, and 49 nm (Figure 3B). Both A53T and E46K mutants had their contour length distributions skewed to shorter L_C values as compared to those of both WT and A30P (Figures 3C,D). The majority of the rupture events for all α -Syn variants appear within the range of ~26 nm to ~50 nm. Although, there were other interactions observed at longer and shorter distances, these had a lower probability than the peaks in the 26–50 nm range. Moreover, other peaks outside this region are not present for all of the variants. For example, the shortest peak observed for E46K at 17 nm is not present in the histogram for other variants. Also, the longest peak at 68 nm does not have a counterpart in histograms for the mutants suggesting that the interaction producing this long rupture event is specific only for WT.

The comparison of contour length distributions shown in Figure 3 and Table 1 demonstrates the discrete distribution of the rupture events as revealed by contour lengths. However, two major peaks, peak 1 at ~30–34 nm and peak 2 at ~40–44 nm, appear in all of the mutants and WT protein. All mutations remove distant interactions (peak ~68 nm for the WT α -Syn sample), but short interactions appear in the A53T and E46K distributions. Also, similar interactions (peak 1 and peak 2) have different contributions to the overall interaction pattern. For example, interactions of WT are dominated by peak 2, which comprises 42% of all the detected interactions. Peak 1 contributes only 15%. In contrast, the A53T mutant has equal contributions of these two peaks at 31% each, and A30P has almost equal contributions of peak 1 and peak 2, 30% and 24%, respectively. The E46K profile is the opposite of the WT α -Syn profile with peak 1 dominating at 61% compared to 15%




Figure 3. Contour length distribution for α -synuclein variants: group B force–distance curves with a single rupture. (A) Wild type (with maxima at 28 ± 2 nm (peak 0), 34 ± 2 nm (peak 1), 44 ± 4 nm (peak 2), 54 ± 3 nm (peak 3), and 68 ± 4 nm (peak 4)), (B) A30P mutant (with maxima at 30 ± 1 , 36 ± 2 , 41 ± 2 , and 49 ± 2 nm), (C) A53T mutant (with maxima at 26 ± 3 , 33 ± 4 , and 44 ± 3 nm), and (D) E46K mutant (with maxima at 17 ± 3 , 31 ± 4 , and 40 ± 2 nm).

Table 1. Relative Contribution of Single (SR)- and Multiple-Rupture (MR) Curves for Different Variants^a

| | SR # of curves | L_C | MR # of curves | L_C (last) | L_C (internal) |
|------|----------------|-----------------|----------------|------------------|------------------|
| A30P | 398 | 30 ± 1 (28%) | 53 | 25 ± 3 (24%) | 17 ± 5 (27%) |
| | | 36 ± 2 (33%) | | 38 ± 3 (62%) | 34 ± 5 (73%) |
| | | 41 ± 2 (22%) | | 48 ± 1 (14%) | |
| | | 49 ± 2 (17%) | | | |
| WT | 421 | 28 ± 2 (16%) | 233 | 25 ± 3 (25%) | 15 ± 2 (20%) |
| | | 34 ± 2 (15%) | | 40 ± 5 (55%) | 23 ± 4 (50%) |
| | | 44 ± 4 (42%) | | 50 ± 1 (11%) | 33 ± 3 (17%) |
| | | 54 ± 3 (14%) | | 63 ± 3 (9%) | 43 ± 5 (13%) |
| | | 68 ± 4 (13%) | | | |
| A53T | 372 | 26 ± 3 (24%) | 674 | 28 ± 6 (28%) | 13 ± 2 (19%) |
| | | 33 ± 4 (41%) | | 38 ± 2 (30%) | 21 ± 4 (32%) |
| | | 44 ± 3 (35%) | | 48 ± 4 (42%) | 30 ± 2 (27%) |
| | | | | | 39 ± 5 (22%) |
| E46K | 317 | 17 ± 3 (24%) | 498 | | 19 ± 3 (39%) |
| | | 31 ± 4 (61%) | | 41 ± 3 (100%) | 26 ± 3 (31%) |
| | | 40 ± 2 (15%) | | | 34 ± 4 (24%) |
| | | | | | 40 ± 2 (6%) |

^aCorresponding contour lengths are also shown with the relative contribution of each peak.

for *peak 2*. These data show preferences of conformations for these variants. E46K has shorter conformations, while WT has longer conformations, and A53T and A30P are somewhere in the middle.

Force–Distance Curves with a Multiple-Rupture Pattern. The available set of data allowed us to analyze group MR force–distance curves characterized by a multiple-rupture pattern similar to the one shown in Figure 2B-iii. We have compared the relative contribution of group SR vs group MR force–distance curves in the overall set of the force–distance curves with specific rupture events. The A30P mutant had a majority of the force curves with single-rupture peaks, which contributed 81% of all force–distance curves with specific rupture events. The relative contribution of SR curves for all of the variants decreased in the following order A30P > WT > A53T > E46K (81% > 63% > 37% > 24%). We also characterized the force curves by the number of multiple events termed curve complexity. Complexity of the curves, defined by the number of rupture peaks in a single force–distance curve, increased in the following order: A30P < WT < A53T < E46K. The E46K mutant had the highest degree of force curve complexity among all of the variants.

Force–Distance Curves with a Multiple-Rupture Pattern. Wild Type Protein. Here, we analyzed force–distance curves with a multiple-rupture pattern. A representative curve of this kind is shown on Figure 2B-iii. We interpreted the last

rupture peak as the one leading to dimer dissociation, and the rest of the rupture peaks were termed internal ruptures. They correspond to the rupture of interactions between protein segments involved in intramolecular or intermolecular α -Syn folding. Both the last rupture in MR curves and the single rupture in SR curves correspond to force-induced dissociation of dimers, and therefore, it is instructive to compare these rupture events with each other. Analysis of contour length for both the last rupture peaks in group MR and the single-rupture peaks in group SR is shown on Figure 4A and B. The




Figure 4. Rupture contour length histograms (A,B,C) and rupture force histograms (D,E,F) for wild type α -Syn. (A and D) Single rupture event group (with contour length maxima at 28 ± 2, 34 ± 2, 44 ± 4, 54 ± 3, and 68 ± 4 nm and a maximal rupture force of 27 ± 8 pN), (B and E) last events from the multiple-rupture group (with contour length maxima at 25 ± 3, 40 ± 5, 50 ± 1, and 63 ± 3 nm and a maximal rupture force of 27 ± 8 pN), and (C and F) internal ruptures from the multiple-rupture group (with contour length maxima at 15 ± 2, 23 ± 4, 33 ± 3, and 43 ± 5 nm and a maximal rupture force of 41 ± 14 pN).

histograms are very similar with minor variations. These variations are manifested in slight shifts of peak maxima to smaller values in the case of the group MR distribution. It appears that the relative contribution of any particular peak is similar for the last rupture histogram (group MR) and the histogram for group SR. For example, a peak at 44 nm (corresponding to *peak 2* of the SR distribution) is dominant in both histograms. Interestingly, a peak at 34 nm (corresponding to the *peak 1* of the SR distribution) is not present in the histogram of contour lengths measured for the last ruptures (Figure 4B). Although, there might be some events registered at that distance, it is obvious that they do not make up a clear peak leaving a gap between the peak at 25 nm and the peak at 40 nm.

Next, we analyzed the contour length for internal ruptures of the MR force–distance curves. The L_C distribution for these ruptures is shown in Figure 4C. In general, the distribution of contour lengths is shifted to smaller values as compared to both the last event of MR curves (Figure 4B) and the single event in SR curves (Figure 4A). Such behavior of the contour length distributions is expected for the sequential rupture of multiple contacts established between polypeptide chains of α -Syn. A peak corresponding to *peak 1* of the group SR distribution (with a maximum contour length of 33 nm) appeared again when we plotted the contour length distribution for internal ruptures as illustrated by Figure 4C. A peak corresponding to *peak 2* of the SR distribution (with a maximum contour length of 43 nm) is also present; however, it no longer dominates the distribution.

Force–Distance Curves with a Multiple-Rupture Pattern. Single Point Mutants. We have also analyzed force–distance curves with a multiple-rupture pattern for the pathological mutants of α -Syn. Comparison of contour lengths for both the last rupture peaks in group MR and the single-rupture peaks in group SR revealed that A30P and A53T mutants behave similarly to WT. Figure 5A and B shows histograms for single-rupture peaks (SR curves) and last rupture peaks (MR curves) for the A53T mutant. Again, *peak 4* (at 63 nm), characteristic for WT, is not present in the histogram for the mutant. *Peak 1* with a maximum at ~ 30 nm




Figure 5. Rupture contour length histograms (A,B,C) and rupture force histograms (D,E,F) for the A53T mutant of α -Syn. (A and D) Single rupture event group (with contour length maxima at 26 ± 3 , 33 ± 4 , and 44 ± 3 nm and a maximal rupture force of 38 ± 9 pN), (B) the last events from the multiple-rupture group (with contour length maxima at 28 ± 6 , 38 ± 2 , and 48 ± 4 nm and a maximal rupture force of 35 ± 7 pN), and (C) internal ruptures from the multiple-rupture group (with contour length maxima at 13 ± 2 , 21 ± 4 , 30 ± 2 , and 39 ± 5 nm and rupture force maxima of 35 ± 8 , 51 ± 3 , 64 ± 6 , and 83 ± 6 pN).

does not completely disappear for A53T as it does for WT, but its contribution is smaller than that in the histogram for single-rupture curves (Figure 5A and B). The A30P mutant has histograms of contour length very similar to those of A53T for both types of analyzed rupture events with only minor variations (Figure S2, Supporting Information). Similar to WT, a peak with a maximum contour length of ~ 30 nm vanishes in the histogram of last rupture peaks for A30P (Figure S2A and B, Supporting Information). Also, despite much complexity in the number of curves with multiple ruptures and the number of ruptures per curve, the contour length distribution for the E46K mutant features only a single peak at 40 nm when the last rupture is analyzed (Figure S3B, Supporting Information). Interestingly, while group SR events for all of the mutants (particularly E46K) are dominated by a peak at ~ 30 nm, a peak at ~ 40 nm becomes dominant in the last rupture events of MR curves for all of the mutants.

The internal rupture events of the group MR force–distance curves were also analyzed. The L_C distribution for these ruptures is shown in Figures 5C and S2C and S3C (Supporting Information). In general, similar to the WT protein, the distributions are shifted to smaller values as compared to both single events in simple curves (Figures 5A and S2A and S3A (Supporting Information)) and the last event of complex curves (Figures 5B and S2B and S3B (Supporting Information)).

Rupture forces measured for all of the mutants are also similar to those of WT (Figures 5 and S2 and S3 (Supporting Information)). Although generally similar, there are differences among the variants. For example, the E46K mutant had a double distribution of rupture forces at 27 and 50 pN for the last event of MR curves compared to a single peaked distribution for SR curves (Figure S3D and E, Supporting Information). The second peak at 50 pN resembles forces detected for internal events of E46K, which peaked at 49 pN (Figure S3F, Supporting Information). This suggests that the last events in the MR curves for E46K have a double nature: one that resembles single events and another one resembling internal ruptures with higher force. The rupture forces of the A53T mutant also had a higher contribution of large forces for internal ruptures. Figure 5 shows that in addition to small forces (35 pN; Figure 5E, first peak) similar to the forces of the single rupture events, we have also observed a contribution from larger forces (72 pN; Figure 5, E second peak) for the A53T mutant. This situation is opposite to that of WT (Figure 4) and A30P (Figure S2, Supporting Information) but similar to that of the E46K mutant (Figure S3, Supporting Information).

To correlate the rupture force values with the contour lengths, we made 2D scatter plots and histograms of the rupture forces and contour lengths in one combined plot shown in Figures S7–S10 (Supporting Information). Such plots illustrate graphically that the majority of ruptures take place at low force and that this dependence is common for all L_C peaks.

DISCUSSION

Single molecule force spectroscopy probing of interactions between α -Syn molecules within a dimeric contact provided new insight into the pattern of α -Syn assembly within the dimers. Reproducible interactions suggest that α -Syn structure within the dimers is characterized by a set of well-defined conformations. We used the approach described in refs 37 and 46 to characterize the protein segments involved in the α -Syn

folding and interactions. In our immobilization procedure, the proteins are covalently bound to the tip and surface at the C-terminus. Therefore, the position of the peak on the force–distance curve is defined by the protein length between the C-terminus and the interacting segment. The closer the segment to the C-terminus, the shorter is the contour length measured with AFM. This is schematically shown in Figure 6A where




Figure 6. Interaction model of α -synuclein molecules. The model describes two (observed for all the variants of α -synuclein) major peaks in the contour length histograms as sites of the same identity interacting with each other with the formation of a dimer. (A) Position of the interacting site further from the C-end (point of attachment) results in longer contour length value. (B) Positions of the beginning of detected interaction sites. Colored arrows correspond to two detected major interaction sites schematically shown in (A), and black arrows show the positions of A30P, E46K, and A53T mutations in α -synuclein. (C and D) Superposition of representative force–distance curves for the detected rupture events corresponding to L_{C1} and L_{C2} .

contour length L_{C1} is defined by the rupture between the segments shown with red arrows. Longer contour length, L_{C2} , corresponds to the dissociation of distantly located segments shown with blue arrows. Using 0.34 nm as a contribution of one amino acid residue to the total contour length,³⁸ we estimated that L_{C1} and L_{C2} correspond to the following C-terminal parts of α -Syn molecules not participating in interactions: ~94–140 and ~79–140 residues, respectively, matching values of 34 nm (*peak1*) and 44 nm (*peak2*) (Figure S4, Supporting Information).

Structure of WT α -Syn in Dimers. According to Figure 4A, the contour length distribution of the SR force–distance curves features five peaks occurring at distinct distances. *Peak 1* (34 nm) and *peak 2* (44 nm) comprise the majority of the contour lengths; other peaks are located at 54, 68, and 28 nm suggesting that protein segments located at these positions are involved in protein folding within dimers. We compared our data on the location of these interacting segments with the known structure of α -Syn within fibrils. A recent solid state NMR study of full-length α -Syn fibrils showed an extended core with multiple β -strands that span residues 38–96.^{47,48} Assuming that the C-terminus is a stretched polypeptide chain acting as a linker, residues 38–96 correspond to the contour length values between 72 and 32 nm. Our observed range of contour length values vary between 23 and 73 nm, which is close to the estimated L_C range based on the NMR data. Figure S4 (Supporting Information) compares the secondary structure of WT α -Syn taken from ref 48 with the positions of interacting segments estimated based on our contour length analysis.

Remarkably, the peaks in the contour length distribution coincide very well with the location of β -strands in the amyloid fibrils of WT α -Syn determined from ssNMR (see table in Figure S4, Supporting Information).⁴⁸ Despite this similarity, there are some differences: (1) a peak at 59 nm predicted from the NMR structure is missing in our measurements, and (2) we observe a single peak at 44 nm instead of the predicted two peaks at 42 and 48 nm. The majority of events were observed at this contour length (*peak 2* = 44 nm) in the L_C distribution of WT α -Syn. Interestingly, this interaction coincides with the position of an 11 aa long segment of the NAC region. This segment comprising residues 73–83 has been suggested to determine α -Syn's ability to form fibrils.^{39,49}

The multiple-rupture events provide additional insight into the α -Syn structure within dimers. Force-induced dissociation of the dimer results in a complete separation of two monomers and thus the AFM tip comes free from the surface (segment 4 in Figure 1). This complete dissociation appears as the last rupture in the force–distance curves with multiple-rupture peaks. Comparison of the last rupture curves in MR curves with SR curves suggested that these two are very similar. For example, *peak 2* at 44 nm is dominant in both histograms. This peak corresponds to the 73–83 aa stretch, which is critically involved in α -Syn folding within fibrils.^{39,49} Three other peaks at 25, 50, and 63 nm have similar positions in the histogram for group MR events compared to the L_C histogram for group SR events. One major difference is the absence of *peak 1* at 34 nm in the histogram of contour lengths of the last ruptures (Figure 4B).

While the L_C distributions of the SR curves and the last-rupture peaks of the MR curves are similar, the L_C histogram for internal ruptures of the MR curves is quite different. The contour length distribution is skewed to shorter contour length values (Figure 4C) with the first peak in the histogram being at $L_C = 15$ nm. This peak corresponds to a very short noninteracting segment ~122–140 aa, which is further in the C-terminal region than expected based on NMR studies of fibrillar structures.^{47,48} The results of the L_C analysis lead to the following assumptions of the structural features of α -Syn within misfolded dimers: (1) the interacting regions which are capable of forming dimers extend beyond the NAC region into the C-terminal part, and (2) the C-terminal part forms a compact configuration which resists the external force applied in the pulling experiments.

The first assumption suggests a simplified model where the observed short interactions occur between the segments beyond the NAC region, and this model is supported by the recent NMR studies of α -Syn in the free state.^{44,50} According to these data, the segment spanning residues 110–120 has a propensity for the formation of β structure. The neutralization of negative charges within the C-terminus at low pH reduces intermolecular repulsion. This effect along with the preference for β structure formation might effectively induce interactions between regions located outside the NAC.

Our second assumption is based on the possibility of the C-terminal region to fold at low pH. Indeed, several studies observed that low pH values induce a collapse of the C-terminal region due to the shielding of negative charges.^{8,43,51,52} The existence of contacts between different segments due to the compactness of the C-terminal region was observed in AFM single-molecule unfolding of monomeric α -Syn.^{27,28} These studies showed that the intramolecular contacts are strong enough to resist mechanical pulling.

Effect of Mutations on the Structure of α -Syn Folding in Dimers. Three single point mutations A30P, E46K, and A53T of α -Syn identified in familial early onset PD alter the interaction pattern between molecules characteristic of the SMFS experiments. The following differences between mutants and WT were found: (1) contour length distributions for all three mutants lack the peak at $L_C = 68$ nm (*peak 4*), and (2) interactions at shorter contour lengths become more dominant.

According to our estimates, *peak 4* corresponds to the interaction segment starting at around the 45th residue. All three mutations are located in the proximity of this interacting site. Thus, we hypothesize that the absence of the peak at this L_C indicates a direct influence of a mutation on the interacting propensity of this segment, thereby reducing its ability to form a stable dimeric contact. This finding suggests that the protein's segment around the 45th residue is structured and involved in protein folding but that mutations destabilize this structure. Therefore, weakening of the protein structure in this region facilitates α -Syn misfolding and aggregation.

The appearance of short interactions in mutants is an indication of the stabilization of interactions within the C-terminal region of the protein. This effect is more pronounced for the E46K and A53T mutants as demonstrated by Figure 3. We hypothesize that pathological mutants, especially A53T and E46K, in the misfolded state have a higher propensity for β structure in the C-terminal part. This assumption is in agreement with ssNMR analysis of fibrils formed by the A53T mutant that detected the formation of an extended β -sheet core in the C-terminal region as compared to the WT α -Syn.⁴⁸

Despite the variability in positions of the short-range interactions, all four variants of α -Syn have two common peaks in contour length distributions: *peak 1* at around ~ 30 nm and *peak 2* at ~ 40 nm. *Peak 2* corresponding to the 73–83 aa segment of the NAC region was expected to dominate the L_C distributions for mutants as it did for WT. Although, all three pathological mutants have this peak, its contribution becomes smaller and decreases in the following order A53T > A30P > E46K. Such behavior is consistent with observations made in NMR studies that fibrils of mutants are structurally perturbed and that the effects of the mutations increase in the order A30P < A53T < E46K.⁴⁷ The decrease in the intensity of *peak 2* is accompanied by an increase in intensity of another peak common to all variants, *peak 1* at ~ 30 nm, which becomes dominant for the mutants. For example, this interaction comprises 61% of all events for the E46K mutant, while *peak 2* contributes only 15%. This indicates that the mutants prefer to assemble via interactions of segment 2 (segment s2 in Figure S4, Supporting Information) in contrast to WT whose preference is segment 3 (segment s3 in Figure S4, Supporting Information).

Similar to WT, the MR force–distance curves with multiple-rupture events were also present for the mutants. The A30P mutant has a lower contribution of the MR curves in comparison to those of WT (19% vs 37%), while A53T and E46K mutants have a higher contribution of these force–distance curves, 63% and 76%, respectively. Internal ruptures of the MR force curves for the mutants appear shifted to short contour length values suggesting that the observed multiple ruptures promote interactions at short distances. Both assumptions described above for the observation of short rupture distances in the case of WT α -Syn are also valid for the mutants. Collapse and compactness of the C-terminal region at

low pH may result in L_C shortening. Indeed, pathological mutants were found to establish stronger contacts than WT,²⁷ thus increasing the compactness of the C-terminus. Also, we cannot exclude that the observed short interactions appear due to the expansion of interacting modalities beyond the NAC region. According to our model of intermolecular interactions, the shortest interaction observed ($L_C = 13$ – 15 nm) corresponds to contacts between monomers upon dimerization extending into the C-terminal region as far as the 125th residue (Figure S5A, Supporting Information). These results suggest a critical involvement of the C-terminal region in α -Syn misfolding and interactions within the dimer.

Possible Pathways for α -Syn Self-Assembly. Comparison of the aggregation pathways suggests that the A53T and E46K mutations accelerate fibrillization rates compared to those of WT,^{15,16} while A30P prefers an oligomerization route.^{17,18} Our SMFS data also show differences in interactions between monomeric molecules of α -Syn. It is therefore tempting to correlate our data with the known aggregation patterns for WT and mutants of α -Syn. According to our data, the major difference for all of the variants of α -Syn is in the number of the SR force–distance curves vs the MR curves. The elevated relative contribution of multiple ruptures correlates very well with the higher rate of amyloid fibril formation for E46K and A53T mutants compared to those of WT.^{17,18,53} The A30P mutant shows a slower fibrillization rate^{17,18} correlating with the fact that the SR curves are mostly detected for this mutant. Multiple contacts between monomers may also significantly slow down the internal reconfiguration rate of α -Syn, which in turn would promote further association steps. Notably, a slow reconfiguration rate of monomers under conditions when aggregation is enhanced has been proposed as a key factor in determining how fast α -Syn molecules associate with each other.^{20–22} Our data suggest that A53T and E46K mutations increase the possibility for monomers to establish multiple contacts, which may stabilize misfolded states of the protein in the dimer and allow for further growth of aggregated species. A similar hypothesis has been recently proposed based on measurements of dimer stability (lifetime) for several amyloidogenic proteins/peptides including α -Syn.²⁹

Although the model shown in Figure 6 depicts interacting regions in shearing geometry which rupture sequentially upon pulling, a more complex situation can be envisioned. For example, multiple contacts between monomers may promote the formation of a folded dimeric structure as illustrated in Figure S6 (Supporting Information). The structure resembles a fold of α -synuclein within fibrils with several layers of parallel in-register β -sheets as recently proposed.^{54,55} Such folded structure establishes inter- and intramolecular contacts between β -sheets.

CONCLUSIONS

In summary, we report novel findings on the pattern of misfolding of α -Syn and the effects of single point mutations on the protein's dimerization, the very first step of its self-assembly. Single molecule force spectroscopy identified local interacting segments within monomeric α -Syn revealing a great deal of heterogeneity in the dimer conformations. Several segments involved in dimer formation are found in the core of amyloid fibrils suggesting that a link exists between structures of α -Syn at first steps of self-assembly and in mature fibrils. Our data demonstrate that the pathogenic mutations A30P, E46K, and A53T do not increase the propensity of α -Syn to misfold but

rather change conformational preferences of α -Syn. There is a direct effect of mutations through destabilization of interactions involving regions located close to the position of the mutations. Additionally, indirect contribution of mutations to interactions between segments located within the C-terminal region distant from the mutation sites is identified. Importantly, E46K and A53T mutations increase dimerization propensity by promoting simultaneous interactions of multiple segments, while the A30P mutant prefers single type interactions compared to those of WT. We hypothesize that such differences in interacting patterns can define the aggregation pathway of the protein, and future studies are needed to clarify this hypothesis. The single molecule force spectroscopy approach described here was used to study the formation of a dimer. This method can be extended to analyze further steps of the aggregation process if modeled dimer structure is probed by monomer or/and designed dimers.

■ ASSOCIATED CONTENT

■ Supporting Information

Results of control experiments probing specificity of detected rupture events; rupture contour length histograms and rupture force histograms for the A30P mutant of α -Syn and the E46K mutant of α -Syn; comparison of force spectroscopy data with ssNMR data; two possible scenarios for observation of short contour length values; rupture of folded monomers; combined 2D plots correlating the rupture force and the contour length for WT α -synuclein, A53T α -synuclein, E46K α -synuclein, and A30P α -synuclein. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Department of Pharmaceutical Sciences, University of Nebraska Medical Center, 986025 Nebraska Medical Center, Omaha, NE 68198-6025, U.S.A. Phone: 402-559-1971. Fax: 402-559-9543. E-mail: ilyubchenko@unmc.edu.

Funding

This work was supported by grants from the National Institutes of Health (SP01GM091743-02 and 5R01 GM096039-02) and National Science Foundation (EPS-1004094).

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank members of the Lyubchenko lab for the discussion of the results and critical comments.

■ ABBREVIATIONS

α -Syn, alpha-synuclein; AFM, atomic force microscopy; NAC, nonamyloid- β component; NR, no rupture; SR, single rupture; MR, multiple rupture; SMFS, single molecule force spectroscopy; WT, wild type; PD, Parkinson's disease

■ REFERENCES

- (1) Bisaglia, M., Mammi, S., and Bubacco, L. (2009) Structural insights on physiological functions and pathological effects of alpha-synuclein. *FASEB J.* 23, 329–340.
- (2) Weinreb, P. H., Zhen, W., Poon, A. W., Conway, K. A., and Lansbury, P. T., Jr. (1996) NACP, a protein implicated in Alzheimer's disease and learning, is natively unfolded. *Biochemistry* 35, 13709–13715.

- (3) Eliezer, D., Kutluay, E., Bussell, R., Jr., and Browne, G. (2001) Conformational properties of alpha-synuclein in its free and lipid-associated states. *J. Mol. Biol.* 307, 1061–1073.
- (4) Conway, K. A., Harper, J. D., and Lansbury, P. T., Jr. (2000) Fibrils formed in vitro from alpha-synuclein and two mutant forms linked to Parkinson's disease are typical amyloid. *Biochemistry* 39, 2552–2563.
- (5) Iwai, A., Yoshimoto, M., Masliah, E., and Saitoh, T. (1995) Non-A beta component of Alzheimer's disease amyloid (NAC) is amyloidogenic. *Biochemistry* 34, 10139–10145.
- (6) Spillantini, M. G., Schmidt, M. L., Lee, V. M. Y., Trojanowski, J. Q., Jakes, R., and Goedert, M. (1997) [alpha]-Synuclein in Lewy bodies. *Nature* 388, 839–840.
- (7) Baba, M., Nakajo, S., Tu, P. H., Tomita, T., Nakaya, K., Lee, V. M., Trojanowski, J. Q., and Iwatsubo, T. (1998) Aggregation of alpha-synuclein in Lewy bodies of sporadic Parkinson's disease and dementia with Lewy bodies. *Am. J. Pathol.* 152, 879–884.
- (8) McClendon, S., Rospigliosi, C. C., and Eliezer, D. (2009) Charge neutralization and collapse of the C-terminal tail of alpha-synuclein at low pH. *Protein Sci.* 18, 1531–1540.
- (9) Uversky, V. N., Li, J., and Fink, A. L. (2001) Evidence for a partially folded intermediate in alpha-synuclein fibril formation. *J. Biol. Chem.* 276, 10737–10744.
- (10) Kruger, R., Kuhn, W., Muller, T., Woitalla, D., Graeber, M., Kosel, S., Przuntek, H., Epplen, J. T., Schols, L., and Riess, O. (1998) Ala30Pro mutation in the gene encoding alpha-synuclein in Parkinson's disease. *Nat. Genet.* 18, 106–108.
- (11) Polymeropoulos, M. H., Lavedan, C., Leroy, E., Ide, S. E., Dehejia, A., Dutra, A., Pike, B., Root, H., Rubenstein, J., Boyer, R., Stenroos, E. S., Chandrasekharappa, S., Athanassiadou, A., Papapetropoulos, T., Johnson, W. G., Lazzarini, A. M., Duvoisin, R. C., Di Iorio, G., Golbe, L. I., and Nussbaum, R. L. (1997) Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science* 276, 2045–2047.
- (12) Zarranz, J. J., Alegre, J., Gómez-Esteban, J. C., Lezcano, E., Ros, R., Ampuero, I., Vidal, L., Hoenicka, J., Rodriguez, O., Atarés, B., Llorens, V., Tortosa, E. G., del Ser, T., Muñoz, D. G., and de Yebenes, J. G. (2004) The new mutation, E46K, of alpha-synuclein causes parkinson and Lewy body dementia. *Ann. Neurol.* 55, 164–173.
- (13) Uversky, V. N., Li, J., and Fink, A. L. (2001) Metal-triggered structural transformations, aggregation, and fibrillation of human alpha-synuclein. A possible molecular link between Parkinson's disease and heavy metal exposure. *J. Biol. Chem.* 276, 44284–44296.
- (14) Krasnoslobodtsev, A. V., Peng, J., Asiago, J. M., Hindupur, J., Rochet, J. C., and Lyubchenko, Y. L. (2012) Effect of spermidine on misfolding and interactions of alpha-synuclein. *PLoS One* 7, e38099.
- (15) Greenbaum, E. A., Graves, C. L., Mishizen-Eberz, A. J., Lupoli, M. A., Lynch, D. R., Englander, S. W., Axelsen, P. H., and Giasson, B. I. (2005) The E46K mutation in alpha-synuclein increases amyloid fibril formation. *J. Biol. Chem.* 280, 7800–7807.
- (16) Conway, K. A., Harper, J. D., and Lansbury, P. T. (1998) Accelerated in vitro fibril formation by a mutant alpha-synuclein linked to early-onset Parkinson disease. *Nat. Med.* 4, 1318–1320.
- (17) Conway, K. A., Lee, S. J., Rochet, J. C., Ding, T. T., Williamson, R. E., and Lansbury, P. T., Jr. (2000) Acceleration of oligomerization, not fibrillization, is a shared property of both alpha-synuclein mutations linked to early-onset Parkinson's disease: implications for pathogenesis and therapy. *Proc. Natl. Acad. Sci. U.S.A.* 97, 571–576.
- (18) Li, J., Uversky, V. N., and Fink, A. L. (2001) Effect of familial Parkinson's disease point mutations A30P and A53T on the structural properties, aggregation, and fibrillation of human alpha-synuclein. *Biochemistry* 40, 11604–11613.
- (19) Mukhopadhyay, S., Krishnan, R., Lemke, E. A., Lindquist, S., and Deniz, A. A. (2007) A natively unfolded yeast prion monomer adopts an ensemble of collapsed and rapidly fluctuating structures. *Proc. Natl. Acad. Sci. U.S.A.* 104, 2649–2654.
- (20) Ahmad, B., Chen, Y., and Lapidus, L. J. (2012) Aggregation of alpha-synuclein is kinetically controlled by intramolecular diffusion. *Proc. Natl. Acad. Sci. U.S.A.* 109, 2336–2341.

- (21) Ahmad, B., and Lapidus, L. J. (2012) Curcumin prevents aggregation in alpha-synuclein by increasing reconfiguration rate. *J. Biol. Chem.* 287, 9193–9199.
- (22) Lapidus, L. J. (2013) Understanding protein aggregation from the view of monomer dynamics. *Mol. Biosyst.* 9, 29–35.
- (23) Rubinstein, A., Lyubchenko, Y. L., and Sherman, S. (2009) Dynamic properties of pH-dependent structural organization of the amyloidogenic beta-protein (1–40). *Prion* 3, 31–43.
- (24) Huang, A., and Stultz, C. M. (2009) Finding order within disorder: elucidating the structure of proteins associated with neurodegenerative disease. *Future Med. Chem.* 1, 467–482.
- (25) Ullman, O., Fisher, C. K., and Stultz, C. M. (2011) Explaining the structural plasticity of alpha-synuclein. *J. Am. Chem. Soc.* 133, 19536–19546.
- (26) Uversky, V. N. (2003) A protein-chameleon: conformational plasticity of alpha-synuclein, a disordered protein involved in neurodegenerative disorders. *J. Biomol. Struct. Dyn.* 21, 211–234.
- (27) Bruciale, M., Sandal, M., Di Maio, S., Rampioni, A., Tessari, I., Tosatto, L., Bisaglia, M., Bubacco, L., and Samori, B. (2009) Pathogenic mutations shift the equilibria of alpha-synuclein single molecules towards structured conformers. *ChemBioChem* 10, 176–183.
- (28) Sandal, M., Valle, F., Tessari, I., Mammi, S., Bergantino, E., Musiani, F., Bruciale, M., Bubacco, L., and Samori, B. (2008) Conformational equilibria in monomeric alpha-synuclein at the single-molecule level. *PLoS Biol.* 6, e6.
- (29) Lyubchenko, Y. L., Kim, B. H., Krasnoslobodtsev, A. V., and Yu, J. (2010) Nanoimaging for protein misfolding diseases. *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.* 2, 526–543.
- (30) Krasnoslobodtsev, A. V., Shlyakhtenko, L. S., Ukraintsev, E., Zaikova, T. O., Keana, J. F., and Lyubchenko, Y. L. (2005) Nanomedicine and protein misfolding diseases. *Nanomedicine* 1, 300–305.
- (31) Rochet, J.-C., and Lansbury, P. T., Jr. (2000) Amyloid fibrillogenesis: themes and variations. *Curr. Opin. Struct. Biol.* 10, 60–68.
- (32) Rief, M., Gautel, M., Oesterhelt, F., Fernandez, J. M., and Gaub, H. E. (1997) Reversible unfolding of individual titin immunoglobulin domains by AFM. *Science* 276, 1109–1112.
- (33) Tskhovrebova, L., Trinick, J., Sleep, J. A., and Simmons, R. M. (1997) Elasticity and unfolding of single molecules of the giant muscle protein titin. *Nature* 387, 308–312.
- (34) Stirnemann, G., Giganti, D., Fernandez, J. M., and Berne, B. J. (2013) Elasticity, structure, and relaxation of extended proteins under force. *Proc. Natl. Acad. Sci. U.S.A.* 110, 3847–3852.
- (35) Sarkar, A., Caamano, S., and Fernandez, J. M. (2007) The mechanical fingerprint of a parallel polyprotein dimer. *Biophys. J.* 92, L36–38.
- (36) Oesterhelt, F., Oesterhelt, D., Pfeiffer, M., Engel, A., Gaub, H. E., and Müller, D. J. (2000) Unfolding pathways of individual bacteriorhodopsins. *Science* 288, 143–146.
- (37) Yu, J., Malkova, S., and Lyubchenko, Y. L. (2008) alpha-Synuclein misfolding: single molecule AFM force spectroscopy study. *J. Mol. Biol.* 384, 992–1001.
- (38) Yang, G., Cecconi, C., Baase, W. A., Vetter, I. R., Breyer, W. A., Haack, J. A., Matthews, B. W., Dahlquist, F. W., and Bustamante, C. (2000) Solid-state synthesis and mechanical unfolding of polymers of T4 lysozyme. *Proc. Natl. Acad. Sci. U.S.A.* 97, 139–144.
- (39) El-Agnaf, O. M., Bodles, A. M., Guthrie, D. J., Harriott, P., and Irvine, G. B. (1998) The N-terminal region of non-A beta component of Alzheimer's disease amyloid is responsible for its tendency to assume beta-sheet and aggregate to form fibrils. *Eur. J. Biochem.* 258, 157–163.
- (40) Yu, J., Warnke, J., and Lyubchenko, Y. L. (2011) Nanoprobng of alpha-synuclein misfolding and aggregation with atomic force microscopy. *Nanomed.: Nanotechnol., Biol., Med.* 7, 146–152.
- (41) McAllister, C., Karymov, M. A., Kawano, Y., Lushnikov, A. Y., Mikheikin, A., Uversky, V. N., and Lyubchenko, Y. L. (2005) Protein interactions and misfolding analyzed by AFM force spectroscopy. *J. Mol. Biol.* 354, 1028–1042.
- (42) Bernstein, S. L., Liu, D., Wyttenbach, T., Bowers, M. T., Lee, J. C., Gray, H. B., and Winkler, J. R. (2004) alpha-Synuclein: stable compact and extended monomeric structures and pH dependence of dimer formation. *J. Am. Soc. Mass Spectrom.* 15, 1435–1443.
- (43) Trexler, A. J., and Rhoades, E. (2010) Single molecule characterization of alpha-synuclein in aggregation-prone states. *Biophys. J.* 99, 3048–3055.
- (44) Wu, K. P., Weinstock, D. S., Narayanan, C., Levy, R. M., and Baum, J. (2009) Structural reorganization of alpha-synuclein at low pH observed by NMR and REMD simulations. *J. Mol. Biol.* 391, 784–796.
- (45) Hoyer, W., Antony, T., Cherny, D., Heim, G., Jovin, T. M., and Subramaniam, V. (2002) Dependence of alpha-synuclein aggregate morphology on solution conditions. *J. Mol. Biol.* 322, 383–393.
- (46) Kim, B. H., Palermo, N. Y., Lovas, S., Zaikova, T., Keana, J. F., and Lyubchenko, Y. L. (2011) Single-molecule atomic force microscopy force spectroscopy study of Abeta-40 interactions. *Biochemistry* 50, 5154–5162.
- (47) Comellas, G., Lemkau, L. R., Nieuwkoop, A. J., Kloepper, K. D., Lador, D. T., Ebisu, R., Woods, W. S., Lipton, A. S., George, J. M., and Rienstra, C. M. (2011) Structured regions of alpha-synuclein fibrils include the early-onset Parkinson's disease mutation sites. *J. Mol. Biol.* 411, 881–895.
- (48) Heise, H., Celej, M. S., Becker, S., Riedel, D., Pelah, A., Kumar, A., Jovin, T. M., and Baldus, M. (2008) Solid-state NMR reveals structural differences between fibrils of wild-type and disease-related A53T mutant alpha-synuclein. *J. Mol. Biol.* 380, 444–450.
- (49) Giasson, B. I., Murray, I. V., Trojanowski, J. Q., and Lee, V. M. (2001) A hydrophobic stretch of 12 amino acid residues in the middle of alpha-synuclein is essential for filament assembly. *J. Biol. Chem.* 276, 2380–2386.
- (50) Rospigliosi, C. C., McClendon, S., Schmid, A. W., Ramlall, T. F., Barre, P., Lashuel, H. A., and Eliezer, D. (2009) E46K Parkinson's-linked mutation enhances C-terminal-to-N-terminal contacts in alpha-synuclein. *J. Mol. Biol.* 388, 1022–1032.
- (51) Cho, M.-K., Nodet, G., Kim, H.-Y., Jensen, M. R., Bernado, P., Fernandez, C. O., Becker, S., Blackledge, M., and Zweckstetter, M. (2009) Structural characterization of alpha-synuclein in an aggregation prone state. *Protein Sci.* 18, 1840–1846.
- (52) Fernandez, C. O., Hoyer, W., Zweckstetter, M., Jares-Erijman, E. A., Subramaniam, V., Griesinger, C., and Jovin, T. M. (2004) NMR of alpha-synuclein-polyamine complexes elucidates the mechanism and kinetics of induced aggregation. *EMBO J.* 23, 2039–2046.
- (53) Choi, W., Zibae, S., Jakes, R., Serpell, L. C., Davletov, B., Crowther, R. A., and Goedert, M. (2004) Mutation E46K increases phospholipid binding and assembly into filaments of human alpha-synuclein. *FEBS Lett.* 576, 363–368.
- (54) Chen, M., Margittai, M., Chen, J., and Langen, R. (2007) Investigation of alpha-synuclein fibril structure by site-directed spin labeling. *J. Biol. Chem.* 282, 24970–24979.
- (55) Vilar, M., Chou, H. T., Luhrs, T., Maji, S. K., Riek-Loher, D., Verel, R., Manning, G., Stahlberg, H., and Riek, R. (2008) The fold of alpha-synuclein fibrils. *Proc. Natl. Acad. Sci. U.S.A.* 105, 8637–8642.