

Effect of *p*-dimethylaminoazobenzene on the nucleic acids : potassium ratios of rat liver

It was observed recently in this laboratory¹ that the growth of the Crocker sarcoma S-180 in Swiss mice caused an increase of the DNA: K ratio in the liver, lung, and kidney of the host, but that the RNA: K ratio remained unchanged. Asimov *et al.*² reported that in human carcinoma there was no change in the DNA: K ratio of the tumor as compared with the tissue surrounding it. In the present experiments, we have studied the effects of *p*-dimethylaminoazobenzene (*p*-DAB), where a well defined tumor can be compared with the tissue from which it arises.

As shown in Table I, the hepatoma is characterized by a DNA: K ratio definitely much higher than that of normal liver. The ratio is also significantly higher for the liver of rats fed *p*-DAB, but which have not yet developed a hepatoma. The RNA: K ratio in the liver decreases significantly, but for the hepatoma the change is not significant at a level of $p = 0.05$.

TABLE I
PERCENTAGE CHANGE IN DNA:K AND RNA:K RATIOS IN THE LIVERS AND HEPATOMAS
OF RATS FED *p*-DIMETHYLAMINOAZOBENZENE *

Group	No. of rats	Change, %	
		DNA: K	RNA: K
Liver, low riboflavin	14	+ 28	— 18
Liver, high riboflavin	8	+ 16	— 9
Hepatoma	12	+ 186	— 4

* Ratios in the livers of the controls: DNK: K, 0.56 ± 0.03 ; RNA: K, 2.74 ± 0.08 .

Riboflavin has long been recognized as a protective factor in animals fed *p*-DAB. In the present study, it has been observed that when the vitamin is fed at a high level, it causes less marked changes in both ratios in the liver tissue of the experimental rats, but no change whatsoever for the hepatoma.

This work was supported in part by grants from The National Institutes of Health, U.S. Public Health Service (C-1370 (C3)), and from The Damon Runyon Memorial Fund.

Department of Biochemistry, Fordham University,
New York 58, N.Y. (U.S.A.)

NESTOR M. RODRIGUEZ
MARY T. BOLTON
LEOPOLD R. CERECEDO

¹ N. M. RODRIGUEZ, H. T. HOCHSTRASSER, J. O. MALBICA AND L. R. CERECEDO, *J. Biol. Chem.*, **211** (1954) 483.

² I. ASIMOV, H. M. LEMON, R. M. REGUERA, M. M. DAVISON AND B. S. WALKER, *J. Cell. Comp. Physiol.*, **37** (1951) 355.

Received January 5th, 1955

A relation between cystine content and ultraviolet sensitivity of proteins

The action of ultraviolet light on proteins has been extensively studied^{1,2}, but the mechanism by which proteins are denatured by light is not understood. An excellent review of possible theories may be found in a recent article by DOTY AND GEIDUSCHEK³. McLAREN⁴ has shown that the quantum yield for the inactivation of proteins decreases as the molecular weight increases. The correlation between the quantum yield and molecular weight, however, is not a particularly good one.

The purpose of this note is to indicate that the quantum yield for the inactivation of enzymes by light of wavelength 2,537 Å is very closely related to the cystine content of the molecule. Table I shows the quantum yield and the per cent cystine composition for all proteins for which data are available. The per cent abundance of the aromatic amino acids is indicated simply to show that no correlation exists between these quantities and the quantum yield. The calculated value of the quantum yield depends upon the value chosen for the molecular weight. The agreement between quantum yield and cystine content is quite remarkable, especially when we consider the many different circumstances upon which the quantum yield may depend.