TITLES OF RELATED PAPERS IN OTHER SECTIONS

The following papers that have recently appeared in other sections of BIOCHIMICA ET BIOPHYSICA ACTA may be of interest to the readers of this specialized section:

BBA - PROTEIN STRUCTURE

Formation of a complex between yeast L-lactate dehydrogenase (cytochrome b_2) and cytochrome c. Ultracentrifugal and gel chromatographic analyses (BBA 37678)	
by T. Yoshimura, A. Matsushima, K. Aki and K. Kakiuchi (Tokushima and Osaka, Japan)	492 (1977) 331–339
by T. Takebayashi, Y. Morita and F. Oosawa (Osaka, Japan) Differential sensitivity to trypsin digestion of purified forms of <i>Micrococcus lysodeikticus</i> ATPase (BF ₁). A study of their structural and confor-	492 (1977) 357–363
mational differences and mechanism of conversion (BBA 37671) by J. Carreira, J. M. Andreu and E. Muñoz (Madrid, Spain) Studies on the amino groups of myosin ATPase. IV. Effect of ATP and its analogs on the spectral properties of trinitrophenylated myosin and its	492 (1977) 387–398
active fragments (BBA 37695) by A. Muhlrad (Jerusalem, Israel)	493 (1977) 154–166
bovine adrenal cortex by affinity chromatography with pregnenolone as ligand (BBA 37726) by B. E. Tilley, N. Watanuki and P. F. Hall (Irvine, Calif., U.S.A.)	493 (1977) 260–271
Actin from the nematode, Ceanorhabditis elegans, is a single electrofocusing species (BBA 37735) by F. H. Schachat, H. E. Harris and H. F. Epstein (Stanford, Calif., U.S.A.)	493 (1977) 304–309
BBA - LIPIDS AND LIPID METABOLISM	, ,
Characteristics of a cytochrome P -450-dependent fatty acid ω -2 hydroxylase from $Bacillus\ megaterium\ (BBA\ 57000)$	
by R. S. Matson, R. S. Hare and A. J. Fulco (Los Angeles, Calif., U.S.A) Phospholipids and detergents as effectors in the liver microsomal hydroxylase system (BBA 57028)	487 (1977) 487–494
by M. Ingelman-Sundberg (Stockholm, Sweden)	488 (1977) 225–234
by B. E. Tilley, M. Watanuki and P. F. Hall (Irvine, Calif., U.S.A.).	488 (1977) 331–340