

BBA 47640

## LINEAR DICHROISM OF LIGHT HARVESTING BACTERIOCHLOROPHYLL PROTEINS FROM *RHODOPSEUDOMONAS* *SPHAEROIDES* IN STRETCHED POLYVINYL ALCOHOL FILMS

JOHN BOLT and KENNETH SAUER

*Department of Chemistry and Laboratory of Chemical Biodynamics, Lawrence Berkeley Laboratory, University of California, Berkeley, CA 94720 (U.S.A.)*

(Received August 22nd, 1978)

**Key words:** *Linear dichroism; Bacteriochlorophyll; Light-harvesting protein; Bacterial photosynthesis; Reaction center; (Rhodopseudomonas sphaeroides)*

### Summary

Light-harvesting bacteriochlorophyll-protein complexes from *Rhodopseudomonas sphaeroides* 2.4.1 and R-26 mutant are solubilized in sodium dodecyl sulfate and imbedded in polyvinyl alcohol. Stretching induces orientation, and the linear dichroism of visible and near infrared absorption is analyzed. Based on a simple model, angles between the particle axis and the transition dipole moments are found. In the near infrared absorption band of the R-26 light-harvesting protein the dichroic ratio varies from 1.30 to 1.57. Using the absorption curves the band is resolved into two exciton components. In the visible absorption band the dichroic ratio has a constant value of 0.43 for the R-26 protein but varies with wavelength for the wild type light-harvesting protein. This variation is attributed to an additional bacteriochlorophyll not present in the R-26 protein.

---

### Introduction

The light-harvesting bacteriochlorophyll *a* and carotenoid of purple photosynthetic bacteria are found complexed with low molecular weight peptides of approximately 10 000 daltons [1–7]. With selective solubilization of the chromatophore membranes these complexes can be isolated. Sauer and Austin separated a complex from the carotenoidless mutant of *Rhodopseudomonas*

*sphaeroides*, R-26 which contained two polypeptides and two bacteriochlorophylls [6]. In the presence of sodium dodecyl sulfate (SDS) this complex is not further aggregated, and it absorbs in the near infrared with a single maximum at 853 nm. The circular dichroism, CD, spectrum shows a double Cotton effect, negative to long wavelengths, and centered on the 853 nm absorption band. A similar complex from the wild type *Rps. sphaeroides* contains an additional bacteriochlorophyll and a carotenoid and exhibits absorption maxima at 799 nm and 850 nm. Cogdell and Thornber classify this complex as B800-B850 [7]. The 800 nm absorption band of the wild type complex can be irreversibly and selectively bleached, leaving a complex with essentially the same absorption and CD properties as the complex isolated from the mutant R-26 [6]. Thus, we believe the complex from R-26 is an altered form of the wild type complex. This paper reports evidence to support this belief.

Recently Rafferty and Clayton reported the linear dichroism of the reaction center bacteriochlorophyll-protein complex in a stretched gelatin film [8]. We use a similar approach, embedding the protein detergent particles in polyvinyl alcohol as the stretching matrix. The information gained is the orientation of the transition dipoles with respect to the stretching axis. Because bacteriochlorophyll has two well-separated electronic transitions,  $Q_x$  and  $Q_y$ , which are believed to be polarized in the monomer bacteriochlorophyll close to the bacteriochlorin symmetry axes [9], and because these pigment protein complexes involve a small number of bacteriochlorophylls, they are suitable for determining the geometry of the pigment molecules from optical spectra.

## Materials and Methods

Bacteria strains R-26 and 2.4.1 of *Rps. sphaeroides* were grown as described by Sauer and Austin [6]. Chromatophores were prepared by the method of Clayton and Wang [10] and were purified on a 0.6–1.2 M discontinuous sucrose gradient. The light-harvesting complex (LH-wt) from wild type chromatophores was prepared according to published procedures [2,11]. The protein complex was dialyzed against 0.01 M Tris-HCl, pH = 7.6, henceforth called buffer, and stored at  $-20^{\circ}\text{C}$ . The light-harvesting complex (LH-R26) from R-26 was prepared as follows: chromatophores were adjusted to an absorbance of  $0.5\text{ cm}^{-1}$  at 860 nm when diluted one hundred-fold. Dodecyl-dimethylamine oxide was added to a final concentration of 1% and the suspension was centrifuged over a 0.6–1.1 M discontinuous sucrose gradient at  $90\,000 \times g$  for 3 h. The green material at the center sucrose interface was collected with a syringe and dialyzed overnight against buffer. The dialyzate was centrifuged 1 h at  $80\,000 \times g$ . The pellet was resuspended in a minimum volume of buffer containing 1% SDS. The protein was then purified on a Sephadex G-150 column at  $13^{\circ}\text{C}$ . The material from 15 ml of chromatophores was applied to a column  $29 \times 2.5\text{ cm}$  and eluted with 0.4% SDS in buffer. Fractions with the ratio of absorbance at 855 nm to absorbance at 275 nm greater than 2.3 were pooled, dialyzed against buffer and stored at  $-20^{\circ}\text{C}$ . Low molecular weight polyvinyl alcohol from Touzart et Matignon was dissolved as a 35% solution in 50 mM Tris-HCl to a final pH = 7.0. The dialyzed proteins were concentrated by centrifugation and resuspended in buffer containing 3.5 mM SDS to an

absorbance at 853 nm of  $25 \text{ cm}^{-1}$ . Due to the labile nature of the 800 nm band, 50 mM dithioerythritol was included in the final solution of LH-wt protein. 0.5–2.0 ml of the protein solution was mixed with 2.5 ml of the polyvinyl alcohol solution at room temperature. After allowing the bubbles to rise overnight, the solution was spread over a glass microscope slide,  $25 \times 76 \text{ mm}$ , and dried in a nitrogen atmosphere at room temperature in the dark. The polyvinyl alcohol film was peeled off the glass slide and hydrated at 100% humidity for 1 h. A center portion was cut, clamped into a simple stretching device and stretched slowly to about half maximum. The film was rehydrated for 30 min before stretching to the final value. The extent of stretching is measured by stretch ratio,  $R_s$ , as defined by Land [12]. Stretch ratio is the axial ratio of the ellipse into which a hypothetical circle on the film is distorted.

Absorption measurements were performed using a Cary 14 recording spectrophotometer with a Hamamatsu R928 photomultiplier. Glan-Thompson prisms were mounted in sample and reference beams. The plane of the polyvinyl alcohol film was mounted perpendicular to the beam with the stretch axis vertical. Absorption spectra with polarizations parallel and perpendicular to the stretch axis were measured by rotating the polarizing prisms for vertical and horizontal polarizations, respectively. Baselines with vertical and horizontal polarizations were measured separately. Absorption of films made and stretched with no proteins present was coincident with the baseline after correcting for the baseline shift at 930 nm. The baseline shift was attributed to reflection and a small amount of light scattering. The plane of the polyvinyl alcohol film was tilted by mounting dry films on a goniometer, with the rotating axis vertical and passing through the center of the beam.

## Results and Discussion

Fig. 1 shows the absorption of the LH-R26 protein parallel and perpendicular to the stretch axis. In the  $Q_y$  region (820–920 nm) there is a distinct change in dichroic ratio across the band. The sigmoid shape of the ratio suggests two overlapping electronic transitions with different dichroic ratios. The midpoint of this sigmoid curve is 6–8 nm to the red of the wavelength of maximum absorbance,  $\lambda_{\text{max}}$ . In the 850 nm absorption band of the LH-wt protein (Fig. 2) this behavior is also observed, suggesting similarity of the two complexes.

In the  $Q_x$  region of the LH-R26 protein (Fig. 1) the dichroic ratio is constant within the precision of the measurement. By contrast, the LH-wt protein has a distinct change across the  $Q_x$  absorption band. When dithioerythritol is omitted from the polyvinyl alcohol film the 797 nm absorption band is bleached (not shown). This also decreases the dichroic ratio on the short wavelength side of the 590 nm absorption band but not on the long wavelength side. In solution, the selective bleaching at 800 nm is accompanied by a shift of the  $Q_x$   $\lambda_{\text{max}}$  from 589 nm to 592 nm. Thus, the increase in  $A_{\parallel}$  at shorter wavelengths in the  $Q_x$  region of the intact LH-wt protein is correlated with the presence of the 800 nm absorption band.

The dichroic ratio across the carotenoid absorption is constant from 440 nm to 520 nm.

Rafferty and Clayton reported orientation of reaction centers in unstretched




Fig. 1. Absorption parallel ( $A_{\parallel}$ ) and perpendicular ( $A_{\perp}$ ) to the stretch axis is shown for the LH-R26 protein in stretched polyvinyl alcohol film. In the  $Q_y$  region  $R_s = 2.5$ . A more concentrated sample was used for better precision in the  $Q_x$  region, and  $R_s = 3.1$ .

gelatin films [8]. This was a result of the shrinkage in thickness by a factor of 30 on drying the gelatin film. The polyvinyl films shrink to no less than one-fourth the original thickness. As a result, no dichroism was detected in unstretched, dried polyvinyl alcohol films when  $A_{\parallel}$  and  $A_{\perp}$  were recorded with the sample tilted by 45 degrees.

To extract quantitative information from the linear dichroism presented in Figs. 1 and 2 it was necessary to establish a model for the orientation phenomenon. This model must include the shape and extent of aggregation of the protein detergent particles and also the extent of orientation achieved by stretching. We will assume that orientation is uniaxial along the direction of stretch and that the particle contains a unique long axis. (However, an oblate ellipsoid model has not been eliminated.) Unfortunately, the aggregation state of the proteins cannot be directly determined in the polyvinyl alcohol glass. The absorption and circular dichroism spectra are unaffected by incorporation into polyvinyl alcohol films. In addition the films show no sign of particle aggregation and are optically clear. For these reasons we feel justified in treating the LH-R26 protein as individual units containing two polypeptides, SDS and two bacteriochlorophylls, and the LH-wt protein as a unit containing two polypeptides, SDS, three bacteriochlorophylls and one carotenoid [6]. We cannot rule out the possible formation of small aggregates; the dichroism observed shows that such aggregation would not be random.

According to the model of Fraser, the degree of orientation can be described by a fraction,  $f$ , of particles aligned perfectly with the stretch axis and a fraction,  $1 - f$ , randomly oriented [13]. The dichroic ratio,  $D$ , can be expressed as




Fig. 2. Absorption parallel and perpendicular to the stretch axis for LH-wt protein in stretched polyvinyl alcohol.  $R_s = 2.9$ .

a function of  $f$  and  $\theta$ , the angle of the transition dipole to the particle axis:

$$D = \frac{A_{||}}{A_{\perp}} = \frac{f \cos^2 \theta + \frac{1}{3}(1-f)}{\frac{1}{2} f \sin^2 \theta + \frac{1}{3}(1-f)} \quad (1)$$

Other choices for plausible distribution functions can be related to Eqn. 1 [14].

Two approaches were used to assess the orientation parameter  $f$ . The first is based on the restrictions inherent in Eqn. 1. In Fig. 3,  $f$  is plotted as a function of  $\theta$  and  $D$ . Typical values for the dichroic ratio are 1.55 ( $Q_y$ ) and 0.4 ( $Q_x$ ), which restrict  $f$  to values greater than 0.15 and 0.5, respectively. Since  $f$  must be the same for all dipoles within the same sample, we restrict it to values greater than 0.5. For dichroic ratios close to 1, the choice of  $f$  is not critical; but for increasingly greater dichroism, the value of  $\theta$  varies with  $f$  more significantly. Another approach is to plot the dichroic ratios as a function of amount of stretching. Fig. 4 plots  $D$  versus  $R_s$  at various wavelengths for a sample of the LH-wt protein. With stretch ratio greater than 3 the dichroic ratios




Fig. 3. The dependence of  $\theta$ , angle between particle axis and transition dipole, on the orientation parameter,  $f$ , for several values of the dichroic ratio,  $D$ . Values of  $D$  are labeled for each curve. Plotted from Eqn. 1; adopted from ref. 15.




Fig. 4. The dependence of dichroic ratio on stretching. The sample is LH-wt protein.  $A_{||}/A_{\perp}$  is plotted versus stretch ratio at several wavelengths.

approach limiting values. This implies that a maximum value of  $f$  has been reached, but it does not imply that  $f$  equals 1. Table I lists values of  $\theta$  based on observed dichroic ratios and using several possible values of the orientation parameter.

The apparent presence of two electronic transitions in the  $Q_y$  region is best interpreted in terms of oscillators with unequal strengths centered at different wavelengths. (Equally weighted oscillators at different wavelengths would result in a CD zero crossing and a dichroic ratio transition with midpoint centered at the peak of the absorption curve.) Exciton coupling of two chromophores predicts splitting of the degenerate transitions into non-degenerate orthogonal transitions, designated plus and minus. Using the simple dipole model of the exciton coupling, the oscillator strengths ( $F_+$  and  $F_-$ ) of the exciton transitions should sum to twice the monomer oscillator strength [16]. The ratio  $F_+/F_-$  gives the angle,  $\beta$ , between the two monomer dipoles by Eqn. 2:

$$\frac{F_+}{F_-} = \frac{1 + \cos \beta}{1 - \cos \beta} \quad (2)$$

The total  $Q_y$  absorption band is resolved into these two exciton absorption bands by writing expressions for  $A_{||}$  and  $A_{\perp}$  including both exciton components,  $A_+$  and  $A_-$ :

$$\begin{aligned} A_{||} &= A_+ \cos^2 \theta_+ + A_- \cos^2 \theta_- \\ A_{\perp} &= \frac{1}{2} A_+ \sin^2 \theta_+ + \frac{1}{2} A_- \sin^2 \theta_- \end{aligned} \quad (3)$$

TABLE I

## ANGLES OF THE TRANSITION DIPOLES TO THE LONG AXIS OF LIGHT-HARVESTING PROTEINS

The values are based on Eqn. 1. The dichroic ratio is given in parentheses. See text for explanation of  $Q_x$  absorption bands  $A_{12x}$  and  $A_{3x}$ .

Wavelength (nm)		$\theta$ (degrees)		
		$f = 0.5$	$f = 0.75$	$f = 1.0$
LH-R26	885	42.3	46.4	48.5
	(1.57)			
	835	47.6	49.9	51.1
	(1.30)			
LH-wt	593	81.8	69.3	65.1
	(0.43)			
	885	42.5	46.5	48.5
	(1.56)			
	835, 800	46.4	49.1	50.5
	(1.36)			
	610	—	71.4	66.4
	(0.38)			
	575	59.5	57.8	57.1
	(0.84)			
	510	61.5	59.1	58.0
	(0.78)			
	$A_{12x}$	—	73.8	67.9
	(0.33)			
	$A_{3x}$	40.9	45.5	47.8
	(1.65)			

The assumption of perfect orientation,  $f = 1$ , does not affect the shape of the resulting bands.  $\theta_+$  and  $\theta_-$  are found from the limiting values of dichroic ratio, 1.30 and 1.57, in Fig. 1. Eqn. 3 can be solved simultaneously for  $A_+$  and  $A_-$  at every wavelength:

$$A_+ = \frac{2A_{\perp} \cos^2 \theta_- - A_{\parallel} \sin^2 \theta_-}{\sin^2 \theta_+ \cos^2 \theta_- - \cos^2 \theta_+ \sin^2 \theta_-} \quad (4)$$

$$A_- = \frac{2A_{\perp} \cos^2 \theta_+ - A_{\parallel} \sin^2 \theta_+}{\sin^2 \theta_- \cos^2 \theta_+ - \cos^2 \theta_- \sin^2 \theta_+}$$

The resolution is plotted in Fig. 5. Taking the peak to peak separation as representative of the exciton splitting, then  $\Delta\nu = 200 \text{ cm}^{-1}$ . Using the relative areas under  $A_+$  and  $A_-$  to estimate the oscillator strengths, then  $F_+/F_- = 1.53$  and  $\beta = 78^\circ$  from Eqn. 2.

In the  $Q_x$  region of the LH-R26 protein no change in dichroic ratio across the band was observed. The CD in this band also shows a double Cotton effect, indicating exciton splitting of this transition. Therefore, the two exciton transition dipoles are tilted at the same angle to the particle axis. For the LH-wt protein, with one additional bacteriochlorophyll/molecule of complex, the change in linear dichroism across the band allows resolution into a band representing the two bacteriochlorophylls which are coupled and a band representing the third bacteriochlorophyll. The ratio of areas under the curves should be 2 : 1.




Fig. 5. Resolution of the  $Q_y$  region of LH-R26 absorption.  $2A_{\perp} + A_{\parallel}$  is the sum of experimental values from Fig. 1.  $A_{+}$  and  $A_{-}$  are the exciton components. Resolution is based on Eqn. 4 with  $\theta_{+} = 51.1^{\circ}$  ( $D_{+} = 1.30$ ) and  $\theta_{-} = 48.5^{\circ}$  ( $D_{-} = 1.57$ ).

The choice of limiting dichroic ratios is more difficult than for the  $Q_y$  band. We define  $A_{12x}$  and  $D_{12x}$  as the absorption and dichroic ratio due to the coupled bacteriochlorophylls;  $A_{3x}$  and  $D_{3x}$  are similarly defined for the third bacteriochlorophyll. On the long wavelength side  $D_{12x} = 0.33$  is chosen in accord with the LH-R26 values. The  $A_{\parallel}$  and  $A_{\perp}$  are then fit to  $A_{12x}$  and  $A_{3x}$  using Eqn. 4 for different values of  $D_{3x}$  until the area under  $A_{3x}$  is half the area




Fig. 6. Resolution of the  $Q_x$  region of LH-wt absorption.  $2A_{\perp} + A_{\parallel}$  is from Fig. 2.  $A_{12x}$  is absorption attributed to the two bacteriochlorophylls absorbing at 850 nm.  $A_{3x}$  is absorption attributed to bacteriochlorophyll absorbing at 800 nm. Resolution is based on Eqn. 4 with  $\theta_{12x} = 67.8^{\circ}$  ( $D_{12x} = 0.33$ ) and  $\theta_{3x} = 47.8^{\circ}$  ( $D_{3x} = 1.65$ ).



under  $A_{12x}$ . The fit is plotted in Fig. 6.  $D_{3x}$  is found to be 1.65. The  $\lambda_{\max}$  for  $A_{12x}$  is 593 nm in agreement with the  $\lambda_{\max}$  of the LH-R26 protein.

## Conclusion

From these simple experiments we have been able to deduce information concerning the geometry of the transition dipoles in the light-harvesting proteins. Fig. 7 summarizes the data. The LH-R26 protein contains two bacteriochlorophylls; both  $Q_x$  and  $Q_y$  absorption transitions are coupled. The  $Q_y$  absorption band is resolved into its exciton components which are tilted at approximately 48 and 51 degrees to the particle axis (Fig. 7a). In the plane containing the two  $Q_y$  exciton dipoles, the monomer dipoles can be found by using the simple point dipole theory for the exciton interaction (Fig. 7b). The  $Q_x$  exciton transitions are tilted at 65 degrees to the particle axis but cannot be resolved. The LH-wt protein contains an additional bacteriochlorophyll and a carotenoid (Fig. 7c). The carotenoid transition dipole is tilted at  $58^\circ$  to the particle axis. The  $Q_x$  and  $Q_y$  transitions of the third bacteriochlorophyll are not coupled to the other two bacteriochlorophylls; they are tilted by 48 and 51




Fig. 7. Orientations of the transition dipoles are sketched with respect to the particle stretch axis, Z, using vector representation of the transition dipoles. (a)  $A_-$  and  $A_+$  represent the two exciton components of the 853 nm band of the LH-R26 protein. They are orthogonal, and the normal to the plane containing them makes an angle of  $43^\circ$  with Z. The exciton components of the  $Q_x$  band make an angle of  $65^\circ$  with Z. (b) In the plane containing  $A_+$  and  $A_-$ , the two monomer transition dipoles,  $A_1$  and  $A_2$ , are shown. (c) Additional components present in the LH-wt protein are shown. The square represents the plane of the third bacteriochlorophyll to which the 800 nm absorption band is attributed.

degrees to the particle axis, such that the normal to the porphyrin plane is tilted by 67 degrees to that axis. More information is needed to obtain a complete picture of the geometry of the pigments. A detailed study of the fluorescence polarization will be used to extend this picture of bacteriochlorophyll arrangement.

## Acknowledgements

We thank Dr. W.W. Parson for suggesting the method of orienting the proteins and Dr. Jacques Breton for many helpful discussions and encouragement. This research was supported by the Division of Biomedical and Environmental Research of the U.S. Department of Energy.

## References

- 1 Thornber, J.P. (1970) *Biochemistry* 9, 2688—2698
- 2 Clayton, R.K. and Clayton, B.J. (1972) *Biochim. Biophys. Acta* 283, 492—504
- 3 Fraker, R.J. and Kaplan, S. (1972) *J. Biol. Chem.* 247, 2732—2737
- 4 Moskalenko, A.A. and Erokhin, Y.E. (1978) *FEBS Lett.* 87, 254—256
- 5 Feick, R. and Drews, G. (1978) *Biochim. Biophys. Acta* 501, 499—513
- 6 Sauer, K. and Austin, L.A. (1978) *Biochemistry* 17, 2011—2019
- 7 Cogdell, R.J. and Thornber, J.P. (1978) Abstracts of the Ciba Foundation Symposium No. 61, 'Chlorophyll Organization and Energy Transfer in Photosynthesis', p. 8
- 8 Rafferty, C.H. and Clayton, R.K. (1978) *Biochim. Biophys. Acta* 502, 51—60
- 9 Sauer, K. (1975) in *Bioenergetics of Photosynthesis* (Govindjee, ed.), pp. 115—181, Academic Press, New York
- 10 Clayton, R.K. and Wang, R. (1971) *Methods Enzymol* 23, 696—704
- 11 Heathcote, P. and Clayton, R.K. (1977) *Biochim. Biophys. Acta* 459, 506—515
- 12 Land, E.H. (1951) *J. Opt. Soc. Am.* 41, 957—963
- 13 Fraser, R.D.B. (1953) *J. Chem. Phys.* 21, 1511—1515
- 14 Fraser, R.D.B. (1958) *J. Chem. Phys.* 28, 1113—1115
- 15 Beer, M. (1956) *Proc. Roy. Soc. (London)* A236, 136—140
- 16 Dratz, E.A., Schultz, A.J. and Sauer, K. (1967) *Brookhaven Symp. Biol.* 19, 303—318