

BBA 41201

PICOSECOND FLUORESCENCE KINETICS IN SPINACH CHLOROPLASTS AT ROOM TEMPERATURE

EFFECTS OF Mg^{2+}

JOHN A. NAIRN *, WOLFGANG HAEHNEL **, PAUL REISBERG and KENNETH SAUER

Department of Chemistry and Laboratory of Chemical Biodynamics, Lawrence Berkeley Laboratory, University of California, Berkeley, CA 94720 (U.S.A.)

(Received April 28th, 1982)

(Revised manuscript received August 17th, 1982)

Key words: Photosynthesis; Fluorescence kinetics; Mg^{2+} ; Excitation transfer; (Spinach chloroplast)

Single-photon timing with picosecond resolution is used to investigate the effect of Mg^{2+} on the room-temperature fluorescence decay kinetics in broken spinach chloroplasts. In agreement with an earlier paper (Haehnel, W., Nairn, J.A., Reisberg, P. and Sauer, K. (1982) *Biochim. Biophys. Acta* 680, 161–173), we find three components in the fluorescence decay both in the presence and in the absence of Mg^{2+} . The behavior of these components is examined as a function of Mg^{2+} concentration at both the F_0 and the F_{max} fluorescence levels, and as a function of the excitation intensity for thylakoids from spinach chloroplasts isolated in the absence of added Mg^{2+} . Analysis of the results indicates that the subsequent addition of Mg^{2+} has effects which occur at different levels of added cation. At low levels of Mg^{2+} (less than 0.75 mM), there appears to be a decrease in communication between Photosystem (PS) II and PS I, which amounts to a decrease in the spillover rate between PS II and PS I. At higher levels of Mg^{2+} (about 2 mM), there appears to be an increase in communication between PS II units and an increase in the effective absorption cross-section of PS II, probably both of these involving the chlorophyll *a/b* light-harvesting antenna.

Introduction

The addition of cations to broken chloroplasts induces changes in the primary processes of photosynthesis. These changes include: (1) a 2-fold increase in the room-temperature fluorescence yield of DCMU-poisoned chloroplasts [1–4], (2) an in-

crease in the 685 nm fluorescence at low temperature relative to the 735 nm fluorescence [2–4], (3) an increase in the PS II quantum efficiency [2–4], and (4) a decrease in the PS I quantum efficiency [2–4]. Murata [2–4] postulated that cations decrease the rate of spillover from PS II to PS I and that cationic regulation of this rate may be the basis of the State 1-to-State 2 transition observed in chloroplasts of intact cells [5,6]. In his model, State 1 (the dark state) is analogous to the state of high cation concentration, with low PS II-to-PS I spillover; State 2 is analogous to the state of low cation concentration, with high PS II-to-PS I spillover. More recent work suggests that the cation effect on energy distribution between PS II and PS

* Present address: E.I. du Pont de Nemours and Co., Inc., Central Research and Development Department, Experimental Station, Wilmington, DE 19898, U.S.A.

** Present address: Ruhr-Universität Bochum, Biochemie der Pflanzen, Postfach 10 21 48, D-4630 Bochum 1, F.R.G.

Abbreviations: DCMU, 3-(3',4'-dichlorophenyl)-1,1-dimethyl urea; Hepes, *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid; Chl, chlorophyll; PS, photosystem.

I is more complicated. Butler and Kitajima [7] concluded from fluorescence induction data at low temperature that, in addition to decreasing the rate of PS II-to-PS I spillover, Mg^{2+} increases the absorption cross-section of PS II. The analysis of fluorescence data by Henkin and Sauer [8] indicated that the only effect of Mg^{2+} is an increase in the absorption cross-section of PS II.

Joliot and Joliot [9] reported that the fluorescence induction curve for intact chloroplasts in whole cells of *Chlorella pyrenoidosa* displays a sigmoidal rise. They attributed the sigmoidicity to the possibility of energy transfer between PS II units. They proposed a theoretical relation between the probability of transfer between PS II units, ρ , and the shape of the fluorescence induction curve; the fluorescence induction curve from intact chloroplasts indicates that $\rho = 0.55$ [9]. Experiments with broken chloroplasts show that the fluorescence induction curve is sigmoidal in the presence of Mg^{2+} , but exponential in the absence of Mg^{2+} ; the two curves correspond to $\rho = 0.5$ – 0.6 in the presence of Mg^{2+} , with ρ decreasing to a low value in the absence of Mg^{2+} [10–13]. The conclusion is that added Mg^{2+} enables energy transfer to occur between PS II units.

Several possibilities have been presented for explaining the mechanism of the cationic regulation of energy distribution. Izawa and Good [14] found that chloroplasts isolated in low-salt medium have unstacked thylakoid membranes and that the addition of salts induces thylakoid stacking. This stacking is correlated with increased light scattering and with the fluorescence increases described above [15,16]. Murakami and Packer [15] and Murata [16] concluded that thylakoid stacking may be the mechanism behind the cation effect and the State 1-to-State 2 transition. More recent experiments show that thylakoid stacking and fluorescence yield changes are separable phenomena; i.e., one effect can be induced independently of the other [17,18]. Experiments with mutants indicate that an Mg^{2+} effect and a State 1-to-State 2 transition are absent in photosynthetic organisms that lack the Chl *a/b* light-harvesting protein [19]. The conclusion is that an interaction between Mg^{2+} and the Chl *a/b* light-harvesting protein induces some change which can control the energy distribution between PS II and PS I [19–21].

In a recent paper [22], we described picosecond resolution of the fluorescence decay kinetics in spinach chloroplasts. The fluorescence decay was found to be characterized by three exponential phases. The three phases can be qualitatively interpreted as follows (see Ref. 22 for a more quantitative analysis; also Reisberg, P., Nairn, J.A., Haehnel, W. and Sauer, K., unpublished observations): (1) One slow phase (1–2 ns) is due to radical-pair recombination in PS II of the oxidized primary electron donor, P-680^+ , and the reduced primary electron acceptor, I^- , which is believed to be a pheophytin molecule [23–26]. The above recombination occurs with a high yield when the secondary electron acceptor Q [27] is reduced [22–26]. (2) Two faster phases are due to excitation that is lost prior to reaching the reaction center. Of these two faster phases, the fastest (50–100 ps) is kinetically controlled by the rate of excitation transfer from the Chl *a* antenna of PS II (Chl a_2) to the reaction center of PS II, and at least a portion of the slower one (400–750 ps) is kinetically controlled by the rate of excitation transfer from the Chl *a/b* light-harvesting antenna to the reaction center of PS II. These two types of chlorophyll antennae make up the PS II antenna in Butler's tripartite model [28]. (3) The room-temperature fluorescence is assumed to be mostly from the PS II antenna; i.e., from the Chl a_2 antenna and the Chl *a/b* light-harvesting antenna. PS I fluorescence may make a contribution, but this should not affect the qualitative interpretation of the above three phases.

In this paper, we report measurements of the three components of the fluorescence decay as a function of the concentration of Mg^{2+} . Experiments were done at the F_0 level and at the F_{max} level in spinach chloroplasts at room temperature. The F_0 level corresponds to the state where all of Q is oxidized (i.e., all PS II reaction centers open), and the F_{max} level corresponds to the state where all of Q is reduced (i.e., all PS II reaction centers closed). To examine the extent of energy transfer between PS II units, it is necessary to look at the fluorescence decay when some of the PS II reaction centers are open and some are closed. For experiments in the partially closed state, we have measured the intensity dependence of the fluorescence decay kinetics for spinach chloroplasts in

the absence of Mg^{2+} . We find that most of our data can be explained by assuming that Mg^{2+} has two effects. The addition of Mg^{2+} to thylakoids from broken spinach chloroplasts isolated in an Mg^{2+} -free buffer first decreases the rate of energy transfer or spillover from PS II to PS I; this first effect saturates at low concentrations of Mg^{2+} (less than 0.75 mM). A second effect, saturating at about 2 mM Mg^{2+} , causes an increase in both the absorption cross-section and the extent of energy transfer between PS II units.

Materials and Methods

Broken spinach chloroplasts were isolated by the method described previously [22]. Fresh spinach leaves, grown either in a growth chamber or in a greenhouse, were ground in a blender for 10 s in 50 mM Hepes-NaOH buffer (pH 7.5) with 0.4 M sucrose and 10 mM NaCl followed by 2 min of centrifugation at $2000 \times g$. After one wash with fresh grinding buffer, the chloroplasts were kept for 20 min at 0°C in 10 mM Hepes-NaOH buffer (pH 7.5) with 0.1 M sucrose and 10 mM NaCl. After centrifugation for 5 min at $2000 \times g$, portions of the pellet were resuspended in several different buffers. Each buffer contained 10 mM Hepes-NaOH (pH 7.5), 0.1 M sucrose and 5 mM NaCl. Each buffer also either had no Mg^{2+} , or contained a concentration of MgCl_2 equal to the concentration desired for the fluorescence measurement. The chloroplasts were allowed to equilibrate in these buffers for at least 1 h. The chlorophyll concentration was adjusted to 18 μg Chl/ml by dilution with the appropriate resuspending buffer. For experiments at the F_0 level, we added 1.25 mM potassium ferricyanide as electron acceptor, 1.25 mM potassium ferrocyanide to control the redox potential, and 2.5 $\mu\text{g}/\text{ml}$ gramicidin D as uncoupler. These levels of added K^+ and polyvalent anions do not by themselves affect the room-temperature fluorescence properties of broken spinach chloroplasts. The uncoupler was added to prevent the slow formation of a pH gradient across the thylakoid membrane, which has been reported to cause a decline in the fluorescence yield [29]. The chloroplast sample was rapidly stirred in a $1 \times 1\text{cm}$ cuvette, and each sample was replaced every 10 min if more data accumulation

was needed. For experiments at the F_{max} level, we added 12.5 μM DCMU and 2 mM hydroxylamine hydrochloride. To close the reaction centers the sample was illuminated with about 10 flashes of saturating light immediately before the lifetime measurement. The intensity-dependence experiment was done like an F_0 experiment, except that ferri- and ferrocyanide were omitted from the resuspending buffer. All measurements were carried out at room temperature (20 – 22°C), and the cuvette was painted black except for a window for the exciting beam and a window in the direction of the photomultiplier. This masking was necessary to eliminate a broadening of the apparent excitation pulse shape due to reflections.

The excitation pulse was provided by a Spectra Physics synchronously pumped mode-locked dye laser which is composed of an SP 171 argon ion laser, an SP 362 mode locker, and a modified SP 375 dye laser. The output pulses of this laser have a full-width half-maximum duration of about 15 ps (as determined by zero background, second-harmonic generation [30]). All experiments used the laser dye rhodamine 6G with excitation pulses at 620 nm.

The single-photon timing detection system and the methods of numerical analysis are described elsewhere [22,31,38]. All of the data analyses presented here result from a resolution of the fluorescence decay kinetics into a sum of exponentials.

TABLE I

LIFETIMES AND RELATIVE YIELDS OF THE FLUORESCENCE FROM SPINACH CHLOROPLASTS

These data show the effects of adding 5 mM Mg^{2+} at both the F_0 and F_{max} levels. All results are from chloroplasts isolated from a single set of spinach leaves. The yield figures are normalized such that $\Sigma\phi$ at F_{max} in the presence of Mg^{2+} equals 100 (data from Ref. 22).

Level	No Mg^{2+}			+ 5 mM Mg^{2+}		
	τ (ps)	ϕ	$\Sigma\phi$	τ (ps)	ϕ	$\Sigma\phi$
F_0	130	4.4	17	100	2.5	25
	360	9.4		420	19.5	
	1500	3.2		1200	3.0	
F_{max}	160	3.5	44	50	1.0	100
	530	20.7		750	17.0	
	1700	19.8		2000	82.0	

That is, the time dependence of the fluorescence decay is given by

$$F(t) = \sum_{i=1}^3 \alpha_i \exp(-t/\tau_i) \quad (1)$$

where α_i and τ_i are the amplitude and the lifetime of the i th component, respectively. The yield of the i th component is equal to $\alpha_i \tau_i$; this yield represents the total number of photons emitted in the i th phase.

We estimate that our fluorescence lifetime measuring system can resolve fluorescence lifetimes as short as 25 ps [22]. Repeated measurements on a given sample result in lifetimes that agree to within 50 ps. Day-to-day variations are larger. Lifetimes and yields sometimes differ as much as 20–30%. The values in Table I were obtained on a single occasion, and comparisons can be made fairly between the results under different experimental conditions. The experiments reported in the figures were done on different occasions, and caution should be used in comparing the exact lifetimes between separate experiments.

Results

Mg^{2+} dependence of F_0 level fluorescence

Table I summarizes the results from our earlier paper [22] which have been renormalized to facilitate comparison of the yields between different experiments. The effects of adding Mg^{2+} to spinach chloroplasts at the F_0 level are: (1) a decrease in the lifetime of the slow phase with no change in its yield, (2) a slight increase in the lifetime of the middle phase accompanied by a doubling of the yield, and (3) a decrease in the yield of the fast phase. The change in the fast phase, however, may be within the uncertainty of our measurements, because the fast phase is the most difficult phase to resolve. The dependence of the three lifetimes on Mg^{2+} concentration is plotted in Fig. 1. The slow phase increases somewhat at low levels of Mg^{2+} and then decreases to its final value by $[Mg^{2+}] = 2$ mM. We note that, like the fast phase, the slow phase is a small part of the total F_0 decay and is difficult to resolve. The precise details of the change in the slow phase lifetime will need confirmation, but we generally




Fig. 1. Lifetimes of the components of the fluorescence decay in spinach chloroplasts at the F_0 level as a function of the concentration of Mg^{2+} .

observe a decrease in the slow phase lifetime upon the addition of 5 mM Mg^{2+} . The lifetimes of the fast and middle phases show only minor changes.

In Fig. 2 are plotted the total yield and the yield of each component vs. Mg^{2+} concentration. The total yield increases about 30%, saturating at $[Mg^{2+}] = 0.75$ mM; this increase is in good agreement with the results of Henkin and Sauer [8]. The changes in the individual decay components are surprisingly complex. The yield of the slow phase




Fig. 2. Total yield and yields of the components of the fluorescence decay in spinach chloroplasts at the F_0 level as a function of the concentration of Mg^{2+} . Yields are normalized such that $\Sigma \phi$ in the absence of Mg^{2+} equals 100.




Fig. 3. Lifetimes of the components of the fluorescence decay in spinach chloroplasts at the F_{\max} level as a function of the concentration of Mg^{2+} .

increases 4-fold, peaking at $[Mg^{2+}] = 0.75$ mM, and then decreases to nearly its original value. The decrease is complete at about $[Mg^{2+}] = 2$ mM. Despite difficulty in resolving the lifetime of the slow phase, the rise and fall of its yield was observed to be similar in each sample investigated. It is, therefore, likely that the rise and fall of the lifetime of the slow phase mentioned above is real. The yield of the middle phase remains approximately constant up to 1 mM and then approxi-




Fig. 4. Total yield and yields of the components of the fluorescence decay in spinach chloroplasts at the F_{\max} level as a function of the concentration of Mg^{2+} . Yields are normalized such that $\Sigma\phi$ in the absence of Mg^{2+} equals 100.




Fig. 5. Lifetimes of the components of the fluorescence decay in spinach chloroplasts isolated in the absence of Mg^{2+} as a function of laser intensity

mately doubles, with the doubling nearly complete by $[Mg^{2+}] = 2$ mM.

Mg^{2+} dependence of the F_{\max} level fluorescence

The effects on the F_{\max} level resulting from increasing the concentration of Mg^{2+} added to broken spinach chloroplasts are: (1) an increase in the lifetime of the slow phase, accompanied by a 4-fold increase in its yield, (2) an increase in the lifetime of the middle phase, accompanied by a slight decrease in its yield, and (3) changes in the fast phase which probably do not lie outside the uncertainty of our measurement. The fast phase in




Fig. 6. Total yield and yields of the components of the fluorescence decay in spinach chloroplasts isolated in the absence of Mg^{2+} as a function of laser intensity. Yields are normalized such that $\Sigma\phi$ at the highest laser intensity equals 100.

the F_{\max} level is especially difficult to resolve because it is a very small component relative to the other phases. The lifetimes of the three components vs. Mg^{2+} concentrations are plotted in Fig. 3. The lifetime of the slow phase increases from 1170 to 1600 ps, saturating at $[\text{Mg}^{2+}] = 0.75$ mM or less. This saturation of the effect at $[\text{Mg}^{2+}] = 0.75$ mM is reproducible for the slow phase lifetime. The lifetime of the middle phase increases only slightly from 380 to 430 ps; the increase occurs at low Mg^{2+} concentration (less than 0.75 mM). We have always seen an increase in this lifetime at the F_{\max} level upon the addition of Mg^{2+} . The increase is sometimes small, and the increase presented in Table I represents about the maximum effect.

The total yield and the yield of each component are plotted vs. Mg^{2+} concentration in Fig. 4. The total yield doubles, saturating at $[\text{Mg}^{2+}] = 2$ mM; this increase is in close agreement with the results of Henkin and Sauer [8]. All of the increase is accounted for by a 4-fold increase in the yield of the slow phase; this increase also saturates at $[\text{Mg}^{2+}] = 2$ mM. The only other effect is a slight decrease in the yield of the middle phase, saturating at about $[\text{Mg}^{2+}] = 1$ mM.

Intensity dependence of the fluorescence decay kinetics in the absence of Mg^{2+}

The effect of intensity on the three kinetic components of spinach chloroplasts in the presence of Mg^{2+} is plotted in Figs. 6–8 in Ref. 22. The results show a smooth transition from the F_0 values to the F_{\max} values given in Table I. Here, we have repeated the same experiment in the absence of Mg^{2+} . The results plotted in Figs. 5 and 6 show smooth transitions between limits similar to the F_0 and F_{\max} values given in Table I. The lifetime of the slow phase is nearly constant at about 1350 ps. The slow phase yield increases 8.3-fold. The lifetime and yield of the middle phase increase somewhat, the yield increasing about 60–70%. The lifetime of the fast phase remains constant, and its yield decreases.

Discussion

Comparison with other Mg^{2+} -dependent fluorescence lifetime measurements

Because we have resolved three fluorescence

decay components where other studies have resolved only two components [32–34] or one component [35], it is difficult to compare our results quantitatively to literature results. A qualitative comparison, however, reveals that our data can be reconciled quite well with other Mg^{2+} -dependent fluorescence lifetime measurements [32–35]. Searle et al. [33] looked at the effect of adding Mg^{2+} to wild-type barley chloroplasts at both the F_0 and F_{\max} levels. At F_0 , they saw very small changes in the lifetimes of two components and a slight increase in the yield of their slow component (600 to 650 ps). Our result for F_0 agrees with this result, if we note that their slow phase is probably an average of our middle and slow phases. The effect of adding Mg^{2+} at the F_{\max} level has been examined for chloroplasts from wild-type barley [33] and from peas [32,34]. All three studies [32–34] recorded increases in yields which predominate in the slow part of the fluorescence. The slow fluorescence lifetime was found either to increase [32,33] or to remain constant [34]. Our results are in essential agreement with these results as well. These previously reported Mg^{2+} effects, however, are generally smaller, because a two-component analysis averages some of the middle phase into the slow phase.

Moya et al. [35], using the technique of phase fluorimetry with a one-component analysis, measured the fluorescence lifetime as a function of intensity both in the presence and absence of Mg^{2+} . A plot of their average lifetime vs. total yield showed that in the absence of Mg^{2+} the average lifetime is proportional to the total yield, and it increases from 0.4 ns in the all-open state to 1.0 ns in the all-closed state. A plot of an average lifetime calculated from our intensity-dependent data in the absence of Mg^{2+} (see Figs. 5 and 6) by the formula:

$$\tau_{\text{mean}} = \sum_{i=1}^3 \tau_i \phi_i / \sum_{i=1}^3 \phi_i \quad (2)$$

is identical to the results of Moya et al. [35] (plot not shown). The deceptively simple linear relation between average lifetime and total yield has influenced the conclusions of several authors.




Fig. 7. Working model for the origin of fluorescence in chloroplasts. k_F , rate constant for fluorescence; k_D , k_d and k_d' , rate constants for radiationless decay; k_{T32} , k_{T23} , k_{T31} , k_{T20} and k_{T02} , rate constants for energy transfer; k_c and k_r , rate constants for electron transfer; P-680, reaction center PS II; Ph, pheophytin; Q, secondary electron acceptor in PS II.

Model for the origin of fluorescence

The results of earlier work in our laboratory [22,38] led to the working model for the origin of fluorescence shown in Fig. 7. The basic structure of the model is derived from the tripartite model proposed by Butler [28]. The major addition is the explicit inclusion of the electron-transfer processes involving pheophytin, which occur in the reaction center of PS II [23–26]. Fig. 7 illustrates the state where the PS II reaction center is closed. Electron transfer beyond pheophytin is blocked, but P-680⁺ and pheophytin[−] (Ph[−]) may undergo recombination resulting in either a fast 'delayed' fluorescence [23–26], triplet formation or radiationless decay to the ground state. If the PS II reaction center is open, electron transfer will continue beyond Q with a high probability and the delayed fluorescence will be quenched. A detailed kinetic analysis of this model (Reisberg, P., Nairn, J.A., Haehnel, W. and Sauer, K., unpublished observations) shows that the total fluorescence which is emitted from the Chl *a/b* proteins and the Chl *a*₂ proteins can be described by a sum of three exponentials. The origin of each phase is a complex interaction among the rate constants in Fig. 7, but they can be qualita-

tively described as follows: (1) The fastest phase (about 100 ps) is kinetically controlled by the decay processes of the Chl *a*₂ antenna, and these processes are dominated by the transfer rate, k_{T20} , from the Chl *a*₂ antenna to the reaction center of PS II. (2) The middle phase (300–750 ps) is kinetically controlled by the decay processes of the Chl *a/b* light-harvesting antenna, and these processes are dominated by transfer rates from the Chl *a/b* to the Chl *a*₂ antenna or to PS I ($k_{T32} + k_{T31}$). (3) The yield of the slow phase is controlled by the presence of Q[−], and its lifetime is affected by two factors. The first is the rate of charge recombination between P-680⁺ and pheophytin[−] and the second is the rate of fluorescence quenching from the chlorophyll antenna.

Interpretation of the Mg²⁺ effects

By investigating the influence of Mg²⁺ on fluorescence induction, Henkin and Sauer [8] found two distinguishable effects of the ion on fluorescence. These effects saturated at 0.5 and 2.5 mM Mg²⁺. Our results presented above also require at least two separate Mg²⁺ effects saturating at approximately these same concentrations. This is most obviously apparent in Fig. 2 where the yield of the slow phase increases between 0.0 and 0.75 mM Mg²⁺ and then decreases until 2.0 mM Mg²⁺. We will discuss these two effects of Mg²⁺ on our data separately.

Effects of high concentrations, [Mg²⁺] > 0.5 mM. In our previous paper [22] we reported a doubling of the slow phase lifetime as PS II reaction centers go from all open to all closed in the presence of Mg²⁺ (see Table I and Ref. 22, Fig. 3). We found it difficult to simulate this doubling without including connections between PS II units. If some PS II reaction centers are open and PS II units are capable of intercommunication, excitation returning to the antenna after a charge recombination may get transferred to an open reaction center and become quenched. The net effect is a shortening of the slow phase lifetime in comparison with the behavior in the absence of intercommunication. Thus, the intensity dependence of the slow phase lifetime can be used to examine the extent of energy transfer between PS II units. In contrast with the results of the experiment in the presence of 5 mM Mg²⁺ [22] (see also Table I), we

find that in the absence of Mg^{2+} the lifetime of the slow phase is essentially constant (Figs. 5 and 6). This suggests that Mg^{2+} is required for communication to occur between PS II units. This conclusion is consistent with the results of fluorescence induction experiments in the presence and absence of Mg^{2+} [10–13].

From the above result, we expect a shortening of the lifetime and a decrease in yield of the slow phase as Mg^{2+} is added at the F_0 level, but no corresponding changes should occur at the F_{max} level. A lifetime shortening and yield decrease in the slow phase do not occur at the F_{max} level because there are no open reaction centers to serve as excitation quenchers, regardless of whether the reaction centers are interconnected. Inspection of Figs. 1–4 shows that in the range 0.5–2.0 mM Mg^{2+} there is indeed a decrease in the lifetimes and yields of the slow phase at the F_0 level, but there are no decreases at the F_{max} level. We believe, therefore, that the transition from separate PS II units to an interconnected state occurs in this concentration range.

Effects of low concentrations, $[\text{Mg}^{2+}] < 0.75$ mM. Low concentrations of Mg^{2+} cause increases in the slow phase lifetimes at F_0 and F_{max} (see Figs. 1 and 3) which are completed by $[\text{Mg}^{2+}] = 0.75$ mM. Two possibilities for this Mg^{2+} effect are either a decrease in a radiationless decay rate (k_D , k_d , and/or k_d') or a decrease in the PS II-to-PS I spillover rate (k_{T31}).

A connection between the behavior of the two photosystems was proposed by Satoh et al. [36], who found an 18% increase in PS I activity when PS II is inhibited. Although the fractional increase of the PS I rate is independent of Mg^{2+} concentration, deconvolution of absolute rates indicates that the spillover rate is about 50% less in the presence of Mg^{2+} than in its absence [28]. Such a decrease would lessen the PS I contribution to the quenching of the excitation in the PS II antenna. This could then account for the increases we observe in the slow phase lifetimes.

An alternative explanation for their results is a direct effect of Mg^{2+} on the electron-transfer kinetics in PS I. In this case, an Mg^{2+} -induced decrease in the radiationless decay rates (k_d and k_d'), as suggested by Melis and Ow [37], would satisfactorily explain our data. It is not possible to

distinguish this effect from changes in spillover rates without parallel measurements of PS I and PS II activities.

A decrease in spillover from PS II to PS I, k_{T31} in Fig. 7, should also be reflected in the middle phase lifetime. This is because, as mentioned above, the middle phase lifetime is approximately equal to $1/(k_{T32} + k_{T31})$; a decrease in k_{T31} should, therefore, result in an increase in the lifetime. At both F_0 and F_{max} in Table I and in Figs. 1 and 3, we see a slight increase in the middle phase lifetime. The changes are usually small and completed by $[\text{Mg}^{2+}] = 0.75$ mM. It is possible, therefore, that the lifetime increases in the middle and slow phases, both of which are completed at low Mg^{2+} concentrations, are due to changes in the PS II-to-PS I spillover rate.

Additional Mg^{2+} effects. Both mechanisms discussed above, by which low levels of Mg^{2+} induce an increase in the slow phase lifetime, should also cause a concomitant increase in the slow phase yield. However, the yield of the slow phase at F_{max} continues to increase up to $[\text{Mg}^{2+}] = 2$ mM, even though the lifetime effects are completed at a lower concentration of Mg^{2+} . A possible explanation for this would be an increase in the absorption cross-section of PS II which would cause the yield of the slow phase to increase without increasing its lifetime. Thus, the Mg^{2+} effects occurring at higher Mg^{2+} levels ($[\text{Mg}^{2+}] = 1$ –2 mM) may result from an Mg^{2+} -induced increase in the absorption cross-section of PS II. The conclusion that both spillover changes and absorption cross-section changes occur upon the addition of Mg^{2+} is in agreement with the work of Butler and Kitajima [7]. The yield of the middle phase in the F_0 experiment increases with $[\text{Mg}^{2+}]$ until about $[\text{Mg}^{2+}] = 2$ mM. This effect is consistent with an increase of the absorption cross-section of PS II. However, we did not observe an increase in the yield of the middle phase in the F_{max} experiment.

Although it is difficult to be precise about changes in the fast phase, we consistently observed a decrease in its lifetime upon the addition of Mg^{2+} . As Mg^{2+} increases communication between PS II units, we postulate that it also strengthens the connection between the Chl a_2 antenna and the PS II reaction center. An Mg^{2+} -induced increase in the rate constant k_{T20} would

then account for the observed shortening of the fast phase.

Conclusion

The addition of Mg^{2+} to broken spinach chloroplasts isolated in the absence of Mg^{2+} has two effects which occur in different concentration ranges. As the Mg^{2+} concentration is increased from 0.0 to 0.75 mM, the rate constant for transfer between PS II and PS I decreases. It is possible that the changes occurring in this concentration range may instead or in addition affect the kinetics of electron transfer or the rate of radiationless decay in the PS II reaction center. As the Mg^{2+} concentration is increased further to 2.0 mM, changes in the Chl *a/b* light-harvesting antenna occur which both increase the absorption cross-section of PS II and bring about communication between PS II units. There is probably also an increase in the transfer rate between the Chl *a*₂ antenna and the PS II reaction center. Most of these correlations can be understood as a consequence of effects of Mg^{2+} on the organization of the Chl *a/b* light-harvesting antenna. This conclusion is in good agreement with the results of Lieberman et al. [19] who concluded that the Chl *a/b* light-harvesting antenna is required for Mg^{2+} effects.

Acknowledgements

This work was supported in part by the Director, Office of Energy Research, Office of Basic Energy Sciences, Division of Biological Energy Conversion and Conservation, U.S. Department of Energy, under contract W-7405-ENG-48, and in part by a National Science Foundation Grant PCM 79-11251. One of us (W.H.) was supported by a grant from the Deutsche Forschungsgemeinschaft. One of us (P.R.) wishes to acknowledge a National Research Service Award from the U.S. Public Health Service. We also thank Dr. Kerry Karukstis for help with the intensity dependence experiment.

References

- 1 Homann, P.H. (1969) *Plant Physiol.* 44, 932–936
- 2 Murata, N. (1969) *Biochim. Biophys. Acta* 189, 171–181

- 3 Murata, N., Tashiro, H. and Takamiya, A. (1970) *Biochim. Biophys. Acta* 197, 250–256
- 4 Murata, N. (1971) *Biochim. Biophys. Acta* 226, 422–432
- 5 Murata, N. (1969) *Biochim. Biophys. Acta* 172, 242–251
- 6 Bonaventura, C. and Myers, J. (1969) *Biochim. Biophys. Acta* 189, 366–383
- 7 Butler, W.L. and Kitajima, M. (1975) *Biochim. Biophys. Acta* 396, 72–85
- 8 Henkin, B.M. and Sauer, K. (1977) *Photochem. Photobiol.* 26, 277–286
- 9 Joliot, A. and Joliot, P. (1964) *C. R. Acad. Sci. Paris* 258D, 4622–4625
- 10 Bennisou, P. (1974) *Biochim. Biophys. Acta* 368, 141–147
- 11 Briantais, J.-M., Verrotte, C. and Moya, I. (1973) *Biochim. Biophys. Acta* 325, 530–538
- 12 Marsho, T.V. and Kok, B. (1970) *Biochim. Biophys. Acta* 223, 240–250
- 13 Joliot, P., Joliot, A. and Kok, B. (1968) *Biochim. Biophys. Acta* 153, 635–652
- 14 Izawa, S. and Good, N.E. (1966) *Plant Physiol.* 41, 544–552
- 15 Murakami, S. and Packer, L. (1971) *Arch. Biochem. Biophys.* 146, 337–347
- 16 Murata, N. (1971) *Biochim. Biophys. Acta* 245, 365–372
- 17 Telfer, A., Nicolson, J. and Barber, J. (1976) *FEBS Lett.* 65, 77–83
- 18 Barber, J. (1979) in *Chlorophyll Organization and Energy Transfer in Photosynthesis*, Ciba Found. Symp. 61, pp. 283–298, Elsevier/North-Holland, Amsterdam
- 19 Lieberman, J.R., Bose, S. and Arntzen, C.J. (1978) *Biochim. Biophys. Acta* 502, 417–429
- 20 Armond, P.A., Arntzen, C.J., Briantais, J.-M. and Verrotte, C. (1976) *Arch. Biochem. Biophys.* 175, 54–63
- 21 Davis, D.J., Armond, P.A., Gross, E.L. and Arntzen, C.J. (1976) *Arch. Biochem. Biophys.* 175, 64–70
- 22 Haehnel, W., Nairn, J.A., Reisberg, P. and Sauer, K. (1982) *Biochim. Biophys. Acta* 680, 161–173
- 23 Klevanik, A.V., Klimov, V.V., Shuvalov, V.A. and Krasnovskii, A.A. (1977) *Dokl. Biophys. (English translation)* 236, 101–104
- 24 Klimov, V.V., Klevanik, A.V., Shuvalov, V.A. and Krasnovskii, A.A. (1977) *FEBS Lett.* 82, 183–186
- 25 Klimov, V.V., Allakhverdiev, S.I. and Pashchenko, V.Z. (1978) *Dokl. Biophys. (English translation)* 242, 166–169
- 26 Shuvalov, V.A., Klimov, V.V., Dolan, E., Parson, W.W. and Ke, B. (1980) *FEBS Lett.* 118, 279–282
- 27 Duysens, L.M.N. and Sweers, H.E. (1963) in *Microalgae and Photosynthetic Bacteria* (Japanese Society of Plant Physiologists, eds.), pp. 353–372, University of Tokyo Press, Tokyo
- 28 Butler, W.L. (1978) *Annu. Rev. Plant. Physiol.* 29, 345–378
- 29 Briantais, J.-M., Verrotte, C., Picaud, M. and Krause, G.H. (1979) *Biochim. Biophys. Acta* 548, 128–138
- 30 Ippen, E.P. and Shank, C.V. (1975) *Appl. Phys. Lett.* 27, 488–490
- 31 Nairn, J.A. (1981) *Orientation and Picosecond Energy Transfer Studies on Chlorophyll in the Photosynthetic Membrane*, Ph. D. Thesis, University of California, Berkeley. Lawrence Berkeley Laboratory Report (LBL-13827)

- 32 Barber, J., Searle, G.F.W. and Tredwell, C.J. (1978) *Biochim. Biophys. Acta* 501, 174–182
- 33 Searle, G.F.W., Tredwell, C.J., Barber, J. and Porter, G. (1979) *Biochim. Biophys. Acta* 545, 496–507
- 34 Beddard, G.S., Fleming, G.R., Porter, G., Searle, G.F.W. and Synowiec, J.A. (1979) *Biochim. Biophys. Acta* 545, 165–174
- 35 Moya, I., Govindjee, Vernotte, C. and Briantais, J.-M. (1977) *FEBS Lett.* 75, 13–18
- 36 Satoh, K., Strasser, R. and Butler, W.L. (1976) *Biochim. Biophys. Acta* 440, 337–345
- 37 Melis, A. and Ow, R.A. (1982) *Biochim. Biophys. Acta* 682, 1–10
- 38 Turko, B., Nairn, J.A. and Sauer, K. (1982) *Rev. Sci. Instrum.*, in the press