

Bioenergetics

Cumulative Contents

Regular papers

- The effect of exchange of bacteriopheophytin *a* with plant pheophytin *a* on charge separation in Y(M210)W mutant reaction centers of *Rhodobacter sphaeroides* at low temperature
A.Y. Shkuropatov, S. Neerken, H.P. Permentier, R. de Wijn, K.A. Schmidt, V.A. Shuvalov, T.J. Aartsma, P. Gast and A.J. Hoff (Russian Federation, The Netherlands) 1
- The respiratory chain of the thermophilic archaeon *Sulfolobus metallicus*: studies on the type-II NADH dehydrogenase
T.M. Bandejas, C.A. Salgueiro, H. Huber, C.M. Gomes and M. Teixeira (Portugal, Germany) 13
- Elimination of high-light-inducible polypeptides related to eukaryotic chlorophyll *a/b*-binding proteins results in aberrant photoacclimation in *Synechocystis* PCC6803
M. Havaux, G. Guedeney, Q. He and A.R. Grossman (France, USA) 21
- Probing connection of PBS with the photosystems in intact cells of *Spirulina platensis* by temperature-induced fluorescence fluctuation
D. Li, J. Xie, Y. Zhao and J. Zhao (People's Republic of China) 35
- Quantitative amino acid analysis of bovine NADH: ubiquinone oxidoreductase (Complex I) and related enzymes. Consequences for the number of prosthetic groups
S.P.J. Albracht, E. van der Linden and B.W. Faber (The Netherlands) 41
- Two EPR-detectable [4Fe–4S] clusters, N2a and N2b, are bound to the NuoI (TYKY) subunit of NADH: ubiquinone oxidoreductase (Complex I) from *Rhodobacter capsulatus*
M. Chevallet, A. Dupuis, J.-P. Issartel, J. Lunardi, R. van Belzen and S.P.J. Albracht (France, The Netherlands) 51
- The *menD* and *menE* homologs code for 2-succinyl-6-hydroxyl-2,4-cyclohexadiene-1-carboxylate synthase and *O*-succinylbenzoic acid–CoA synthase in the phylloquinone biosynthetic pathway of *Synechocystis* sp. PCC 6803
T. Wade Johnson, S. Naithani, C. Stewart Jr., B. Zybailov, A. Daniel Jones, J.H. Golbeck and P.R. Chitnis (USA) 67
- ¹⁸O isotope effect in the photosynthetic water splitting process
K. Burda, K.P. Bader and G.H. Schmid (Poland, Germany) 77
- HiPIP in *Rubrivivax gelatinosus* is firmly associated to the membrane in a conformation efficient for electron transfer towards the photosynthetic reaction centre
C. Lieutaud, W. Nitschke, A. Verméglio, P. Parot and B. Schoepp-Cothenet (France) 83
- The photochemical trapping rate from red spectral states in PSI–LHCI is determined by thermal activation of energy transfer to bulk chlorophylls
R.C. Jennings, G. Zucchelli, R. Croce and F.M. Garlaschi (Italy) 91
- Computer simulation of water in cytochrome *c* oxidase
X. Zheng, D.M. Medvedev, J. Swanson and A.A. Stuchebrukhov (USA) 99

Cytochrome <i>c</i> is transformed from anti- to pro-oxidant when interacting with truncated oncoprotein prothymosin α <i>O.V. Markova, A.G. Evstafieva, S.E. Mansurova, S.S. Moussine, L.A. Palamarchuk, M.O. Pereverzev, A.B. Vartapetian and V.P. Skulachev (Russia)</i>	109	<i>Short sequence paper</i>	
Passive penetration of nitrate through the plasma membrane of <i>Paracoccus denitrificans</i> and its potentiation by the lipophilic tetraphenylphosphonium cation <i>I. Kučera (Czech Republic)</i>	119	<i>QcrCAB</i> operon of a nocardia-form actinomycete <i>Rhodococcus rhodochrous</i> encodes cytochrome reductase complex with diheme cytochrome <i>cc</i> subunit <i>N. Sone, M. Fukuda, S. Katayama, A. Jyoudai, M. Syugyou, S. Noguchi and J. Sakamoto (Japan)</i>	125
		<i>Author Index</i>	133
		<i>Cumulative Contents</i>	135