

Rapid report

Water-gated mechanism of proton translocation by cytochrome *c* oxidase

 Mårten Wikström^{a,*}, Michael I. Verkhovsky^a, Gerhard Hummer^b
^a*Helsinki Bioenergetics Group, Institute of Biotechnology and Biocentrum Helsinki, University of Helsinki, PB 65 (Viikinkaari 1), FI-00014 Helsinki, Finland*
^b*Laboratory of Chemical Physics, NIDDK, National Institutes of Health, Room 132, Building 5, Bethesda, MD 20892-0520, USA*

Received 25 March 2003; accepted 25 March 2003

Abstract

Cytochrome *c* oxidase is essential for aerobic life as a membrane-bound energy transducer. O₂ reduction at the haem *a*₃–Cu_B centre consumes electrons transferred via haem *a* from cytochrome *c* outside the membrane. Protons are taken up from the inside, both to form water and to be pumped across the membrane (M.K.F. Wikström, *Nature* 266 (1977) 271 [1]; M. Wikström, K. Krab, M. Saraste, *Cytochrome Oxidase, A Synthesis*, Academic Press, London, 1981 [2]). The resulting electrochemical proton gradient drives ATP synthesis (P. Mitchell, *Chemiosmotic Coupling in Oxidative and Photosynthetic Phosphorylation*, Glynn Research, Bodmin, UK, 1966 [3]). Here we present a molecular mechanism for proton pumping coupled to oxygen reduction that is based on the unique properties of water in hydrophobic cavities. An array of water molecules conducts protons from a conserved glutamic acid, either to the Δ-propionate of haem *a*₃ (pumping), or to haem *a*₃–Cu_B (water formation). Switching between these pathways is controlled by the redox-state-dependent electric field between haem *a* and haem *a*₃–Cu_B, which determines the water–dipole orientation, and therefore the proton transfer direction. Proton transfer via the propionate provides a gate to O₂ reduction. This pumping mechanism explains the unique arrangement of the metal cofactors in the structure. It is consistent with the large body of biochemical data, and is shown to be plausible by molecular dynamics simulations. © 2003 Elsevier Science B.V. All rights reserved.

Keywords: Electron transfer; Proton pumping; Proton–electron coupling; Cell respiration

Subunit I of all haem–copper oxidases has a unique arrangement of redox cofactors [4–8]. Haem *a* lies about 1/3 into the membrane domain from the positively charged (P) side. Haem *a*₃ lies very close, at the same depth, and almost at a right angle to haem *a*. Cu_B is within 5 Å from the haem iron of *a*₃; together, they form a binuclear site where O₂ is bound and reduced to water (Fig. 1a). All protons pumped across the membrane are transferred across the so-called D-pathway [9], which starts with D132¹ near the negatively charged (N) side of the membrane, and continues into the membrane along polar residues and bound water molecules [9–11], finally connecting to the conserved E286 next to a hydrophobic cavity 10 Å from the binuclear site (Fig. 1b). E286 is essential for proton translocation and for enzyme activity [12,13]. At least three out of the four substrate

protons consumed at the haem *a*₃–Cu_B site per cycle are also transferred through this pathway [9,13,14].

How is chemical energy used for the mechanical translocation of individual protons across the membrane against an electrochemical gradient, in addition to charging up the membrane by selective uptake of electrons and substrate protons from opposite sides? To answer this question, we focus on how protons are transferred further from E286, how pumped protons are distinguished from consumed substrate protons, and how proton translocation is coupled to the driving redox reaction. No proton connectivity beyond E286 is seen in the X-ray structures. The Δ-propionate of haem *a*₃ (Fig. 1b) has been suggested to initiate the exit path of the pumped protons [4,10], and this has been experimentally supported [15]. Water molecules have been shown to have a key role in light-driven proton pumping by bacteriorhodopsin [16]. Statistical–mechanical calculations [11] predicted that proton transfer could be mediated by three to four water molecules transiently present inside the hydrophobic cavity between E286, the propionate, and the binuclear site (Fig. 1b). During each catalytic cycle, two water molecules are, in fact, produced at

* Corresponding author. Tel.: +358-9-191-58000; fax: +358-9-191-58001.

E-mail address: marten.wikstrom@helsinki.fi (M. Wikström).

¹ All numbering of amino acid residues refers to subunit I of cytochrome *c* oxidase from *Rhodobacter sphaeroides*. Amino acids are abbreviated by the one-letter code.




Fig. 1. (a) Overall function of cytochrome *c* oxidase. The blue arrows depict how the chemical reaction of O_2 reduction to water is orientated with respect to the positively charged P-side and the negatively charged N-side of the membrane. The red arrows show proton translocation (pumping) coupled to the chemistry. (b) A view, parallel to the membrane, of key components of the cytochrome *c* oxidase structure (adapted from the crystal structure [8] of cytochrome *c* oxidase from *R. sphaeroides*). The D-pathway transfers protons from the N-side of the membrane (a) via E286, either to be pumped across the membrane (red arrows), or to be consumed to form water at the $haem\ a_3$ – Cu_B site (blue arrows; see text). Light blue spheres depict water molecules seen in the X-ray structure. The hydrophobic cavity above E286 is predicted to transiently contain three to four water molecules by computational methods [11]. The Δ -propionate of haem a_3 (Δ) is proposed to convey pumped protons to the hydrophilic domain above the haem groups. The programme VMD was used for the molecular graphics [32].

this site as the product of O_2 reduction. Water molecules move in and out of the enzyme during the cycle and this has been shown to limit electron transfer [17]. More recent work indicates that water molecules might indeed provide proton conductivity from E286 to both the propionate and the binuclear site [10,18], consistent with the role of the D-pathway in both translocation and consumption of protons.

Water molecules behave uniquely in narrow hydrophobic cavities [19], and may not only serve as ‘proton wires’, but also as the switch between proton transfer for ‘pumping’ and water formation. Due to the scarcity of interactions with the surrounding structure, and the limited space, they tend to form orientated single file arrays where each water molecule only interacts with its two neighbours. The hydrogen bond orientation of the array determines the allowed direction of Grotthuss proton transfer [20]. Orientated water arrays are excellent unidirectional conductors of single protons [21]; the proton mobility exceeds that in bulk water by a factor of

40. Any reorientation of such an array occurs collectively to maintain the strong water–water hydrogen bonds, hence changing the allowed direction of proton transfer. Thus, alternative water chains from E286 to the propionate, or to the binuclear site, might provide a switch of the proton current to either destination (cf. Ref. [22]). We propose here that the direction of such a switch may be determined by the redox states of the electron donor (haem *a*) and the acceptor (the binuclear centre). This would provide the important link between proton translocation and the redox reaction, and make proton pumping a prerequisite for completion of the oxygen reduction chemistry, assuring tight coupling between the two. The effect of a unit charge at either redox site on a single water molecule in the cavity yields a charge–dipole interaction energy of 0.4–0.8 kcal/mol at an Fe–water distance of 8–10 Å (with $\epsilon=4$). Thus, an electric field between haem *a* and the $haem\ a_3$ – Cu_B site, parallel to the membrane, may orientate the water dipoles in

the cavity between them to form a proton transfer path from the carboxylic group of E286, either to the Δ -propionate of haem a_3 , or to the binuclear site.

Fig. 2 summarises the proposed mechanism based on redox-linked switching between proton ‘wires’ for pumping and oxygen reduction. Electron transfer from haem a to the binuclear site does not occur unless accompanied by a charge-compensating proton transfer [23,24]; however, initially, with haem a reduced and the binuclear site oxidised, there is no proton connectivity to the latter. Instead, the electric field orientates the water molecules for proton transfer from E286 to the Δ -propionate of haem a_3 (Fig. 2a). Transfer of the proton to the propionate (Fig. 2b) raises the redox potential of the binuclear centre to allow transfer of the electron (Fig. 2c). The electron transfer stabilises the ‘pumped’ proton electrostatically and inverts the electric field between the redox sites. This reorientates the water molecules to provide a proton path from E286 to the binuclear site (Fig. 2d). The conserved R481 strongly interacts with the propionate [25] (Fig. 1b). Its protonation will therefore cause deprotonation of the arginine, with transfer of the pumped proton to the hydrophilic domain on the P-side of haem a_3 (Fig. 1b), followed by reprotonation of the arginine by the propionic acid (Fig. 2d). Reprotonation of E286 via the D-pathway blocks back-transfer of the pumped proton, and if it fails, as it does in some mutants [9], proton translocation is compromised. The proton on

E286 is transferred into the binuclear site to form water where it fully neutralises the charge of the electron (Fig. 2e). As a result, the initially transferred proton is expelled towards the P-side of the membrane by electrostatic repulsion (Fig. 2f).

This mechanism is in accordance with the proton dependence of reduction of the binuclear site [23,24], the apparent dependence of the redox states of the two haems on the same proton-accepting site [2], and the dependence of electron transfer between haem a and the binuclear site on the water occupancy of the enzyme [17]. It further explains why haem a is positioned very close to the a_3 -Cu_B site, which is required for haem a to exert its electrostatic effect on the water orientation. Moreover, the short distance ensures a high electron transfer rate constant, which compensates for the presumably rate-limiting initial proton transfer via the propionate. It is noteworthy that the redox-dependent orientation of the water dipoles transfers positive charge from the proton donor (E286) to the acceptor (propionate or binuclear centre) through a displacement current [21], which further enhances the coupling between electron and proton transfer. Involvement of the K-pathway in transferring one of the substrate protons to the binuclear centre [9,13,14] is not contradictory, because this is also expected to cause release of a proton pumped via the D-pathway by the mechanism described here. Finally, the unique positioning of haem a and haem a_3 -Cu_B in the




Fig. 2. Schematic of the proposed mechanism. In (a), haem a is reduced and the haem a_3 -Cu_B site is oxidised. The electric field between the two orientates the water molecules to form a Grotthuss array for proton transfer from glutamic acid-286 to the Δ -propionate of haem a_3 . Transfer of a proton from E286 to the propionate (b) is coupled to transfer of the electron to the binuclear site (c). Due to the electron transfer, the electric field switches orientation, and the water dipoles reorientate to provide a proton transfer path from E286 to the binuclear site (d). Uptake of a substrate proton via E286 (d) into the binuclear site to form water (e) forces the first proton out of the membrane, and E286 is again reprotonated (f).

same membrane plane (Fig. 1a) ensures that the transmembrane potential will not affect the electric field between them.

To test the energetic feasibility of the proposed redox-dependent switching of water arrays, we performed a series of molecular dynamics simulations. A 30-Å diameter sphere of structure was chosen around the haem groups in subunit I of cytochrome *c* oxidase from *R. sphaeroides* (access code 1M56, Ref. [8]). The covalent link between H284 and Y288 [5], and the OH[−] ligand of Cu_B [26], were constructed using the HyperChem 7.03 programme (HyperCube Inc., Gainesville, FL), and energy-minimised using the MM+ force field. Molecular dynamics simulations were done with Amber99 or Charmm27, as applied in HyperChem. The TIP3P model [27] of water was used. The partial charges of the metal centres and their ligands were from semiempirical ZINDO/1 calculations using HyperChem. The following groups were allowed to move during the simulations: three to five water molecules in the cavity above E286, E286 itself, M107, W172, W280, V287, G283, the Δ-propionate of haem *a*₃, and the Cu_B–OH structure. Five water molecules were initially arbitrarily positioned in the cavity. Minimisation and initial simulations expelled one water molecule. In most cases, four water molecules were present, of which one was sometimes expelled in 5–200-ps simulations at 300 K with different redox states of haem *a* and Cu_B, keeping haem *a*₃ oxidised. In many simulations, the E286 carboxyl group was turned to a position towards the P-side [10,28], and tethered to the closest water molecule by constraining the distance between the carboxyl OH and water oxygens to 3.0 Å, applying a harmonic force constant of 7 kcal/mol Å², and the carboxyl OH...water O angle at 180° with a force constant of 7 kcal/mol deg². This prevented excursion of the glutamic acid side chain away from the P-side waters to contact the D-pathway on the N-side (Fig. 1b) [28]. Alternatively, and with similar results, the carboxylic side chain was held in the P-side position by constraining its distance from the haem groups. Similar results were obtained from molecular dynamics simulations including subunits I, II, and III of bovine cytochrome *c* oxidase [5] (access code 2OCC) using AMBER 4.1 (University of California, San Francisco, CA) and the AMBER94 force field [29] without constraints. In these latter simulations, several different charge parameterisations for the redox centres were tested and gave consistent results. Hence, the behaviour of the water arrays seems to be robust and not very sensitive to the precise structure, the force field, or the exact partial charges of the redox centres.

With haem *a* reduced and haem *a*₃–Cu_B oxidised, two water molecules formed a hydrogen-bonded path within picoseconds from E286 to the propionate (Fig. 3a). The second water molecule donates a hydrogen bond either directly to the propionate, or more frequently to the NH group of W172, which in turn is stably hydrogen-bonded to the propionate. This path remained stable for at least 100




Fig. 3. Proton-conducting, hydrogen-bonded water arrays from E286 to the Δ-propionate of haem *a*₃ (a), or to the OH[−] ligand of Cu_B (b), as obtained from molecular dynamics simulations with (a) haem *a* reduced and Cu_B oxidised, and (b) with haem *a* oxidised and Cu_B reduced (see text). The structure is shown from a slightly different angle in (a) and (b). Note in (a) the orientation of the water molecules next to the OH[−] ligand of Cu_B, and the hydrogen bond to G283, and in (b) the orientation of the water molecule below W172, which forms hydrogen bonds to the sulfur of M107, and the water molecule nearest to E286. Colour code: carbon (blue green); copper (Cu_B, green); oxygen (red); nitrogen (dark blue); sulfur (yellow), and hydrogen (white).

ps. In contrast, when haem *a* is oxidised and Cu_B is reduced, a hydrogen-bonded water array formed within picoseconds from E286 to the OH[−] ligand of Cu_B (Fig. 3b). Despite their limitations, these simulations clearly show that the control of water array orientation by redox state is plausible in the way required by the proposed mechanism.

It is not surprising that many features of the suggested mechanism are reminiscent of proton translocation models proposed in the past [4,10,18,24,30,31]. However, the unique gating of proton transfer by a redox-state-controlled orientation of water arrays proposed here provides a simple solution to many of the key problems that were previously unsolved. In particular, it explains how the same proton-conducting pathway may be used both for protons to be consumed in formation of water and protons to be translocated.

Acknowledgements

M.W. wishes to thank Alexei Stuchebrukhov and Janos K. Lanyi for disclosing their unpublished data. Mikael P. Johansson and William A. Eaton are acknowledged for helpful discussions. The Academy of Finland (programme 44895), Biocentrum Helsinki, the University of Helsinki, and the Sigrid Jusélius Foundation are acknowledged for financial support.

References

- [1] M.K.F. Wikström, Proton pump coupled to cytochrome *c* oxidase in mitochondria, *Nature* 266 (1977) 271–273.
- [2] M. Wikström, K. Krab, M. Saraste, Cytochrome Oxidase, A Synthesis, Academic Press, London, 1981.
- [3] P. Mitchell, Chemiosmotic Coupling in Oxidative and Photosynthetic Phosphorylation, Glynn Research, Bodmin, UK, 1966.
- [4] S. Iwata, C. Ostermeier, B. Ludwig, H. Michel, Structure at 2.8 Å resolution of cytochrome *c* oxidase from *Paracoccus denitrificans*, *Nature* 376 (1995) 660–669.
- [5] S. Yoshikawa, K. Shinzawa-Itoh, R. Nakashima, R. Yaono, E. Yamashita, N. Inoue, M. Yao, M.J. Fei, C. Peters Libeu, T. Mizushima, H. Yamaguchi, T. Tomizaki, T. Tsukihara, Redox-coupled crystal structural changes in bovine heart cytochrome *c* oxidase, *Science* 280 (1998) 1723–1729.
- [6] J. Abramson, S. Riistama, G. Larsson, A. Jasaitis, M. Svensson-Ek, L. Laakkonen, A. Puustinen, S. Iwata, M. Wikström, The structure of the heme-copper oxidase from *Escherichia coli* and its binding site for ubiquinone, *Nat. Struct. Biol.* 7 (2000) 910–917.
- [7] T. Soulimane, G. Buse, G.P. Bourenkov, H.D. Bartunik, R. Huber, M.E. Than, Structure and mechanism of the aberrant *ba*₃-cytochrome *c* oxidase from *Thermus thermophilus*, *EMBO J.* 19 (2000) 1766–1776.
- [8] M. Svensson-Ek, J. Abramson, G. Larsson, S. Törnroth, P. Brzezinski, S. Iwata, The X-ray crystal structures of wild-type and EQ(I-286) mutant cytochrome *c* oxidases from *Rhodobacter sphaeroides*, *J. Mol. Biol.* 321 (2002) 329–339.
- [9] R.B. Gennis, Multiple proton-conducting pathways in cytochrome oxidase and a proposed role for the active-site tyrosine, *Biochim. Biophys. Acta* 1365 (1998) 241–248.
- [10] I. Hofacker, K. Schulten, Oxygen and proton pathways in cytochrome *c* oxidase, *Proteins Struct. Funct. Genet.* 30 (1998) 100–107.
- [11] S. Riistama, G. Hummer, A. Puustinen, R.B. Dyer, W.H. Woodruff, M. Wikström, Bound water in the proton translocation mechanism of the haem-copper oxidases, *FEBS Lett.* 414 (1997) 275–280.
- [12] M.L. Verkhovskaya, J.A. García-Horsman, A. Puustinen, J.-L. Rigaud, J.E. Morgan, M.I. Verkhovsky, M. Wikström, Glutamic acid 286 in subunit I of cytochrome *bo*₃ is involved in proton translocation, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 10128–10131.
- [13] P. Brzezinski, P. Ädelroth, Pathways of proton transfer in cytochrome *c* oxidase, *J. Bioenerg. Biomembranes* 30 (1998) 99–107.
- [14] M. Wikström, A. Jasaitis, C. Backgren, A. Puustinen, M.I. Verkhovsky, The role of the D- and K-pathways of proton transfer in the function of the haem-copper oxidases, *Biochim. Biophys. Acta* 1459 (2000) 514–520.
- [15] A. Puustinen, M. Wikström, Proton exit from the heme-copper oxidase of *Escherichia coli*, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 35–37.
- [16] G. Varo, J.K. Lanyi, Effects of hydrostatic pressure on the kinetics reveal a volume increase during the bacteriorhodopsin photocycle, *Biochemistry* 34 (1995) 12161–12169.
- [17] J.A. Kornblatt, The water channel of cytochrome *c* oxidase: inferences from inhibitor studies, *Biophys. J.* 75 (1998) 3127–3134.
- [18] X. Zheng, D.M. Medvedev, J. Swanson, A.A. Stuchebrukhov, Computer simulation of water in cytochrome *C* oxidase, *Biochim. Biophys. Acta* 1557 (2003) 99–107.
- [19] G. Hummer, J.C. Rasaiah, J.P. Noworyta, Water conduction through the hydrophobic channel of a carbon nanotube, *Nature* 414 (2001) 188–190.
- [20] N. Agmon, The Grotthuss mechanism, *Chem. Phys. Lett.* 244 (1995) 456–462.
- [21] C. Dellago, M.M. Naor, G. Hummer, Proton transport through water-filled carbon nanotubes, *Phys. Rev. Lett.* 90 (2003) 105902.
- [22] A.S. Pawate, J. Morgan, A. Namslauer, D. Mills, P. Brzezinski, S. Ferguson-Miller, R.B. Gennis, A Mutation in subunit I of cytochrome oxidase from *Rhodobacter sphaeroides* results in an increase in steady-state activity but completely eliminates proton pumping, *Biochemistry* 41 (2002) 13417–13423.
- [23] M.I. Verkhovsky, J.E. Morgan, M. Wikström, Control of electron delivery to the oxygen reduction site of cytochrome *c* oxidase. A role for protons, *Biochemistry* 34 (1995) 7483–7491.
- [24] P.R. Rich, Towards an understanding of the chemistry of oxygen reduction and proton translocation in the iron-copper respiratory oxidases, *Aust. J. Plant Physiol.* 22 (1996) 479–486.
- [25] A. Kannt, C.R.D. Lancaster, H. Michel, The coupling of electron transfer and proton translocation: electrostatic calculations on *Paracoccus denitrificans* cytochrome *c* oxidase, *Biophys. J.* 74 (1998) 708–721.
- [26] Y.C. Fann, I. Ahmed, N.J. Blackburn, J.S. Boswell, M.L. Verkhovskaya, B.M. Hoffman, M. Wikström, Structure of Cu_B in the binuclear heme-copper center of the cytochrome *aa*₃-type quinol oxidase from *Bacillus subtilis*: an ENDOR and EXAFS study, *Biochemistry* 34 (1995) 10245–10255.
- [27] W.L. Jorgensen, J. Chandrasekhar, J.D. Madura, R.W. Impey, M.L. Klein, Comparison of simple potential functions for simulating liquid water, *J. Chem. Phys.* 79 (1983) 926–935.
- [28] R. Pomès, G. Hummer, M. Wikström, Structure and dynamics of a proton shuttle in cytochrome *c* oxidase, *Biochim. Biophys. Acta* 1365 (1998) 255–260.
- [29] W.D. Cornell, P. Cieplak, C.I. Bayley, I.R. Gould, K.M. Merz Jr., D.M. Ferguson, D.C. Spellmeyer, T. Fox, J.W. Caldwell, P.A. Kollman, A second generation force field for the simulation of proteins, nucleic acids, and organic molecules, *J. Am. Chem. Soc.* 117 (1995) 5179–5197.
- [30] J.E. Morgan, M.I. Verkhovsky, M. Wikström, The histidine cycle: a new model for proton translocation in the respiratory heme-copper oxidases, *J. Bioenerg. Biomembranes* 26 (1994) 599–608.
- [31] H. Michel, Cytochrome *c* oxidase: catalytic cycle and mechanisms of proton pumping—A discussion, *Biochemistry* 38 (1999) 15129–15140.
- [32] W. Humphrey, A. Dalke, K. Schulten, VMD—Visual molecular dynamics, *J. Mol. Graph.* 14 (1996) 33–38.