

Mitochondrial biogenesis in fast skeletal muscle of CK deficient mice

A. Vaarmann^{a,b,c}, D. Fortin^{a,b}, V. Veksler^{a,b}, I. Momken^{a,b},
R. Ventura-Clapier^{a,b}, A. Garnier^{a,b,*}

^a INSERM, U769, Châtenay-Malabry, F-92296, France

^b Univ Paris-Sud, UMR-S 769, IFR141, Châtenay-Malabry, F-92296, France

^c Department of Pharmacology, Center of Molecular and Clinical Medicine, University of Tartu, Estonia

Received 8 October 2007; received in revised form 5 November 2007; accepted 5 November 2007

Available online 12 November 2007

Abstract

Creatine kinase (CK) is a phosphotransfer kinase that catalyzes the reversible transfer of a phosphate moiety between ADP and creatine and that is highly expressed in skeletal muscle. In fast glycolytic skeletal muscle, deletion of the cytosolic M isoform of CK in mice (M-CK^{-/-}) leads to a massive increase in the oxidative capacity and of mitochondrial volume. This study was aimed at investigating the transcriptional pathways leading to mitochondrial biogenesis in response to CK deficiency. Wild type and M-CK^{-/-} mice of eleven months of age were used for this study. Gastrocnemius muscles of M-CK^{-/-} mice exhibited a dramatic increase in citrate synthase (+120%) and cytochrome oxidase (COX, +250%) activity, and in mitochondrial DNA (+60%), showing a clear activation of mitochondrial biogenesis. Similarly, mRNA expression of the COXI (mitochondria-encoded) and COXIV (nuclear-encoded) subunits were increased by +103 and +94% respectively. This was accompanied by an increase in the expression of the nuclear respiratory factor (NRF2α) and the mitochondrial transcription factor (mtTFA). Expression of the co-activator PGC-1α, a master gene in mitochondrial biogenesis was not significantly increased while that of PGC-1β and PRC, two members of the same family, was moderately increased (+45% and +55% respectively). While the expression of the modulatory calcineurin-interacting protein 1 (MCIP1) was dramatically decreased (−68%) suggesting inactivation of the calcineurin pathway, the metabolic sensor AMPK was activated (+86%) in M-CK^{-/-} mice. These results evidence that mitochondrial biogenesis in response to a metabolic challenge exhibits a unique pattern of regulation, involving activation of the AMPK pathway.

© 2007 Elsevier B.V. All rights reserved.

Keywords: Skeletal muscle; CK-KO mice; Energy metabolism; AMPK; Transcription

1. Introduction

The family of creatine kinase isoenzymes catalyzes the reversible transfer of a phosphate moiety between creatine and ATP. It is highly expressed in striated muscles and is a key player in intracellular energy storage and transport. The major isoenzymes of CK in muscle are the cytosolic isoform (MM-CK) and the mitochondrial isoform (mi-CK). Fast gastrocnemius muscle expresses almost exclusively MM-CK [1] that is free in the cytosol or structurally associated with myofibrils and membranes of the sarcoplasmic reticulum (SR), and function-

ally coupled to ATPases for optimal functioning of the contractile machinery and SR calcium uptake [2].

Fast skeletal muscle fibers have very low mitochondrial content and mainly rely on quickly mobilisable energy sources (mainly phosphocreatine (PCr) and glycogen) to develop strong and fast contractions but this is possible only for short periods of time because of limited reserves. These muscles are thus quickly fatigable and should recover their energy reserves through anaerobic glycolysis and less importantly mitochondrial oxidations.

Engineered mice with invalidated expression of M-CK (M-CK^{-/-}) were developed by the group of Bé Wieringa [3] to understand the effects of altered energy metabolism. Functional tests revealed that fast skeletal muscle of CK^{-/-} mice has abnormal calcium transient, lacks burst activity at the onset of stimulation but exhibits paradoxical decreased fatigability [4,5]. This is accompanied by a marked increase in the relative

* Corresponding author. INSERM, U769, Université Paris-Sud, 5 rue Jean-Baptiste Clément, F-92296 Châtenay-Malabry, France. Tel.: +33 1 46 83 5249; fax: +33 1 46 83 54 75.

E-mail address: anne.garnier@u-psud.fr (A. Garnier).

mitochondrial volume, the mitochondrial enzyme content, and the muscle oxidative capacity together with a relocation of mitochondria towards myofibrils [3,6–8]. Proteomic and mRNA analysis [9,10] confirmed the metabolic remodeling of the CK^{−/−} gastrocnemius muscle towards a more oxidative phenotype. This metabolic remodeling is thought to compensate for the lack of creatine kinase by switching energy metabolism of gastrocnemius muscle from the use of PCr stores and anaerobic glycolysis to oxidative metabolism. However, the signaling and molecular events governing this metabolic shift are currently unknown.

Mitochondrial biogenesis depends on the coordinated expression of the nuclear and mitochondrial genomes. Mitochondrial DNA (mtDNA), encodes 13 subunits of the oxidative phosphorylation system (OXPHOS). The remaining OXPHOS subunits as well as other mitochondrial proteins are encoded by the nuclear DNA. An inducible transcriptional co-activator termed peroxisome proliferator-activated receptor gamma co-activator 1 α (PGC-1 α) has emerged as a critical factor coordinating the activation of metabolic genes required for substrate utilization and mitochondrial biogenesis [11–13]. Effects of PGC-1 α on mitochondrial biogenesis could be explained via its interaction with several DNA-binding transcription factors, such as the nuclear respiratory factors (NRFs). These factors in turn up-regulate the expression of nuclear genes encoding respiratory chain components and other mitochondrial proteins, as well as of the mitochondrial transcription factor A (mtTFA), a factor required for mtDNA replication and transcription [14]. In rodent skeletal muscles, regularly performed exercise induces an increase in PGC-1 α , coincidentally with an increase in NRFs and mtTFA mRNA and/or protein expression (for recent reviews see [15–17]). In humans, PGC-1 α and its transcription cascade correlate with exercise capacity and vastus lateralis muscle oxidative capacity [18].

In response to changes in environment, muscle activity or energy state, mitochondrial biogenesis is controlled by upstream signaling events. Calcineurin, a calcium sensitive phosphatase, involved in the transcriptional response of skeletal muscle to endurance training through the dephosphorylation and the nuclear import of the nuclear transcription factor of activated T cell (NFAT) family [19], was proposed to control the expression of PGC-1 α [20]. In human skeletal muscle, the transcriptional activity of calcineurin correlates with exercise capacity and muscle oxidative capacity [18]. However, calcineurin inhibition fails to block the exercise-induced PGC-1 α expression and activation of mitochondrial biogenesis [21]. Moreover, calcineurin inhibition increases rather than decreases oxidative capacity in soleus muscle consistent with the partial transition from type I to the more oxidative type IIa fiber in this muscle [22]. Other pathways have been implicated in the control of PGC-1 α expression and activity. Among those, the p38 mitogen-activated protein kinase (p38 MAPK) and the calcium–calmodulin dependent protein kinases (CaMKs), also activated during exercise, could contribute to increase muscle oxidative capacity [23,24].

On the other hand, skeletal muscle mitochondrial biogenesis can be induced by energetic deficiency. Energetic deficiency can

activate the AMP-activated protein kinase (AMPK). Under conditions of high (exercise) or disturbed (energetic deficiency) energy turnover, AMP concentration increases and induces the phosphorylation of the Thr172 of the α subunit thus activating the catalytic activity of AMPK. Chronic depletion of creatine by feeding an analog (β -guanidino propionic acid (β -GPA), activates AMPK, increases mitochondrial content and up-regulates expression of genes involved in mitochondrial biogenesis among which NRF1 and PGC-1 α [25,26]. Thus by sensing the metabolic state of the muscle, AMPK appears to be an important regulator of mitochondrial biogenesis. However, deletion of one or the other catalytic subunit of AMPK did not impair the exercise-induced activation of mitochondrial gene expression [27]. Moreover at present, the way by which AMPK activation increases PGC-1 α gene transcription in skeletal muscle is currently unknown.

The aim of the present work has been to determine whether and how mitochondrial biogenesis occurs in response to cytosolic creatine kinase deficiency. The superficial part of the gastrocnemius muscle of M-CK deficient mice was used for this study because this muscle mainly rely on creatine kinase for contractile activity and because it is known to undergo thorough remodeling towards increased oxidative capacity. The results show that mitochondrial biogenesis seems to be triggered by energy depletion-induced AMPK activation and increased PGC-1 α transcriptional activity, rather than by calcium-dependent increase in PGC-1 α expression.

2. Materials and methods

2.1. Animals

Procedures involved in the generation and genotyping of M-CK^{−/−} (kind gift from Drs B. Wieringa and F. Oerlemans University of Nijmegen, Netherlands) have been described in detail elsewhere [3]. Eleven-month old C57BL6 wild type (WT, $n=6$), and M-CK^{−/−} ($n=6$) mice were used for this study. Animals were anaesthetized with an intraperitoneal pentobarbital injection (0.15 mg/g BW^{−1}), sacrificed and the superficial parts of gastrocnemius (fast-twitch, glycolytic) muscles were isolated, rapidly frozen and kept at -80°C . The investigation conforms to Inserm Institution guidelines defined by the European Community guiding principles in the care and use of animals and the French decree no. 87/848 of October 19, 1987.

2.2. Enzyme analysis

Frozen tissue samples were weighed, homogenized in ice-cold buffer (50 mg wet weight per ml) containing: HEPES 5 mM (pH 8.7), ethyleneglycol-bis (β -aminoethyl ether) N , N , N' , N' -tetraacetic acid (EGTA) 1 mM, dithiothreitol 1 mM, and Triton X-100 (0.1%) and incubated for 60 min at 4°C for complete enzyme extraction. The cytochrome c oxidase (COX), and citrate synthase (CS) were assayed by standard spectrophotometric methods at 30°C and pH 7.5 [28].

2.3. Real-time quantitative RT-PCR analysis

Total muscle RNA was extracted using standard procedures. Oligo-dT first strand cDNA was synthesized from 5 μg total RNA using superscript II reverse transcriptase (Invitrogen). Real-time RT-PCR was performed using the SYBR[®]Green method on a LightCycler rapid thermal cycler (Roche Diagnostics) as previously described [28]. Primers were designed in a different exon of the target gene sequence, eliminating the possibility of amplifying genomic DNA (Table 1). A Basic Local Alignment Search Tool (BLAST) search

Table 1
Primers used for real-time PCR amplification

Gene	GenBank AN	Forward primer Reverse primer (5'–3')	PCR product size (pb)	Annealing temperature (°C)
PGC-1 α	NM_008904	CAC CAA ACC CAC AGA GAA CAG GCA GTT CCA GAG AGT TCC ACA	210	58
PGC-1 β	NM_133249	TGG AAA GCC CCT GTG AGA GT TTG TAT GGA GGT GTG GTG GG	202	60
PRC	XM_358330	AGG AAA CTC AGG CAG CAT TG GGC GGT GGA TTT AGG AGA TT	178	60
NRF2 α	NM_008065	AGGTGACGAGATGGGCTGC CGTTGTCCCCATTTTGCG	604	65
mtTFA	NM_009360	GCT GAT GGG TAT GGA GAA G GAG CCG AAT CAT CCT TTG C	161	56
COX I	NC_006914	CAC TAA TAA TCG GAG CCC CA TTC ATC CTG TTC CTG CTC CT	129	60
COX IV	NM_053091	TGG GAG TGT TGT GAA GAG TGA GCA GTG AAG CCG ATG AAG AAC	273	58
Drp1	NM_152816	CTG ACG CTT GTG GAT TTA CC CCC TTC CCA TCA ATA CAT CC	277	58
Mfn2	NM_133201	ACG AGC AAT GGG AAG AGC AC TCC ATC AGC ACG AGG TCA TC	284	60
MCIP1	NM_019466	CAG CGA AAG TGA GAC CAG GG ACG GGG GTG GCA TCT TCT AC	309	60
MEF2C	NM_025282	CAG GGA ACG GGT ATG GCA ATC CAA TGA CTG AGC CGA CTG GGA	239	60
PPAR δ	NM_011145	GCC TCC ATC GTC AAC AAA GA TCT ACC TGG GGC ACA TTC AT	230	60
GCB	NM_008094	CCC ATT TCA CTC TTT GCC AG AGG TTC ATT CTC CGC TGT CA	198	60

PGC-1 α and PGC-1 β : peroxisome proliferator-activated receptor gamma co-activator-1 α and β ; PRC: PGC-1 α related co-activator; NRF2 α : nuclear respiratory factor-2 α ; mtTFA, mitochondrial transcription factor A; COX I and IV: cytochrome oxidase subunits I and IV; Drp1: dynamin-related protein 1; Mfn2: mitofusin 2; MCIP1: modulatory calcineurin-interacting protein 1; MEF2C: myocyte-specific enhancer factor 2; PPAR δ : peroxisome proliferator-activated receptor δ ; GCB: glucocerebrosidase.

performed for each set of primers revealed that sequence homology was obtained only for the target gene. Glucocerebrosidase (GCB) was chosen as housekeeping gene for normalization as its expression did not differ between the two groups. Values for each gene were normalized to GCB mRNA content in order to compensate for variation in input RNA amounts and efficiency of reverse transcription, then they were multiplied by total RNA per amount of tissue (g wet weight⁻¹) to compare expression level in different conditions [28].

2.4. Southern blot analysis of total DNA

Total cellular DNA was extracted by standard methods including successive steps of proteinase K digestion, organic extraction and ethanol precipitation. To measure mtDNA levels, a Southern blot analysis was performed using concomitant hybridization with a cDNA probe for mtDNA and a cDNA probe for nDNA as control for the amount of nuclear DNA as previously described [28]. Signals were detected by chemiluminescent reagents (CDP-StarTM, Amersham) and quantified using an image analyzer (Bio-Rad) to determine mtDNA to nDNA ratio.

2.5. Western blot analysis

Specific antibodies were used to measure the protein content of phosphorylated and non-phosphorylated AMPK (Upstate Biotechnology Inc., Lake Placid, New York, USA) and phospho- and total p38 MAPK (Cell Signaling) in control and M-CK^{-/-} gastrocnemius muscles. Blots were revealed with enhanced chemiluminescent substrate (ECL, Amersham, France). Light emission was detected by autoradiography using an image-analysis system (Bio-Rad Geldoc 1000). Quantification was performed using Quantity One software (Biorad) and expressed as a ratio of the signal obtained with the phosphorylated protein of interest relative to the non-phosphorylated protein.

2.6. Statistical analysis

All data are expressed as means \pm sem and were compared using a Student's *t*-test. Values of $p \leq 0.05$ were considered significant.

3. Results

3.1. Mitochondrial activity and protein expression

Mitochondrial content of gastrocnemius muscles was estimated by measuring the activity of two markers of mitochondrial activity, citrate synthase (CS) an enzyme of the Krebs cycle and cytochrome oxidase (COX), the complex IV of the respiratory chain. M-CK deletion resulted in significant 3.5-fold and 2.2-fold increases in COX and CS activities respectively (Fig. 1A). The COX/CS ratio was significantly increased by 1.55-fold, showing an excess increase in COX specific activity. Moreover, a 1.6-fold increase in mitochondrial DNA over nuclear DNA ratio (from 0.78 ± 0.06 to 1.26 ± 0.15 , $p < 0.05$) confirmed the increase in mitochondrial mass. The total mRNA content was significantly increased from 0.469 ± 0.028 to 0.642 ± 0.043 mg \cdot gww⁻¹ ($p < 0.05$) in M-CK^{-/-} gastrocnemius muscle. Thus, this increase in mRNA was taken into account to calculate the concentration of mRNAs in muscles.

In order to confirm that both the mitochondrial and the nuclear genomes were activated, we measured expression of two subunits of COX, one encoded by the mitochondrial genome




Fig. 1. Mitochondrial enzyme activity and expression in gastrocnemius muscle of WT ($n=6$) and M-CK^{-/-} ($n=6$) mice. A. Increase in cytochrome oxidase and citrate synthase activity measured by spectrophotometry. B. Increased mRNA expression of subunit I (COXI, mitochondria-encoded) and subunit IV (COXIV, nuclear-encoded) of the cytochrome oxidase and of dynamin-related protein 1 (Drp1) but not mitofusin 2 (Mfn2) two proteins that are involved in mitochondrial dynamics. Values are means \pm sem. ** $p < 0.01$ versus WT.

(COXI) and one encoded by the nuclear genome (COXIV). When expressed as per mg of tissue, the amounts of mRNA encoding COXI and COXIV were both significantly increased by 203% and 194% respectively (Fig. 1B), showing that both genomes were coordinately activated in M-CK^{-/-} gastrocnemius muscle. Mitochondrial biogenesis also involves mitochondrial dynamics. Shape and size of mitochondria are regulated by a complex process of fusion and fission. Two proteins, the dynamin-related protein 1 (Drp1) involved in fission and the mitofusin 2 (Mfn2) involved in fusion are implicated in this process. In M-CK^{-/-} mice, Drp1 expression was increased 1.9-fold while Mfn2 exhibited a non-significant increase (Fig. 1B).

3.2. Mitochondrial transcription cascade

We next examined the transcription cascade involved in mitochondrial biogenesis (Fig. 2A). It is well accepted that the mitochondrial transcription factor mtTFA is involved in both transcription and replication of mitochondrial DNA. In M-CK^{-/-} mice mtTFA expression was significantly increased by 50%, in accordance with increased COXI mRNA expression and mtDNA content. In rodents, mtTFA promoter contains NRF2 α but not

NRF1 recognition sites [29]. Accordingly, NRF2 α expression was also increased by 43% in M-CK^{-/-} gastrocnemius muscle.

Upstream of NRFs, PGC-1 α transcriptional co-activator family plays a major role in mitochondrial biogenesis induced by diverse physiological stimuli [13,30]. We investigated whether PGC-1 α , PGC-1 β and PGC-1 α related co-activator (PRC) were increased in gastrocnemius muscle (Fig. 2B). Surprisingly, PGC-1 α expression was not increased while PGC-1 β and PRC were slightly increased with PRC only reaching significance. As PPAR δ has also been involved in skeletal muscle biogenesis [31] we evaluated its mRNA expression. No significant change was observed in M-CK^{-/-} muscles.

3.3. Mitochondrial biogenesis signaling

We then attempted to elucidate the signaling pathways involved in mitochondrial biogenesis in gastrocnemius muscle of M-CK^{-/-} mice. In order to assess calcineurin transcriptional activity, we measured the level of transcription of the myocyte-enriched calcineurin-interacting protein 1 (MCIP1), which contains 15 repeats of the NFAT binding site and thus has been shown to be the most sensitive indicator of calcineurin transcriptional




Fig. 2. Mitochondrial transcription cascade in gastrocnemius muscle of WT ($n=6$) and M-CK^{-/-} ($n=6$) mice. A. Real-time RT-PCR analysis of the nuclear respiratory factor 2 (NRF2) and the mitochondrial transcription factor A (mtTFA). B. mRNA expression of the peroxisome proliferator-activated receptor (PPAR) gamma co-activator 1 α (PGC-1 α) or 1 β (PGC-1 β) isoforms, the PGC-1-related co-activator (PRC), and the delta isoform of PPAR. Values are means \pm sem and are shown as arbitrary units normalized to the glucocorticoid and to the amount of mRNA. * $p < 0.05$, ** $p < 0.01$ versus WT.




Fig. 3. Calcium signaling pathways of mitochondrial biogenesis in gastrocnemius muscle of WT ($n=6$) and M-CK^{-/-} ($n=6$) mice. A. Calcineurin activation was assessed by the level of expression of the modulatory calcineurin-interacting protein 1 (MCIP1) and the myocyte-specific enhancer factor 2 (MEF2C). B. Activation of p38 MAPK has been assessed by the ratio of signals obtained by immunoblots of the phosphorylated and total forms; upper part representative immunoblots; lower part mean values. Values are means \pm sem. * $p < 0.05$ versus WT.

activity [18]. MCIP1 expression exhibited a 3-fold decrease suggesting decreased rather than increased transcriptional activity of calcineurin in M-CK^{-/-} mice (Fig. 3A). Expression level of the MEF2C transcription factor which acts in synergy with NFAT to regulate muscle fiber type [32] was slightly but not significantly increased in M-CK^{-/-} mice. On the other hand, we examined whether the p38 MAPK pathway was activated in gastrocnemius muscle of M-CK^{-/-} mice by western blotting with antibodies specific for the phosphorylated or the total p38 MAPK (Fig. 3B). No increase in p38 MAPK phosphorylation was observed in gastrocnemius muscle of M-CK^{-/-} mice suggesting that this pathway does not participate in mitochondrial biogenesis.

Finally AMPK was also shown to induce mitochondrial biogenesis in muscle, in response to metabolic stress and exercise. AMPK phosphorylation was significantly increased in M-CK^{-/-} gastrocnemius muscle (Fig. 4).

4. Discussion

This study was aimed at investigating the transcriptional cascade involved in mitochondrial biogenesis induced by cre-

atine kinase depletion in fast skeletal muscle of mice (Fig. 5). The results show that the 2-fold increase in mitochondrial mass was accounted for by an increase in both mitochondria and nucleus-encoded mitochondrial proteins, accompanied by a 60% increase in mitochondrial DNA. This could be explained by an increase in mtTFA and NRF2 transcription factors. However, while neither expression of PGC-1 α nor PPAR δ , factors implicated in muscle mitochondrial biogenesis was increased, a slight increase in PRC expression was observed. Moreover, AMPK but not calcineurin or p38 MAPK was activated, evidencing a unique pattern of mitochondrial biogenesis in response to M-CK depletion.

4.1. Mitochondrial activity and content

Creatine kinase deficiency is a unique model of metabolic deficiency. It leads to a massive metabolic remodeling especially in the fast gastrocnemius muscle. Indeed, a doubling of oxidative capacity and mitochondrial volume density was described in gastrocnemius of M-CK or M-CK and mi-CK deficient mice [6,8,33]. As mi-CK is almost absent from fast fibers [1], this mitochondrial remodeling results from M-CK deficiency. At the same time, mitochondrial DNA is increased by only 60%, showing the complexity of mitochondrial biogenesis. On the other hand, as a more than 2-fold increase in CS activity was measured, this suggests that not only mitochondrial mass but also mitochondrial specific activity was increased, in line with increased state 3 respiration rates in mitochondria isolated from




Fig. 4. AMPK activation in gastrocnemius muscle of WT ($n=6$) and M-CK^{-/-} ($n=6$) mice. Activation of AMP-activated protein kinase (AMPK) has been assessed by the ratio of signals obtained by immunoblots of the phosphorylated and total forms; upper part representative immunoblots; lower part mean values. Values are means \pm sem. * $p < 0.05$ versus WT.




Fig. 5. A model for mitochondrial biogenesis in gastrocnemius muscle of CK deficient mice. Energy deficiency induced by M-CK deletion lead to increased mitochondrial biogenesis. This effect was not accompanied by increased activity of calcineurin or phosphorylation of p38 MAPK suggesting that neither calcium signaling through calcineurin or CaMK, nor p38 signaling was involved. AMPK appeared phosphorylated suggesting a direct link between energy depletion, change in AMP/ATP ratio and mitochondrial biogenesis. A slight increase in PGC-1 β and PRC but no change in PGC-1 α or PPAR δ expression, the two main regulators of mitochondrial biogenesis in muscle, was observed. However, PGC-1 α may be involved in mitochondrial biogenesis by direct phosphorylation and activation by AMPK.

CK $^{-/-}$ gastrocnemius muscle [34]. Mitochondrial function depends on the proper assembly of complexes of the electron transport chain (ETC) that are embedded within the inner mitochondrial membrane. The most widely studied ETC complex is COX which converts oxygen to water and provides part of protons required for ATP synthesis. COX assembly depends on COX subunits that are both nuclear- and mitochondria-encoded. Both the mitochondria-encoded COX I and the nuclear-encoded COX IV subunits were up-regulated by a factor of 2 in M-CK deficient mice. This led to a 3.5 increase in COX activity. Moreover this increase in COX activity exceeded that of CS as a significant 1.5 increase in COX/CS activity was observed. Indeed the COX/CS activity ratio can vary in a muscle type specific manner being much higher in oxidative (2.3 in heart, 2.6 in soleus) [35] than in fast glycolytic muscle (0.3 in gastrocnemius), showing the complex regulation of mitochondrial protein expression in muscles. Interestingly, in M-CK $^{-/-}$ mice the COX/CS ratio was significantly increased to 0.5 toward that of an oxidative muscle, but still much lower than WT soleus muscle.

The structural and functional adaptations of the mitochondrial network in challenged skeletal muscles result not only from changes in the mitochondrial protein expression and proper assembly but also involve mitochondrial dynamics. Training induced mitochondrial biogenesis in humans involve the coordinate increase in both Mfn2, a protein involved in fusion, and Drp1, a protein involved in mitochondrial fission, in vastus lateralis muscle [18]. Here we see a significant 50% increase in expression of Drp1, while the expression of Mfn2 remained unchanged. This suggests that in this model, the increase in

mitochondrial number results from growing and fission of preexisting mitochondria.

4.2. Mitochondrial biogenesis transcription cascade

It is now well documented that mitochondrial biogenesis involves the integration of multiple transcriptional regulatory pathways controlling both the activation of nuclear and mitochondrial genomes. The mitochondrial genome however should not only be expressed but also replicated. This can be achieved by the nuclear-encoded transcription factor mtTFA that binds to the mitochondrial DNA and induces both replication and transcription [13]. Accordingly mtTFA expression was significantly enhanced in M-CK $^{-/-}$ gastrocnemius muscle. Upstream of mtTFA, the NRFs are thought to increase the expression of multiple mitochondrial proteins by binding to specific recognition sites. The expression of NRF2 α , the main isoform involved in rodent mitochondrial biogenesis [36] was also increased in M-CK $^{-/-}$ mice, but with lower significance ($p < 0.057$). Upstream of NRFs, PGC-1 α is thought to orchestrate mitochondrial biogenesis in most tissues [37]. Moreover, PGC-1 α expression correlates with oxidative capacity of cardiac and skeletal muscles [28]. It was thus surprising that its expression was not increased in gastrocnemius muscle of M-CK $^{-/-}$ mice, despite massive increase in mitochondrial content. Loss-of-function experiments have established that PGC-1 α is not an absolute requirement for mitochondrial biogenesis. KO mice have preserved volume density of mitochondria in both heart and skeletal muscles [38], but reduced mRNA for mitochondrial proteins [38]. In a similar study however, Leone et al described slightly different muscle

phenotype with a 30% decrease in mitochondrial volume and mRNA expression of mitochondrial proteins in soleus muscle and decrease in respiration rate [39]. Although results differ between the two studies, they show that the absence of PGC-1 α can be partly overcome by other mitochondrial biogenesis pathways. PGC-1 α is a member of an important family of transcriptional co-activators that includes PGC-1 β and PGC-related co-activator (PRC) [40]. Those transcription factors share a high degree of homology and redundancy with PGC-1 α with respect to the manner they activate mitochondrial biogenesis although they seem to respond differently to activators of mitochondrial biogenesis. Thus the slight increase in both PGC-1 β ($p < 0.069$) and PRC ($p < 0.05$) could participate in increased mitochondrial biogenesis in M-CK $^{-/-}$ gastrocnemius. Finally, PPAR δ whose invalidation was also shown to alter mitochondrial content [31,41] was unchanged in M-CK mice. However, the slight increase in transcription of members of PPAR and PGC transcription factor families may not be sufficient to explain the robust increase in gastrocnemius mitochondrial content (Fig. 5).

4.3. Signaling pathways

It is likely that Ca²⁺ and energy depletion signals act in concert to promote metabolic plasticity in CK $^{-/-}$ fast skeletal muscle. Indeed, altered calcium movements have been described in skeletal muscle of M-CK and CK $^{-/-}$ gastrocnemius muscle. CK $^{-/-}$ fast muscle fibers have increased resistance to fatigue and prolonged calcium transient [5]. It was thus proposed that calcium homeostasis impairment due to CK deficiency may have played a role in directing the increase in mitochondrial capacity [34]. The calcium-regulated serine/threonine protein phosphatase calcineurin has been implicated in the transduction of motor neuron signals to alter gene expression programs in skeletal muscle. However, assessment of the transcriptional activity of calcineurin by MCIP1 expression showed a decreased rather than increased calcineurin activity. Moreover, MEF2C was not up-regulated to significant level in these mice. This suggests that calcium is not primarily involved in gastrocnemius increase in mitochondrial biogenesis. The absence of increased calcineurin activation could indeed be linked to the low level of physical activity and thus of contraction-induced calcium increase in M-CK $^{-/-}$ mice [35].

Evidences are accumulating that the transcriptional activity of PGC-1 α can be regulated by posttranslational modifications including phosphorylations. Transcriptional activity of PGC-1 α can be enhanced by phosphorylation and disruption of a PGC-1 α /p160MBP interaction through activation of the p38 MAPK pathway [42,43]. However, no activation of p38 MAPK was observed in this study suggesting that p38 MAPK is not involved in mitochondrial biogenesis in this model.

Finally, AMPK can activate mitochondrial biogenesis during long term exercise or energetic deficiency in muscle. It has been shown that AMPK can activate the transcription of PGC-1 α indicating that its mitochondriogenic action can partly originate from enhancement of PGC-1 α expression [26,44]. Moreover, skeletal muscle of transgenic mice expressing a dominant-negative mutant form of AMPK have blunted response to energy

deprivation by β -GPA feeding, showing that AMPK is necessary for mitochondrial biogenesis in response to metabolic stress [26]. Low frequency stimulation that mimics exercise training induces AMPK-PGC-1 α signaling pathway [45]. AMPK is phosphorylated and activated in response to the increase in the AMP/ATP ratio. Indeed, both a decrease in ATP and an increase in AMP have been shown in hindlimb muscle of M-CK-KO mice [3]. The present results show that AMPK phosphorylation is robustly increased in the gastrocnemius muscle of M-CK $^{-/-}$ mice. However, AMPK activation did not lead to increased expression of PGC-1 α . It was shown very recently that AMPK can directly interact with and phosphorylate PGC-1 α showing a posttranslational regulation of mitochondrial biogenesis by AMPK; moreover, AMPK phosphorylation of PGC-1 α initiates many of the important gene regulatory functions of AMPK in skeletal muscle [46]. This strongly suggests that M-CK deficiency creates an energy deprivation state in gastrocnemius muscle of M-CK $^{-/-}$ mice that induces basal activation of AMPK, AMPK-dependent phosphorylation of PGC-1 α and increased transcriptional activity of PGC-1 α . Interestingly, activation of AMPK induces decreased skeletal muscle fiber size and increased mitochondrial biogenesis [47] in S6K deficient mice. Basal activation of AMPK in M-CK $^{-/-}$ mice could be a major factor involved in muscle atrophy and decreased exercise capacity of CK deficient mice [35].

Thus, following M-CK deletion, fast skeletal muscle undergoes a metabolic remodeling and fiber atrophy. These events seem to be triggered by energy depletion-induced AMPK activation, rather than by calcium-dependent processes (Fig. 5).

Acknowledgments

We thank Dr. R. Fischmeister for the continuous support. RV-C is supported by “Centre National de la Recherche Scientifique”. We thank Dr. B. Wieringa for the kind support with the mice. We thank Claudine Delomé and the “Plateforme transcriptome” for their help in RT-PCR experiments and Valérie Domergue and the animal care center for taking care of the animals.

References

- [1] W.N. Qin, Z. Khuchua, J. Boero, R.M. Payne, A.W. Strauss, Oxidative myocytes of heart and skeletal muscle express abundant sarcomeric mitochondrial creatine kinase, *Histochem. J.* 31 (1999) 357–365.
- [2] R. Ventura-Clapier, A. Kuznetsov, V. Veksler, E. Boehm, K. Anfous, Functional coupling of creatine kinases in muscles: species and tissue specificity, *Mol. Cell. Biochem.* 184 (1998) 231–247.
- [3] J. van Deursen, A. Heerschap, F. Oerlemans, W. Ruitenbeek, P. Jap, H. ter Laak, B. Wieringa, Skeletal muscles of mice deficient in muscle creatine kinase lack burst activity, *Cell* 74 (1993) 621–631.
- [4] M. Gorselink, M.R. Drost, W.A. Coumans, G.P. van Kranenburg, R.P. Hesselink, G.J. van der Vusse, Impaired muscular contractile performance and adenine nucleotide handling in creatine kinase-deficient mice, *Am. J. Physiol., Endocrinol Metabol.* 281 (2001) E619–E625.
- [5] A.J. Dahlstedt, H. Westerblad, Inhibition of creatine kinase reduces the rate of fatigue-induced decrease in tetanic Ca(2+), *J. Physiol.* 533 (2001) 639–649.
- [6] V.I. Veksler, A.V. Kuznetsov, K. Anfous, P. Mateo, J. van Deursen, B. Wieringa, R. Ventura-Clapier, Muscle creatine kinase-deficient mice.2.

- Cardiac and skeletal muscles exhibit tissue-specific adaptation of the mitochondrial function. *J. Biol. Chem.* 270 (1995) 19921–19929.
- [7] A. Kaasik, V. Veksler, E. Boehm, M. Novotova, R. Ventura-Clapier, From energy store to energy channeling: a study in creatine kinase deficient fast skeletal muscle, *FASEB J.* 17 (2003) 708–710.
 - [8] M. Novotova, M. Pavlovicova, V. Veksler, R. Ventura-Clapier, I. Zahradnik, Ultrastructural remodeling of fast skeletal muscle fibers induced by invalidation of creatine kinase, *Am. J. Physiol., Cell Physiol.* 291 (2006) C1279–C1285.
 - [9] A.J. de Groof, B. Smeets, M.J. Groot Koerkamp, A.N. Mul, E.E. Janssen, H.F. Tabak, B. Wieringa, Changes in mRNA expression profile underlie phenotypic adaptations in creatine kinase-deficient muscles, *FEBS Lett.* 506 (2001) 73–78.
 - [10] A.J. de Groof, F.T. Oerlemans, C.R. Jost, B. Wieringa, Changes in glycolytic network and mitochondrial design in creatine kinase-deficient muscles, *Muscle Nerve* 24 (2001) 1188–1196.
 - [11] Z.D. Wu, P. Puigserver, U. Andersson, C.Y. Zhang, G. Adelmant, V. Mootha, A. Troy, S. Cinti, B. Lowell, R.C. Scarpulla, B.M. Spiegelman, Mechanisms controlling mitochondrial biogenesis and respiration through the thermogenic coactivator PGC-1, *Cell* 98 (1999) 115–124.
 - [12] D. Knutti, A. Kralli, PGC-1, a versatile coactivator, *Trends Endocrinol. Metab.* 12 (2001) 360–365.
 - [13] R.C. Scarpulla, Transcriptional activators and coactivators in the nuclear control of mitochondrial function in mammalian cells, *Gene* 286 (2002) 81–89.
 - [14] N.G. Larsson, J. Wang, H. Wilhelmsson, A. Oldfors, P. Rustin, M. Lewandoski, G.S. Barsh, D.A. Clayton, Mitochondrial transcription factor A is necessary for mtDNA maintenance and embryogenesis in mice, *Nat. Genet.* 18 (1998) 231–236.
 - [15] M. Fluck, H. Hoppeler, Molecular basis of skeletal muscle plasticity—from gene to form and function, *Rev. Physiol., Biochem. Pharmacol.* 146 (2003) 159–216.
 - [16] R.M. Reznick, G.I. Shulman, The role of AMP-activated protein kinase in mitochondrial biogenesis, *J. Physiol.* 574 (2006) 33–39.
 - [17] D.A. Hood, I. Irrcher, V. Ljubicic, A.M. Joseph, Coordination of metabolic plasticity in skeletal muscle, *J. Exp. Biol.* 209 (2006) 2265–2275.
 - [18] A. Garnier, D. Fortin, J. Zoll, B. N'Guessan, B. Mettauer, E. Lampert, V. Veksler, R. Ventura-Clapier, Coordinated changes in mitochondrial function and biogenesis in healthy and diseased human skeletal muscle, *FASEB J.* 19 (2005) 43–52.
 - [19] E.R. Chin, E.N. Olson, J.A. Richardson, Q. Yang, C. Humphries, J.M. Shelton, W. Wu, R. Bassel-Duby, R.S. Williams, A calcineurin-dependent transcriptional pathway controls skeletal muscle fiber type, *Genes Dev.* 12 (1998) 2499–2509.
 - [20] J. Lin, H. Wu, P.T. Tarr, C.Y. Zhang, Z. Wu, O. Boss, L.F. Michael, P. Puigserver, E. Isotani, E.N. Olson, B.B. Lowell, R. Bassel-Duby, B.M. Spiegelman, Transcriptional co-activator PGC-1 α drives the formation of slow-twitch muscle fibres, *Nature* 418 (2002) 797–801.
 - [21] P.M. Garcia-Roves, J. Huss, J.O. Holloszy, Role of calcineurin in exercise-induced mitochondrial biogenesis, *Am. J. Physiol., Endocrinol. Metab.* 290 (2006) E1172–E1179.
 - [22] X. Bigard, H. Sanchez, J. Zoll, P. Mateo, V. Rousseau, V. Veksler, R. Ventura-Clapier, Calcineurin co-regulates contractile and metabolic components of slow muscle phenotype, *J. Biol. Chem.* 275 (2000) 19653–19660.
 - [23] T. Akimoto, S.C. Pohnert, P. Li, M. Zhang, C. Gumbs, P.B. Rosenberg, R.S. Williams, Z. Yan, Exercise stimulates Pgc-1 α transcription in skeletal muscle through activation of the p38 MAPK pathway, *J. Biol. Chem.* 280 (2005) 19587–19593.
 - [24] H. Wu, S.B. Kanatous, F.A. Thurmond, T. Gallardo, E. Isotani, R. Bassel-Duby, R.S. Williams, Regulation of mitochondrial biogenesis in skeletal muscle by CaMK, *Science* 296 (2002) 349–352.
 - [25] R. Bergeron, J.M. Ren, K.S. Cadman, I.K. Moore, P. Perret, M. Pypaert, L.H. Young, C.F. Semenkovich, G.I. Shulman, Chronic activation of AMP kinase results in NRF-1 activation and mitochondrial biogenesis, *Am. J. Physiol.* 281 (2001) E1340–E1346.
 - [26] H. Zong, J.M. Ren, L.H. Young, M. Pypaert, J. Mu, M.J. Birnbaum, G.I. Shulman, AMP kinase is required for mitochondrial biogenesis in skeletal muscle in response to chronic energy deprivation, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 15983–15987.
 - [27] S.B. Jorgensen, J.F. Wojtaszewski, B. Viollet, F. Andreelli, J.B. Birk, Y. Hellsten, P. Schjerling, S. Vaulont, P.D. Neuffer, E.A. Richter, H. Pilegaard, Effects of α -AMPK knockout on exercise-induced gene activation in mouse skeletal muscle, *FASEB J.* 19 (2005) 1146–1148.
 - [28] A. Garnier, D. Fortin, C. Delomenie, I. Momken, V. Veksler, R. Ventura-Clapier, Depressed mitochondrial transcription factors and oxidative capacity in rat failing cardiac and skeletal muscles, *J. Physiol.* 551 (2003) 491–501.
 - [29] Y.S. Choi, H.K. Lee, Y.K. Pak, Characterization of the 5'-flanking region of the rat gene for mitochondrial transcription factor A (Tfam), *Biochim. Biophys. Acta* 1574 (2002) 200–204.
 - [30] P. Puigserver, Z. Wu, C.W. Park, R. Graves, M. Wright, B.M. Spiegelman, A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis, *Cell* 92 (1998) 829–839.
 - [31] Y.X. Wang, C.L. Zhang, R.T. Yu, H.K. Cho, M.C. Nelson, C.R. Bayuga-Ocampo, J. Ham, H. Kang, R.M. Evans, Regulation of muscle fiber type and running endurance by PPAR δ , *PLoS Biol.* 2 (2004) e294.
 - [32] H. Wu, F.J. Naya, T.A. McKinsey, B. Mercer, J.M. Shelton, E.R. Chin, A.R. Simard, R.N. Michel, R. Bassel-Duby, E.N. Olson, R.S. Williams, MEF2 responds to multiple calcium-regulated signals in the control of skeletal muscle fiber type, *EMBO J.* 19 (2000) 1963–1973.
 - [33] A. Kaasik, V. Veksler, E. Boehm, M. Novotova, A. Minajeva, R. Ventura-Clapier, Energetic crosstalk between organelles: architectural integration of energy production and utilization, *Circ. Res.* 89 (2001) 153–159.
 - [34] F. ter Veld, J.A. Jeneson, K. Nicolay, Mitochondrial affinity for ADP is twofold lower in creatine kinase knock-out muscles. Possible role in rescuing cellular energy homeostasis, *FEBS J.* 272 (2005) 956–965.
 - [35] I. Momken, P. Lechene, N. Koulmann, D. Fortin, P. Mateo, B.T. Doan, J. Hoerter, X. Bigard, V. Veksler, R. Ventura-Clapier, Impaired voluntary running capacity of creatine kinase-deficient mice, *J. Physiol.* 565 (2005) 951–964.
 - [36] R.C. Scarpulla, Nuclear activators and coactivators in mammalian mitochondrial biogenesis, *Biochim. Biophys. Acta* 1576 (2002) 1–14.
 - [37] P. Puigserver, B.M. Spiegelman, Peroxisome proliferator-activated receptor- γ coactivator 1 α (PGC-1 α): transcriptional coactivator and metabolic regulator, *Endocr. Rev.* 24 (2003) 78–90.
 - [38] Z. Arany, H. He, J. Lin, K. Hoyer, C. Handschin, O. Toka, F. Ahmad, T. Matsui, S. Chin, P.H. Wu, I.I. Rybkin, J.M. Shelton, M. Manieri, S. Cinti, F.J. Schoen, R. Bassel-Duby, A. Rosenzweig, J.S. Ingwall, B.M. Spiegelman, Transcriptional coactivator PGC-1 α controls the energy state and contractile function of cardiac muscle, *Cell Metab.* 1 (2005) 259–271.
 - [39] T.C. Leone, J.J. Lehman, B.N. Finck, P.J. Schaeffer, A.R. Wende, S. Boudina, M. Courtois, D.F. Wozniak, N. Sambandam, C. Bernal-Mizrachi, Z. Chen, J.O. Holloszy, D.M. Medeiros, R.E. Schmidt, J.E. Saffitz, E.D. Abel, C.F. Semenkovich, D.P. Kelly, PGC-1 α deficiency causes multi-system energy metabolic derangements: muscle dysfunction, abnormal weight control and hepatic steatosis, *PLoS Biol.* 3 (2005) 101.
 - [40] J. Lin, C. Handschin, B.M. Spiegelman, Metabolic control through the PGC-1 family of transcription coactivators, *Cell Metab.* 1 (2005) 361–370.
 - [41] S. Luquet, J. Lopez-Soriano, D. Holst, A. Fredenrich, J. Melki, M. Rassoulzadegan, P.A. Grimaldi, Peroxisome proliferator-activated receptor δ controls muscle development and oxidative capability, *FASEB J.* 17 (2003) 2299–2301.
 - [42] P. Puigserver, J. Rhee, J. Lin, Z. Wu, J.C. Yoon, C.Y. Zhang, S. Krauss, V.K. Mootha, B.B. Lowell, B.M. Spiegelman, Cytokine stimulation of energy expenditure through p38 MAP kinase activation of PPAR γ coactivator-1, *Mol. Cell.* 8 (2001) 971–982.
 - [43] M. Fan, J. Rhee, J. St-Pierre, C. Handschin, P. Puigserver, J. Lin, S. Jaeger, H. Erdjument-Bromage, P. Tempst, B.M. Spiegelman, Suppression of mitochondrial respiration through recruitment of p160 myb binding protein to PGC-1 α : modulation by p38 MAPK, *Genes Dev.* 18 (2004) 278–289.
 - [44] S.B. Jorgensen, E.A. Richter, J.F. Wojtaszewski, Role of AMPK in skeletal muscle metabolic regulation and adaptation in relation to exercise, *J. Physiol.* 574 (2006) 17–31.
 - [45] P.J. Atherton, J. Babraj, K. Smith, J. Singh, M.J. Rennie, H. Wackerhage, Selective activation of AMPK-PGC-1 α or PKB-TSC2-mTOR signaling

- can explain specific adaptive responses to endurance or resistance training-like electrical muscle stimulation, *FASEB J.* 19 (2005) 786–788.
- [46] S. Jager, C. Handschin, J. St-Pierre, B.M. Spiegelman, AMP-activated protein kinase (AMPK) action in skeletal muscle via direct phosphorylation of PGC-1 α , *Proc. Natl. Acad. Sci. U. S. A.* 104 (2007) 12017–12022.
- [47] V. Aguilar, S. Alliouachene, A. Sotiropoulos, A. Sobering, Y. Athea, F. Djouadi, S. Miraux, E. Thiaudiere, M. Foretz, B. Viollet, P. Diolez, J. Bastin, P. Benit, P. Rustin, D. Carling, M. Sandri, R. Ventura-Clapier, M. Pende, S6 kinase deletion suppresses muscle growth adaptations to nutrient availability by activating AMP kinase, *Cell Metab.* 5 (2007) 476–487.