

We have recently shown that the  $H^+/ATP$  ratio can significantly decrease during ATP hydrolysis by the ATPsynthase of *Rb. capsulatus*, when the concentration of either ADP or Pi is maintained at a low level. This same phenomenon has then been observed in isolated membranes of *E. coli*. We have now purified the ATPsynthase of *E. coli* and reconstituted it into liposomes, in order to verify whether the same behavior could be observed in the isolated enzyme. The ATP hydrolysis and proton pumping activity were measured under the same experimental conditions. The hydrolysis was measured either with the colorimetric pH indicator Phenol Red or with an ATP regenerating enzymatic assay, and the proton pumping was evaluated by a calibrated ACMA assay. The hydrolysis activity was inhibited by Pi with an apparent  $K_d$  of 400  $\mu M$ , while the steady state  $\Delta pH$  was stimulated up to 200  $\mu M$  Pi and was only slightly inhibited up to 1000  $\mu M$  Pi. Both the inhibition of ATP hydrolysis and the stimulation of proton pumping as a function of Pi were lost upon ADP removal by an ADP trap. We conclude that the isolated and reconstituted ATPsynthase of *E. coli* can vary its degree of coupling as a function of Pi and ADP.

doi:10.1016/j.bbabo.2008.05.062

### S1.25 Immobilization of the $H^+$ -ATPsynthase on glass surface and single molecule fluorescence spectroscopy

Peter Oswald<sup>a</sup>, Irene Schütz<sup>a</sup>, Kathrin Förster<sup>a</sup>, Eva Galvez<sup>b</sup>, Peter Gräber<sup>a</sup>

<sup>a</sup>Albert-Ludwigs-University of Freiburg, Department of Physical Chemistry, Albertstr. 23a, 79104 Freiburg, Germany

<sup>b</sup>Instituto de Carboquímica (CSIC), Zaragoza, Spain

E-mail: peter.oswald@physchem.uni-freiburg.de

During ATP synthesis the rotor subunits of the  $H^+$ -ATPsynthase  $\gamma\epsilon c_{10}$  rotate versus stator subunits  $\alpha_3\beta_3 \delta ab_2$ . This rotation has been shown by single pair Fluorescence Resonance Energy Transfer (spFRET). The double labeled enzyme was integrated into liposomes and the diffusion time through the confocal volume of the microscope limited the observation time. In order to extend the observation time the proteoliposomes were immobilized on a glass surface. To immobilize the proteoliposomes two techniques have been used. First, the enzyme was reconstituted into liposomes containing biotinylated lipids. They are immobilized with streptavidin on a biotinylated surface. Second, the  $H^+$ -ATPsynthase carried a his-tag on the  $\beta$ -subunits. After modification of the surface with BSA or Silan proteoliposomes were immobilized via the his-tagged enzyme. With this method the observation time of a single enzyme was significantly increased, however also photobleaching of the organic fluorophores was increased. Therefore, quantum dots (QD<sub>580</sub> and QD<sub>600</sub>) with hydrophilic shells have been used as fluorescence donors. They were bound covalently to the  $\beta$ -subunits of the  $H^+$ -ATPsynthase and Atto 647N was used as acceptor. This allows the observation of a single enzyme in the second to minute time range.

doi:10.1016/j.bbabo.2008.05.063

### S1.26 Structure of the C-terminal domain of the $\epsilon$ subunit of chloroplast-type $F_1$

Hiromasa Yagi<sup>a</sup>, Hiroki Konno<sup>b</sup>, Tomoe Murakami-Fuse<sup>b</sup>, Hideo Akutsu<sup>a</sup>, Toru Hisabori<sup>b</sup>

<sup>a</sup>Institute for Protein Research, Osaka University, 3-2 Yamadaoka, Suita 565-0871, Japan

<sup>b</sup>Chemical Resources Laboratory, Tokyo Institute of Technology, Nagatsuta 4259-R1-8, Midori-Ku, Yokohama 226-8503, Japan  
E-mail: hkonno@res.titech.ac.jp

The chloroplast  $F_1$ -ATPase  $\epsilon$  subunit inhibits ATP hydrolysis with ATP-independent manner, whereas the bacterial  $\epsilon$  is ATP-dependent. To understand the structure–function relationship of  $\epsilon$ , we prepared the chimera  $\epsilon$  subunit combining the N-terminal domain from *Thermosynechococcus elongatus* and the C-terminal  $\alpha$ -helical domain from spinach  $CF_1$  ( $\epsilon_{NB\_CC}$ ) or  $EF_1$  ( $\epsilon_{NB\_CE}$ ). The  $\epsilon_{WT}$  from *T. elongatus* and  $\epsilon_{NB\_CC}$  inhibited the ATP hydrolysis activity of cyanobacteria  $F_1$  to a similar extent, whereas  $\epsilon_{NB\_CE}$  was less potent. The solution structures of  $\epsilon_{WT}$  and  $\epsilon_{NB\_CC}$  solved by NMR were so-called “retracted-state”, which was similar to that of  $EF_1$ - $\epsilon$  or  $TF_1$ - $\epsilon$ . However, the length of  $\alpha$ -helices in the C-terminal domain of these  $CF_1$ -type  $\epsilon$  was longer than that of the bacterial  $\epsilon$ . Interestingly, the loop between two helices of the  $\epsilon_{NB\_CC}$  was formed by only four residues and the retracted position of the helices was quite different from the previous reports. Significance of the unique conformation of the C-terminal  $\alpha$ -helices is discussed.

doi:10.1016/j.bbabo.2008.05.064

### S1.27 Step size of proton-driven c ring rotation in single $F_0F_1$ -ATP synthase by FRET

M.G. Düser<sup>a</sup>, N. Zarrabi<sup>a</sup>, S. Ernst<sup>a</sup>, D.J. Cipriano<sup>b</sup>, G.D. Glick<sup>c</sup>, S.D. Dunn<sup>b</sup>, M. Börsch<sup>a</sup>

<sup>a</sup>3rd Institute of Physics, University of Stuttgart, Stuttgart, Germany

<sup>b</sup>Department of Biochemistry, University of Western Ontario, London, Canada

<sup>c</sup>Department of Chemistry, University of Michigan, Ann Arbor, USA

E-mail: m.boersch@physik.uni-stuttgart.de

A mean ratio of 4.0 protons transported per synthesized ATP has been determined for the *E. coli*  $F_0F_1$ -ATP synthase recently. However, the  $F_0$  part likely contains 10  $c$  subunits corresponding to 3.3  $H^+/ATP$ . Synthesis of ATP is performed by a stepwise internal rotation of subunits in  $F_0F_1$ . Sequential conformational changes of single enzymes are monitored in real time by fluorescence resonance energy transfer, FRET. Therefore two different fluorophores have to be attached to those protein domains, which move during function. We investigated the step size of proton-driven  $c$  subunit rotation in  $F_0F_1$  by single-molecule FRET between  $a$  and  $c$ . ‘Duty cycle optimized alternating laser excitation’ minimized FRET artefacts. Rotary movements with stochastic single step sizes between 36° and 144° were determined by Hidden Markov Models. As the two coupled motors of  $F_0F_1$  showed apparently different step sizes, contributions of rotor and stator subunits for transient energy storage can be located using FRET. Monitoring  $c$  rotation we identified the action mode of the allosteric inhibitor aurovertin B, which modulates single  $F_0F_1$  activity by slowing down rotation upon ATP hydrolysis, but acts differently during ATP synthesis.

doi:10.1016/j.bbabo.2008.05.065

### S1.28 Mechanochemical coupling of $F_1$ -ATPase and intracellular ATP imaging

Hiroyuki Noji<sup>a</sup>, Rikiya Watanabe<sup>a</sup>, Daichi Okuno<sup>a</sup>, Hiromi Imamura<sup>a,b</sup>

<sup>a</sup>Institute of Scientific and Industrial Research, Osaka University, Japan

<sup>b</sup>PRSTO, JST, Osaka, Japan

E-mail: hnoji@sanken.osaka-u.ac.jp