

^aLaboratory of Intracellular Ion Channels, Nencki Institute of Experimental Biology, Polish Academy of Sciences, 3 Pasteur Street, 02-093 Warsaw, Poland

^bDepartment of Biophysics, University Life of Sciences—SGGW, 159 Nowoursynowska Street, 02-776 Warsaw, Poland

^cLaboratory of Bioenergetics, Institute of Molecular Biology and Biotechnology, Adam Mickiewicz University, Fredry 10, 61-701 Poznań, Poland

^dDepartment of Epileptology, Bonn University, Bonn, Germany

E-mail: a.szewczyk@nencki.gov.pl

Potassium channels (ATP-regulated, calcium activated and voltage dependent potassium channels) present in inner mitochondrial membranes were implicated in cytoprotective phenomenon in various tissues. These channels modulate mitochondrial matrix volume, mitochondrial respiration and membrane potential, and generation of reactive oxygen species. In this paper we describe the biophysical and pharmacological properties of new mitochondrial potassium channels recorded in *Acanthamoeba castellanii* and potato tuber mitochondria. Additionally, properties of mitochondrial potassium channels present in neuronal, cardiac tissue and endothelial cells will be described.

This work was supported by grants from MNiSW P-N/031/2006, N30105331/1676 and Polish Mitochondrial Network MitoNet.pl.

doi:10.1016/j.bbabo.2008.05.105

S3/7 Intramitochondrial signaling – Interactions among mitoK_{ATP}, PKCε, ROS, and MPT

Keith D. Garlid, Alexandre D.T. Costa

Department of Biology, Portland State University, Portland OR, 97201, USA

E-mail: garlid@pdx.edu

Our aim was to apprehend the pathways by which mitoK_{ATP} opening leads to inhibition of the mitochondrial permeability transition (MPT), thereby reducing ischemia–reperfusion injury. We showed previously that mitoK_{ATP} is opened by activation of a mitochondrial PKCε, designated PKCε1, that is closely associated with mitoK_{ATP}. MitoK_{ATP} opening causes an increase in ROS production by Complex I of the respiratory chain. This ROS activates a second pool of PKCε, designated PKCε2, which inhibits the mitochondrial permeability transition (MPT). We measured mitoK_{ATP}-dependent changes in mitochondrial matrix volume to further investigate the relationships among PKCε, mitoK_{ATP}, ROS, and MPT. We present evidence that (1) H₂O₂ and NO cause mitoK_{ATP} opening that is mediated by PKCε1 and not by direct actions on mitoK_{ATP}; (2) superoxide has no effect on mitoK_{ATP} opening; (3) H₂O₂ or NO inhibits MPT opening, and both compounds do so independently of mitoK_{ATP} activity via activation of PKCε2; (4) mitoK_{ATP} opening induced by PKG, PMA or diazoxide is not mediated by ROS; and (5) mitoK_{ATP}-generated ROS activates PKCε1 and induces phosphorylation-dependent mitoK_{ATP} opening *in vitro* and *in vivo*. Thus, mitoK_{ATP}-dependent mitoK_{ATP} opening constitutes a positive feedback loop capable of maintaining the channel open after the stimulus is no longer present. This feedback pathway may be responsible for the lasting protective effect of preconditioning, colloquially known as the memory effect.

doi:10.1016/j.bbabo.2008.05.106

(S3) Membrane transporters symposium abstracts (poster and raised abstracts)

S3.8 Effects of inhibitors on the unfolding of the mitochondrial ADP/ATP carrier by single-molecule force spectroscopy

Alex Hellawell^a, Alexej Kedrov^b, Adam Klosin^b, R. Bill Broadhurst^c, Edmund R.S. Kunji^a, Daniel Müller^b

^aMedical Research Council, Dunn Human Nutrition Unit, Cambridge, UK

^bThe Group of Cellular Machines, BioTEC, Technical University of Dresden, Germany

^cDepartment of Biochemistry, University of Cambridge, Cambridge, UK

E-mail: ek@mrc-dunn.cam.ac.uk

The mitochondrial ADP/ATP carrier exchanges cytosolic ADP for ATP synthesised in the mitochondrial matrix and replenishes the eukaryotic cell with metabolic energy. Two specific inhibitors of the carrier are known; atractyloside (ATR) and carboxyatractyloside (CATR), which differ in one carboxylate. Reconstituted histidine-tagged yeast ADP/ATP carrier AAC3 with either ATR or CATR bound was subjected to single-molecule force spectroscopy. The amino-terminal end of the protein was pulled out of the α-helical bundle in pairs of helices, reflecting the tripartite structure of the carrier. Additional resistance to unfolding was observed on helix H2 when CATR was bound rather than ATR. Two-dimensional NMR spectroscopy was used to confirm the stereochemistry of ATR, showing that the additional carboxylate of CATR is in the equatorial position. We interpret the extra resistance to be caused by the removal of the inhibitor together with the first two α-helices of the carrier, as the inhibitor is bound most strongly to these α-helices. The single-molecule force spectroscopy studies explain why CATR confers additional structural stability to the carrier.

doi:10.1016/j.bbabo.2008.05.107

S3.9 Effect of single gene deletions of *mrpA–G* and *mrpE* point mutations on activity of the Mrp Na⁺/H⁺ antiporter of alkaliphilic *Bacillus* and formation of hetero-oligomeric Mrp complex

Masato Morino^a, Shinsuke Natsu^a, Talia H. Swartz^b, Terry A. Krulwich^b, Masahiro Ito^a

^aGraduate School of Life Sciences, Toyo University, Japan

^bDepartment of Pharmacology and Systems Therapeutics, Mount Sinai School of Medicine, New York, USA

E-mail: ito@itakura.toyo.ac.jp

The putative “multi-subunit” Mrp family of secondary monovalent cation proton antiporters is physiologically important in diverse bacteria. The aim of this study was to examine structure–function of the product of the seven-gene *mrp* operon from an alkaliphilic *Bacillus*. The cloned operon was engineered so that each of the Mrp proteins (MrpA–G) could be detected. When expressed in an antiporter-deficient strain of *Escherichia coli*, Mrp-dependent Na⁺(Li⁺)/H⁺ antiport was observed. Analyses by combined Blue Native electrophoresis and SDS-PAGE demonstrated complexes that contain all 7 gene products in size ranges that could be monomers and dimers. Analyses of single, non-polar *mrp* gene deletion mutants showed that: all Mrp proteins were required for significant antiport activity; MrpD is required for stable membrane incorporation of all other Mrp proteins;