

#### S4.29 Catalytic properties of Na<sup>+</sup>-translocating NADH:quinone oxidoreductases from *Vibrio harveyi*, *Klebsiella pneumoniae*, and *Azotobacter vinelandii*

Maria S. Fadeeva<sup>a</sup>, Yulia V. Bertsova<sup>b</sup>, Cinthia Núñez<sup>c</sup>, Guadalupe Espín<sup>c</sup>, Alexander V. Bogachev<sup>b</sup>

<sup>a</sup>Faculty of Bioengineering and Bioinformatics, Moscow State University, Russia

<sup>b</sup>Department of Molecular Energetics of Microorganisms, A.N. Belozersky Institute of Physico-Chemical Biology, Moscow State University, Russia

<sup>c</sup>Departamento de Microbiología Molecular, Instituto de Biotecnología, Universidad Nacional Autónoma de México. Av. Universidad 2001. Col. Chamilpa. Cuernavaca, Morelos. 62210, México

E-mail: [masha@genebee.msu.ru](mailto:masha@genebee.msu.ru)

The main goal of our study was a comparative analysis of the catalytic properties of sodium-translocating NADH:quinone oxidoreductases (Na<sup>+</sup>-NQRs) from marine bacterium *Vibrio harveyi*, enterobacterium *Klebsiella pneumoniae*, and soil microorganism *Azotobacter vinelandii*. It is shown that their enzymes drastically differ in their affinity to sodium ions with apparent  $K_M$  values 2.7 mM, 0.67 mM and  $\approx$ 0.1 mM respectively. The enzymes also possess different sensitivity to inhibitors. Na<sup>+</sup>-NQR from *A. vinelandii* is not sensitive to low HQNO concentrations, while Na<sup>+</sup>-NQRs from *V. harveyi* and *K. pneumoniae* can be inhibited with  $I_{0.5}$  values 0.13  $\mu$ M and 0.55  $\mu$ M respectively. Also Na<sup>+</sup>-NQR from *K. pneumoniae* is fully resistant to either Ag<sup>+</sup> (which is considered to be specific inhibitor of Na<sup>+</sup>-NQR from *V. harveyi*) or *N*-ethylmaleimide. Na<sup>+</sup>-NQR from *A. vinelandii* possess transitional sensitivity to these modifiers of SH-groups. All the Na<sup>+</sup>-NQR-type enzymes are sensitive to diphenyliodonium. So the main unique characteristic of Na<sup>+</sup>-NQR is its specific requirement for sodium ions, which can be not readily detectable, since the affinity of Na<sup>+</sup>-NQR to Na<sup>+</sup> can be very high.

doi:10.1016/j.bbabbio.2008.05.160

### (S5) Mitochondrial biogenesis symposium lecture abstracts

#### S5/1 Control of the synthesis of uncoupling and coupling proteins in brown adipose tissue

Barbara Cannon<sup>a</sup>, Tatiana V. Kramarova<sup>a</sup>, Natasa Petrovic<sup>a</sup>, Irina G. Shabalina<sup>a</sup>, Josef Houstek<sup>b</sup>, Jan Nedergaard<sup>a</sup>

<sup>a</sup>The Wenner-Gren Institute, The Arrhenius Laboratories F3, Stockholm University, SE-10691 Stockholm, Sweden

<sup>b</sup>Czech Academy of Sciences, Prague, Czech Republic

E-mail: [barbara.cannon@wgi.su.se](mailto:barbara.cannon@wgi.su.se)

In active brown adipose tissue, the balance between coupled and uncoupled respiration is the opposite of that seen elsewhere. This is accomplished through two features: low content of coupling proteins (the ATP-synthase complex) and high content of uncoupling protein (UCP1). In the tissue, very high expression (mRNA levels) of all subunits of ATP-synthase is seen — except for subunit c, implying that ATP-synthase assembly is under control of sub-c amount. We have now demonstrated that artificially-induced overexpression of sub-c results in increased amounts of fully competent ATP-synthase. In wildtype, despite high mRNA levels for the other subunits, no unassembled ATP-synthase subunits are observed in blue-native gels. This implies translation control of the other components of the ATP-synthase. Concerning the  $\beta$ -subunit, the control in different tissues may be related to formation of an RNA/protein complex that is dependent on a stem-loop structure in the 3'UTR mRNA. Brown adipose tissue recruitment and UCP1 expression are normally considered to be under sympathetic

control. There is physiological reason for a nonsympathetic recruitment pathway. We find that chronic treatment of brown (pre)adipocytes with PPAR $\gamma$ -agonists activates mitochondriogenesis and UCP1 expression, leading to thermogenically competent brown-fat cells, i.e. cells that although naive to norepinephrine respond thermogenically to norepinephrine.

doi:10.1016/j.bbabbio.2008.05.161

#### S5/2 Regulation of mitochondrial dynamics by nitric oxide is a key event in myogenesis

Clara De Palma<sup>a</sup>, Sestina Falcone<sup>b</sup>, Maria T. Bassi<sup>b</sup>, Luca Scorrano<sup>c</sup>, Giulio Cossu<sup>d</sup>, Salvador Moncada<sup>e</sup>, Silvia Brunelli<sup>d,f</sup>, Emilio Clementi<sup>a,b</sup>

<sup>a</sup>Department of Preclinical Sciences, University of Milano, Italy

<sup>b</sup>E. Medea Scientific Institute, Bosisio Parini, Italy

<sup>c</sup>Venetian Institute of Molecular Medicine, Padova, Italy

<sup>d</sup>San Raffaele Scientific Institute, Milano, Italy

<sup>e</sup>The Wolfson Institute for Biomedical Research, University College London

<sup>f</sup>Department of Experimental Medicine, University of Milano-Bicocca, Milano, Italy

E-mail: [emilio.clementi@unimi.it](mailto:emilio.clementi@unimi.it)

Mitochondria exist in two interconverting forms, i.e. as small isolated particles, and as extended filaments, networks or clusters. Here we provide evidence that in differentiating myoblasts endogenous nitric oxide (NO) generation controls mitochondrial shape: in the absence of NO mitochondrial fission occurs rapidly. The action of NO is specifically addressed to mitochondrial fission since in PEG fusion assay organelle fusion was not modified by the treatment with the NO synthase-inhibitor L-NAME. A key protein involved in mitochondrial fission is the large GTPase DRP-1. DRP-1 translocation to the mitochondria promotes mitochondrial fission. DRP-1 translocation and mitochondrial fission were stimulated by L-NAME and inhibited by exogenous NO. In addition, NO inhibited DRP-1 GTPase activity. We also found that in differentiating myoblasts NO is required for the expression of differentiation markers including myogenin and muscle specific myosin since L-NAME inhibited myogenic differentiation, and exogenous NO restored it. Overexpression of a dominant negative DRP-1 reversed the inhibitory effect of L-NAME on myogenesis. Our results indicate that NO controls a key event in mitochondrial dynamics that may have relevant implications for both myogenesis and control of energy metabolism in developing skeletal muscle.

doi:10.1016/j.bbabbio.2008.05.162

#### S5/3 Bioenergetics of mitochondrial protein topogenesis

Walter Neupert

Adolf-Butenandt-Institute, University of Munich, Germany

E-mail: [Neupert@med.uni-muenchen.de](mailto:Neupert@med.uni-muenchen.de)

The TIM23 translocase is involved in the topogenesis of the vast majority of mitochondrial proteins. Preproteins pass through the TOM complex of the outer membrane and are then transferred across or laterally inserted into the inner membrane (IM). The electrical membrane potential  $\Delta\psi$  is required for the translocation of the targeting signal across the IM where it can be reached by the chaperone mtHsp70. Further translocation does not require  $\Delta\psi$ , but instead matrix ATP. ATP hydrolysis drives cycles of binding of mtsp70 to incoming unfolded preproteins. mtHsp70 is part of the mitochondrial import motor which comprises further components,