

result of a point mutation (T to C) at nucleotide position 8851 of the mitochondrial DNA (mtDNA). This mutation changes a highly conserved tryptophane residue to arginine at amino acid position 109 of the subunit *a* of mitochondrial F_1F_0 -ATP synthase, a complex which provides most of the cellular ATP in humans. Nothing is known on the consequences of the T8851C mutation on the mitochondrial ATP synthase. To gain insight into the primary pathogenic mechanisms induced by T8851C, we have investigated the consequences of this mutation on the ATP synthase of yeast where the protein homologous to subunit *a* (referred to as Atp6p) is also encoded by the mtDNA. The modified yeast exhibited a very slow growth phenotype on non-fermentable carbon sources, both at 28 °C (the optimal temperature for growing yeast) and at 36 °C. *In vitro*, mitochondria from T8851C yeast grown at 28 °C showed a 60% deficit in ATP production, while those prepared from the mutant grown at 36 °C had an ATP synthesis activity below 5% that of the wild type. The mutated F_1F_0 complex was correctly assembled, at both temperatures, and had a very poor ATPase activity (10% that of the wild type), both in mitochondria and after purification. Electron microscopy revealed that many of the mitochondrial matrices in T8851C yeast grown at 36 °C exhibited septae made of apposed inner mitochondrial membranes. Another anomaly was an increased mitophagic activity, presumably in response to the T8851C-induced damaging of mitochondria. Thus, in addition to a bioenergetic deficit, alterations in mitochondrial dynamics and homeostasis may also participate in the pathogenic mechanism induced by T8851C.

doi:10.1016/j.bbabbio.2010.04.169

4P.8 Iron deficiency in children with mitochondrial disease

Young-Mock Lee, Hae Eun Kwon, Junh Hun Lee, Hoon Chul Kang, Joon Soo Lee, Heung Dong Kim

Department of Pediatrics, Gangnam Severance Hospital, Severance Children's Hospital, Yonsei University College of Medicine, Seoul, Republic of Korea

E-mail: ymleemd@yuhs.ac

Mitochondrial disease is an energy metabolic disorder with various organ involvements. Iron is widely known to be one of the most important nutrients required for normal brain development and several essential metabolic functions. We retrospectively studied the laboratory data on iron deficiency (ID) in 69 children with mitochondrial respiratory chain complex (MRC) defects by biochemical enzyme assay using muscle tissue. We analyzed the differences between groups of mitochondrial disease based on the presence of ID. ID has higher prevalence in children with mitochondrial disease than in the normal population. There were 6 (9%) patients with low hemoglobin, 12 (17%) with low serum ferritin, and 22 (32%) with low transferrin saturation levels among children with MRC defects. In comparisons between the ID and the non-ID group of MRC-defect patients, the frequency of MRC I defect was significantly higher in the ID group while that of MRC IV defect was higher in the non-ID group. Abnormal brain magnetic resonance imaging (MRI) findings were more frequently detected in the ID group. The incidence of failure to thrive and gastrointestinal symptoms were significantly higher in the ID group. Early diagnosis and proper treatment of ID are recommended. Especially in cases with risk factors such as failure to thrive or gastrointestinal manifestation, active evaluation of ID should be encouraged.

doi:10.1016/j.bbabbio.2010.04.170

4P.9 The new molecular p.M177T identified in two unrelated patients with clinical features of SCO2-dependent cytochrome c oxidase deficiency

Sylvia Łuczak, Dorota Piekutowska-Abramczuk, Paweł Kowalski, Elżbieta Ciara, Dorota Jurkiewicz, Maria Borucka-Mankiewicz, Anna Tańska, Magdalena Pelc, Joanna Trubicka, Małgorzata Krajewska-Walasek, Ewa Pronicka

The Children's Memorial Health Institute, Warsaw, Poland

E-mail: s.luczak@czd.pl

Cytochrome c oxidase (COX, CIV) is one of the complexes of the OXPHOS system located in the inner mitochondrial membrane and catalyzing the last step of electron's transfer from cytochrome c to molecular oxygen. It is composed of 13 subunits encoded by mitochondrial and nuclear DNA. A correct assembly and function of COX require a substantial number of the nuclear, ancillary proteins, including SCO2, which is involved in the transport and incorporation of the copper ions to the CuA enzymatic site on COXII subunit. Human SCO2 gene is located on the chromosome 22q13, and contains two exons. Only the 801 bp fragment of exon 2 undergoes translation into 266 amino acid protein. Mutations in the SCO2 gene lead to serious damage of the protein resulting in severe COX deficiency observed mainly in muscle, heart and brain. The common substitution, g.1541G>A (p.E140K) was identified at least on one allele in all so far reported patients with COX deficiency. In the group of 23 Polish patients, the common substitution was found on 84% of the studied alleles. The clinical features of the disease associated with SCO2 deficiency include early onset, fatal hypertrophic cardiomyopathy with respiratory insufficiency, encephalopathy, hypotonia and metabolic acidosis. The aim of this study was to characterize the molecular background of the disease in three patients from two unrelated families with clinically and biochemically recognized cytochrome c oxidase deficiency. Here we present patients with the same genotype, comprising the common mutation, g.1541G>A and a new, not described in the literature, molecular variant g.1653T>C. The new variant affects the highly conserved methionine at 177 position of the SCO2 protein (p.M177T) and was not found on 600 control alleles. Additionally, g.1653T>C substitution was predicted by SIFT BLink programme as a pathogenic mutation. Our findings indicate that the compound heterozygous genotype, p.M177T/p.E140K, is responsible for clinical manifestation of destroyed SCO2 protein.

The study was partly supported by the CMHI project no. S11/2009 and Polish Ministry of Science Project no. PB 0890/P05/2005/29.

doi:10.1016/j.bbabbio.2010.04.171

4P.10 Impaired mitochondrial energetic in patients harbouring SURF1 mutations is caused by uncoupling of cytochrome c oxidase

Petr Pecina, Alena Pecinova, Nikola Kovarova, Martin Kalous, Ivan Miksik, Josef Houstek

Institute of Physiology and Center for Applied Genomics, AS CR, v.v.i., Prague, Czech Republic

E-mail: petr.pecina@gmail.com

Leigh syndrome is most frequently caused by mutations of *SURF1* gene, which encodes cytochrome c oxidase (COX) specific assembly factor. Our previous studies suggested that fibroblasts from patients harbouring *SURF1* mutations accumulate incomplete forms of COX lacking several small nuclear-encoded subunits with decreased H^+ /e stoichiometry. In experiments aimed at detailed characterization of the mitochondrial energetics, we observed 30–50% decreased respiratory capacity available for ATP synthesis (RCR_p). When using COX-specific substrates ascorbate + TMPD, the RCR_p was 7-fold lower than in controls, suggestive of deficient proton pumping of COX in patient