

can be, at least partially, due to their interactions with the endogenous pool of fatty acids.

This work was supported by RFBR (grants 06-04-49687 and 09-04-01238).

doi:[10.1016/j.bbabbio.2010.04.206](https://doi.org/10.1016/j.bbabbio.2010.04.206)

5P.21 The effect of Ca^{2+} on reactive oxygen species generation in brain mitochondria in the absence of permeability transition

Laszlo Tretter, Zsolt Komar, Vera Adam-Vizi

Semmelweis University, Department of Medical Biochemistry, Hungary

Hungarian Academy of Sciences, Neurobiochemical Group, Hungary

E-mail: tretter@eok.sote.hu

Glutamate excitotoxicity is a key element in the pathomechanism of acute (ischemia-reperfusion) and chronic (Alzheimer's disease, Parkinson's disease, Huntington's disease) neurological disorders. Stimulation of glutamate receptors results in the elevation of intracellular $[\text{Ca}^{2+}]$ which can activate reactive oxygen species (ROS) generation. The aim of our experiments was to study the effects of high micromolar Ca^{2+} concentrations on the H_2O_2 generation in isolated guinea pig brain mitochondria, supported by NADH-generating substrates; glutamate plus malate. H_2O_2 formation was detected extramitochondrially by Amplex red assay. In parallel with ROS formation NAD(P)H autofluorescence was detected. Mitochondrial membrane potential ($\Delta\psi_m$) was measured by safranin O and TMRM fluorescence respectively. Swelling of mitochondria was detected by light scattering. Permeability transition pore (PTP) opening was measured by calcium induced calcium release and by quenching of calcein fluorescence. PTP was prevented by ADP, a very efficient inhibitor of mitochondrial permeability transition. In the presence of ADP 50 μM Ca^{2+} (500 nmol/mg protein) did not induce PTP opening but enhanced mitochondrial H_2O_2 release by $81 \pm 18\%$. Mitochondria were able to take up calcium; after a transient depolarization $\Delta\psi_m$ was restored, even hyperpolarization was detected and parallel with these NAD(P)H fluorescence was increased. With 300 μM Ca^{2+} membrane potential collapsed without recovery and H_2O_2 release was unchanged. At 300 μM $[\text{Ca}^{2+}]$ in the presence of ADP, mitochondria were unable to complete Ca^{2+} -uptake but no signs of PTP were detected in the time-frame of the experiments. In highly polarized mitochondria, in the presence of ATP or oligomycin ROS production was elevated, Ca^{2+} failed to stimulate mitochondrial ROS generation and hyperpolarization did not follow the Ca^{2+} -induced depolarization. It is suggested that in the presence of nucleotides the effect of Ca^{2+} on mitochondrial ROS release is related to changes in $\Delta\psi_m$. The increased ROS release evoked by Ca^{2+} in the presence of ADP in isolated mitochondria is unrelated to PTP and would not explain the extensive cellular ROS production observed during glutamate excitotoxicity.

doi:[10.1016/j.bbabbio.2010.04.207](https://doi.org/10.1016/j.bbabbio.2010.04.207)

5P.22 Cardiolipin: Altered content and fatty acid composition in mitochondria from mtDNA mutator mice

Mikhail Yu. Vyssokikh^{1,2}, Irina G. Shabalina², Alexandra Trifunovic³, Barbara Cannon², Vladimir P. Skulachev¹, Jan Nedergaard²

¹Belozersky Institute of Physico-Chemical Biology,

Lomonosov Moscow State University, Vorobyevy Gory 1, Moscow 119991, Russia

²The Wenner-Gren Institute, the Arrhenius Laboratories F3, Stockholm University, SE-106 91 Stockholm, Sweden

³Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases, University of Cologne, Germany

E-mail: mikhail.vyssokikh@gmail.com

We have recently demonstrated that the assembly and turnover of the mitochondrial respiratory chain complexes I, III and IV (but not complexes II and V) are altered in mitochondria from mtDNA mutator mice [1]. Since it is known that cardiolipin is essential for assembly and stability of respiratory chain complexes, we have elected to study this phospholipid in mitochondria from mtDNA mutator mice. The content of mitochondrial phospholipids was analysed by two-dimensional high performance thin layer chromatography (2D-HPTLC). The content of cardiolipin was significantly lower in liver and skeletal muscle mitochondria from mtDNA mutator mice than in wild-type mitochondria. To analyse the fatty acid composition of cardiolipin, gas chromatography/flame ionization detection or electron ionization-mass spectrometry (GC/FID or EI MS) was applied. The content of the polyunsaturated *n*-6 fatty acids was remarkably lowered in the cardiolipin fraction from skeletal muscle and liver mitochondria of mtDNA mutator mice, as compared with wild-type mice. Mitochondrial phospholipids were also studied in mice chronically treated with mitochondria-targeted antioxidant (plastoquinone derivative, SkQ1), added to drinking water. The content of cardiolipin and its fatty acid composition were normalised in mtDNA mutator mitochondria after treatment with SkQ1. We conclude that cardiolipin content is decreased and its fatty acid composition markedly altered in mitochondria from mtDNA mutator mice. The cause and significance of these alterations are of interest, considering the special role of cardiolipin in mitochondrial bioenergetics.

Reference

[1] Edgar D *et al.* (2009) *Cell. Metab.* **10**: 131–138.

doi:[10.1016/j.bbabbio.2010.04.208](https://doi.org/10.1016/j.bbabbio.2010.04.208)

5P.23 Antioxidant defence systems and generation of reactive oxygen species in osteosarcoma cells with defective mitochondria: Effect of selenium

Marta Wojewoda, Joanna Szczepanowska, Jerzy Duszyński

Nencki Institute of Experimental Biology,

Department of Biochemistry, Warsaw, Poland

E-mail: m.wojewoda@nencki.gov.pl

Mitochondrial diseases originate from mutations in mitochondrial or nuclear genes encoding for mitochondrial proteome. Neurogenic muscle weakness, ataxia and retinitis pigmentosa (NARP) syndrome is associated with the T8993G transversion in *ATP6* gene which results in substitution at the very conservative site in the subunit 6 of mitochondrial ATP synthase. Defects in the mitochondrial respiratory chain and the ATPase are considered to be accompanied by changes in the generation of reactive oxygen species (ROS). This study was aimed to elucidate effects of selenium on ROS and antioxidant system of NARP cybrid cells with 98% of T8993G mutation load. We found that selenium decreased ROS generation and increased the level and activity of antioxidant enzymes such as glutathione peroxidase (GPx) and thioredoxin reductase (TrxR). Therefore, we propose selenium to be a promising therapeutic agent not only in the case of NARP syndrome but also other diseases associated with mitochondrial dysfunctions and oxidative stress.

doi:[10.1016/j.bbabbio.2010.04.209](https://doi.org/10.1016/j.bbabbio.2010.04.209)

5P.24 Response of *Acanthamoeba castellanii* mitochondria to hydrogen peroxide stress

Andrzej Woyda-Ploszczyca, Jarosław Haremza, Wojciech Michalak, Nina Antos-Krzeminska, Wiesława Jarmuszkiewicz

Laboratory of Bioenergetics, Faculty of Biology,

Adam Mickiewicz University, Poznań, Poland

E-mail: awoy@amu.edu.pl