

Udine, Italy

E-mail: strecker@zbc.kgu.de

Apurinic apyrimidinic endonuclease 1/redox effector factor 1 (APE1/Ref-1) is a vital protein that acts as a master regulator of cellular response to oxidative stress conditions and contributes to the maintenance of genome stability [1, 2]. APE1 is involved in both the base excision repair (BER) pathways of DNA lesions, acting as the major apurinic/apyrimidinic (AP) endonuclease, and in transcriptional regulation of gene expression as a redox coactivator of different transcription factors. In different mammalian cell types the APE1 subcellular distribution is mainly nuclear and is critical for controlling cellular proliferative rate [3]. However, cytoplasmic, mitochondrial, and endoplasmic reticulum localization has also been reported [4] but the specific mechanism for redirecting APE1 into the different subcellular compartments is not known. Mitochondrial DNA is constantly exposed to oxidative injury. Due to its location close to the main source of reactive oxygen species, the inner mitochondrial membrane, mtDNA is more susceptible than nuclear DNA to oxidative damage. Because BER is the main DNA repair pathway coping with oxidative lesions, most of the investigations on mitochondrial DNA repair focus on mtBER. To verify mitochondrial localization of APE1/Ref-1, a flag-tagged protein was expressed in HeLa cells and its intracellular localization was tracked by immunocytochemistry. Subfractioned bovine heart mitochondria were also used to analyze the localization of mtAPE1. Mitoplasts were prepared by low digitonin treatment of mitochondria and separated from fraction containing outer membranes and intermembrane space soluble proteins by differential centrifugation. After submitochondrial fractionation, mtAPE1 was predominantly found in the soluble protein rich fraction of the intermembrane space. Notably, in agreement with a previous study [5], only the full-length protein was detected in mitochondrial extracts, raising doubts about the mitochondrial targeting mechanism previously suggested [6]. The mitochondrial import pathway and the potential targeting signals of mtAPE1 are currently under investigation.

References

- [1] Tell G *et al.* (2009) *Antioxid Redox Signal* **11**: 601–620.
- [2] Tell G *et al.* (2010) *Mol. Cell. Biol.* **30**: 366–371.
- [3] Izumi T *et al.* (2005) *Proc. Natl. Acad. Sci. U.S.A.* **102**: 5739–5743.
- [4] Mitra S *et al.* (2007) *DNA Repair (Amst.)* **6**: 1064–1070.
- [5] Tell G *et al.* (2001) *Mutat. Res.* **485**: 143–152.
- [6] Chattopadhyay R *et al.* (2006) *Nucleic Acids Res.* **34**: 2067–2076.

doi:[10.1016/j.bbabbio.2010.04.436](https://doi.org/10.1016/j.bbabbio.2010.04.436)

7P.3 Effects of dietary methylmercury on the Zebrafish (*Danio rerio*) liver proteome expression and mitochondrial function

Brésart David¹, Fourdrillis Séverine¹,
Mathy Gregory^{1,2}, Sluse Francis¹

¹Laboratory of Bioenergetics and Cellular Physiology, B6,
Allée de la Chimie 3, 4000 Liège, Belgium

²Laboratoire de biochimie végétale, Bât. B22 Biochimie végétale,
boulevard du Rectorat 27, 4000 Liège, Belgium

E-mail: hotback7@hotmail.com

In the present work we have studied the chronic and the acute effects of methylmercury (MeHg) in a model fish (*Danio rerio*). The chronic effects of methylmercury intoxication were showed by a comparative proteomics approach realized on the cellular and mitochondrial soluble proteome in the liver. At the mitochondrial proteome level 56 proteins exhibiting a significant difference in their abundance were identified. The proteins involved in the Krebs cycle and the amino acid metabolism were down-regulated while the

enzymes implicated in the β -oxidation and the OXPHOS apparatus were up-regulated as well as the antioxidant systems and the ROS scavenger enzymes. At the cellular proteome level, 74 proteins were identified and exhibited a significant difference. The results suggested that glycolysis is down-regulated, as well as the pentose phosphate pathway. The acute effect of methylmercury was studied on isolated liver mitochondria using a Clark electrode oxymeter. The state 3 mitochondrial respiration was readily inhibited by MeHg, on the contrary the state 4 respiration exhibited a concentration threshold in the inhibition, suggesting an inhibition at the level of the dicarboxylate transport. This proposal was confirmed by comparing the effects of both n-butylmalonate, a well-known inhibitor of the dicarboxylate carrier, and MeHg. Thus the proteomic approach allows us to understand the metabolic changes that permit the cell to ensure its viability during dietary methylmercury intoxication. According to respiratory results it could be proposed that those changes were the consequences of substrate transport impairment by MeHg.

doi:[10.1016/j.bbabbio.2010.04.437](https://doi.org/10.1016/j.bbabbio.2010.04.437)

7P.4 Src-mediated tyrosine phosphorylation: A new way of regulating mitochondrial bioenergetics?

Etienne Hébert Chatelain, Thierry Letellier, Jeanne Dachary-Prigent
INSERM-U688 Physiopathologie Mitochondriale,
Université Victor Ségalen-Bordeaux 2, 146 rue Léo Saignat,
F-33076 Bordeaux-Cedex, France
E-mail: etienne.hebertchatelain@etud.uordeaux2.fr

Mitochondria produce the most part of the energy consumed by cells through oxidative phosphorylation. Since energetic demands of cells depend on their functions, energy production must be tightly regulated. Phosphorylation (or dephosphorylation) is one of the major regulation pathways of enzymes. Numerous proteins are tyrosine-phosphorylated among electron transport system components in rat brain mitochondria [1]. Several tyrosine kinases are also present in those mitochondria but the tyrosine Src kinase seems the most active of them [2]. Our goal was to study how the regulation of this kinase influences oxidative phosphorylation in rat brain mitochondria. First, we demonstrated that tyrosine phosphatase PTP1B, a well-known Src regulator in cytosol [3], seems to regulate Src activity in rat brain mitochondria as it does in cytosol (i.e. it activates Src by dephosphorylating its terminal pY527 residue). Moreover, *in vitro* ATP addition caused activation of Src and global increase of tyrosine-phosphorylated mitochondrial proteins. The global increase of tyrosine-phosphorylated mitochondrial proteins was accompanied by an increase of COX activities and decrease of complexes III and V. Besides, activation of Src, increase of tyrosine phosphorylation and changes of enzymatic activities were reduced when mitochondria were preincubated with pharmacological inhibitors of Src or PTP1B. Finally, mitochondrial ADP-stimulated respiration decreased in the presence of each inhibitor. These results clearly show that Src kinase activity (partly regulated by PTP1B) influences mitochondrial physiology. Therefore, Src-mediated tyrosine phosphorylation seems to represent a major regulation pathway of oxidative phosphorylation, at least in rat brain mitochondria.

References

- [1] Augereau O *et al.* (2005) *CMLS* **62**: 1478–1488.
- [2] Tibaldi E, *et al.* (2008) *J. Cell. Biochem.* **104**: 840–849.
- [3] Roskoski R (2004) *Bioch. Biophys. Res. Comm.* **324**: 1154–11164.

doi:[10.1016/j.bbabbio.2010.04.438](https://doi.org/10.1016/j.bbabbio.2010.04.438)