



## Review

## Biosynthesis and physiology of coenzyme Q in bacteria ☆


Laurent Aussel <sup>a</sup>, Fabien Pierrel <sup>b</sup>, Laurent Loiseau <sup>a</sup>, Murielle Lombard <sup>c</sup>, Marc Fontecave <sup>c</sup>, Frédéric Barras <sup>a,\*</sup>
<sup>a</sup> Laboratoire de Chimie Bactérienne, UMR 7283 Aix-Marseille Université - CNRS, Institut de Microbiologie de la Méditerranée, 31 Chemin Joseph Aiguier 13009 Marseille, France

<sup>b</sup> Laboratoire de Chimie et Biologie des Métaux, UMR 5249 CEA - Université Grenoble I - CNRS, 17 Rue des Martyrs, 38054 Grenoble Cedex France

<sup>c</sup> Laboratoire de Chimie des Processus Biologiques, UMR 8229 CNRS, UPMC, Collège de France, 11 Place Marcellin Berthelot, 75231 Paris Cedex 05 France

## ARTICLE INFO

## Article history:

Received 9 December 2013

Received in revised form 23 January 2014

Accepted 24 January 2014

Available online 28 January 2014

## Keywords:

Coenzyme Q

ubi genes

Escherichia coli

Q<sub>8</sub> biosynthesis

Aerobic respiration

Salmonella

## ABSTRACT

Ubiquinone, also called coenzyme Q, is a lipid subject to oxido-reduction cycles. It functions in the respiratory electron transport chain and plays a pivotal role in energy generating processes. In this review, we focus on the biosynthetic pathway and physiological role of ubiquinone in bacteria. We present the studies which, within a period of five decades, led to the identification and characterization of the genes named *ubi* and involved in ubiquinone production in *Escherichia coli*. When available, the structures of the corresponding enzymes are shown and their biological function is detailed. The phenotypes observed in mutants deficient in ubiquinone biosynthesis are presented, either in model bacteria or in pathogens. A particular attention is given to the role of ubiquinone in respiration, modulation of two-component activity and bacterial virulence. This article is part of a Special Issue entitled: 18th European Bioenergetic Conference.

© 2014 Elsevier B.V. All rights reserved.

## 1. Introduction

In most living organisms, catalytic reactions involved in cell energization generate electrons which are funneled to the quinone pool. Thereafter, the reduced quinones (quinols) serve as substrates for reduction of the terminal acceptors. In *Escherichia coli*, three kinds of quinones are involved in this process: ubiquinone, also known as coenzyme Q (Q), menaquinone (MK) and demethylmenaquinone (DMK) [1–3]. MK and DMK have low midpoint potentials ( $E^\circ = -74$  mV and  $+36$  mV, respectively, [4]) and are involved in anaerobic respiration while Q, which has a higher midpoint potential ( $E^\circ = +100$  mV, [4]), is involved in aerobic respiration [5,6]. The *E. coli* genome contains gene clusters for three cytochrome oxidase enzymes: cytochrome *bo* oxidase (*cyoABCD*), cytochrome *bd-I* oxidase (*cydABX*) and cytochrome *bd-II* oxidase (*appCD*). The three enzymes function as the major terminal oxidases in the aerobic respiratory chain of *E. coli* catalyzing electron transfer from ubiquinol to oxygen [7].

Ubiquinone is a widespread redox-active lipid which consists of a conserved aromatic ring and a polyprenyl hydrophobic tail, with the number of isoprenyl units varying among species: six in *Saccharomyces cerevisiae*, eight in *E. coli* and ten in humans [8–10]. Therefore, *E. coli* ubiquinone is designated Q<sub>8</sub>. Its biosynthesis is a highly conserved pathway, which involves a large number of genes, named *ubi*, that have been identified from genetic studies [11,12]. Q<sub>8</sub> is located in the bacterial plasma membrane and was described to be an essential element for aerobic respiratory growth, gene regulation, oxidative stress adaptation, and various processes depending upon proton motive force (PMF) [13–16]. In this review, we present the genes involved in the Q<sub>8</sub> biosynthetic pathway in bacteria, with a particular attention on those recently identified and on the remaining gaps in current knowledge. We focus on the enzymatic actors, i.e. those involved in the decoration of the aromatic ring leading to Q<sub>8</sub>, as well as the accessory ones. A few structures of Ubi proteins are presented and the phenotypes associated with *ubi* mutations are described and discussed.

2. The Q<sub>8</sub> biosynthetic pathway in bacteria


The *ubi* genes have been extensively studied over a period of five decades since the pioneering work of Cox and Gibson [17]. Biosynthesis of Q<sub>8</sub> in *E. coli* requires nine *ubi* genes, most of them encoding enzymes that decorate the aromatic ring of the 4-hydroxybenzoate (4-HB) universal precursor (Fig. 1). Noticeably in *E. coli*, *ubi* genes are all scattered around the chromosome (Fig. 2A and B). It is important to stress that in a few cases, the mutation of genes required for Q<sub>8</sub> production led to the accumulation of 3-octaprenylphenol (OPP), an early intermediate of Q<sub>8</sub> biosynthetic pathway (Fig. 1 and Table 1). As a consequence, these

Abbreviations: SAM, S-Adenosylmethionine; Q/Q<sub>8</sub>, Coenzyme Q/Q<sub>8</sub>; DMK/DMK<sub>8</sub>, Demethylmenaquinone; DDMQ<sub>8</sub>, Demethyldemethoxy coenzyme Q<sub>8</sub>; DMQ<sub>8</sub>, Demethoxy coenzyme Q<sub>8</sub>; DMSO, Dimethyl sulfoxide; FAD, Flavine adenine dinucleotide; FMN, Flavin mononucleotide; H<sub>2</sub>O<sub>2</sub>, Hydrogen peroxide; 3-HB, 3-Hydroxybenzoate; 4-HB, 4-Hydroxybenzoate; Fe-S, Iron-sulfur; MK/MK<sub>8</sub>, Menaquinone; OHB, 3-Octaprenyl-4-hydroxybenzoate; 4-HP<sub>8</sub>, 3-Octaprenyl-4-hydroxyphenol; OPP, 3-Octaprenylphenol; PHBH, Para-hydroxybenzoate hydroxylase; PMF, Proton motive force; O<sub>2</sub><sup>-</sup>, Superoxide anion; TMAO, Trimethylamine N-oxide

☆ This article is part of a Special Issue entitled: 18th European Bioenergetic Conference.

\* Corresponding author at: Laboratoire de Chimie Bactérienne, 31 Chemin Joseph Aiguier, 13402 Marseille Cedex 20, France. Tel.: +33 491 16 45 79; fax: +33 491 71 89 14.

E-mail address: [barras@imm.cnrs.fr](mailto:barras@imm.cnrs.fr) (F. Barras).



**Fig. 1.** Biosynthetic pathway of ubiquinone in *Escherichia coli*. The numbering of the aromatic carbon atoms is shown on coenzyme  $Q_8$ , and the octaprenyl tail is represented by R on C-3 of the different biosynthetic intermediates. The name of the enzymes catalyzing the reactions (each labeled with a lowercase letter) is indicated. Abbreviations used for 4-hydroxybenzoate (4-HB), 3-octaprenyl-4-hydroxybenzoate (OHB), 3-octaprenylphenol (OPP), coenzyme  $Q_8$  ( $Q_8$ ), C1-demethyl-C6-demethoxy- $Q_8$  (DDMQ $_8$ ), and C6-demethoxy- $Q_8$  (DMQ $_8$ ) are underlined. The XanB2 protein, present in some prokaryotes but not in *E. coli*, catalyzes the production of 4-HB from chorismate. The biosynthetic intermediates that accumulate in mutants affected in the different steps are listed in Table 1.

genetics studies sometimes failed at identifying precisely the biosynthetic step altered by the mutation.

In 1968, a screen based on the selection of mutants unable to grow on malate and examination of the quinone content of these strains


grown on glucose led to the identification of *ubiA* as the first gene involved in  $Q_8$  biosynthesis [18]. Poole and colleagues found the *ubiA* mutant to be unable to grow aerobically on non-fermentable substrates but able to grow anaerobically on glycerol with alternative electron acceptors such as fumarate [19]. The *ubiA* gene was predicted to encode a membrane-bound 4-hydroxybenzoate octaprenyltransferase. Within the same operon, *ubiA* lies downstream the *ubiC* gene (Fig. 2B). The UbiC protein catalyzes the first committed step in the biosynthesis of  $Q_8$ , the conversion of chorismate to 4-HB (Reaction a, Fig. 1) [20]. Interestingly, in *Xanthomonas campestris* (which does not contain any *ubiC* gene or homologue), the XanB2 protein was reported to convert chorismate into 3-HB or 4-HB using two distinct catalytic domains not related to UbiC (Reaction a, Fig. 1) [21].

Mutation in the *E. coli ubiG* gene yields strains unable to grow aerobically on nonfermentable substrates [22]. UbiG is one of the few Ubi enzymes that has been purified and assayed in vitro. Activity assays showed that UbiG is a S-adenosylmethionine (SAM)-dependent methyltransferase, which catalyzes the two O-methylation steps of  $Q_8$  biosynthesis (Reactions e and i, Fig. 1) [23].

The *ubiE* gene encodes the second, also SAM-dependent, methyltransferase of the  $Q_8$  biosynthetic pathway (Reaction g, Fig. 1) [24]. The *ubiE* mutant is deficient for growth on succinate and accumulates demethyldemethoxy-coenzyme  $Q_8$  (DDMQ $_8$ ) and demethylmenaquinone (DMK $_8$ ) as predominant intermediates (Table 1), leading to the conclusion that *ubiE* is required for C-methylation in both ubiquinone and menaquinone synthesis [24].

An *E. coli ubiD* mutant accumulates 3-octaprenyl-4-hydroxybenzoate (OHB) (Reaction c, Fig. 1) [25,26]. It was shown that a partially purified membrane-bound UbiD protein was able to convert OHB into OPP, strongly suggesting UbiD involvement in the decarboxylation step of  $Q_8$  biosynthesis [26]. However, a *ubiD* mutant retains the ability to produce about 25% of the wild-type levels of  $Q_8$ , consistent with the existence of a second decarboxylase [26]. Inactivation of the *ubiX* gene leads to low levels of  $Q_8$ , a reduced growth on succinate and accumulation of OHB (Table 1), suggesting a decarboxylase activity for UbiX [27]. A hypothesis is that UbiD and UbiX, which share no sequence similarity, function together during the decarboxylation of OHB (Reaction c, Fig. 1) [27]. Moreover, it might be worth noting that in *Salmonella enterica* serovar Paratyphi, *ubiX* and *ubiD* are organized as a single fusion gene *ubiX-ubiD* (SPA2778 gene). In *E. coli* O157:H7, in addition to UbiD and UbiX, a probable aromatic acid decarboxylase called Pad1 was identified but its role in  $Q_8$  biosynthesis has not been investigated so far.

Hydroxylations represent three of the nine reactions required for  $Q_8$  biosynthesis in *E. coli*. These reactions introduce hydroxyl groups at positions C-5, C-1, and C-6 of the aromatic ring of  $Q_8$  (Reactions d, f and h, Fig. 1), and the genes encoding these enzymes were proposed to be *ubiB*, *ubiH*, and *ubiF*, respectively (Fig. 2A and B) [28]. *ubiH* was the first of these genes to be identified and, based on sequence comparison, was proposed to be a flavin-containing monooxygenase [29]. A *ubiF* mutant was shown to accumulate demethoxy coenzyme  $Q_8$



**Fig. 2.** Genetic organization and location of *ubi* genes on the *Escherichia coli* chromosome. A. Location of the eleven *ubi* genes on the 100 min map of the *E. coli* chromosome. B. Genetic organization of the *ubi* genes in *E. coli*. The location of each gene (or the first gene of the operon) is indicated on the left. The genes required for ubiquinone biosynthesis are indicated in bold. Genes encoding monooxygenases are symbolized by black arrows and genes encoding methyltransferases are symbolized by gray arrows.

**Table 1**  
Accumulated intermediates from the Q<sub>8</sub> biosynthetic pathway in *Escherichia coli* mutants.

Gene affected by the mutation <sup>a</sup>	Deficient reaction	Accumulated Q <sub>8</sub> -intermediates <sup>b</sup>	Reference
<i>ubiB</i>	Not known	OPP	[31]
<i>ubiC</i>	a	None <sup>c</sup>	[73]
<i>ubiD</i>	c	OHB <sup>c</sup>	[27]
<i>ubiX</i>	c	OHB <sup>c</sup>	[27]
<i>ubil</i>	d	4-HP <sub>8</sub> <sup>c</sup>	[32]
<i>ubiG</i>	e and i	OPP and compound 3 (Fig. 1)	[74]
<i>ubiE</i>	g	DDMQ <sub>8</sub>	[24]
<i>ubiF</i>	h	DDMQ <sub>8</sub>	[30]
<i>ubiA</i>	b	No prenylated products formed	[19]
<i>ubij</i>	Not known	No intermediate characterized	[34]
<i>ubiH</i>	f	OPP, compounds 1 and 2 (Fig. 1)	[75]

<sup>a</sup> The mutations that affect the *ubi* genes reported in this table vary from point mutation to transposon insertion or complete deletion. See references for details.

<sup>b</sup> As detected by the techniques used in the reported studies. For example, OPP is adequately detected by using radiolabeled 4-HB but is not evidenced when using electrochemical detection coupled to HPLC.

<sup>c</sup> A low level of Q<sub>8</sub> is observed in these strains.

(DDMQ<sub>8</sub>) (Table 1, Fig. 1) indicating that UbiF, also related to flavin-containing monooxygenases, was responsible for introducing a hydroxyl group at C-6 of the aromatic ring [30]. UbiB was initially proposed to be involved in the C-5 hydroxylation because a *ubiB* mutant was shown to accumulate OPP and failed to produce Q<sub>8</sub> (Table 1) [25,31]. However, the UbiB protein does not contain any signature sequence for monooxygenases but rather shares identity with a large family of eukaryotic-type protein kinases [31]. Actually, we discovered that the C-5 hydroxylase was the product of the *visC* gene [32]. Accordingly, the *visC* gene, locating immediately downstream of *ubiH* within the same operon, was renamed *ubil* (Fig. 2B). A *ubil* mutant has a low level of Q<sub>8</sub> and accumulates 3-octaprenyl-4-hydroxyphenol (4-HP<sub>8</sub>), an intermediate that results from a C5-hydroxylation defect (Table 1) [32]. In fact, 4-HP<sub>8</sub> is formed by hydroxylation of OPP on C1 by UbiH without reaction d and e taking place (Fig. 1). Therefore, the UbiH protein is able to perform the C1-hydroxylation in the absence of the methoxyl group on C5. The UbiI protein displays the typical domains of flavin-containing monooxygenases and shares 30 and 39% sequence identity with UbiH and UbiF, respectively [32]. Thus, we proposed to assign to UbiI the first hydroxylation step in Q<sub>8</sub> biosynthesis (Reaction d, Fig. 1) [32]. The role of the *ubiB* gene product remains to be investigated.

In *E. coli*, the *yigP* gene (called *ubij* in Fig. 2B) lies in between the *ubiE* and *ubiB* genes [31]. Recently, *yigP* was proposed to encode a small RNA of 252 nucleotides (referred to as *esrE*), which was proposed to be essential in *E. coli* [33]. We also characterized the *yigP* gene in *E. coli* and *Salmonella* and our results did not support these conclusions as *yigP* deletions were obtained in both bacteria. We found *yigP* to be required for Q<sub>8</sub> biosynthesis and we changed the name of *yigP* into *ubij* [34]. Regarding the small RNA issue, a “scrambled” *ubij* allele including the mutation of 30% of the nucleotides without changing the amino acid sequence restored Q<sub>8</sub> biosynthesis in *Salmonella ubij* mutant [34]. We therefore believe that the biological function of *ubij* is mediated by a protein. We also demonstrated that a *ubij* mutant was impaired for growth under aerobic conditions, but did not present any growth defect anaerobically, either with glucose or glycerol supplemented with different electron acceptors [34].

In conclusion, all enzymatic reactions necessary for Q<sub>8</sub> biosynthesis in aerobic conditions have been assigned to a Ubi protein (Fig. 1). However, the role of UbiB and UbiJ proteins remains to be established, as they are also essential for this process.


### 3. Structural analysis

The *E. coli* chorismate lyase UbiC structure was solved at 1.4 Å resolution [35,36]. This monomeric enzyme is composed of 164 amino acid

residues with a molecular weight of 19 kDa [19]. The fold involves a 6-stranded antiparallel β-sheet with no spanning helices and novel connectivity [36]. Moreover, the 4-HB is completely sequestered from solvent in a largely hydrophobic environment behind two helix–turn–helix loops [36]. Ultrahigh resolution (1.0 Å) crystal structure of the *E. coli* UbiC product complex (PDB ID: 1TT8) reveals details of a substrate-sized internal cavity, also behind the flaps, near the product site (Fig. 3) [37]. A common feature of the UbiC structures is the presence of two internal-binding pockets connected by a short tunnel [37].


Structural information about the UbiX decarboxylase are available from studies carried out in *Pseudomonas aeruginosa* and *E. coli* O157:H7. The *P. aeruginosa* UbiX structure was determined to 1.5 Å resolution [38]. It shows that the enzyme assembles into a dodecamer, each subunit displaying a typical Rossmann fold, and contains one Flavin MonoNucleotide (FMN) at the interface between two subunits (PDB ID: 3ZQU) [38]. A paralog of UbiX named Pad1 (52% identity) was identified in *E. coli* O157:H7. Its three-dimensional structure has been determined and refined at 2.0 Å resolution (PDB ID: 1SBZ) (Fig. 4A) [39]. Each Pad1 monomer of the dodecameric assembly consists of a typical Rossmann fold and contains a non-covalently bound molecule of FMN (Fig. 4A) [39]. The FMN cofactor also lies at the interface between two monomers. As expected, the structures of UbiX from *P. aeruginosa* and Pad1 from *E. coli* O157:H7 are highly comparable as shown by the root-mean-square deviation of aligned Cα atoms in the range 1.1–1.5 Å (Fig. 4B).

Structural information for the UbiD decarboxylase derived from studies carried out in *P. aeruginosa* and *E. coli*. Two genes, PA0254 and PA5237, predicted to encode UbiD homologs arise in *P. aeruginosa* [40]. The three-dimensional structure of PA0254 has been determined in two different crystal forms to resolutions of 1.95 and 2.3 Å, respectively, showing a dimeric assembly (PDB ID: 4IP2) (Fig. 5A). Strikingly, the quaternary structure of *P. aeruginosa* PA0254 (25% identity with *E. coli* UbiD) differs from the hexameric organization of UbiD from *E. coli* (PDB ID: 2IDB, unpublished, Fig. 5B) and of PA5237 (76% identity with *E. coli* UbiD) [40]. Each subunit of PA0254, whose overall fold is



**Fig. 3.** Structure of the chorismate lyase UbiC from *Escherichia coli*. Crystal structure of the chorismate lyase UbiC from *E. coli* in complex with the 4-hydroxybenzoate, 1.0 Å resolution (PDB ID: 1TT8) [37]. The core of the protein fold is a 6-stranded antiparallel β-sheet, with two helix–turn–helix loops colored in yellow and orange. The two internal cavities are colored in gray. The 4-HB product is located in one of these cavities, behind the two flaps in a hydrophobic pocket.





**Fig. 4.** Structures of the UbiX decarboxylase from *Pseudomonas aeruginosa* and the Pad1 paralogue from *Escherichia coli* O157:H7. A. Crystal structure of the dodecameric Pad1 from *E. coli* O157:H7 complexed with a flavin mononucleotide (FMN), 2.0 Å resolution (PDB ID: 1SBZ) [39]. The view along the threefold axis shows the trimeric arrangement of Pad1. Each subunit is shown in a different color. The FMN cofactor is shown as yellow sticks and its binding site is located at the interface between two monomers. B. Superimposition of one subunit of the dodecameric Pad1 from *E. coli* O157:H7 (in green) with the corresponding subunit of the dodecameric UbiX from *Pseudomonas aeruginosa* (PDB ID 3ZQU) (in red). The flavin cofactors FMN of the two proteins are colored in green in UbiX and in red in Pad1.

similar to that of UbiD from *E. coli*, consists of three domains. The N-terminal part of the molecule is built up of two domains that pack tightly to each other and a C-terminal  $\alpha/\beta$  domain which displays a topology characteristic for the UbiD protein family [40]. The middle domain shows significant structural similarity to the FMN-binding split barrel from a family of flavoproteins, including the NADH: FMN oxidoreductase from *Methylobacillus flagelates* (PDB ID: 3E4V), the flavin reductase from *Shewanella baltica* (PDB ID: 3HMZ), or the flavoredoxin from *Desulfovibrio vulgaris* (PDB ID: 2D5M). This middle domain also contains a metal binding site, with a magnesium ion coordinated by two histidine and glutamate residues, which are conserved in the corresponding metal binding site of some FMN binding proteins. However, no evidence for the incorporation of a flavin cofactor in PA0254 is available so far.


Recently, we determined the crystal structure of a truncated form of UbiI, involved in the aerobic C5-hydroxylation reaction (Fig. 6A) [32]. Only this form, lacking the 35C-terminal residues, and not the full-length protein, too prone to precipitate, could crystallize. Nevertheless, this structure provides some useful information, in particular as it shares many features with that of the canonical FAD-containing parahydroxybenzoate hydroxylase (PHBH) [41]. In particular, comparison with PHBH reveals a FAD binding site in UbiI (Fig. 6B) which has been validated by site directed mutagenesis, thus supporting that UbiI and probably UbiH and UbiF are indeed FAD-containing monooxygenases [32].

As shown from this brief survey, it is obvious that a lot is still missing regarding the structural biology of  $Q_8$  pathway: few proteins have been structurally characterized and rather incomplete information has been gained from those which have been crystallized. This opens a large field of research for the future.


#### 4. Physiological role of $Q_8$

##### 4.1. Interplay between $Q_8$ and redox sensing two-component systems

The Arc (anoxic redox control) two-component system of *E. coli* comprises the ArcB transmembrane sensor kinase and the cytosolic ArcA response regulator. It plays a major role in a transcriptional regulatory network that allows facultative anaerobic bacteria to sense various respiratory growth conditions [42]. Georgellis and colleagues showed that quinones act as direct negative signals that inhibit



**Fig. 5.** Structures of the UbiD decarboxylases from *Pseudomonas aeruginosa* and *Escherichia coli*. A. Crystal structure of the dimeric PA0254 from *P. aeruginosa*, a putative aromatic acid decarboxylase bearing a C-terminal domain characteristic of the UbiD protein family, 1.95 Å resolution (PDB ID: 4IP2) [40]. The three domains of each subunit are shown in blue (N-terminal domain), green (middle domain) and red (C-terminal domain). The magnesium site is indicated by a black sphere in the middle domain. B. Crystal structure of the hexameric UbiD from *E. coli* (PDB ID: 2IDB). Each subunit is shown in a different color.



**Fig. 6.** Structure of the Ubil monooxygenase from *Escherichia coli*. A. Crystal structure of the C-terminal truncated form of *E. coli* C5-hydroxylase Ubil involved in  $Q_8$  biosynthesis, 2.0 Å resolution (PDB ID: 4 K22) [32]. B. Proposed molecular docking model of Ubil with FAD. The flavin cofactor is shown as yellow sticks.

autophosphorylation of ArcB during aerobiosis [43]. They demonstrated that the oxidative power provided by the  $Q_8$  pool induced the formation of two intermolecular cysteine disulfide bonds within the cytosolic domains of an ArcB dimer, leading to the inactivation of ArcB kinase activity [43,44]. Bekker and colleagues provided evidence that additional regulation of ArcB kinase activity by the redox state of menaquinone ( $MK_8$ ) was prevalent under microaerobic to anaerobic conditions [45]. They concluded that activation of the ArcBA system was controlled by both  $MK_8$  and  $Q_8$  pools [45]. Moreover, the extent of ArcA phosphorylation was shown to be modulated by the oxygen supply rate in an *ubiE* mutant, leading to the conclusion that the oxidized form of  $DMK_8$  could inactivate ArcB [46]. However, the authors did not take in consideration the  $DDMK_8$  which also accumulates in a *ubiE* mutant [24,34] and could play a role in ArcB regulation in the *ubiE* mutant. A complementary study demonstrated that (D) $MK_8$  was required for activation of ArcB upon a shift to anoxic conditions [4]. The midpoint redox potential of the cysteines of ArcB was determined to be approximately  $-41$  mV, which is in agreement with the proposed model in which the  $Q_8$  pool can oxidize these cysteines under aerobic conditions and the disulfide bonds can be reduced by the (D) $MK_8$  pool upon a shift to anaerobic conditions [4].

Another example of redox regulation is illustrated by the RegBA two-component system of *Rhodobacter capsulatus*. RegB is a membrane-spanning sensor kinase that autophosphorylates and transfers the phosphoryl group to its cognate response regulator RegA [47]. The RegB/RegA system regulates the synthesis of numerous energy-generating and energy-utilizing systems [48]. Recently, RegB was shown to bind weakly both oxidized and reduced  $Q_8$  with nearly equal affinity, although oxidized  $Q_8$  alone inhibited kinase activity [49]. The current view is that interaction of RegB with both the reduced and oxidized forms of  $Q_8$  allows the kinase to monitor and tune the cellular energy state [49].

In conclusion, in both *E. coli* and *R. capsulatus* models, and presumably many other bacteria, a connection between the pool, and the redox state, of  $Q_8$  and two-component systems endows the cell with the capacity to modulate the expression of multiple genes in response to changes in redox conditions.

#### 4.2. $Q_8$ and oxidative stress

The involvement of ubiquinone in oxidative stress resistance was first addressed in eukaryotes, where it was suggested that its reduced

form, i.e. ubiquinol, was able to function as a lipid-soluble anti-oxidant [50]. Ubiquinol was shown to scavenge lipid peroxyl radicals and thereby prevents a chain reaction causing oxidative damage to polyunsaturated fatty acids of biological membranes, a process known as lipid peroxidation [51]. In yeast, the biosynthetic intermediate  $DMQ_6$  was concluded to lack antioxidant activity because it failed to protect cells against oxidative stress generated by hydrogen peroxide ( $H_2O_2$ ) or linolenic acid [52]. In prokaryotes, an exhaustive study aimed at investigating the role of  $Q_8$  in resistance against oxidative stress was carried out by Søballe and Poole [53]. Using a *ubiCA* mutant deficient for  $Q_8$  production, they observed an accumulation of superoxide ( $O_2^{\bullet-}$ ) in the membranes of the mutant compared to wild-type membranes, reflecting the lack of superoxide-scavenging ubiquinol [53]. Moreover, intracellular  $H_2O_2$  concentration was increased 1.8-fold in the *ubiCA* mutant, which was also found to be hypersensitive to an oxidative stress mediated by  $H_2O_2$  [53]. Expression of *katG* gene, encoding a catalase, and intracellular catalase activity, were both increased in the *ubiCA* mutant [53]. These observations are consistent with the hypothesis that  $Q_8$  limits  $O_2^{\bullet-}$  and  $H_2O_2$  accumulation in scavenging reactive oxygen species. However, given the multiplicity of catalases, peroxidases and superoxide dismutases present in most bacteria, the relative contribution of  $Q_8$  to the overall anti-oxidative stress defenses system remains to be assessed.

#### 4.3. $Q_8$ and respiration

The respiratory chain of *E. coli* contains many enzymes allowing the organism to transfer electrons to oxygen or to use alternative terminal acceptors such as nitrate, dimethyl sulfoxide (DMSO), trimethylamine N-oxide (TMAO) or fumarate [1,2,7].  $Q_8$  is generally depicted as the aerobic quinone since it is more abundant than  $MK_8$  during aerobic growth, whereas  $MK_8$  is essential for anaerobic respiration using fumarate, DMSO and TMAO as electron acceptors [13,14]. In this way, a *ubiCA* mutant showed severely diminished growth yields aerobically but not anaerobically using nitrate or fumarate as terminal electron acceptors [54]. Conversely, a *ubiE* mutant, which contains  $DMK_8$  but no  $MK_8$ , was able to grow with fumarate, TMAO and DMSO, but not with nitrate as electron acceptor [55]. It was then concluded that  $DMK_8$  (in addition to  $MK_8$ ) could serve as a redox mediator in fumarate, TMAO and DMSO respiration, but not in nitrate respiration [55]. Moreover, it was



demonstrated that nitrate reductase A could accept electrons from both Q<sub>8</sub> and MK<sub>8</sub> pools, the coupling being more effective with ubiquinol than with menaquinol [56].

Recently, Bekker and colleagues reported that DMK<sub>8</sub> played a role not only in TMAO –, DMSO- and fumarate-dependent respiration, but also in oxidation of succinate [15]. They further concluded that all three quinol oxidases of *E. coli* accepted electrons from DMK<sub>8</sub> based on the residual aerobic respiration observed in a *ubiE* mutant [15]. As already mentioned, another hypothesis is that DDMQ<sub>8</sub> might provide electron towards the quinol oxidases in the *ubiE* mutant. In the future, it will be interesting to know whether these remaining uncertainties will challenge the classic view associating ubiquinone to aerobic growth and menaquinone to anaerobic growth.

#### 4.4. Q<sub>8</sub> and antibiotics resistance

The proton-pumping NADH:ubiquinone oxidoreductase, also called complex I, catalyzes the electron transfer from NADH to Q<sub>8</sub> linked with a proton translocation from the negative inner to the positive outer side of the membrane [57]. Thus, a proton-motive (PMF) force is generated, which is utilized mainly for ATP synthesis. PMF is instrumental in allowing either import or export of antibiotics [58–60]. As a consequence, *ubi* mutations are likely to modify levels of resistance to antibiotics sustained by bacteria. In fact, resistance to low levels of aminoglycoside antibiotics has been used for long for the isolation of menaquinone-deficient mutants in *Bacillus subtilis* since the MK deficiency depresses the rate of accumulation of aminoglycosides, which accounts for the resistance phenotype [61]. Thereby, a phenotype of increased aminoglycoside resistance led to the identification of the *aarE* gene, the *ubiA* homologue in *Providencia stuartii* [62]. In *E. coli*, a *ubiF* mutation was associated with pleomycin resistance [63] while a *ubiD* mutation was found to be associated with decreased transport of streptomycin and gentamicin, and increased resistance to those antibiotics [64]. This mutant also exhibited increased resistance to several other aminoglycoside antibiotics, but not spectinomycin [64]. A series of *E. coli* mutants located in the vicinity of *ubiB* and *ubiD* and consistent with a disruption of Q<sub>8</sub> synthesis were found to present zwittermicin A resistance, a novel broad-spectrum antibiotics produced by *Bacillus cereus* [65]. Rationally, a *ubiD* mutant was also resistant to zwittermicin A [65]. All together, these studies highlight that connections between Q<sub>8</sub> production and antibiotics resistance occur in many bacteria.

#### 4.5. Q<sub>8</sub> and bacterial virulence

In the *Caenorhabditis elegans* model, Clarke and colleagues revealed that nematodes fed diets of respiratory deficient *E. coli* lacking Q<sub>8</sub> lived significantly longer than worms fed the wild-type parental strain [66]. Moreover, these *E. coli* mutants were degraded in the early adult worm and did not accumulate in the intestinal tract [66]. These data led the authors to propose that bacterial respiration might act as a virulence factor, influencing the ability of bacteria to colonize – and subsequently harm – the animal host [66]. As a matter of fact, the role of respiration in host–pathogen interactions has recently been underlined in several studies, including those dedicated to *Shigella* pathogenicity [67] and more recently to *Salmonella* [34].

Indeed, early studies brought in light the requirement of a functional Q<sub>8</sub> biosynthesis pathway for the flagellation of *Salmonella* [68]. Interestingly, the increase in motility response occurred within a narrow range of the increase in Q<sub>8</sub> content [68]. More recently, we showed that *Salmonella ubiJ* mutant, a strain deficient for Q<sub>8</sub> production under aerobic conditions, was killed within macrophages [34]. When macrophages were infected with an anaerobic inoculum of the *ubiJ* mutant (in which Q<sub>8</sub> was still produced), the *ubiJ* mutant recovered its proliferation capacity, clearly establishing the requirement for Q<sub>8</sub> for efficient intracellular proliferation [34]. Several possibilities can be considered to connect Q<sub>8</sub> defects and virulence attenuation: (i) the necessity for the bacterium

to use aerobic respiration – thus Q<sub>8</sub> – to grow intracellularly; (ii) the antioxidant role of Q<sub>8</sub> to avoid the oxidative burst produced by the host; (iii) the requirement of a PMF-dependent process for virulence. Thereby, our recent data combined to results from other groups highlighted the importance of Q<sub>8</sub> biosynthesis in the context of the host–pathogen interplay and revealed an unexpected link between Q<sub>8</sub> and bacterial virulence.

### 5. Conclusion and perspectives

In this review, we have presented the actors involved in Q<sub>8</sub> biosynthetic pathway. Phenotypic characterizations of various mutants coupled with the examination of Q<sub>8</sub> content led to the identification of genes involved in the decoration of the aromatic ring, from the 4-HB precursor to the final ubiquinone. When available, the three-dimensional structures of these enzymes were presented. Last, the role of Q<sub>8</sub> in diverse cellular processes was detailed: genetic regulation through the two-component systems ArcBA and RegBA, adaptation to oxidative stress, respiration, antibiotics resistance and bacterial virulence.

Whereas nine *ubi* genes were found to be directly involved in Q<sub>8</sub> biosynthesis, the biological function of *ubiB* and *ubiJ* gene products remain unknown. The involvement of UbiB in the C-5 hydroxylation step has to be definitively abandoned. Instead, UbiB might play a regulatory role through a kinase activity, even though it is still unknown if it displays such an activity and what substrate it acts on. Based on sequence similarities between *ubiJ* and genes involved in lipid metabolism, UbiJ could serve as a carrier of the isoprenoid hydrophobic tail prior the action of monooxygenases and methyltransferases. Alternatively, it might chaperone prenylated intermediates during the biosynthetic process.

Intriguingly, we found that the *ubiJ* mutant synthesizes almost wild-type levels of Q<sub>8</sub> under anaerobic growth conditions [34]. Actually, *ubiH*, *ubiF* and *ubiL* mutants also synthesize significant levels of Q<sub>8</sub> anaerobically [28,32]. The three latter genes encode flavin-dependent monooxygenases, which use molecular oxygen to catalyze their respective hydroxylation reactions. In consequence, these proteins will not participate in Q<sub>8</sub> biosynthesis under anaerobic conditions because of the absence of molecular oxygen. Therefore, alternative anaerobic hydroxylases must carry out the hydroxylation reactions in anaerobic conditions as proposed by Alexander and Young [28], but so far the identity of these proteins has remained elusive. The involvement of UbiJ only in aerobic Q<sub>8</sub> biosynthesis is more difficult to rationalize than that of the aerobic monooxygenases and may reflect a functional link with those enzymes.

It is important to note that with the exception of the *E. coli* methyltransferase UbiG [23], the partially purified membrane-bound UbiD [26] and the UbiA octaprenyltransferase [69], none of the other enzymes (the decarboxylases, the putative kinase and the hydroxylases) that participate to Q<sub>8</sub> biosynthesis have been assayed under in vitro conditions using pure proteins. As a consequence, most of the predicted activities associated with Ubi enzymes remain to be fully established by straight biochemical analyses. This is not surprising considering the difficulties to purify these proteins, their probable association to the membranes and their instability, as well as the difficulties to get access to the substrates, not commercially available.

Another issue is whether Ubi proteins may belong to a large multiprotein complex as suggested by an early report [70]. By a combination of sonication, gel filtration and equilibrium sedimentation with a sucrose gradient, the author purified a mega complex of at least 12 proteins, ranging from 40 to 80 kDa and exhibiting a molecular weight estimated to 2.10<sup>6</sup> Da. This complex had the capacity to process OPP into Q<sub>8</sub> in the presence of NADPH, SAM and O<sub>2</sub>. It is intriguing that, in spite of this fascinating observation, no further studies concerning this bacterial complex have been reported then. The existence of a high-molecular weight complex has also been documented in *Saccharomyces cerevisiae*, mostly by genetic and partly by biochemical studies [71,72]. The complex may

actually consist of several subcomplexes likely associated with the mitochondrial inner membrane [72], but neither its complete composition nor its structural organization is known. In particular, the purification of this complex has not been reported. Thus, the question of the existence of a Q<sub>8</sub> biosynthetic complex in bacteria remains open and needs further investigation.

## Acknowledgements

Thanks are due to the members of the FB group for fruitful discussions. We acknowledge Ludovic Pecqueur for the help in the preparation of molecular graphics figures with PyMol. This work was funded by the CNRS, Aix-Marseille Université (AMU), the Institut Universitaire de France (IUF), and the French State Program “Investissements d’Avenir” (Grant “LABEX DYNAMO”, ANR-11-LABX-0011).

## References

- [1] G. Uden, J. Bongaerts, Alternative respiratory pathways of *Escherichia coli*: energetics and transcriptional regulation in response to electron acceptors, *Biochim. Biophys. Acta* 1320 (1997) 217–234.
- [2] R.K. Poole, G.M. Cook, Redundancy of aerobic respiratory chains in bacteria? Routes, reasons and regulation, *Adv. Microb. Physiol.* 43 (2000) 165–224.
- [3] I. Belevich, V.B. Borisov, M.I. Verkhovsky, Discovery of the true peroxy intermediate in the catalytic cycle of terminal oxidases by real-time measurement, *J. Biol. Chem.* 282 (2007) 28514–28519.
- [4] A.F. Alvarez, C. Rodriguez, D. Georgellis, Ubiquinone and menaquinone electron carriers represent the yin and yang in the redox regulation of the ArcB sensor kinase, *J. Bacteriol.* 195 (2013) 3054–3061.
- [5] B. Schoep-Cothenet, C. Lieutaud, F. Baymann, A. Vermeiglio, T. Friedrich, D.M. Kramer, W. Nitschke, Menaquinone as pool quinone in a purple bacterium, *Proc. Nat. Acad. Sci. U. S. A.* 106 (2009) 8549–8554.
- [6] A. Nakamura, T. Suzawa, Y. Kato, T. Watanabe, Species dependence of the redox potential of the primary electron donor p700 in photosystem I of oxygenic photosynthetic organisms revealed by spectroelectrochemistry, *Plant Cell Physiol.* 52 (2011) 815–823.
- [7] V.B. Borisov, R.B. Gennis, J. Hemp, M.I. Verkhovsky, The cytochrome bd respiratory oxygen reductases, *Biochim. Biophys. Acta* 1807 (2011) 1398–1413.
- [8] B. Nowicka, J. Kruk, Occurrence, biosynthesis and function of isoprenoid quinones, *Biochim. Biophys. Acta* 1797 (2010) 1587–1605.
- [9] M. Bentinger, M. Tekle, G. Dallner, Coenzyme Q-biosynthesis and functions, *Biochem. Biophys. Res. Commun.* 396 (2010) 74–79.
- [10] M. Turunen, J. Olsson, G. Dallner, Metabolism and function of coenzyme Q, *Biochim. Biophys. Acta* 1660 (2004) 171–199.
- [11] R. Meganathan, Ubiquinone biosynthesis in microorganisms, *FEMS Microbiol. Lett.* 203 (2001) 131–139.
- [12] U.C. Tran, C.F. Clarke, Endogenous synthesis of coenzyme Q in eukaryotes, *Mitochondrion* 7 (2007) S62–S71.
- [13] B.J. Wallace, I.G. Young, Role of quinones in electron transport to oxygen and nitrate in *Escherichia coli*. Studies with a ubiA- menA- double quinone mutant, *Biochim. Biophys. Acta* 461 (1977) 84–100.
- [14] B. Søballe, R.K. Poole, Microbial ubiquinones: multiple roles in respiration, gene regulation and oxidative stress management, *Microbiology* 145 (1999) 1817–1830.
- [15] P. Sharma, M.J. Teixeira de Mattos, K.J. Hellingwerf, M. Bekker, On the function of the various quinone species in *Escherichia coli*, *FEBS J.* 279 (2012) 3364–3373.
- [16] S. Grimaldi, B. Schoep-Cothenet, P. Ceccaldi, B. Guigliarelli, A. Magalon, The prokaryotic Mo/W-bisPGD enzymes family: a catalytic workhorse in bioenergetic, *Biochim. Biophys. Acta* 1827 (2013) 1048–1085.
- [17] G.B. Cox, F. Gibson, Biosynthesis of vitamin K and ubiquinone. Relation to the shikimic acid pathway in *Escherichia coli*, *Biochim. Biophys. Acta* 93 (1964) 204–206.
- [18] G.B. Cox, F. Gibson, J. Pittard, Mutant strains of *Escherichia coli* K-12 unable to form ubiquinone, *J. Bacteriol.* 95 (1968) 1591–1598.
- [19] G. Wu, H.D. Williams, F. Gibson, R.K. Poole, Mutants of *Escherichia coli* affected in respiration: the cloning and nucleotide sequence of ubiA, encoding the membrane-bound p-hydroxybenzoate:octaprenyltransferase, *J. Gen. Microbiol.* 139 (1993) 1795–1805.
- [20] B.P. Nichols, J.M. Green, Cloning and sequencing of *Escherichia coli* ubiC and purification of chorismate lyase, *J. Bacteriol.* 174 (1992) 5309–5316.
- [21] L. Zhou, J.Y. Wang, J.H. Wang, A. Poplawsky, S.J. Lin, B.S. Zhu, C.Q. Chang, T.L. Zhou, L.H. Zhang, Y.W. He, Mol. Microbiol. 87 (2013) 80–93.
- [22] G. Wu, H.D. Williams, M. Zamanian, F. Gibson, R.K. Poole, Isolation and characterization of *Escherichia coli* mutants affected in aerobic respiration: the cloning and nucleotide sequence of ubiG. Identification of an S-adenosylmethionine-binding motif in protein, RNA, and small-molecule methyltransferases, *J. Gen. Microbiol.* 138 (1992) 2101–2112.
- [23] W.W. Poon, R.J. Barkovich, A.Y. Hsu, A. Frankel, P.T. Lee, J.N. Shepherd, D.C. Myles, C.F. Clarke, Yeast and rat Coq3 and *Escherichia coli* UbiG polypeptides catalyze both O-methyltransferase steps in coenzyme Q biosynthesis, *J. Biol. Chem.* 274 (1999) 21665–21672.
- [24] P.T. Lee, A.Y. Hsu, H.T. Ha, C.F. Clarke, A C-methyltransferase involved in both ubiquinone and menaquinone biosynthesis: isolation and identification of the *Escherichia coli* ubiE gene, *J. Bacteriol.* 179 (1997) 1748–1754.
- [25] G.B. Cox, I.G. Young, L.M. McCann, F. Gibson, Biosynthesis of ubiquinone in *Escherichia coli* K-12: location of genes affecting the metabolism of 3-octaprenyl-4-hydroxybenzoic acid and 2-octaprenylphenol, *J. Bacteriol.* 99 (1969) 450–458.
- [26] R.A. Leppik, I.G. Young, F. Gibson, Membrane-associated reactions in ubiquinone biosynthesis in *Escherichia coli*. 3-Octaprenyl-4-hydroxybenzoate carboxy-lyase, *Biochim. Biophys. Acta* 436 (1976) 800–810.
- [27] M. Gulmezian, K.R. Hyman, B.N. Marbois, C.F. Clarke, G.T. Javor, The role of UbiX in *Escherichia coli* coenzyme Q biosynthesis, *Arch. Biochem. Biophys.* 467 (2007) 144–153.
- [28] K. Alexander, I.G. Young, Alternative hydroxylases for the aerobic and anaerobic biosynthesis of ubiquinone in *Escherichia coli*, *Biochemistry* 17 (1978) 4750–4755.
- [29] K. Nakahigashi, K. Miyamoto, K. Nishimura, H. Inokuchi, Isolation and characterization of a light-sensitive mutant of *Escherichia coli* K-12 with a mutation in a gene that is required for the biosynthesis of ubiquinone, *J. Bacteriol.* 174 (1992) 7352–7359.
- [30] O. Kwon, A. Kotsakis, R. Meganathan, Ubiquinone (coenzyme Q) biosynthesis in *Escherichia coli*: identification of the ubiF gene, *FEMS Microbiol. Lett.* 186 (2000) 157–161.
- [31] W.W. Poon, D.E. Davis, H.T. Ha, T. Jonassen, P.N. Rather, C.F. Clarke, Identification of *Escherichia coli* ubiB, a gene required for the first monooxygenase step in ubiquinone biosynthesis, *J. Bacteriol.* 182 (2000) 5139–5146.
- [32] M. Hajj Chehade, L. Loiseau, M. Lombard, L. Pecqueur, A. Ismail, M. Smadja, B. Golinelli-Pimpianeau, C. Mellot-Drazniak, O. Hamelin, L. Aussel, S. Kieffer-Jaquinet, N. Labessan, F. Barras, M. Fontecave, F. Pierrel, A new gene involved in coenzyme Q biosynthesis in *Escherichia coli*: UbiI functions in aerobic C5-hydroxylation, *J. Biol. Chem.* 288 (2013) 20085–20092.
- [33] Z. Chen, Y. Wang, Y. Li, Y. Li, N. Fu, J. Ye, H. Zhang, Esre: a novel essential non-coding RNA in *Escherichia coli*, *FEBS Lett.* 586 (2012) 1195–1200.
- [34] L. Aussel, L. Loiseau, M. Hajj Chehade, B. Pocachard, M. Fontecave, F. Pierrel, F. Barras, Coenzyme Q biosynthesis in *Escherichia coli* and *Salmonella typhimurium*: ubiJ, a new gene required for aerobic growth and proliferation in macrophage, *J. Bacteriol.* 196 (2014) 70–79.
- [35] C. Stover, M.P. Mayhew, M.J. Holden, A. Howard, D.T. Gallagher, Crystallization and 1.1-Å diffraction of chorismate lyase from *Escherichia coli*, *J. Struct. Biol.* 129 (2000) 96–99.
- [36] D.T. Gallagher, M. Mayhew, M.J. Holden, A. Howard, K.J. Kim, V.L. Vilker, The crystal structure of chorismate lyase shows a new fold and a tightly retained product, *Proteins* 44 (2001) 304–311.
- [37] N. Smith, A.E. Roitberg, E. Rivera, A. Howard, M.J. Holden, M. Mayhew, S. Kaistha, D.T. Gallagher, Structural analysis of ligand binding and catalysis in chorismate lyase, *Arch. Biochem. Biophys.* 445 (2006) 72–80.
- [38] J. Kopeck, R. Schnell, G. Schneider, Structure of PA4019, a putative aromatic acid decarboxylase from *Pseudomonas aeruginosa*, *Acta Crystallogr. Sect. F: Struct. Biol. Cryst. Commun.* 67 (2011) 1184–1188.
- [39] E.S. Rangarajan, Y. Li, P. Iannuzzi, A. Tocilj, L.W. Hung, A. Matte, M. Cygler, Crystal structure of a dodecameric FMN-dependent UbiX-like decarboxylase (Pad1) from *Escherichia coli* O157: H7, *Protein Sci.* 13 (2004) 3006–3016.
- [40] A. Jaciewicz, A. Izumi, K. Brunner, R. Schnell, G. Schneider, Structural insights into the UbiD protein family from the crystal structure of PA0254 from *Pseudomonas aeruginosa*, *PLoS One* 8 (2013) e63161.
- [41] H.A. Schreuder, P.A. Prick, R.K. Wierenga, G. Vriend, K.S. Wilson, W.G. Hol, J. Drenth, Crystal structure of the p-hydroxybenzoate hydroxylase-substrate complex refined at 1.9 Å resolution. Analysis of the enzyme-substrate and enzyme-product complexes, *J. Mol. Biol.* 208 (1989) 679–696.
- [42] A.S. Lynch, E.C. Lin, *Escherichia coli* and *Salmonella*: cellular and molecular biology, in: F.C. Neidhardt, R. Curtis III, J.L. Ingraham, E.C. Lin, K.B. Low, B. Magasanik, W.S. Reznikoff, M. Riley, M. Schaechter, H.E. Umberger (Eds.), *Am. Soc. Microbiol.*, 1996, pp. 1526–1538, (Washington, DC).
- [43] D. Georgellis, O. Kwon, E.C. Lin, Quinones as the redox signal for the arc two-component system of bacteria, *Science* 292 (2001) 2314–2316.
- [44] R. Malpica, B. Franco, C. Rodriguez, O. Kwon, D. Georgellis, Identification of a quinone-sensitive redox switch in the ArcB sensor kinase, *Proc. Nat. Acad. Sci. U. S. A.* 101 (2004) 13318–13323.
- [45] M. Bekker, A. Alexeeva, W. Laan, G. Sawers, J. Teixeira de Mattos, K. Hellingwerf, The ArcBA two-component system of *Escherichia coli* is regulated by the redox state of both the ubiquinone and the menaquinone pool, *J. Bacteriol.* 192 (2010) 746–754.
- [46] P. Sharma, S. Stagg, M. Bekker, K. Bettenbrock, K.J. Hellingwerf, Kinase activity of ArcB from *Escherichia coli* is subject to regulation by both ubiquinone and demethylmenaquinone, *PLoS One* 8 (2013) e75412.
- [47] T.H. Bird, S. Du, C.E. Bauer, Autophosphorylation, phosphotransfer, and DNA-binding properties of the RegB/RegA two-component regulatory system in *Rhodobacter capsulatus*, *J. Biol. Chem.* 274 (1999) 16343–16348.
- [48] S. Elsen, L.R. Swem, D.L. Swem, C.E. Bauer, RegB/RegA, a highly conserved redox-responding global two-component regulatory system, *Microbiol. Mol. Biol. Rev.* 68 (2004) 263–279.
- [49] J. Wu, C.E. Bauer, RegB kinase activity is controlled in part by monitoring the ratio of oxidized to reduced ubiquinones in the ubiquinone pool, *MBio* 1 (2010) e00272–10.
- [50] L. Ernster, G. Dallner, Biochemical, physiological and medical aspects of ubiquinone function, *Biochim. Biophys. Acta* 1271 (1995) 195–204.
- [51] P. Forsmark-Andree, G. Dallner, L. Ernster, Endogenous ubiquinol prevents protein modification accompanying lipid peroxidation in beef heart submitochondrial particles, *Free Radic. Biol. Med.* 19 (1995) 749–757.
- [52] S. Padilla, T. Jonassen, M.A. Jiménez-Hidalgo, D.J. Fernández-Ayala, G. López-Lluch, B. Marbois, P. Navas, C.F. Clarke, C. Santos-Ocaña, Demethoxy-Q, an intermediate of

- coenzyme Q biosynthesis, fails to support respiration in *Saccharomyces cerevisiae* and lacks antioxidant activity, *J. Biol. Chem.* 279 (2004) 25995–26004.
- [53] B. Søballe, R.K. Poole, Ubiquinone limits oxidative stress in *Escherichia coli*, *Microbiology* 146 (2000) 787–796.
- [54] B. Søballe, R.K. Poole, Requirement for ubiquinone downstream of cytochrome(s) b in the oxygen-terminated respiratory chains of *Escherichia coli* K-12 revealed using a null mutant allele of *ubiCA*, *Microbiology* 144 (1998) 361–373.
- [55] U. Wissenbach, D. Ternes, G. Unden, An *Escherichia coli* mutant containing only demethylmenaquinone, but no menaquinone: effects on fumarate, dimethylsulfoxide, trimethylamine N-oxide and nitrate respiration, *Arch. Microbiol.* 158 (1992) 68–73.
- [56] T.H. Brondijk, D. Fiegen, D.J. Richardson, J.A. Cole, Roles of NapF, NapG and NapH, subunits of the *Escherichia coli* periplasmic nitrate reductase, in ubiquinol oxidation, *Mol. Microbiol.* 44 (2002) 245–255.
- [57] T. Friedrich, K. Steinmüller, H. Weiss, The proton-pumping respiratory complex I of bacteria and mitochondria and its homologue in chloroplasts, *FEBS Lett.* 367 (1995) 107–111.
- [58] H.W. Taber, J.P. Mueller, P.F. Miller, A.S. Arrow, Bacterial uptake of aminoglycoside antibiotics, *Microbiol. Rev.* 51 (1987) 439–457.
- [59] B.D. Davis, L.L. Chen, P.C. Tai, Misread protein creates membrane channels: an essential step in the bactericidal action of aminoglycosides, *Proc. Nat. Acad. Sci. U. S. A.* 83 (1986) 6164–6168.
- [60] B. Ezraty, A. Vergnes, M. Banzhaf, Y. Duverger, A. Huguenot, A.R. Brochado, S.Y. Su, L. Espinosa, L. Loiseau, B. Py, A. Typas, F. Barras, Fe–S cluster biosynthesis controls uptake of aminoglycosides in a ROS-less death pathway, *Science* 340 (2013) 1583–1587.
- [61] H.W. Taber, E.A. Dellers, L.R. Lombardo, Menaquinone biosynthesis in *Bacillus subtilis*: isolation of men mutants and evidence for clustering of men genes, *J. Bacteriol.* 145 (1981) 321–327.
- [62] M.R. Paradise, G. Cook, R.K. Poole, P.N. Rather, Mutations in *aarE*, the *ubiA* homolog of *Providencia stuartii*, result in high-level aminoglycoside resistance and reduced expression of the chromosomal aminoglycoside 2'-N-acetyltransferase, *Antimicrob. Agents Chemother.* 42 (1998) 959–962.
- [63] C.M. Collis, G.W. Grigg, An *Escherichia coli* mutant resistant to phleomycin, bleomycin, and heat inactivation is defective in ubiquinone synthesis, *J. Bacteriol.* 171 (1989) 4792–4798.
- [64] L.E. Bryan, H.M. Van Den Elzen, Effects of membrane-energy mutations and cations on streptomycin and gentamicin accumulation by bacteria: a model for entry of streptomycin and gentamicin in susceptible and resistant bacteria, *Antimicrob. Agents Chemother.* 12 (1977) 163–177.
- [65] E.V. Stabb, J. Handelsman, Genetic analysis of zwittermicin A resistance in *Escherichia coli*: effects on membrane potential and RNA polymerase, *Mol. Microbiol.* 27 (1998) 311–322.
- [66] F. Gomez, G.C. Monsalve, V. Tse, R. Saiki, E. Weng, L. Lee, C. Srinivasan, A.R. Frand, C.F. Clarke, Delayed accumulation of intestinal coliform bacteria enhances life span and stress resistance in *Caenorhabditis elegans* fed respiratory deficient *E. coli*, *BMC Microbiol.* 12 (2012) 300.
- [67] B. Marteyn, N.P. West, D.F. Browning, J.A. Cole, J.G. Shaw, F. Palm, J. Mounier, M.C. Prévost, P. Sansonetti, C.M. Tang, Modulation of *Shigella* virulence in response to available oxygen in vivo, *Nature* 465 (2010) 355–358.
- [68] J. Bar-Tana, B.J. Howlett, R. Hertz, Ubiquinone synthetic pathway in flagellation of *Salmonella typhimurium*, *J. Bacteriol.* 143 (1980) 637–643.
- [69] L.A. Wessjohann, B. Sonntag, Prenylation of benzoic acid derivatives catalyzed by a transferase from *Escherichia coli* overproduction: method development and substrate specificity, *Angew. Chem. Int. Ed. Engl.* 35 (1996) 1697–1699.
- [70] H.E. Knochel, Isolation of a soluble enzyme complex comprising the ubiquinone-8 synthesis apparatus from the cytoplasmic membrane of *Escherichia coli*, *Biochem. Biophys. Res. Commun.* 91 (1979) 919–925.
- [71] P. Gin, C.F. Clarke, Genetic evidence for a multi-subunit complex in coenzyme Q biosynthesis in yeast and the role of the Coq1 hexaprenyl diphosphate synthase, *J. Biol. Chem.* 280 (2005) 2676–2681.
- [72] C.H. He, L.X. Xie, C.M. Allan, U.C. Tran, C.F. Clarke, Coenzyme Q supplementation or over-expression of the yeast Coq8 putative kinase stabilizes multi-subunit Coq polypeptide complexes in yeast coq null mutants, *Biochim. Biophys. Acta* (2013), <http://dx.doi.org/10.1016/j.bbap.2013.12.017>.
- [73] M. Siebert, K. Severin, L. Heide, Formation of 4-hydroxybenzoate in *Escherichia coli*: characterization of the *ubiC* gene and its encoded enzyme chorismate pyruvate-lyase, *Microbiology* 140 (1994) 897–904.
- [74] P. Stroobant, I.G. Young, F. Gibson, Mutants of *Escherichia coli* K-12 blocked in the final reaction of ubiquinone biosynthesis: characterization and genetic analysis, *J. Bacteriol.* 109 (1972) 134–139.
- [75] I.G. Young, P. Stroobant, C.G. Macdonald, F. Gibson, Pathway for ubiquinone biosynthesis in *Escherichia coli* K-12: gene-enzyme relationships and intermediates, *J. Bacteriol.* 114 (1973) 42–52.