SUBJECT INDEX

| Acetylcholinesterase Cholinesterase activities in subcellular fractions of rat liver: Association of — with the surface membrane, and other properties of the enzyme (Wheeler et al.) (255) 917 Acetylcholinesterase Human erythrocyte membrane bound enzyme: — (Heller, Hanahan) (255) 251 Acholeplasma laidlawii cell membranes The effect of cholesterol and epicholesterol incorporation on the permeability and on the phase transition of intact — and derived liposomes (de Kruyff et al.) (255) 331 Active transport | and alkaline phosphatase in the microvilli of isolated brush border of rat kidney (Thomas, Kinne) (255) 114 Anesthetic The action and adsorption of local ———————————————————————————————————— |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Coupling in secondary ——. Activation | (255) 171 |
| of transport by co-transport and/or counter-transport with the fluxes of other solutes (Heinz et al.) (255) 442 Adenylate kinase | Anesthetics The hydrophobic expansion of erythrocyte membranes by the phenol ——— (Machleidt et al.) (255) 178 |
| Erythrocyte membrane-bound enzymes: | Anesthetics |
| ATPase, phosphatase and ——— in human, bovine and porcine erythrocytes (Heller, Hanahan) (255) 239 Adsorption | The membrane concentrations of neutral and positive ———————————————————————————————————— |
| Relation between —— at an oil-water | (Roth, Seeman) (255) 207 |
| interface and membrane permeability (Wang | Antibiotics |
| et al.) (255) 691 | Interaction of —— with membranes: |
| Air-water interface | Chlorothricin (Pache, Chapman) (255) 348 |
| Structural requirements of sterol for the | Antibiotics |
| interaction with lecithin at the | Interaction of ——— with membranes: |
| (Demel et al.) (255) 311 | Polymyxin B and gramicidin S (Pache et al.) |
| Alamethicin | (255) 358 |
| The unit conductance channel of ——— | Aqueous pores |
| (Gordon, Haydon) (255) 1014 | ——— in lipid bilayers and erythrocyte |
| Albumin | membranes (Solomon, Gary-Bobo) (255)1019 |
| Reaction of human serum ——— and | ATP |
| human erythrocytes with tritiated 2,4,6- | The effect of ——— on the CD spectrum of |
| trinitrobenzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Algae | membrane fraction from oxyntic cells (Masotti et al.) (255) 420 ATP |
| General cable theory for cells of —— | Sodium transfer in bullfrog small intestine. |
| Characeae (Volkov) (255) 709 | Stimulation by exogenous ——— (Ge- |
| Alkaline phosphatase | |
| Studies on the arrangement of aminopep- | rencser, Armstrong) (255) 663 ATPase |
| tidase and ———in the microvilli of isolated | A comparison of the sodium-potassium- |
| brush border of rat kidney (Thomas, Kinne) | |
| (255) 114 | activated ———— activities found in isolated brush border and plasma membrane of the |
| Amino acid accumulation | rat intestinal mucosa (Quigley, Gotterer) |
| In vitro ——— by intestinal strips from | (255) 107 |
| normal and essential fatty acid-deficient | ATPase |
| rats (Ahmed, Walker) (255) 815 | Erythrocyte membrane-bound enzymes: |
| α-Aminoisobutyric acid uptake | , phosphatase and adenylate kinase |
| —— by cultured beating heart cells | in human, bovine and porcine erythrocytes |
| (Dunand et al.) (255) 462 | (Heller, Hanahan) (255) 239 |
| Aminopeptidase | ATPase |
| Studies on the arrangement of —— | Functions of the E-ATP and E-P complexes |
| | |

| Brodsky) (255) 220 ATPase Phosphorylation of bovine brain sodium, potassium-stimulated — by adenosine [32P]triphosphate studied by a rapid-mixing technique (Mardh, Zetterqvist) (255) 231 ATPase The role of phospholipid in calcium-stimulated — activity of sarcoplasmic | Calcium release ATP synthesis coupled to from sarcoplasmic reticulum vesicles (Panet, Selinger) (255) 34 Calcium-stimulated ATPase The role of phospholipid in activity of sarcoplasmic reticulum (Meissner, Fleischer) (255) 19 Cardiolipin forms hexagonal structure with |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| reticulum (Meissner, Fleischer) (255) 19 ATPase activity | divalent cations (Rand, SenGupta) (255) |
| Changes in erythrocyte membranes during preparation, as expressed by (Hanahan, Ekholm) (255) 413 ATPase activity | Casein-lysolecithin interactions I. Nuclear magnetic resonance and spin label studies (Barratt, Rayner) (255) 974 |
| The effect of the calcium/magnesium concentration ratio on placental (Ca ²⁺ -Mg ²⁺)- ———————————————————————————————————— | Casein-lysolecithin interactions II. Gel filtration, gel electrophoresis and density-gradient ultracentriguation of the lysolecithin-α ₈₁ -casein system (Partrick et al.) (255) 981 |
| sarcoplasmic reticulum vesicles (Paner, Selinger) (255) 34 | Cation complexes Detection of ionophore ——— on phospho- |
| Axon Intracellular pH electrode. Experiments on the giant squid ——— (Bicher, Ohki) (255) | lipid membranes (Haynes) (255) 406 Cation permeability The effect of pH at hemolysis on the reconstitution of law in human another |
| 900 Bacterial cell membranes | stitution of low ——— in human erythrocyte ghosts (Lepke, Passow) (255) 696 |
| Molecular organization in — I. Sodium dodecyl sulphate solubilization and fractionation of the components of a depleted membrane from Micrococcus lysodeikticus (Estrugo et al.) (255) 960 | Cation transport The relationship between the glucose transport and ——————————————————————————————————— |
| Biotin absorption The sodium activation of ——— in hamster | glucose transport and glucose in rat soleus muscle (Kohn, Vlausen) (255) 798 |
| small intestine in vitro (Berger et al.) (255) 873 | Cell division ———————————————————————————————————— |
| Brush border A comparison of the sodium-potassium- | Protein leakage from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 |
| activated ATPase activities found in iso- lated ——— and plama membrane of the rat intestinal mucosa (Quigley, Gotterer) | Cell membrane Preparation of fat ——— with high sensitivity to lipolytic hormones (Laudat et al.) |
| (255) 107 Brush border | (255) 1005 Cell membranes |
| Iron influx across the ——— of rabbit duodenum: Effects of anemia and iron | Molecular organization in bacterial ———. I. Sodium dodecyl sulphate solubilization |
| loading (Acheson, Schultz) (255) 479 Brush borders Sodium-dependent D-glucose binding to a filamentous fraction of Tris-disrupted | and fractionation of the components of a depleted membrane from Micrococcus lyso- deikticus (Estrugo et al.) (255) 960 Cell membranes |
| from hamster jejunum (Faust et al.) (255) 685 | The effect of cholesterol and epicholesterol incorporation on the permeability and on |
| Brush border Studies on the arrangement of aminopeptidase and alkaline phosphatase in the | the phase transition of intact Acholeplasma laidlawii ——— and derived liposomes (de Kruyff et al.) (255) 331 |
| microvilli of isolated — of rat kidney (Thomas, Kinne) (255) 114 Cable theory General — for cells of algae Characeae (Volkov) (255) 709 | Cell membranes General anesthetics expand — at surgical concentrations (Seeman, Roth) (255) 171 Cells of algae |
| Calcium/magnesium concentration ratio The effect of the ——— on placental (Ca²+— | General cable theory for ——— Characeae (Volkov) (255) 709 |
| Mg ²⁺)-ATPase activity (Shami, Radde) (255) 675 | Cell wall mannan Metabolism of 2-deoxy-p-glucose by baker's |

| yeast. IV. Incorporation of 2-deoxy-D-glu- | Coupling in secondary active transport. |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| cose into ——— (Biely et al.) (255) 631 | Activation of transport by co-transport and/ |
| Characeae | or ——— with the fluxes of other solutes |
| General cable theory for cells of algae ——— | (Heinz et al.) (255) 442 |
| (Volkov) (255) 709 | Coupled transport |
| Chloride transfer | A theory of ——— in cells (Verhoff, |
| Stimulation by anoxia of active — in | Sundaresan) (255) 425 |
| isolated bullfrog small intestine (Armstrong | 2',3'-Cyclic nucleotide 3'-phosphohydrolase |
| et al.) (255) 647 | in human erythrocyte membranes |
| Chloroplasts | (Sudo et al.) (255) 640 |
| Isolation of different classes of spinach | Electroconductivity |
| by counter-current distribution. A | changes during the mitotic cycle in |
| | |
| methodological study (Karlstam, Alberts- | Ehrlich ascites tumour cells (Malenkov et |
| son) (255) 539 | al.) (255) 304 |
| Chlorothricin | Electrolyte transport |
| Interaction of antibiotics with membranes: | by bullfrog colon in vitro (Perheen- |
| ——— (Pache, Chapman) (255) 348 | tupa et al.) (255) 858 |
| Chlorpromazine | Endoplasmic reticulum |
| The membrane concentrations of neutral | The lipids of the plasma membranes and |
| positive anesthetics (alcohols, ———, mor- | ——— from cultured baby hamster kidney |
| phine) fit the Meyer-Overton rule of | cells (BHK21) (Renkonen et al.) (255) 66 |
| anesthesia; negative narcotics do not (Roth, | Epicholesterol |
| Seeman) (255) 207 | The effect of cholesterol and — in- |
| Cholesterol | corporation on the permeability and on the |
| The effect of — and epicholesterol in- | phase transition of intact Acholeplasma |
| corporation on the permeability and on the | laidlawii cell membranes and derived lipo- |
| phase transition of intact Acholeplasma | somes (de Kruyff et al.) (255) 331 |
| laidlawii cell membranes and derived lipo- | Erythrocytes |
| somes (de Kruyff et al.) (255) 331 | Differential effects of lipids on the osmotic |
| Cholesterol replacement | fragility of hamster ——— (Livne et al.) |
| Rate of —— processes in rat. III. Modifi- | (255) 744 |
| cation and critical study of the isotope | Erythrocytes |
| equilibrium technique (Lutton, Chevallier) | Kinetic parameters of glucose efflux from |
| | |
| | |
| (255) 762 | human —— under zero-trans conditions |
| (255) 762 Cholinesterase activities | human ——— under zero-trans conditions (Karlish et al.) (255) 126 |
| (255) 762 Cholinesterase activities ———————————————————————————————————— | human ——— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes |
| (255) 762 Cholinesterase activities ———————————————————————————————————— | human ——— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and |
| (255) 762 Cholinesterase activities ———————————————————————————————————— | human ——— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human ——— with tritiated 2,4,6-trinitro- |
| (255) 762 Cholinesterase activities ———————————————————————————————————— | human —— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human —— with tritiated 2,4,6-trinitrobenzenesulfonic acid and tritiated picryl |
| (255) 762 Cholinesterase activities ———————————————————————————————————— | human —— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human —— with tritiated 2,4,6-trinitrobenzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 |
| (255) 762 Cholinesterase activities ———————————————————————————————————— | human —— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human —— with tritiated 2,4,6-trinitrobenzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes |
| (255) 762 Cholinesterase activities ———————————————————————————————————— | human ——— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human ——— with tritiated 2,4,6-trinitrobenzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate |
| (255) 762 Cholinesterase activities ———————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) |
| (255) 762 Cholinesterase activities ———————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitrobenzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 |
| (255) 762 Cholinesterase activities ———————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitrobenzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts |
| Cholinesterase activities —— in subcellular fractions of rat liver: Association of acetylcholinesterase with the surface membrane, and other properties of the enzyme (Wheeler et al.) (255) 917 Chromatophores The effect of aging resolved —— of Rhodospirillum rubrum on the capacity to reconstitute the energy-linked transhydrogenation (Guber et al.) (255) 167 Cobra venom Activation of enzymes in erythrocyte mem- | human —— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human —— with tritiated 2,4,6-trinitrobenzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human —— (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the recon- |
| Cholinesterase activities —— in subcellular fractions of rat liver: Association of acetylcholinesterase with the surface membrane, and other properties of the enzyme (Wheeler et al.) (255) 917 Chromatophores The effect of aging resolved —— of Rhodospirillum rubrum on the capacity to reconstitute the energy-linked transhydrogenation (Guber et al.) (255) 167 Cobra venom Activation of enzymes in erythrocyte membranes by a basic protein isolated from | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in |
| Cholinesterase activities —— in subcellular fractions of rat liver: Association of acetylcholinesterase with the surface membrane, and other properties of the enzyme (Wheeler et al.) (255) 917 Chromatophores The effect of aging resolved —— of Rhodospirillum rubrum on the capacity to reconstitute the energy-linked transhydrogenation (Guber et al.) (255) 167 Cobra venom Activation of enzymes in erythrocyte mem- | human —— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human —— with tritiated 2,4,6-trinitrobenzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human —— (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the recon- |
| Cholinesterase activities ———————————————————————————————————— | human ———— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human ———— with tritiated 2,4,6-trinitrobenzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human ———— (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human ————————— (Lepke, Passow) (255) 696 Erythrocyte membranes |
| Cholinesterase activities ———————————————————————————————————— | human ——— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human ——— with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human ———— (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human ———— (Lepke, Passow) (255) 696 |
| Cholinesterase activities —— in subcellular fractions of rat liver: Association of acetylcholinesterase with the surface membrane, and other properties of the enzyme (Wheeler et al.) (255) 917 Chromatophores The effect of aging resolved —— of Rhodospirillum rubrum on the capacity to reconstitute the energy-linked transhydrogenation (Guber et al.) (255) 167 Cobra venom Activation of enzymes in erythrocyte membranes by a basic protein isolated from —— (Fajnholc et al.) (255) 850 Conductance channel The unit —— of alamethicin (Gordon, | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome |
| Cholinesterase activities —— in subcellular fractions of rat liver: Association of acetylcholinesterase with the surface membrane, and other properties of the enzyme (Wheeler et al.) (255) 917 Chromatophores The effect of aging resolved —— of Rhodospirillum rubrum on the capacity to reconstitute the energy-linked transhydrogenation (Guber et al.) (255) 167 Cobra venom Activation of enzymes in erythrocyte membranes by a basic protein isolated from —— (Fajnholc et al.) (255) 850 Conductance channel The unit —— of alamethicin (Gordon, Haydon) (255) 1014 Connective tissue membrane | human —— under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human —— with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human —— (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human —— (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthe- |
| Cholinesterase activities —— in subcellular fractions of rat liver: Association of acetylcholinesterase with the surface membrane, and other properties of the enzyme (Wheeler et al.) (255) 917 Chromatophores The effect of aging resolved —— of Rhodospirillum rubrum on the capacity to reconstitute the energy-linked transhydrogenation (Guber et al.) (255) 167 Cobra venom Activation of enzymes in erythrocyte membranes by a basic protein isolated from —— (Fajnholc et al.) (255) 850 Conductance channel The unit —— of alamethicin (Gordon, Haydon) (255) 1014 | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 Erythrocyte membranes |
| Cholinesterase activities —— in subcellular fractions of rat liver: Association of acetylcholinesterase with the surface membrane, and other properties of the enzyme (Wheeler et al.) (255) 917 Chromatophores The effect of aging resolved —— of Rhodospirillum rubrum on the capacity to reconstitute the energy-linked transhydrogenation (Guber et al.) (255) 167 Cobra venom Activation of enzymes in erythrocyte membranes by a basic protein isolated from —— (Fajnholc et al.) (255) 850 Conductance channel The unit —— of alamethicin (Gordon, Haydon) (255) 1014 Connective tissue membrane | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 |
| Cholinesterase activities ———————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitrobenzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 Erythrocyte membranes Activation of enzymes in — by a basic |
| Cholinesterase activities ——————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 Erythrocyte membranes |
| Cholinesterase activities ———————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 Erythrocyte membranes Activation of enzymes in — by a basic protein isolated from cobra venom (Fajnholc et al.) (255) 850 Erythrocyte membranes |
| Cholinesterase activities ——————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 Erythrocyte membranes Activation of enzymes in — by a basic protein isolated from cobra venom (Fajnholc et al.) (255) 850 Erythrocyte membranes |
| Cholinesterase activities —— in subcellular fractions of rat liver: Association of acetylcholinesterase with the surface membrane, and other properties of the enzyme (Wheeler et al.) (255) 917 Chromatophores The effect of aging resolved —— of Rhodospirillum rubrum on the capacity to reconstitute the energy-linked transhydrogenation (Guber et al.) (255) 167 Cobra venom Activation of enzymes in erythrocyte membranes by a basic protein isolated from —— (Fajnholc et al.) (255) 850 Conductance channel The unit —— of alamethicin (Gordon, Haydon) (255) 1014 Connective tissue membrane A —— as a molecular sieve (Elsbach, Pettis) (255) 149 Co-transport Coupling in secondary active transport. Activation of transport by —— and/or counter-transport with the fluxes of other | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 Erythrocyte membranes Activation of enzymes in — by a basic protein isolated from cobra venom (Fajnholc et al.) (255) 850 Erythrocyte membranes Anesthetics expand — without causing |
| Cholinesterase activities ——————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 Erythrocyte membranes Activation of enzymes in — by a basic protein isolated from cobra venom (Fajnholc et al.) (255) 850 Erythrocyte membranes Anesthetics expand — without causing potassium loss (Roth, Seeman) (255) 190 |
| Cholinesterase activities ——————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 Erythrocyte membranes Activation of enzymes in — by a basic protein isolated from cobra venom (Fajnholc et al.) (255) 850 Erythrocyte membranes Anesthetics expand — without causing potassium loss (Roth, Seeman) (255) 190 Erythrocyte membranes |
| Cholinesterase activities ——————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 Erythrocyte membranes Activation of enzymes in — by a basic protein isolated from cobra venom (Fajnholc et al.) (255) 850 Erythrocyte membranes Anesthetics expand — without causing potassium loss (Roth, Seeman) (255) 190 |
| Cholinesterase activities ———————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitro-benzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 Erythrocyte membranes Activation of enzymes in — by a basic protein isolated from cobra venom (Fajnholc et al.) (255) 850 Erythrocyte membranes Anesthetics expand — without causing potassium loss (Roth, Seeman) (255) 190 Erythrocyte membranes Aqueous pores in lipid bilayers and — |
| Cholinesterase activities ——————————————————————————————————— | human — under zero-trans conditions (Karlish et al.) (255) 126 Erythrocytes Reaction of human serum albumin and human — with tritiated 2,4,6-trinitrobenzenesulfonic acid and tritiated picryl chloride (Arrotti, Garvin) (255) 79 Erythrocytes Temperature dependence of phosphate entry into stored human — (Peterson) (255) 844 Erythrocyte ghosts The effect of pH at hemolysis on the reconstitution of low cation permeability in human — (Lepke, Passow) (255) 696 Erythrocyte membranes The action and adsorption of local anesthetic enantiomers on — and synaptosome membranes (Roth et al.) (255) 199 Erythrocyte membranes Activation of enzymes in — by a basic protein isolated from cobra venom (Fajnholc et al.) (255) 850 Erythrocyte membranes Anesthetics expand — without causing potassium loss (Roth, Seeman) (255) 190 Erythrocyte membranes Aqueous pores in lipid bilayers and — (Solomon, Gary-Bobo) (255) 1019 |

| expressed by ATPase activity (Hanahan, | surface membrane of normal and trans- |
|------------------------------------------------|---------------------------------------------------------------------------------|
| Ekholm) (255) 413 | formed mammalian cells (Inbar et al.) (255) |
| Erythrocyte membranes | 703 |
| 2',3'-Cyclic nucleotide 3'-phosphohydrolase | Ganglion |
| in human ——— (Sudo et al.) (255) 640 | Movement in a ——— (Shaw, Newby) (255) |
| Erythrocyte membranes | 411 |
| The hydrophobic expansion of ——— by | y-Globulins |
| the phenol anesthetics (Machleidt et al.) | Intestinal absorption and degradation of |
| (255) 178 | rat and bovine in the suckling rat |
| Erythrocyte membranes | (Jones) (255) 530 |
| Micromorphologic consequences following | Glucose |
| perturbation of ——by trypsin, phospho- | The effect of sterol structure on the per- |
| lipase A, lysolecithin, sodium dodecylsulfate | meability of liposomes to, glycerol |
| and saponin: A correlated freeze-etching | and rubidium (Demel et al.) (255) 321 |
| and biochemical study (Speth et al.) (255) 386 | 2-deoxy-D-Glucose |
| Erythrocyte membranes | Metabolism of ——— by baker's yeast. IV. |
| Studies on the succinylation of — | Incorporation of 2-deoxy-D-glucose into |
| (Moldow et al.) (255) 133 | |
| | cell wall mannan (Bielp et al.) (255) 631 |
| Erythrocyte membrane-bound enzymes | D-Glucose binding |
| : ATPase, phosphatase and adenyl- | Sodium-dependent ———— to a filamentous |
| ate kinase in human, bovine and porcine | fraction of Tris-disrupted brush borders |
| erythrocytes (Heller, Hanahan) (255) 239 | from hamster jejunum (Faust et al.) (255) |
| Erythrocyte membrane bound enzyme | 685 |
| Human ——: Acetylcholinesterase (Hel- | Glucose efflux |
| ler, Hanahan) (255) 251 | Kinetic parameters of ——— from human |
| Erythrocyte membrane protein | erythrocytes under zero-trans conditions |
| The interaction of fluorescence probes and | (Karlish et al.) (255) 126 |
| nuclear magnetic resonance probes with | Glucose 6-phosphate |
| fractions (Randall et al.) (255) 888 | Properties of the ——-solubilized brain |
| Erythrocyte membrane proteins | hexokinase: Evidence for a lipoprotein |
| Selective solubilization of ——— with | complex (Craven, Basford) (255) 620 |
| guanidine hydrochloride (Steck) (255) 553 | Glucose transport |
| Erythrocyte volume | The relationship between the ——— and |
| Perturbation of ———: Constancy of mem- | cation transport across cell membranes in |
| brane transport parameters for certain slow | isolated tissues. VII. The effects of extra- |
| penetrants (Farmer, Macey) (255) 502 | cellular Na+ and K+ on the 3-O-methyl- |
| Escherichia coli K ₁₂ | glucose transport and glucose in rat soleus |
| A mutant of ——— energy-uncoupled for | muscle (Kohn, Clausen) (255) 798 |
| lactose transport (Wilson, Kusch) (255) 786 | Glycerol |
| Essential fatty acid | The effect of sterol structure on the per- |
| In vitro amino acid accumulation by in- | meability of liposomes to glucose, |
| testinal strips from normal and | and rubidium (Demel et al.) (255) 321 |
| deficient rats (Ahmed, Walker) (255) 815 | Gramicidin |
| Fat cell membrane | Solution and interfacial properties of |
| Preparation of ——— with high sensitivity | pertinent to its effect on membranes (Kemp |
| to lipolytic hormones (Laudat et al.) (255) | et al.) (255) 493 |
| 1005 | Gramicidin S |
| Fatty acid transport | Interaction of antibiotics with membranes. |
| into the brain (Dhopeshwarkar et | Polymyxin B and ——— (Pache et al.) |
| al.) (255) 572 | |
| Fibroblast | (255) 35 ⁸ Hemolysis |
| Studies of surface membranes of two mouse | |
| cell lines (Scher, Barland) (255) 580 | The effect of pH at —— on the reconsti- |
| Filamentous cells | tution of low cation permeability in human |
| Cell division in a species of Erwinia. XIII. | erythrocyte ghosts (Lepke, Passow) (255) |
| | 696 |
| Protein leakage from the peripheral area of | Hexokinase |
| (Grula, Hopfer) (255) 822 | Properties of the glucose 6-phosphate- solubilized brain ———: Evidence for a |
| Fluorescence probes | Solublinzed Drain ———: Evidence for a |
| The interaction of and nuclear | lipoprotein complex (Craven, Basford) (255) |
| magnetic resonance probes with erythrocyte | 620 |
| membrane protein fractions (Randall et al.) | Hexose transport |
| (255) 888 | and phosphorylation by hamster |
| L-Fucose | kidney cortex slices and everted jejunal |
| Availability of ———-like sites on the | rings (Elsas, MacDonell, Jr.) (255) 948 |

| | 75 1 224 . (116 1 |
|-------------------------------------------------------|----------------------------------------------------------|
| Interfacial properties | Permeability of a modified ——— to |
| Solution and ——— of gramicidin pertinent | sodium (Petkau, Chelack) (255) 161 |
| to its effect on membranes (Kemp et al.) | Lipolytic hormones |
| (255) 493 | Preparation of fat cell membrane with high |
| | • • • • • • • • • • • • • • • • • • • • |
| Intestinal absorption | sensitivity to ——— (Laudat et al.) (255) |
| ——— and degradation of rat and bovine | 1005 |
| γ -globulins in the suckling rat (Jones) (255) | Lipopolysaccharide-protein complex |
| 530 | Protein surface films and a ——— extracted |
| Intestinal strips | from Pseudomonas aeruginosa (Suzuki, |
| In vitro amino acid accumulation by ——— | Goto) (255) 734 |
| | |
| from normal and essential fatty acid-defi- | Lipoprotein complex |
| cient rats (Ahmed, Walker) (255) 815 | Properties of the glucose 6-phosphate-solu- |
| Intestine | bilized brain hexokinase: Evidence for a |
| The sodium activation of biotin absorption | ——— (Craven, Basford) (255) 620 |
| in hamster small ——— in vitro (Berger et | Lipoprotein subunits |
| al.) (255) 873 | Studies on plasma membranes from liver |
| 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | |
| Ionophore complexes | cells. Separation and characterization of |
| Detection of cation on phospholipid | of the isolated plasma membranes |
| membranes (Haynes) (255) 406 | (Barclay et al.) (255) 931 |
| Iron influx | Liposomes |
| ——— across the brush border of rabbit | Changes in permeability of Staphylococcus |
| duodenum: Effects of anemia and iron | aureus and derived — with varying |
| | lipid composition (Haest et al.) (255) 720 |
| loading (Acheson, Schultz) (255) 479 | |
| Isotope equilibrium technique | Liposomes |
| Rate of cholesterol replacement processes in | The effect of cholesterol and epicholesterol |
| rat. III. Modification and critical study of | incorporation on the permeability and on |
| the — (Lutton, Chevallier) (255) 762 | the phase transition of intact Achole- |
| Jejunal rings | plasma laidlawii cell membranes and derived |
| Hexose transport and phosphorylation by | ——— (de Kruyff et al.) (255) 331 |
| | |
| hamster kidney cortex slices and everted | Liposomes |
| (Elsas, MacDonell, Jr.) (255) 948 | The effect of sterol structure on the per- |
| Kidney cortex slices | meability of ——— to glucose, glycerol and |
| Hexose transport and phosphorylation by | rubidium (Demel et al.) (255) 321 |
| hamster — and everted jejunal rings | Liposome permeability |
| (Elsas, MacDonell, Jr.) (255) 948 | Effect of haemoglobin on ——— to rubi- |
| Lactose transport | dium and other solutes (Calissano et al.) |
| A mutant of Escherchia coli K ₁₂ energy- | (255) 1009 |
| | |
| uncoupled for ——— (Wilson, Kusch) (255) | Local anesthetic |
| 786 | The action and adsorption of ——— enan- |
| Lecithins | tiomers on erythrocyte membranes and |
| High resolution proton relaxation studies | synaptosome membranes (Roth et al.) (255) |
| of ——— (Lee et al.) (255) 43 | 199 |
| Lecithin | Lysolecithin |
| Structural requirements of sterols for the | Micromorphologic consequences following |
| interaction with ——— at the air—water | |
| | perturbation of erythrocyte membranes by |
| interface (Demel et al.) (255) 311 | trypsin, phospholipase A, ——, sodium |
| Lipid bilayers | dodecylsulfate and saponin: A correlated |
| Aqueous pores in ——— and erythrocyte | freeze-etching and biochemical study (Speth |
| membranes (Solomon, Gary-Bobo) (255) 1019 | et al.) (255) 386 |
| Lipid composition | Lysolecithin-casein interactions |
| Changes in permeability of Staphylococcus | I. Nuclear magnetic resonance and |
| aureus and derived liposomes with varying | spin label studies (Barratt, Rayner) (255) |
| | |
| ——— (Haest et al.) (255) 720 | 974 |
| Lipids | Lysolecithin-casein interactions |
| Circular dichroism of mitochondrial mem- | ——. II. Gel filtration, gel electrophoresis |
| branes before and after extraction of ——— | and density-gradient ultracentrifugation of |
| and surface proteins (Zahler et al.) (255) | the lysolecithin- α_{s1} -casein system (Partrick |
| 365 | et al.) (255) 981 |
| Lipids | Magnesium/calcium concentration ratio |
| The — of the plasma membranes and | The effect of the ——— on placental (Ca ²⁺ — |
| | |
| endoplasmic reticulum from cultured baby | Mg ²⁺)-ATPase activity (Shami, Radde) |
| hamster kidney cells (BHK12) (Renkonen | (255) 675 |
| et al.) (255) 66 | Mannan |

et al.) (255) 66 Lipid membrane

Mannan
Metabolism of 2-deoxy-D-glucose by baker's

| yeast. IV. Incorporation of 2-deoxy-D-glu- | Membrane transport parameters |
|----------------------------------------------|-------------------------------------------------|
| cose into cell wall ——— (Biely et al.) (255) | Perturbation of erythrocyte volume: Con- |
| 631 | stancy of ——— for certain slow penetrants |
| Membrane | (Farmer, Macey) (255) 502 |
| Availability of L-fucose-like sites on the | 3-O-Methylglucose transport |
| surface ——— of normal and transformed | The relationship between the glucose trans- |
| mammalian cells (Inbar et al.) (255) 703 | port and cation transport across cell mem- |
| Membrane | |
| | branes in isolated tissues. VII. The effects of |
| A connective tissue ——— as a molecular | extracellular Na+ and K+ on the ——— and |
| sieve (Elsbach, Pettis) (255) 149 | glucose in rat soleus muscle (Kohn, Clausen) |
| Membranes | (255) 798 |
| Hindrance of solute diffusion within ——— | Meyer-Overton rule of anesthesia |
| as measured with microporous membranes | The membrane concentrations of neutral |
| of known pore geometry (Beck, Schultz) | and positive anesthetics (alcohols, chlor- |
| (255) 273 | promazine, morphine) fit the ———; nega- |
| Membranes | tive narcotics do not (Roth, Seeman) (255) 207 |
| Interaction of antibiotics with ———: | Micrococcus lysodeikticus |
| Chlorothricin (Pache, Champan) (255) 348 | Molecular organization in bacterial cell |
| Membranes | membranes. I. Sodium dodecyl sulphate |
| Interaction of antibiotics with ——: | solubilization and fractionation of the com- |
| Polymyxin B and gramicidin S (Pache et | ponents of depleted membrane from ——— |
| al.) (255) 358 | (Estrugo et al.) (255) 960 |
| Membrane | Microporous membranes |
| Solution of the equations describing the | Hindrance of solute diffusion within mem- |
| sodium flow in a semi-infinite composite | branes as measured with ——— of known |
| | |
| system containing a ——— (Fleming, | pore geometry (Beck, Schultz) (255) 273 |
| Castleden) (255) I | Microsomal membranes |
| Membranes | Structural properties of rough and smooth |
| Studies of surface ——— of two mouse | A study with fluorescence probes |
| fibroblast cell lines (Scher, Barland) (255) | (Dallner, Azzi) (255) 589 |
| 580 | Microvilli |
| Membrane ATPase | Studies on the arrangement of aminopep- |
| Functions of the E-ATP and E-P com- | tidase and alkaline phosphatase in the |
| plexes in the ——— reaction (Shamoo, | ——— of isolated brush border of rat |
| Brodsky) (255) 220 | kidney (Thomas, Kinne) (255) 114 |
| Membrane fraction | Mitochondrial membranes |
| The effect of ATP on the CD spectrum of | Circular dichroism of ——— before and |
| ——— from oxyntic cells (Masotti et al.) | after extraction of lipids and surface pro- |
| (255) 420 | teins (Zahler et al.) (255) 365 |
| Membrane from Micrococcus lysodeikticus | Mitochondrial membranes |
| Molecular organisation in bacterial cell | Fractionation of ——— with sodium deoxy- |
| membranes. I. Sodium dodecyl sulphate | cholate (Hall, Crane) (255) 602 |
| solubilisation and fractionation of the com- | Mitochondrial membrane protein |
| ponents of a depleted ——— (Estrugo et | Physical characteristics of ——— dissolved |
| al.) (255) 960 | in chloroform—methanol (Curtis) (255) 833 |
| Membrane permeability | Mitotic cycle |
| Relation between adsorption at an oil- | Electroconductivity changes during the |
| water interface and ——— (Wang et al.) | —— in Ehrlich ascites tumour cells |
| | |
| (255) 691 Mambana matantial | (Malenkov et al.) (255) 304 |
| Membrane potential | Morphine |
| The buffer value and trans—— of | The membrane concentrations of neutral |
| Escherichia coli (White, O'Brien) (255) 780 | and positive anesthetics (alcohols, chlor- |
| Membrane protein | promazine, ———) fit the Meyer-Overton |
| Physical characteristics of mitochondrial | rule of anesthesia: negative narcotics do not |
| dissolved in chloroform-methanol | (Roth, Seeman) (255) 207 |
| (Curtis) (255) 833 | Narcotics |
| Membrane protein | The membrane concentrations of neutral |
| The interaction of fluorescence probes and | and positive anesthetics (alcohols, chlor- |
| nuclear magnetic resonance probes with | promazine, morphine) fit the Meyer-Over- |
| erythrocyte — fractions (Randall et | ton rule of anesthesia: negative ——— do |
| al.) (255) 888 | not (Roth, Seeman) (255) 207 |
| Membrane resistance | Nuclear envelopes |
| Low — in sucrose gap — a parallel | The isolation of the ——— of rat liver |
| leakage path (Pooler, Oxford) (255) 681 | (Agutter) (255) 397 |

| | *** |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Nuclear magnetic resonance robes | Phospholipid membranes |
| The interaction of fluorescence probes and | Detection of ionophore-cation complexes on |
| ——— with erythrocyte membrane protein | ——— (Haynes) (255) 406 |
| fractions (Randall et al.) (255) 888 | Picryl chloride |
| Nucleoside transport | Reaction of human serum albumin and |
| Membrane —— in rabbit polymorpho- | human erythrocytes with tritiated 2,4,6,- |
| nuclear leukocytes (Taube, Berlin) (255) 6 | trinitrobenzenesulfonic acid and tritiated |
| Oil-water interface | ——— (Arrotti, Garvin) (255) 79 |
| Relation between adsorption at an ——— | Plasma membrane |
| and membrane permeability (Wang et al.) | A comparison of the sodium-potassium- |
| | activated ATPase activities found in iso- |
| (255) 691 Ouabain | lated brush border and ——— of the rat |
| | |
| uninhibited sodium transport in | intestinal mucosa (Quigley, Gotterer) (255) |
| human erythrocytes: The effects of tri- | 107 |
| flocin (Dunn) (255) 567 | Plasma membranes |
| Osmotic fragility | The lipids of the ——— and endoplasmic |
| Differential effects of lipids on the ——— of | reticulum from cultured baby hamster |
| hamster erythrocytes (Livne et al.) (255) | kidney cells (BHK21) (Renkonen et al.) |
| 744 | (255) 66 |
| Oxyntic cells | Plasma membranes |
| The effect of ATP on the CD spectrum of | Studies on ——— from liver cells. Separa- |
| membrane fraction from (Masotti | tion and characterization of lipoprotein sub- |
| et al.) (255) 420 | units of the isolated plasma membranes |
| Permeability of liposomes | (Barclay et al.) (255) 931 |
| The effect of sterol structure on the ——— | |
| | Polymorphonuclear leukocytes |
| to glucose, glycerol and rubidium (Demel | Membrane nucleoside transport in rabbit |
| et al.) (155) 321 | ——— (Taube, Berlin) (255) 6 |
| pH electrode | Polymyxin B |
| Intracellular — Experiments on the | Interaction of antibiotics with membranes. |
| giants squid axon (Bicher, Ohki) (255) 900 | —— and gramicidin S (Pache et al.) (255) |
| Phosphate entry | 358 |
| Temperature dependence of ——— into | Pores |
| stored human erythrocytes (Peterson) (255) | Aqueous — in lipid bilayers and ery- |
| 844 | throcyte membranes (Solomon, Gary-Bobo) |
| Phosphatase | (255) 1019 |
| Erythrocyte membrane-bound enzymes: | Potassium flux |
| ATPase, — and adenylate kinase in | —— and sodium transport in the isolated |
| human, bovine and porcine erythrocytes | frog skin (Candia, Zadunaisky) (255) 517 |
| (Heller, Hanahan) (255) 239 | Potassium loss |
| Phosphatase | Anesthetics expand erythrocyte membranes |
| | Anesthetics expand of yulf ocyte membranes |
| Studies on the arrangement of aminopep- | without squaing (Both Somen) |
| tidage and allealine in the mismorilli | without causing ——— (Roth, Seeman) |
| tidase and alkaline —— in the microvilli | (255) 190 |
| of isolated brush borders of rat kidney | (255) 190 Probenecid |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 | (255) 190 Probenecid ———: Effects on water permeability in |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles | (255) 190 Probenecid |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous ——— | (255) 190 Probenecid ———: Effects on water permeability in |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles | (255) 190 Probenecid ———: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous ——— | (255) 190 Probenecid ——: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous (Redwood et al.) (255) 557 | (255) 190 Probenecid : Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous ———————————————————————————————————— | (255) 190 Probenecid : Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. from the peripheral area of filamen- |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous (Redwood et al.) (255) 557 Phospholipase A Micromorphologic consequences following perturbation of erythrocyte membranes by | (255) 190 Probenecid —: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. —— from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous (Redwood et al.) (255) 557 Phospholipase A Micromorphologic consequences following perturbation of erythrocyte membranes by trypsin, ———, lysolecithin, sodium dode- | (255) 190 Probenecid ——: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. ——— from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous (Redwood et al.) (255) 557 Phospholipase A Micromorphologic consequences following perturbation of erythrocyte membranes by trypsin, ———, lysolecithin, sodium dode- cylsulfate and saponium: A correlated | (255) 190 Probenecid ——: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. ——— from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex Protein surface films and a ——— extracted |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous (Redwood et al.) (255) 557 Phospholipase A Micromorphologic consequences following perturbation of erythrocyte membranes by trypsin, ———, lysolecithin, sodium dode- cylsulfate and saponium: A correlated freeze-etching and biochemical study (Speth | (255) 190 Probenecid —: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. —————————————————————————————————— |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous (Redwood et al.) (255) 557 Phospholipase A Micromorphologic consequences following perturbation of erythrocyte membranes by trypsin, ———, lysolecithin, sodium dodecylsulfate and saponium: A correlated freeze-etching and biochemical study (Speth et al.) (255) 386 | (255) 190 Probenecid ——: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. ——— from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex Protein surface films and a ——— extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous (Redwood et al.) (255) 557 Phospholipase A Micromorphologic consequences following perturbation of erythrocyte membranes by trypsin, ————, lysolecithin, sodium dodecylsulfate and saponium: A correlated freeze-etching and biochemical study (Speth et al.) (255) 386 Phospholipid | (255) 190 Probenecid —: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. ——from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex Protein surface films and a — extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Protein surface films |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous ———————————————————————————————————— | (255) 190 Probenecid —: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. ——from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex Protein surface films and a ——extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Protein surface films ——and a lipopolysaccharide protein |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous ———————————————————————————————————— | Probenecid —: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. —— from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex Protein surface films and a —— extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Protein surface films —— and a lipopolysaccharide protein complex extracted from Pseudomonas |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous ———————————————————————————————————— | Probenecid —: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. —— from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex Protein surface films and a —— extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Protein surface films —— and a lipopolysaccharide protein complex extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous ———————————————————————————————————— | Probenecid ——: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. ——— from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex Protein surface films and a ——— extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Protein surface films ——— and a lipopolysaccharide protein complex extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Prymnesium parvum toxin |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous (Redwood et al.) (255) 557 Phospholipase A Micromorphologic consequences following perturbation of erythrocyte membranes by trypsin, ——, lysolecithin, sodium dodecylsulfate and saponium: A correlated freeze-etching and biochemical study (Speth et al.) (255) 386 Phospholipid The role of —— in calcium-stimulated ATPase activity of sarcoplasmic reticulum (Meissner, Fleischer) (255) 19 Phospholipids Surface properties of synthetic ——— | Probenecid ——: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. ——— from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex Protein surface films and a ——— extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Protein surface films ———— and a lipopolysaccharide protein complex extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Prymnesium parvum toxin Nature of initial damage to Ehrlich ascites |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous — (Redwood et al.) (255) 557 Phospholipase A Micromorphologic consequences following perturbation of erythrocyte membranes by trypsin, — , lysolecithin, sodium dodecylsulfate and saponium: A correlated freeze-etching and biochemical study (Speth et al.) (255) 386 Phospholipid The role of — in calcium-stimulated ATPase activity of sarcoplasmic reticulum (Meissner, Fleischer) (255) 19 Phospholipids Surface properties of synthetic — (Hayashi et al.) (255) 98 | Probenecid ——: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. ——— from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex Protein surface films and a ——— extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Protein surface films ———— and a lipopolysaccharide protein complex extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Prymnesium parvum toxin Nature of initial damage to Ehrlich ascites cells caused by ———— (Dafni, Giberman) |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous (Redwood et al.) (255) 557 Phospholipase A Micromorphologic consequences following perturbation of erythrocyte membranes by trypsin, ———————————————————————————————————— | Probenecid —: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. ——from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex Protein surface films and a — extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Protein surface films ——and a lipopolysaccharide protein complex extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Prymnesium parvum toxin Nature of initial damage to Ehrlich ascites cells caused by —— (Dafni, Giberman) (255) 380 |
| of isolated brush borders of rat kidney (Thomas, Kinne) (255) 114 Phosphatidylcholine vesicles Dielectric studies on homogeneous — (Redwood et al.) (255) 557 Phospholipase A Micromorphologic consequences following perturbation of erythrocyte membranes by trypsin, — , lysolecithin, sodium dodecylsulfate and saponium: A correlated freeze-etching and biochemical study (Speth et al.) (255) 386 Phospholipid The role of — in calcium-stimulated ATPase activity of sarcoplasmic reticulum (Meissner, Fleischer) (255) 19 Phospholipids Surface properties of synthetic — (Hayashi et al.) (255) 98 | Probenecid ——: Effects on water permeability in frog urinary bladder (Ripoche et al.) (255) 91 Protein leakage Cell division in a species of Erwinia. XIII. ——— from the peripheral area of filamentous cells (Grula, Hopfer) (255) 822 Protein-lipopolysaccharide complex Protein surface films and a ——— extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Protein surface films ———— and a lipopolysaccharide protein complex extracted from Pseudomonas aeruginosa (Suzuki, Goto) (255) 734 Prymnesium parvum toxin Nature of initial damage to Ehrlich ascites cells caused by ———— (Dafni, Giberman) |

| meability to ——— and other solutes | Changes in permeability of ——— and |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (Calissano et al.) (255) 1009 | derived liposomes with varying lipid com- |
| Rubidium | position (Haest et al.) (255) 720 |
| The effect of sterol structure on the per- | Sterols |
| meability of liposomes to glucose, glycerol | Structural requirements of ——— for the |
| and — (Demel et al.) (255) 321 | interaction with lecithin at the air-water |
| Saponin | interface (Demel et al.) (255) 311 |
| Micromorphologic consequences following | Sterol structure |
| perturbation of erythrocyte membranes by | The effect of —— on the permeability of |
| trypsin, phospholipase A, lysolecithin, so- | liposomes to glucose glycerol and rubidum |
| dium dodecylsulfate and ——: A cor- | (Demel et al.) (255) 321 |
| | Sucrose antagonism |
| related freeze-etching and biochemical study | |
| (Speth et al.) (255) 386 | of drug and temperature effects on |
| Sarcoplasmic reticulum | cardiac sarcoplasmic reticulum (Solaro et |
| Sucrose antagonism of drug and tempera- | al.) (255) 751 |
| ture effects on cardiac ——— (Solaro et al.) | Sucrose gap |
| (255) 75 ^I | Low membrane resistance in ——————————————————————————————————— |
| Sarcoplasmic reticulum | parallel leakage path (Pooler, Oxford) (255) |
| The role of phospholipid in calcium-stium- | 681 |
| lated ATPase activity of ——— (Meissner, | Surface membrane |
| Fleischer) (255) 19 | Availability of L-fucose-like sites on the |
| Sarcoplasmic reticulum vesicles | — of normal and transformed mamma- |
| ATP synthesis coupled to calcium release | lian cells (Inbar et al.) (255) 703 |
| from ——— (Panet, Selinger) (255) 34 | Surface membrane |
| Sodium | Cholinesterase activities in subcellular frac- |
| Permeability of a modified lipid membrane | tions of rat liver: Association of acetyl- |
| to — (Petkau, Chelack) (255) 161 | cholinesterase with the —, and other |
| Sodium dodecylsulfate | properties of the enzyme (Wheeler et al.) |
| Micromorphologic consequences following | (255) 917 |
| perturbation of erythrocyte membranes by | Surface membranes |
| trypsin, phopholipase A, lysolecithin, | Studies of — of two mouse fibroblast |
| and saponin: A correlated freeze-etching | cell lines (Scher, Barland) (255) 580 |
| and biochemical study (Speth et al.) (255) | Surface proteins |
| 386 | Circular dichroism of mitochondrial mem- |
| Sodium-potassium-activated ATPase | branes before and after extraction of lipids |
| A comparison of the ——— activities | and ——— (Zahler et al.) (255) 365 |
| found in isolated brush border and plasma | Synaptosome membranes |
| membrane of the rat intestinal mucosa | The action and adsorption of local anesthe- |
| (Quigley, Gotterer) (255) 107 | tic enantiomers on erythrocyte membranes |
| Sodium, potassium-stimulated ATPase | and ——— (Roth et al.) (255) 199 |
| Phosphorylation of bovine brain — by | Thiamine triphosphatase |
| adenosine [32P]-triphosphate studied by a | - in the nervous system (Barchi, |
| rapid-mixing technique (Mårdh, Zetterqvist | Braun) (255) 402 |
| | Transhydrogenation |
| (255) 231 Sodium transfer | The effect of aging resolved chromatophores |
| ——— in bullfrog small intestine. Stimu- | of Rhodospirillum rubrum on the capacity |
| | or imodospirmum rubrum on the capacity |
| | to reconstitute the energy-linked |
| lation by exogenous ATP (Gerencser, | to reconstitute the energy-linked ———————————————————————————————————— |
| Armstrong) (255) 663 | (Guber et al.) (255) 167 |
| Armstrong) (255) 663 Sodium transport | (Guber et al.) (255) 167 Transmembrane potential |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited ———————————————————————————————————— | (Guber et al.) (255) 167 Transmembrane potential The buffer value and ——— of Escherichia |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited ———————————————————————————————————— | (Guber et al.) (255) 167 Transmembrane potential The buffer value and ——————————————————————————————————— |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited ———————————————————————————————————— | (Guber et al.) (255) 167 Transmembrane potential The buffer value and ——————————————————————————————————— |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited ———————————————————————————————————— | (Guber et al.) (255) 167 Transmembrane potential The buffer value and — of Escherichia coli (White, O'Brien) (255) 780 Transport Coupling in secondary active — |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited ———————————————————————————————————— | (Guber et al.) (255) 167 Transmembrane potential The buffer value and of Escherichia coli (White, O'Brien) (255) 780 Transport Coupling in secondary active Activation of transport by co-transport and/or |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited ———————————————————————————————————— | (Guber et al.) (255) 167 Transmembrane potential The buffer value and of Escherichia coli (White, O'Brien) (255) 780 Transport Coupling in secondary active Activation of transport by co-transport and/or counter-transport with the fluxes of other |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited —— in human erythrocytes: The effects of triflocin (Dunn) (255) 567 Sodium transport Potassium flux and —— in the isolated frog skin (Candia, Zadunaisky) (255) 517 Solute diffusion | (Guber et al.) (255) 167 Transmembrane potential The buffer value and of Escherichia coli (White, O'Brien) (255) 780 Transport Coupling in secondary active Activation of transport by co-transport and/or counter-transport with the fluxes of other solutes (Heinz et al.) (255) 442 |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited ———————————————————————————————————— | (Guber et al.) (255) 167 Transmembrane potential The buffer value and ——————————————————————————————————— |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited ———————————————————————————————————— | (Guber et al.) (255) 167 Transmembrane potential The buffer value and ——————————————————————————————————— |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited ———————————————————————————————————— | (Guber et al.) (255) 167 Transmembrane potential The buffer value and of Escherichia coli (White, O'Brien) (255) 780 Transport Coupling in secondary active Activation of transport by co-transport and/or counter-transport with the fluxes of other solutes (Heinz et al.) (255) 442 Transport in cells A theory of coupled (Verhoff, Sundaresan) (255) 425 |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited —— in human erythrocytes: The effects of triflocin (Dunn) (255) 567 Sodium transport Potassium flux and —— in the isolated frog skin (Candia, Zadunaisky) (255) 517 Solute diffusion Hindrance of —— within membranes as measured with microporous membranes of known pore geometry (Beck, Schultz) (255) 273 | (Guber et al.) (255) 167 Transmembrane potential The buffer value and of Escherichia coli (White, O'Brien) (255) 780 Transport Coupling in secondary active Activation of transport by co-transport and/or counter-transport with the fluxes of other solutes (Heinz et al.) (255) 442 Transport in cells A theory of coupled (Verhoff, Sundaresan) (255) 425 Triflocin |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited —— in human erythrocytes: The effects of triflocin (Dunn) (255) 567 Sodium transport Potassium flux and —— in the isolated frog skin (Candia, Zadunaisky) (255) 517 Solute diffusion Hindrance of —— within membranes as measured with microporous membranes of known pore geometry (Beck, Schultz) (255) 273 Squid axon | (Guber et al.) (255) 167 Transmembrane potential The buffer value and of Escherichia coli (White, O'Brien) (255) 780 Transport Coupling in secondary active Activation of transport by co-transport and/or counter-transport with the fluxes of other solutes (Heinz et al.) (255) 442 Transport in cells A theory of coupled (Verhoff, Sundaresan) (255) 425 Triflocin Ouabain-uninhibited sodium transport in |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited ———————————————————————————————————— | (Guber et al.) (255) 167 Transmembrane potential The buffer value and of Escherichia coli (White, O'Brien) (255) 780 Transport Coupling in secondary active Activation of transport by co-transport and/or counter-transport with the fluxes of other solutes (Heinz et al.) (255) 442 Transport in cells A theory of coupled (Verhoff, Sundaresan) (255) 425 Triflocin Ouabain-uninhibited sodium transport in human erythrocytes: The effects of |
| Armstrong) (255) 663 Sodium transport Ouabain-uninhibited —— in human erythrocytes: The effects of triflocin (Dunn) (255) 567 Sodium transport Potassium flux and —— in the isolated frog skin (Candia, Zadunaisky) (255) 517 Solute diffusion Hindrance of —— within membranes as measured with microporous membranes of known pore geometry (Beck, Schultz) (255) 273 Squid axon | (Guber et al.) (255) 167 Transmembrane potential The buffer value and of Escherichia coli (White, O'Brien) (255) 780 Transport Coupling in secondary active Activation of transport by co-transport and/or counter-transport with the fluxes of other solutes (Heinz et al.) (255) 442 Transport in cells A theory of coupled (Verhoff, Sundaresan) (255) 425 Triflocin Ouabain-uninhibited sodium transport in |

SUBJECT INDEX 1001

Trypsin

Micromorphologic consequences following perturbation of erythrocyte membranes by ______, phospholipase A, lysolecithin, sodium dodecylsulfate and saponin: A corre-

lated freeze-etching and biochemical study (Speth et al.) (255) 386

Uridine uptake

Studies on the mechanism of serum stimulation of ——— in serum-less mouse cells (Hare) (255) 905

Water permeability

Probenecid: Effects on ____ in frog urinary bladder (Ripoche et al.) (255) 91