

## ERRATA

## BIOCHIMICA ET BIOPHYSICA ACTA Vol. 433 (1976)

- p. 81 and 86, Figs. 1 and 3: for "The bar represents 1  $\mu\text{m}$ " read "The bar represents 3  $\mu\text{m}$ "
- p. 82, Fig. 2: for "The bar represents 0.5  $\mu\text{m}$ " read "The bar represents 1.5  $\mu\text{m}$ "
- p. 219, line 10: for " $(2/e^2 \text{ diameter} \approx 6 \mu\text{m})$ " read " $(1/e^2 \text{ diameter} \approx 6 \mu\text{m})$ "
- p. 220, legend to Table I: for "... was 20  $\mu\text{m}$  or less" read "... was 2.0  $\mu\text{m}$  or less"  
last column of Table I: for " $(= K_B T / \sigma \pi \eta a')$ " read " $(= k_B T / 6 \pi \eta a')$ "

## EXTRA ERRATA

## BIOCHIMICA ET BIOPHYSICA ACTA Vol. 426 (1976)

- p. 63, replace fig. 2 and legend by the following




Fig. 2. Dual-wavelength traces of calcium uptake by cardiac relaxing system: Rates are expressed as nmol calcium accumulated/mg cardiac sarcoplasmic reticulum protein/min and are illustrated by numbers and arrows above the traces. Reaction conditions: 40 mM Tris/maleate (pH 6.8), 10 mM KCl, 10 mM  $\text{MgCl}_2$ , 5 mM sodium oxalate, 0.2 mM murexide and 125  $\mu\text{g}$  cardiac sarcoplasmic reticulum protein/ml; when present, 50  $\mu\text{g}$  protein kinase,  $10^{-6}$  M cyclic AMP, 30  $\mu\text{g}$  phosphorylase *b* kinase. Cuvette was equilibrated at 30  $^\circ\text{C}$  for 5 min, then 3.3 mM disodium ATP was added to the reaction. After 5 min of incubation, 66  $\mu\text{M}$  calcium was added to initiate the uptake reaction. — = control; -- = cyclic AMP; · — = phosphorylase *b* kinase; . . . = protein kinase and cyclic AMP.