

Walter, H., Krob, E.J. and Ascher, G.S.
Aging of erythrocytes results in altered cell surface properties in the rat, but not in the human. Studies by partitioning in two-polymer aqueous phase systems (641) 202

Wohlhueter, R.M., see Graff, J.C. (641) 320

Yoshida, S., see Hayashi, K. (641) 106

Zahler, P., see Kempf, C. (641) 88

Zimmerman, U. and Scheurich, P.

Fusion of *Avena sativa* mesophyll cell protoplasts by electrical breakdown (641) 160

ERRATA

BIOCHIMICA ET BIOPHYSICA ACTA, VOL. 640 (1981)

p. 365, second paragraph, line 8 should read:
tion reports that the aldehyde derivatives of lysine, α -aminoadipic acid- δ -semi-

p. 423, legend to Fig. 1, second line should read:

of 8.5 . 1 at different pH values: pH 7 (● — ● — ●), pH 6 (· · · · ·), pH 5 (— — —) and pH 4 (○ — ○ — ○)

p. 630, third paragraph, line 7 should read:
factor for the j th transition, normalized to unity. K is the local field factor

p. 736, fourth paragraph, line 4 should read:
vesicles were made by sonicating the multilamellar preparation (10 μ mol/ml) for

p. 809, Legend to Fig. 1 should read as follows:

Fig. 1. (a) Phosphotungstic acid negative-stain preparation of resealed singly lysed white ghosts showing the protrusions (arrows) and halo substructures (arrowhead) on the membrane surface. Magnification, $\times 40\,000$. (b) Freeze-fracture micrograph of singly lysed resealed white ghosts. The arrows indicate protrusions from the membrane. The protoplasmic fractured faces (PF) are covered by many 80–100 Å particles while the extracellular (EF) surfaces have fewer particles. Similar membrane fragmentation is observed for doubly lysed resealed ghosts. Arrow at the lower right-hand corner indicates the direction of shadowing. The membrane evaginations viewed from the cytoplasmic space are shown in the insert. Magnification, $\times 58\,500$.

BIOCHIMICA ET BIOPHYSICA ACTA, VOL. 603 (1980)

p. 322, 13th line of Summary, for 'increased' read 'decreased'