

ADDENDUM

BIOCHIMICA ET BIOPHYSICA ACTA, VOL
731 (1983)

pp. 97–108

NONELECTROLYTE SUBSTITUTION FOR WATER IN PHOSPHATIDYLCHOLINE BI- LAYERS

R.V. McDANIEL, S.A. SIMON AND T.J. McINTOSH

(Durham, NC, U.S.A.)

The following complementary information to this
article has been received:

The interlayer spacings in the non-aqueous lamel-

lar liquid crystals of phosphatidylcholine and (a)
ethylene glycol [1]; (b) alkane diols [2] and (c)
poly(ethyleneglycol) dodecyl ethers [3] are availa-
ble in the literature. In addition, the dynamics of
ethylene glycol have been described [4].

References

- 1 Moucharafieh, N. and Friberg, S.E. (1979) *Mol. Cryst. Liq. Cryst.* 49, 231
- 2 El Nokaly, M., Ford, L., Friberg, S. and Larsen, D. (1981) *J. Colloid Interface Sci.* 84, 228 (1982)
- 3 Ganzuo, L., El Nokaly, M. and Friberg, S. (1982) *Mol. Cryst. Liq. Cryst.* 72, 183
- 4 Larsen, D.W., Friberg, S. and Christenson, H. (1980) *J. Am. Chem. Soc.* 102, 6565

ERRATA

BIOCHIMICA ET BIOPHYSICA ACTA, VOL.
769 (1984)

p. 87, left-hand column, line 5:
for 'concentration' read 'concentrator'

p. 91, right-hand column, line 35:
for '(1,4)-MSL' read '(1,14)-MSL'

p. 93, left-hand column, lines 27–29 should read:
served at physiological temperature. Only in sys-
tems where protein-protein associations occur
[1,2,6,9] has this component been observed.