



Cumulative Contents

Information for Contributors

Regular papers

Effect of 17 β -estradiol on chondrocyte membrane fluidity and phospholipid metabolism is membrane-specific, sex-specific, and cell maturation-dependent

Z. Schwartz, P.A. Gates, E. Nasatzky, V.L. Sylvia, J. Mendez, D.D. Dean and B.D. Boyan (Israel, USA)

Stable GDP analog-induced inactivation of G_i proteins promotes cardiac adenylyl cyclase inhibition by guanosine 5'-($\beta\gamma$ -imino)triphosphate and muscarinic acetylcholine receptor

L. Piacentini, R. Mura, K.H. Jakobs and F. Nieroomand (Germany)

Inhibition of ATP-diphosphohydrolase (apyrase) of *Torpedo* electric organ by 5'-p-fluorosulfonylbenzoyl-adenosine

E. Martí, I. Gómez de Aranda and C. Solsona (Spain)

Arginine uptake through a novel cationic amino acid:K⁺ symporter, System R⁺, in brush border membrane vesicles from larval *Manduca sexta* midgut

Z. Liu and W.R. Harvey (USA)

Cationic lysine uptake by System R⁺ and zwitterionic lysine uptake by System B in brush border membrane vesicles from larval *Manduca sexta* midgut

Z. Liu and W.R. Harvey (USA)

Monomeric human red cell glucose transporter (Glut1) in non-ionic detergent solution and a semi-elliptical torus model for detergent binding to membrane proteins

L. Haneskog, L. Andersson, E. Brekkan, A.-K. Englund, K. Kameyama, L. Liljas, E. Greijer, J. Fischbarg and P. Lundahl (Sweden, Japan, USA)

A kinetic characterization of putrescine and spermidine uptake and export in human erythrocytes

G.H. Fukumoto and C.V. Byus (USA)

Cryptic receptors for insulin-like growth factor II in the plasma membrane of rat adipocytes – a possible link to cellular insulin resistance

Z.-W. Yu, A. Wickman and J.W. Eriksson (Sweden)

Caffeine as an antioxidant: inhibition of lipid peroxi-

vii

dation induced by reactive oxygen species

T.P.A. Devasagayam, J.P. Kamat, H. Mohan and P.C. Kesavan (India)

63

Effect of vanadate on glucose transporter (GLUT4) intrinsic activity in skeletal muscle plasma membrane giant vesicles

S. Kristiansen, J. Youn and E.A. Richter (Denmark)

71

Membrane changes associated with the temperature-sensitive P85^{bag-mos}-dependent transformation of rat kidney cells as determined by dielectrophoresis and electrorotation

Y. Huang, X.-B. Wang, F.F. Becker and P.R.C. Gascoyne (USA)

76

Lindane suppresses the lipid-bilayer permeability in the main transition region

M.C. Sabra, K. Jørgensen and O.G. Mouritsen (Denmark)

85

Effect of glycation of albumin on its binding to renal brush-border membrane vesicles: influence of aging in rats

P. Verbeke, M. Perichon, J. Schaefferbeke and H. Bakala (France)

93

The measurement of transmembrane electrical potential with lipophilic cations

P.D. Hockings and P.J. Rogers (Australia)

101

Association of hepatic system A amino acid transporter with the membrane-cytoskeletal proteins ankyrin and fodrin

M.E. Handlogten, E.E. Dudenhausen, W. Yang and M.S. Kilberg (USA)

107

Sulfate transport in human placental brush-border membrane vesicles

S.M. Grassl (USA)

115

Interactions of alkylglucosides with the renal sodium/D-glucose cotransporter

H. Kipp, J.-T. Lin and R.K.H. Kinne (Germany)

124

Effects of extracellular pH on intracellular pH-regulation and growth in a human colon carcinoma cell-line

G. Bischof, E. Cosentini, G. Hamilton, M. Riegler, J. Zacherl, B. Teleky, W. Feil, R. Schiessel, T.E. Machen and E. Wenzl (Austria, USA)

131

Pathway for Ca²⁺ influx into cells by trichosporin-B-VIa, an α -aminoisobutyric acid-containing peptide, from the fungus *Trichoderma polysporum*

E. Tachikawa, K. Nogimori, S. Takahashi, K.

57

Mizuma, K. Itoh, T. Kashimoto, Y. Nagaoka, A. Iida and T. Fujita (Japan)	140	tions. Indications for kinetic similarities between H^+/K^+ -ATPase and Na^+/K^+ -ATPase	207
Inhibition of calcium channel activation in GH3 cells by static magnetic fields		J. Rettinger (Germany)	
A.D. Rosen (USA)	149	Isolation of mitochondrial porin of the fly <i>Protophormia</i> : porin modification by the pesticide CGA 140'408 studied in lipid bilayer membranes	
Exogenous heat shock protein hsp70 activates potassium channels in U937 cells		P. Wiesner, B. Popp, A. Schmid, R. Benz and H. Kayser (Germany, Switzerland)	216
Yu.A. Negulyaev, E.A. Vedernikova, A.V. Kinev and A.P. Voronin (Russia)	156	Uptake of neutral α - and β -amino acids by human proximal tubular cells	
Iron transport into erythroid cells by the Na^+/Mg^{2+} antiport		H. Jessen, H. Røigaard and C. Jacobsen (Denmark)	225
L.M. Stonell, D.L. Savigni and E.H. Morgan (Australia)	163	Membrane fluidity response to odorants as seen by 2H -NMR and infrared spectroscopy	
Information for Contributors	v	M. Bouchard, N. Boudreau and M. Auger (Canada)	233
Rapid report		Alteration of Na^+ -coupled transport in site-directed mutants of the melibiose carrier of <i>Escherichia coli</i>	
Temperature dependence of endocytosis in renal epithelial cells in culture		P.J. Franco and T.H. Wilson (USA)	240
Z. Mamdoui, M.-C. Giocondi, R. Laprade and C. Le Grimellec (France, Canada)	171	Receptor affinity of neurotensin message segment immobilized on liposome	
Short sequence-papers		J. Zhao, S. Kimura and Y. Imanishi (Japan)	249
Characterization of the 3' UTR sequence encoded by the AQP-1 gene in human retinal pigment epithelium		Human placental ATP diphosphohydrolase is a highly N-glycosylated plasma membrane enzyme	
A. Ruiz and D. Bok (USA)	174	S. Christoforidis, T. Papamarcaki and O. Tsolas (Greece)	257
cDNAs from <i>Onchocerca</i> sp. encoding members of the MRS3/MRS4 class of mitochondrial solute carriers		Pathways for flip-flop of mono- and di-anionic phospholipids in the erythrocyte membrane	
J. Catmull and D.J. Miller (Australia, USA)	179	M.V. Serra, D. Kamp and C.W.M. Haest (Italy, Germany)	263
Regular papers		Sodium-dependent high-affinity binding of carnitine to human placental brush border membranes	
Biochemical characterization of a V-ATPase of tracheal smooth muscle plasma membrane fraction		A.S. Roque, P.D. Prasad, J.S. Bhatia, F.H. Leibach and V. Ganapathy (USA)	274
G. Pacheco, I. Lippo de Bécemberg, R. Gonzalez de Alfonzo and M.J. Alfonzo (Venezuela)	182	Vesicles made of glycopospholipids with homogeneous (two fluorocarbon or two hydrocarbon) or heterogeneous (one fluorocarbon and one hydrocarbon) hydrophobic double chains	
H^+ , K^+ , and Na^+ transport across phospholipid vesicular membrane by the combined action of proton uncoupler 2,4-dinitrophenol and valinomycin		F. Guillod, J. Greiner and J.G. Riess (France)	283
B.S. Prabhananda and M.H. Kombrabail (India)	193	Monovalent cations differentially affect membrane surface properties and membrane curvature, as revealed by fluorescent probes and dynamic light scattering	
The mechanism of L-glutamate transport by lactating rat mammary tissue		R. Kraayenhof, G.J. Sterk, H.W. Wong Fong Sang, K. Krab and R.M. Epand (The Netherlands, Canada)	293
I.D. Millar, D.T. Calvert, M.A. Lomax and D.B. Shennan (UK)	200	Cumulative Contents, Vol. 1282	303
Characteristics of Na^+/K^+ -ATPase mediated proton current in Na^+ - and K^+ -free extracellular solu-		Author index	305