

Biomembranes

Cumulative Contents

Rapid reports

Two fatty acids can replace one phospholipid in condensed complexes with cholesterol
T.M. Okonogi, A. Radhakrishnan and H.M. McConnell (USA)

Transdermal insulin delivery using lipid enhanced electroporation
A. Sen, M.E. Daly and S.W. Hui (USA)

Phosphatidyl ethanolamine is essential for targeting the arginine transporter Canlp to the plasma membrane of yeast
M. Opekarová, I. Robl and W. Tanner (Czech Republic, Germany)

Regular papers

Modulation of acetylcholine receptor channel kinetics by hydrocortisone
E. Nurowska and F. Ruzzier (Italy, Poland)

Ultradeflexible lipid vesicles can penetrate the skin and other semi-permeable barriers unfragmented. Evidence from double label CLSM experiments and direct size measurements
G. Cevc, A. Schätzlein and H. Richardsen (Germany)

A novel pH-sensitive liposome formulation containing oleyl alcohol
J.J. Sudimack, W. Guo, W. Tjarks and R.J. Lee (USA)

Insertion of carrier proteins into hydrophilic loops of the *Escherichia coli* lactose permease
C.K. Engel, L. Chen and G.G. Privé (Canada)

Stability of the lactose permease in detergent solutions
C.K. Engel, L. Chen and G.G. Privé (Canada)

Conformational transitions of membrane-bound HIV-1 fusion peptide
A. Sáez-Ciri6n and J.L. Nieva (Spain)

Cubic phase is induced by cholesterol in the dispersion of 1-palmitoyl-2-oleoyl-phosphatidylethanolamine
X. Wang and P.J. Quinn (UK)

Cloaking cytolytic peptides for liposome-based detection and potential drug delivery

H.S. Aojula, S. Offerman, R.R. Aojula, A.P. Hutchinson, S. Nicklin and D.J. Clarke (UK) 73

CTP:phosphocholine cytidyltransferase and protein kinase C recognize different physical features of membranes: differential responses to an oxidized phosphatidylcholine

A.E. Drobnies, S.M.A. Davies, R. Kraayenhof, R.F. Epand, R.M. Epand and R.B. Cornell (Canada, The Netherlands) 82

High-throughput screening of bacteriorhodopsin mutants in whole cell pastes

L.C. Martinez and G.J. Turner (USA) 91

Asp³⁴⁴ and Thr³⁴⁵ are critical for cation exchange mediated by NhaD, Na⁺/H⁺ antiporter of *Vibrio cholerae*

E. Ostroumov, J. Dzioba, P.C. Loewen and P. Dibrov (Canada) 99

Interaction of nominally soluble proteins with phospholipid monolayers at the air–water interface

W.H. Pitcher III, S.L. Keller and W.H. Huestis (USA) 107

Molecular analysis of the mechanism of potassium uptake through the TRK1 transporter of *Saccharomyces cerevisiae*

R. Haro and A. Rodríguez-Navarro (Spain) 114

Amphiphilic effects of local anesthetics on rotational mobility in neuronal and model membranes

I. Yun, E.-S. Cho, H.-O. Jang, U.-K. Kim, C.-H. Choi, I.-K. Chung, I.-S. Kim and W.G. Wood (South Korea, USA) 123

A study of L-leucine, L-phenylalanine and L-alanine transport in the perfused rat mammary gland: possible involvement of LAT1 and LAT2

D.B. Shennan, D.T. Calvert, M.T. Travers, Y. Kudo and C.A.R. Boyd (UK) 133

Differential effects induced by α - and β -endosulfan in lipid bilayer organization are reflected in proton permeability

R.A. Videira, M.C. Antunes-Madeira and V.M.C. Madeira (Portugal) 140

- Ionic strength has a greater effect than does transmembrane electric potential difference on permeation of tryptamine and indoleacetic acid across Caco-2 cells
M. Sugawara, M. Kurosawa, K. Sakai, M. Kobayashi, K. Iseki and K. Miyazaki (Japan) 149
- Towards high-resolution ^1H -NMR in biological membranes: magic angle spinning of bicelles
C. Carlotti, F. Aussenac and E.J. Dufourc (France) 156
- Micromanipulation of phospholipid bilayers by atomic force microscopy
N. Maeda, T.J. Senden and J.-M. Di Meglio (Australia, France) 165
- La^{3+} and Gd^{3+} induce shape change of giant unilamellar vesicles of phosphatidylcholine
T. Tanaka, Y. Tamba, S.Md. Masum, Y. Yamashita and M. Yamazaki (Japan) 173
- Functional characterization of the mouse organic-anion-transporting polypeptide 2
J.E. van Montfoort, T.E. Schmid, I.-D. Adler, P.J. Meier and B. Hagenbuch (Switzerland, Germany) 183
- Bioluminescence reaction catalyzed by membrane-bound luciferase in the “firefly squid,” *Watasenia scintillans*
F.I. Tsuji (USA) 189
- The hepatocyte glucose-6-phosphatase subcomponent T3: its relationship to GLUT2
J.S. Hah, J. Ryu, W. Lee, C.Y. Jung and M. Lachaal (USA) 198
- Photodynamic activation of ion transport through lipid membranes and its correlation with an increased dielectric constant of the membrane
F. Killig and G. Stark (Germany) 207
- Band-3 protein function in human erythrocytes: effect of oxygenation–deoxygenation
A. Galtieri, E. Tellone, L. Romano, F. Misiti, E. Bellocco, S. Ficarra, A. Russo, D. Di Rosa, M. Castagnola, B. Giardina and I. Messina (Italy) 214
- Binding of bilirubin with albumin-coupled liposomes: implications in the treatment of jaundice
A.K. Masood, S.M. Faisal, M.K. Mushahid, A. Nadeem, M.U. Siddiqui and M. Owais (India, USA) 219
- Superoxide dismutase entrapped in long-circulating liposomes: formulation design and therapeutic activity in rat adjuvant arthritis
M. Luisa Corvo, J.C.S. Jorge, R. Van't Hof, M.E.M. Cruz, D.J.A. Crommelin and G. Storm (Portugal, Netherlands) 227
- Interaction of human serum albumin with membranes containing polymer-grafted lipids: spin-label ESR studies in the mushroom and brush regimes
R. Bartucci, M. Pantusa, D. Marsh and L. Sportelli (Italy, Germany) 237
- Phenotypic subunit composition of the tobacco (*Nicotiana tabacum* L.) vacuolar-type H^+ -translocating ATPase
M. Drobny, M. Schnölzer, S. Fiedler, U. Lüttge, E. Fischer-Schliebs, A.-L. Christian and R. Ratajczak (Germany) 243
- Synergistic permeability enhancing effect of lysophospholipids and fatty acids on lipid membranes
J. Davidsen, O.G. Mouritsen and K. Jørgensen (Denmark) 256
- Modulation of the expression of GABA_A receptors in rat cerebellar granule cells by protein tyrosine kinases and protein kinase C
R. Balduzzi, A. Cupello and M. Robello (Italy) 263
- Roles of peptide–peptide charge interaction and lipid phase separation in helix–helix association in lipid bilayer
D. Shigematsu, M. Matsutani, T. Furuya, T. Kiyota, S. Lee, G. Sugihara and S. Yamashita (Japan) 271
- Partitioning of Triton X-100, deoxycholate and C_{10}EO_8 into bilayers composed of native and hydrogenated egg yolk sphingomyelin
F. Ollila and J.P. Slotte (Finland) 281
- Amphotericin B interactions with a DOPC monolayer. Electrochemical investigations
R. Stoodley, J. Shepherd, K.M. Wasan and D. Bizzotto (Canada) 289
- Review*
- Genes for calcium-permeable channels in the plasma membrane of plant root cells
P.J. White, H.C. Bowen, V. Demidchik, C. Nichols and J.M. Davies (UK) 299
- Regular papers*
- Angiotensin II stimulates renal proximal tubule Na^+ -ATPase activity through the activation of protein kinase C
L.B.A. Rangel, C. Caruso-Neves, L.S. Lara and A.G. Lopes (Brazil) 310

Spectroscopic characterization of nanoErythroosomes in the absence and presence of conjugated polyethyleneglycols: an FTIR and ^{31}P -NMR study <i>R. Pouliot, A. Saint-Laurent, C. Chypre, R. Audet, I. Vitté-Mony, R.C. Gaudreault and M. Auger (Canada)</i>	317	Thiamin transport in <i>Escherichia coli</i> : the mechanism of inhibition by the sulfhydryl-specific modifier <i>N</i> -ethylmaleimide <i>A.D. Hollenbach, K.A. Dickson and M.W. Washabaugh (USA)</i>	421
Calcium uptake and calcium transporter expression by trophoblast cells from human term placenta <i>R. Moreau, G. Daoud, R. Bernatchez, L. Simoneau, A. Masse and J. Lafond (Canada)</i>	325	Cholesterol markedly reduces ion permeability induced by membrane-bound amphotericin B <i>S. Matsuoka and M. Murata (Japan)</i>	429
Analysis of membrane topology of the human reduced folate carrier protein by hemagglutinin epitope insertion and scanning glycosylation insertion mutagenesis <i>X.Y. Liu and L.H. Matherly (USA)</i>	333	Determination of membrane protein topology by red-edge excitation shift analysis: application to the membrane-bound colicin E1 channel peptide <i>M.C. Tory and A.R. Merrill (Canada)</i>	435
Effect of liposome charge and PEG polymer layer thickness on cell–liposome electrostatic interactions <i>N. Dan (USA)</i>	343	Detection of cellular responses to toxicants by dielectrophoresis <i>K. Ratanachoo, P.R.C. Gascoyne and M. Ruchirawat (Thailand, USA)</i>	449
Transfection with fluorinated lipoplexes based on fluorinated analogues of DOTMA, DMRIE and DPPE <i>J. Gaucheron, C. Santaella and P. Vierling (France)</i>	349	Cationic triple-chain amphiphiles facilitate vesicle fusion compared to double-chain or single-chain analogues <i>A.V. Koulov, L. Vares, M. Jain and B.D. Smith (USA)</i>	459
Antibacterial peptide pleurocidin forms ion channels in planar lipid bilayers <i>N. Saint, H. Cadiou, Y. Bessin and G. Molle (France)</i>	359	Alterations of wheat root plasma membrane lipid composition induced by copper stress result in changed physicochemical properties of plasma membrane lipid vesicles <i>A.H. Berglund, M.F. Quartacci, L. Calucci, F. Navari-Izzo, C. Pinzino and C. Liljenberg (Sweden, Italy)</i>	466
A quantitative validation of fluorophore-labelled cell-permeable peptide conjugates: fluorophore and cargo dependence of import <i>R. Fischer, T. Waizenegger, K. Köhler and R. Brock (Germany)</i>	365	Cholesterol-dependent insertion of glycosylphosphatidylinositol-anchored enzyme <i>S. Morandat, M. Bortolato and B. Roux (France)</i>	473
Behavior of silkworm yolk protein on phospholipid membranes <i>Y. Toshima and K. Kawasaki (Japan)</i>	375	Hypochlorous acid inhibits glutathione <i>S</i> -conjugate export from human erythrocytes <i>M. Soszyński, I.B. Zavodnik, L.B. Zavodnik, L. Żylinska, G. Bartosz and M. Bryszewska (Poland, Belarus)</i>	479
Cholesterol as stabilizer of the oxytocin receptor <i>G. Gimpl and F. Fahrenholz (Germany)</i>	384	Cadmium transport through type II alveolar cell monolayers: contribution of transcellular and paracellular pathways in the rat ATII and the human A549 cells <i>C. Jumarie (Canada)</i>	487
Evaluation and optimization of DNA delivery into gliosarcoma 9L cells by a cholesterol-based cationic liposome <i>D. Lesage, A. Cao, D. Briane, N. Lievre, R. Coudert, M. Raphael, J.L. Salzmänn and E. Taillandier (France)</i>	393		
Brain spectrin exerts much stronger effect on anionic phospholipid monolayers than erythroid spectrin <i>W. Diakowski and A.F. Sikorski (Poland)</i>	403	<i>Short sequence-paper</i>	
Membrane dielectric changes indicate induced apoptosis in HL-60 cells more sensitively than surface phosphatidylserine expression or DNA fragmentation <i>X. Wang, F.F. Becker and P.R.C. Gascoyne (USA)</i>	412	Self-regulation of rat liver GAP junction by phosphorylation <i>P. Ghosh, S. Ghosh and S. Das (India)</i>	500
		<i>Author Index</i>	505
		<i>Cumulative Contents</i>	509